


# Dynamics of plastic networks of excitatory and inhibitory spiking neurons

Clayton Seitz

July 13, 2021


# Inferring the transfer function from ITC data



Measuring the *static* transfer function from novel images assuming that input currents are Gaussian variables

$$\phi(\xi) = \frac{r_{max}}{1 + \exp \beta(\xi - \xi_0)}$$


# Inferring the learning rule from ITC data



Inferring the change in input current  $\xi_{in}$  from the change in firing rate in **novel** relative to **familiar** stimuli

2

# Inferring the learning rule from ITC data



The change in input current to a neuron can then be read from the firing rate of that neuron when presented a novel stimulus

$$\Delta \xi_i(r) \propto (2q + 1 - \tanh(\beta(r - x)))$$

# A Hebbian update for synaptic weights

Assuming that  $\Delta W_{ij} \propto f(r_i)g(r_j)$ , the change in input current is related to synaptic plasticity by

$$\Delta \xi_i \propto f(r_i) \sum_j g(r_j) r_j$$

which we have fit from the data as

$$\Delta \xi_i(r) \propto (2q + 1 - \tanh(\beta(r - x)))$$

so we can write

$$f(r_i) = \frac{(2q + 1 - \tanh(\beta(r - x)))}{\sum_j g(r_j) r_j}$$

# Presenting novel and familiar stimuli to the network

