

Homework 10

Quantum Mechanics

December 4, 2022

C SEITZ

Problem 1. 4.7

Solution.

The wave function in three dimensions for a free particle ($V = 0$), is

$$\begin{aligned}\psi(\mathbf{x}, t) &= u(\mathbf{x})e^{-iE_nt/\hbar} \\ \psi^*(\mathbf{x}, -t) &= u^*(\mathbf{x})e^{-iE_nt/\hbar}\end{aligned}$$

where $u(\mathbf{x}) = e^{i\vec{p}\cdot\vec{k}}$. Note that the phase remains unchanged under complex conjugation and time reversal. Now if we reverse the direction of momentum i.e. $|p\rangle \rightarrow |p'\rangle$ for $\vec{p}\cdot\vec{p}' = -1$,

$$\psi'(\mathbf{x}, t) = u'(\mathbf{x})e^{-iE_nt/\hbar}$$

Notice that $u'(\mathbf{x}) = e^{-i\vec{p}'\cdot\vec{k}} = u^*(\mathbf{x})$. Therefore $\psi'(\mathbf{x}, t) = \psi^*(\mathbf{x}, -t)$

$$\chi_+(\hat{n}) = \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} e^{i\gamma} \end{pmatrix}$$

It is also known that

$$\chi_-(\hat{n}) = \begin{pmatrix} -\sin \frac{\theta}{2} e^{-i\gamma} \\ \cos \frac{\theta}{2} \end{pmatrix}$$

So we just need to prove that the given transformation gives this result, which it does

$$-i\sigma_2\chi^*(\hat{n}) = -i \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} e^{-i\gamma} \end{pmatrix} = \begin{pmatrix} -\sin \frac{\theta}{2} e^{-i\gamma} \\ \cos \frac{\theta}{2} \end{pmatrix}$$

■

Problem 2. 4.8

Solution.

First note that

$$H\Theta|n\rangle = \Theta H|n\rangle = E_n\Theta|n\rangle$$

so $|n\rangle$ and $\Theta|n\rangle$ have the same energy. If the states are nondegenerate then $|n\rangle$ and $\Theta|n\rangle$ represent the same state. Their wavefunctions are then the same:

$$\langle x'|n\rangle = \langle n|x'\rangle^*$$

which occurs if they are real, or have a phase difference independent of x . For this reason the wavefunction $\psi = e^{ip\cdot x/\hbar}$ does not violate time reversal invariance, because it is degenerate with $e^{ip\cdot x/\hbar}$.

■

Problem 3. 4.9

Solution.

$$\begin{aligned} \Theta|\alpha\rangle &= \int d^3\mathbf{p} \Theta|\mathbf{p}\rangle \langle\mathbf{p}|\alpha\rangle^* \\ &= \int d^3\mathbf{p} |-\mathbf{p}\rangle \langle\mathbf{p}|\alpha\rangle^* \\ &= \int d^3\mathbf{p} |\mathbf{p}\rangle \langle-\mathbf{p}|\alpha\rangle^* \\ &= \phi^*(-p) \end{aligned}$$

■

Problem 4. 4.10

Solution.



Problem 5. *4.11*

Solution.



Problem 6. *4.12*

Solution.

