

Homework 6

Quantum Mechanics

October 28th, 2022

C SEITZ

Problem 1. *Problem 3.12 from Sakurai*

Solution.

In general the ensemble average of an operator $[A]$ is defined as

$$[A] = \text{Tr}(\rho A)$$

where $\hat{\rho} = \sum_i w_i \rho_i$ and $\rho_i = |\alpha_i\rangle \langle \alpha_i|$

$$\begin{aligned}\hat{\rho} &= a |+\rangle \langle +| + (1-a) |-\rangle \langle -| \\ &= \begin{pmatrix} \frac{1-a}{2} + a & \frac{1}{2}i(1-a) \\ -\frac{1}{2}i(1-a) & \frac{1-a}{2} \end{pmatrix}\end{aligned}$$

$$\begin{aligned}[S_x] &= \text{Tr}(\hat{\rho} S_x) \\ &= \frac{\hbar}{2} \text{Tr} \left(\begin{pmatrix} \frac{1}{2}i(1-a) & \frac{1-a}{2} + a \\ \frac{1-a}{2} & -\frac{1}{2}i(1-a) \end{pmatrix} \right) = 0\end{aligned}$$

$$\begin{aligned}[S_y] &= \text{Tr}(\hat{\rho} S_y) \\ &= \frac{\hbar}{2} \text{Tr} \left(\begin{pmatrix} \frac{a-1}{2} & -i \left(\frac{1-a}{2} + a \right) \\ \frac{1}{2}i(1-a) & \frac{a-1}{2} \end{pmatrix} \right) = \frac{\hbar}{2}(a-1)\end{aligned}$$

$$\begin{aligned}[S_z] &= \text{Tr}(\hat{\rho} S_z) \\ &= \frac{\hbar}{2} \text{Tr} \left(\begin{pmatrix} \frac{1-a}{2} + a & -\frac{1}{2}i(1-a) \\ -\frac{1}{2}i(1-a) & \frac{a-1}{2} \end{pmatrix} \right) = \frac{\hbar}{2}a\end{aligned}$$

This is what we would expect for the pure ensemble cases $a = 1$ and $a = 0$. ■

Problem 2. *Problem 3.13 from Sakurai*

Solution.

The state vector in the S_z basis has the form

$$|\alpha\rangle = c_+ |+\rangle + c_- |-\rangle$$

First note that

$$\langle S_z \rangle = |c_+|^2 - |c_-|^2 \quad |c_+|^2 + |c_-|^2 = 1$$

Together, these equations tell us the magnitude of each complex component.

$$|c_+|^2 = \frac{\langle S_z \rangle + 1}{2} \quad |c_-|^2 = \frac{1 - \langle S_z \rangle}{2}$$

$$\begin{aligned} \langle S_x \rangle &= \langle \alpha | (|+\rangle \langle -| + |-\rangle \langle +|) (c_+ |+\rangle + c_- |-\rangle) \\ &= \langle \alpha | (c_- |+\rangle + c_+ |-\rangle) \\ &= (c_+^* \langle +| + c_-^* \langle -|) (c_- |+\rangle + c_+ |-\rangle) \\ &= c_+^* c_- + c_-^* c_+ \\ &= |c_+| |c_-| (e^{i(\theta-\phi)} + e^{i(\phi-\theta)}) \\ &= 2|c_+| |c_-| \cos(\theta - \phi) \end{aligned}$$

$$\begin{aligned} \langle S_y \rangle &= \langle \alpha | ((i |+\rangle \langle -| - i |-\rangle \langle +|) (c_+ |+\rangle + c_- |-\rangle) \\ &= i \langle \alpha | (c_- |+\rangle - c_+ |-\rangle) \\ &= i(c_+^* \langle +| + c_-^* \langle -|) (c_- |+\rangle - c_+ |-\rangle) \\ &= c_+^* c_- - c_-^* c_+ \\ &= |c_+| |c_-| (e^{i(\theta-\phi)} - e^{i(\phi-\theta)}) \\ &= 2i|c_+| |c_-| \sin(\theta - \phi) \end{aligned}$$

So $\langle S_x \rangle$ gives us the phase difference of c_+ and c_- . Then the sign of $\langle S_y \rangle$ tells us whether θ or ϕ is larger, since sine is odd. This is all we can hope to extract from the expectation values, since multiplying by a global phase $e^{i\delta} |\alpha\rangle$ has no effect on the expectation values.



Problem 3. *Problem 3.14 from Sakurai*

Solution.

$$\begin{aligned}\hat{\rho} &= \sum_i w_i |\psi_i\rangle \langle \psi_i| \\ &= \frac{1}{3} (|\alpha\rangle \langle \alpha| + |\beta\rangle \langle \beta| + |2\rangle \langle 2|)\end{aligned}$$

We can write this out explicitly in the subspace spanned by $|0, 1, 2\rangle$

$$|\alpha\rangle \langle \alpha| = \frac{1}{2} \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad |\beta\rangle \langle \beta| = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix} \quad |2\rangle \langle 2| = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

$$\hat{\rho} = \frac{1}{6} \begin{pmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 3 \end{pmatrix}$$

Now recall that $H = \hbar\omega(N + \frac{1}{2})$ which reads

$$H = \hbar\omega \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix} + \left(\frac{\hbar\omega}{2}\right) \mathbb{I}_{3 \times 3}$$

$$\begin{aligned}[H] &= \text{Tr}(\rho H) = \hbar\omega \text{Tr}(\rho N + \rho/2) \\ &= \hbar\omega (\text{Tr}(\rho N) + \text{Tr}(\rho/2)) \\ &= \frac{11}{6} \hbar\omega\end{aligned}$$

■

Problem 4. *Problem 3.15 from Sakurai*

Solution.

$$\rho(t_0) = \sum_i w_i |\psi_i; t_0\rangle \langle \psi_i; t_0|$$

In the Schrodinger picture, the coefficients of the state vectors evolve. Therefore,

$$\begin{aligned}\rho(t) &= \sum_i w_i \mathcal{U}(t, t_0) |\psi_i; t_0\rangle \langle \psi_i; t_0| \mathcal{U}^\dagger(t, t_0) \\ &= \mathcal{U}(t, t_0) \left(\sum_i w_i |\psi_i; t_0\rangle \langle \psi_i; t_0| \right) \mathcal{U}^\dagger(t, t_0) \\ &= \mathcal{U}(t, t_0) \rho(t_0) \mathcal{U}^\dagger(t, t_0)\end{aligned}$$

For a pure ensemble in state $|\psi_i\rangle$, the density matrix is

$$\rho(t_0) = |\psi_i; t_0\rangle \langle \psi_i; t_0|$$

At a later time, the density matrix is

$$\begin{aligned}\rho(t) &= \mathcal{U}(t, t_0) \rho(t_0) \mathcal{U}^\dagger(t, t_0) \\ &= \mathcal{U}(t, t_0) |\psi_i; t_0\rangle \langle \psi_i; t_0| \mathcal{U}^\dagger(t, t_0) \\ &= |\psi_i; t_0\rangle \langle \psi_i; t_0|\end{aligned}$$

■

Problem 5. *Problem 3.16 from Sakurai*

Solution. A 3x3 matrix has 9 parameters, but we only need

■

Problem 6. *Problem 3.40 from Sakurai*

Solution. The singlet state is

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|+-\rangle + |-+\rangle)$$

If B doesn't make a measurement, B will have no effect on A 's measurement. So the probability for A to obtain $s_{1z} = \hbar/2$ is of course $1/2$. The probability that A measures $s_{1x} = \hbar/2$ in this state is also $1/2$. This is because obtaining $s_{1x} = \hbar/2$ is equiprobable for the two states in the singlet superposition.

If observer B has determined that $s_{2z} = \hbar/2$, then observer A must observe $s_{1z} = -\hbar/2$ since the measurement made by B collapses $|\psi\rangle$ to $|-+\rangle$.

Furthermore, if observer B has measured $s_{2z} = \hbar/2$, then particle 1 must be in the $|+\rangle$ state (as stated before) which means $s_{1x} = \pm\hbar/2$ with equal probability.

■