

# Homework 3

Quantum Mechanics

February 12, 2023

C SEITZ

## Problem 1. Problem 2.48

### Solution.

The polar decomposition of a matrix is  $A = UJ = KU$  where  $U$  is a unitary operator and  $J, K$  are positive operators that satisfy  $J = \sqrt{A^\dagger A}$  and  $K = \sqrt{AA^\dagger}$ . If  $P$  is itself a positive matrix we can immediately say that its polar decomposition is  $P = IP = PI$ . If the matrix to decompose is unitary then of course  $UU^\dagger = U^\dagger U = I$  so its decomposition is itself. If the matrix to decompose is Hermitian, then  $H = H^\dagger$  and

$$J = K = \sqrt{H^2} = \sqrt{\sum_i \lambda_i^2 |i\rangle \langle i|} = \sum_i |\lambda_i| |i\rangle \langle i|$$

and its polar decomposition is therefore  $U \sum_i |\lambda_i| |i\rangle \langle i|$  or  $\sum_i |\lambda_i| |i\rangle \langle i| U$ . ■

## Problem 2. Problem 2.49

### Solution.

The polar decomposition is  $A = UJ$ . The spectral decomposition of  $J$  is

$$J = \sqrt{\sum_i \lambda_i \lambda_i^* |i\rangle \langle i|} = \sum_i |\lambda_i| |i\rangle \langle i|$$

For the unitary matrix  $U$ , we have

$$U = \sum_j \lambda_j |j\rangle \langle j|$$

Therefore the product  $UJ$  reads

$$\begin{aligned}
 UJ &= \left( \sum_j \lambda_j |j\rangle \langle j| \right) \left( \sum_i |\lambda_i| |i\rangle \langle i| \right) \\
 &= \sum_{ij} |\lambda_i| \lambda_j |i\rangle \langle i|j\rangle \langle j|
 \end{aligned}$$

■

**Problem 3.** *Problem 2.50*

**Solution.**

$$A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$$

First, consider

$$A^\dagger A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \quad AA^\dagger = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$$

$$J = \sqrt{A^\dagger A} =$$

■

**Problem 4.** *Problem 2.51*

**Solution.** The Hadamard gate  $H$  is unitary if  $H^\dagger = H^{-1}$ . It is easy to see that

$$H^\dagger = H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

It's inverse is

$$H^{-1} = -\frac{1}{\sqrt{2}} \begin{pmatrix} -1 & -1 \\ -1 & 1 \end{pmatrix} = H$$

■

**Problem 5.** *Problem 2.52*

**Solution.**

$$H^2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

■

**Problem 6.** *Problem 2.53*

**Solution.** Writing out the characteristic equation gives that the eigenvalues are  $\lambda = \pm\sqrt{2}$ . ■

**Problem 7.** *Problem 2.54*

**Solution.** Since the two operators commute, they are simultaneously diagonalizable. Consider the following spectral decompositions

$$A = \sum_n a_n |n\rangle \langle n|$$

$$B = \sum_n b_n |n\rangle \langle n|$$

Therefore, it must be true that

$$A + B = \sum_n (a_n + b_n) |n\rangle \langle n|$$

Now these matrices are Hermitian so their eigenvectors are orthogonal, and the product of matrix exponentials is just

$$\begin{aligned} \exp(A) \exp(B) &= \left( \sum_n \exp(a_n) |n\rangle \langle n| \right) \left( \sum_m \exp(b_m) |m\rangle \langle m| \right) \\ &= \sum_{m,n} \delta_{mn} \exp(a_n) \exp(b_m) |n\rangle \langle m| \\ &= \sum_n \exp(a_n) \exp(b_n) |n\rangle \langle n| \\ &= \sum_n \exp(a_n + b_n) |n\rangle \langle n| \\ &= \exp(A + B) \end{aligned}$$

■

**Problem 8.** *Problem 2.55***Solution.**

$$\begin{aligned}
UU^\dagger &= \exp\left(\frac{-iH(t_2 - t_1)}{\hbar}\right) \exp\left(\frac{iH(t_2 - t_1)}{\hbar}\right) \\
&= \left(\sum_n \exp\left(\frac{-iE_n(t_2 - t_1)}{\hbar}\right) |n\rangle \langle n|\right) \left(\sum_m \exp\left(\frac{iE_m(t_2 - t_1)}{\hbar}\right) |m\rangle \langle m|\right) \\
&= \sum_{m,n} \delta_{mn} |n\rangle \langle m| \\
&= \sum_n |n\rangle \langle n| = I
\end{aligned}$$

where  $H$  is a Hermitian operator. ■

**Problem 9.** *Problem 2.56***Solution.**

$U$  is unitary so its eigenvalues  $u_n$  have unit norm, which means

$$K = -i \log(U) = -i \sum_n \log(u_n) |n\rangle \langle n| = \sum_n \theta |n\rangle \langle n|$$

since

$$\log(u_n) = \log(|u_n| e^{i\theta}) = \log(|u_n|) + i\theta = i\theta$$

Therefore,  $K = K^\dagger$  since  $\theta \in \mathbb{R}$ . ■

**Problem 10.** *Problem 2.57***Solution.**

$$L_l |\alpha\rangle = \frac{\ell |l\rangle}{|\ell|}$$

$$M_m \frac{\ell |l\rangle}{|\ell|} = \frac{m\ell}{|m||\ell|} |m\rangle$$

which is equivalent to

$$\begin{aligned}
 N_{m\ell} |\alpha\rangle &= M_m L_\ell |\alpha\rangle \\
 &= \frac{|m\rangle \langle m|\ell\rangle \langle \ell|}{|m||\ell|} |\alpha\rangle \\
 &= \frac{\ell |m\rangle \langle m|}{|m||\ell|} |\ell\rangle \\
 &= \frac{m\ell}{|m||\ell|} |m\rangle
 \end{aligned}$$

■

**Problem 11.** *Problem 2.58*

**Solution.**

Since the system is in an eigenstate of  $M$  with eigenvalue  $m$ , the average will be  $m$

$$\langle M \rangle = \langle m| M |m\rangle = \langle m| m |m\rangle = m$$

The variance must then be zero

$$\begin{aligned}
 (\Delta M)^2 &= \langle M^2 \rangle - \langle M \rangle^2 \\
 &= m^2 - m^2 = 0
 \end{aligned}$$

■

**Problem 12.** *Problem 2.59*

**Solution.**

$$\langle 0| X |0\rangle = \langle 0|1\rangle = 0$$

$$\begin{aligned}
 (\Delta X)^2 &= \langle X^2 \rangle - \langle X \rangle^2 \\
 &= \langle X^2 \rangle \\
 &= \langle 0| X^2 |0\rangle \\
 &= 1
 \end{aligned}$$

■

**Problem 13.** *Problem 2.60*

**Solution.**

$$\vec{v} \cdot \sigma = \begin{pmatrix} v_z & v_x - iv_y \\ v_x + iv_y & -v_z \end{pmatrix}$$

The corresponding characteristic equation is

$$\lambda^2 - (v_z^2 + v_y^2 + v_x^2) = 0$$

If  $\vec{v}$  is normalized then  $\lambda = \pm 1$ .

■

**Problem 14.** *Problem 2.61*

**Solution.**

■