

Bell's Inequality

Clayton W. Seitz

April 28, 2023

CHSH and Tsirelson's Inequalities

Alice: Q, R Bob: S, T

Classical observables distributed according to $P(Q, R, S, T)$. Combination of correlations between Alice and Bobs measurements are bounded according to the CHSH inequality

$$|E(QS) + E(RS) + E(RT) - E(QT)| \leq 2$$

For the quantum version, define 4 spin operators along arbitrary directions $Q = \vec{q} \cdot \sigma, R = \vec{r} \cdot \sigma, S = \vec{s} \cdot \sigma, T = \vec{t} \cdot \sigma$.

$$|\langle Q \otimes S \rangle + \langle R \otimes S \rangle + \langle R \otimes T \rangle - \langle Q \otimes T \rangle| \leq 2\sqrt{2}$$

The Tsirelson bound

Solution to Problem 2.3 in the book:

$$(Q \otimes S + R \otimes S + R \otimes T - Q \otimes T)^2 = 4I + [Q, R] \otimes [S, T]$$

which can be used to derive

$$\begin{aligned}\langle (Q \otimes S + R \otimes S + R \otimes T - Q \otimes T)^2 \rangle &\leq \langle (Q \otimes S + R \otimes S + R \otimes T - Q \otimes T)^2 \rangle \\ &= \langle 4I + [Q, R] \otimes [S, T] \rangle \\ &= 4 + \langle [Q, R] \otimes [S, T] \rangle\end{aligned}$$

for fixed Q, R, S, T .

The book uses $\vec{q} = (0, 0, 1)$, $\vec{r} = (1, 0, 0)$, $\vec{s} = (-\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}})$, $\vec{t} = (-\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}})$

$$\vec{r} \cdot \sigma \otimes \vec{s} \cdot \sigma = \begin{pmatrix} 0 & \vec{s} \cdot \sigma \\ \vec{r} \cdot \sigma & 0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & 1 \\ -1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \end{pmatrix}$$

$$\vec{r} \cdot \sigma \otimes \vec{t} \cdot \sigma = \begin{pmatrix} 0 & \vec{t} \cdot \sigma \\ \vec{r} \cdot \sigma & 0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 & -1 \\ 0 & 0 & -1 & -1 \\ 1 & -1 & 0 & 0 \\ -1 & -1 & 0 & 0 \end{pmatrix}$$

$$\vec{q} \cdot \sigma \otimes \vec{t} \cdot \sigma = \begin{pmatrix} \vec{t} \cdot \sigma & 0 \\ 0 & -\vec{t} \cdot \sigma \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

$$\vec{q} \cdot \sigma \otimes \vec{s} \cdot \sigma = \begin{pmatrix} \vec{s} \cdot \sigma & 0 \\ 0 & -\vec{s} \cdot \sigma \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & -1 \end{pmatrix}$$

$$\langle \vec{q} \cdot \sigma \otimes \vec{s} \cdot \sigma \rangle = \frac{1}{\sqrt{2}} (-\alpha^*(\alpha + \beta) + \beta^*(\beta - \alpha) + \gamma^*(\gamma + \delta) + \delta^*(\gamma - \delta))$$

$$\langle \vec{r} \cdot \sigma \otimes \vec{s} \cdot \sigma \rangle = \frac{1}{\sqrt{2}} (-\alpha^*(\gamma + \delta) + \beta^*(\delta - \gamma) - \gamma^*(\alpha + \beta) + \delta^*(\beta - \alpha))$$

$$\langle \vec{r} \cdot \sigma \otimes \vec{t} \cdot \sigma \rangle = \frac{1}{\sqrt{2}} (\alpha^*(\gamma - \delta) - \beta^*(\delta + \gamma) + \gamma^*(\alpha - \beta) - \delta^*(\beta + \alpha))$$

$$\langle \vec{q} \cdot \sigma \otimes \vec{t} \cdot \sigma \rangle = \frac{1}{\sqrt{2}} (\alpha^*(\alpha - \beta) - \beta^*(\beta + \alpha) + \gamma^*(\delta - \delta) + \delta^*(\gamma + \delta))$$

Quantifying entanglement: partial traces

$$\begin{aligned}\mathrm{Tr}_A(\rho_{AB}) &= \sum_{ijkl} \rho_{ij}^{kl} \mathrm{Tr}_A(|i\rangle \langle k|) \otimes |j\rangle \langle l| \\ &= \sum_i \left(\sum_{jl} \rho_{ij}^{il} |j\rangle \langle l| \right) \\ &= (\rho_{00}^{00} + \rho_{10}^{10}) |0\rangle \langle 0| + (\rho_{00}^{01} + \rho_{10}^{11}) |0\rangle \langle 1| + (\rho_{01}^{00} + \rho_{11}^{10}) |1\rangle \langle 0| + (\rho_{01}^{01} + \rho_{11}^{11}) |1\rangle \langle 1|\end{aligned}$$

$$\begin{aligned}\mathrm{Tr}_B(\rho_{AB}) &= \sum_{ijkl} \rho_{ij}^{kl} |i\rangle \langle k| \otimes \mathrm{Tr}_B(|j\rangle \langle l|) \\ &= \sum_j \left(\sum_{ik} \rho_{ij}^{kj} |i\rangle \langle k| \right) \\ &= (\rho_{00}^{00} + \rho_{01}^{01}) |0\rangle \langle 0| + (\rho_{00}^{10} + \rho_{01}^{11}) |0\rangle \langle 1| + (\rho_{10}^{00} + \rho_{11}^{01}) |1\rangle \langle 0| + (\rho_{10}^{10} + \rho_{11}^{11}) |1\rangle \langle 1|\end{aligned}$$

Reduced density matrices for an arbitrary state

$$\text{Tr}_A(\rho_{AB}) = \begin{pmatrix} \rho_{00}^{00} + \rho_{10}^{10} & \rho_{00}^{01} + \rho_{10}^{11} \\ \rho_{01}^{00} + \rho_{11}^{10} & \rho_{01}^{01} + \rho_{11}^{11} \end{pmatrix} = \begin{pmatrix} |\alpha|^2 + |\gamma|^2 & \alpha\beta^* + \gamma\delta^* \\ \beta\alpha^* + \delta\gamma^* & |\beta|^2 + |\delta|^2 \end{pmatrix}$$

$$\text{Tr}_B(\rho_{AB}) = \begin{pmatrix} \rho_{00}^{00} + \rho_{01}^{01} & \rho_{00}^{10} + \rho_{01}^{11} \\ \rho_{10}^{00} + \rho_{11}^{01} & \rho_{10}^{10} + \rho_{11}^{11} \end{pmatrix} = \begin{pmatrix} |\alpha|^2 + |\beta|^2 & \alpha\gamma^* + \beta\delta^* \\ \gamma\alpha^* + \delta\beta^* & |\gamma|^2 + |\delta|^2 \end{pmatrix}$$

Definition of entanglement entropy

The entanglement entropy of a bipartite system is the Von Neumann entropy of either reduced density matrix (it doesn't matter which one we choose)

$$S(\rho_A) = -\text{Tr}(\rho_A \log \rho_A) = -\sum_x \lambda_x \log \lambda_x$$

for eigenvalues λ_x of ρ_A . This tells us: do the reduced states ρ_A and ρ_B contain all the *information* in ρ_{AB} ?

Entanglement entropy and Tsirelson's bound

Sanity check:

Draw random reals $a, b, c, d, e, f, g, h \sim U([0, 1]^8)$

Construct $|\psi\rangle = (a + ib)|00\rangle + (c + id)|01\rangle + (e + if)|10\rangle + (g + ih)|11\rangle$

Normalize $|\psi\rangle \rightarrow \frac{|\psi\rangle}{\sum_n |c_n|^2}$

Compute $S(\rho_A)$ and $\langle\psi|(Q \otimes S + R \otimes S + R \otimes T - Q \otimes T)|\psi\rangle$