

TTIC 31230, Fundamentals of Deep Learning

David McAllester, Winter 2020

Stochastic Gradient Descent (SGD)

Heat Capacity with

Loss as Energy

and Learning Rate as Temperature

MCMC models of SGD

These Plots are from the original ResNet paper. Left plot is for CNNs without residual skip connections, the right plot is ResNet.

Thin lines are training error, thick lines are validation error.

In all cases η is reduced twice, each time by a factor of 2.

Converged Loss as a Function of η

For each value of η we converge at a loss $\mathcal{L}(\eta)$.

$$\begin{aligned}\mathcal{L}(0) &\doteq \lim_{\eta \rightarrow 0} \mathcal{L}(\eta) \\ &= \mathcal{L}(\Phi^*) \quad \Phi^* \text{ a local optimum}\end{aligned}$$

Can we do a Taylor expansion of $\mathcal{L}(\eta)$?

$$\mathcal{L}(\eta) = \mathcal{L}(\Phi^*) + \left(\frac{d\mathcal{L}}{d\eta} \bigg|_{\eta=0} \right) \eta + \dots$$

Heat Capacity

$$\mathcal{L}(\eta) = \mathcal{L}(\Phi^*) + \left(\frac{d\mathcal{L}}{d\eta} \Big|_{\eta=0} \right) \eta + \dots$$

Let i index a training example and let g_i denote $\nabla_{\Phi} \mathcal{L}_i(\Phi)$ at $\Phi = \Phi^*$.

Heat Capacity Theorem:

$$\frac{\partial \mathcal{L}(\eta)}{\partial \eta} \Big|_{\eta=0} = \frac{1}{4} E_i \|g_i\|^2$$

Proof Step 1

Let i index a training example and let $\mathcal{L}_i(\Phi^* + \Delta\Phi)$ be the loss on training example i with model parameters $\Phi^* + \Delta\Phi$.

We take a second order Taylor expansion.

$$\mathcal{L}(\Phi) = E_i \mathcal{L}_i(\Phi)$$

$$\mathcal{L}_i(\Phi^* + \Delta\Phi) = \mathcal{L}_i(\Phi^*) + g_i \Delta\Phi + \frac{1}{2} \Delta\Phi^\top H_i \Delta\Phi$$

$$E_i g_i = 0$$

$$E_i H_i \quad \text{is positive definite}$$

Proof: Step 2

Let Q_η be the stationary distribution on Φ defined by the SGD stochastic process.

Let P_η be the distribution on $\Delta\Phi = \Phi - \Phi^*$ with $\Phi \sim Q_\eta$.

$$\begin{aligned}\mathcal{L}(\eta) &= E_{\Delta\Phi \sim P_\eta} E_i \mathcal{L}_i + g_i \Delta\Phi + \frac{1}{2} \Delta\Phi^\top H_i \Delta\Phi \\ &= E_i \mathcal{L}_i(\Phi^*) + E_{\Delta\Phi \sim P_\eta} (E_i g_i) \Delta\Phi + \frac{1}{2} \Delta\Phi^\top (E_i H_i) \Delta\Phi \\ &= \mathcal{L}(\Phi^*) + E_{\Delta\Phi \sim P_\eta} \frac{1}{2} \Delta\Phi^\top (E_i H_i) \Delta\Phi\end{aligned}$$

Proof: Step 3

Because P_η is a stationary distribution on $\Delta\Phi$ we must have

$$E_{\Delta\Phi \sim P_\eta} E_i ||\Delta\Phi - \eta(g_i + H_i\Delta\Phi)||^2 = E_{\Delta\Phi \sim P_\eta} ||\Delta\Phi||^2$$

$$E_{\Delta\Phi \sim P_\eta} E_i - 2\eta\Delta\Phi^\top (g_i + H_i\Delta\Phi) + \eta^2 ||(g_i + H_i\Delta\Phi)||^2 = 0$$

$$E_{\Delta\Phi \sim P_\eta} \left(\frac{1}{2} \Delta\Phi^\top (E_i \ H_i) \Delta\Phi \right) = \frac{\eta}{4} E_{\Delta\Phi \sim P_\eta} E_i ||(g_i + H_i\Delta\Phi)||^2$$

$$\mathcal{L}(\eta) = \mathcal{L}(\Phi^*) + \frac{\eta}{4} E_{\Delta\Phi \sim P_\eta} E_i ||(g_i + H_i\Delta\Phi)||^2$$

Proof Step 4

$$\mathcal{L}(\eta) = \mathcal{L}(\Phi^*) + \frac{\eta}{4} E_{\Delta\Phi \sim P_\eta} E_i \|(g_i + H_i \Delta\Phi)\|^2$$

$$\left. \frac{\partial \mathcal{L}(\eta)}{\partial \eta} \right|_{\eta=0} = \frac{1}{4} \lim_{\eta \rightarrow 0} E_{\Delta\Phi \sim P_\eta} E_i \|(g_i + H_i \Delta\Phi)\|^2$$

$$= \frac{1}{4} E_i \|g_i\|^2$$

END