

Measures

Condom Sabotage

1. Agreeing to use a condom, but intentionally breaking the condom when putting it on
2. Agreeing to use a condom, but intentionally breaking the condom after it was on
3. Agreeing to use a condom, but removing it before or during sex without telling her

Event-Level Drinking

1. How many drinks did you consume on that day before having sex?
2. What was her highest level of intoxication while she was with you?

Alcohol Expectancies for Condom Use

1. When I drink, it is hard to use a condom.
2. Drinking makes me forget about using condoms.
3. After a few drinks it is more difficult for me to talk to a potential sexual partner about using a condom.

Sex-Related Drinking Motives

1. To feel less depressed about sexual activity
2. To forget your worries about sexual activity
3. To feel less nervous about sexual activity
4. To feel more self-confident and sure of yourself during sexual activity
5. To help you forget your problems related to sexual activity
6. To feel less upset about sexual activity
7. To make sexual activity more exciting
8. To make sexual activity more fun
9. To make you more relaxed during sexual activity

10. So that you will enjoy sexual activity more

11. To feel sexier

12. To increase your chances of having sexual activity

Enhancement subscale (mean of: SRDM4, SRDM7, SRDM8, SRDM9, SRDM10, SRDM11, SRDM12). Coping subscale (mean of: SRDM1, SRDM2, SRDM3, SRDM5, SRDM6)

