

The all-seeing eye of resonant Auger electron spectroscopy: a study on aqueous KCl

Tsveta Miteva,^{*,†} Nikolai V. Kryzhevoi,[‡] Nicolas Sisourat,[†] Christophe Nicolas,[¶]
Wandared Pokapanich,[§] Thanit Saisopa,^{||} Prayoon Songsiriritthigul,^{||} Yuttakarn
Rattanachai,[⊥] Andreas Dreuw,[#] Jan Wenzel,[#] Jérôme Palaudoux,[†] Gunnar
Öhrwall,[@] Ralph Püttner,[△] Lorenz S. Cederbaum,[‡] Jean-Pascal Rueff,^{†,¶} and
Denis Céolin^{*,¶}

[†]*Sorbonne Université, CNRS, Laboratoire de Chimie Physique Matière et Rayonnement,
UMR 7614, F-75005 Paris, France*

[‡]*Theoretische Chemie, Physikalisch-Chemisches Institut, Universität Heidelberg, Im
Neuenheimer Feld 229, D-69120 Heidelberg, Germany*

[¶]*Synchrotron SOLEIL, l'Orme des Merisiers, Saint-Aubin, F-91192 Gif-sur-Yvette Cedex,
France*

[§]*Faculty of Science, Nakhon Phanom University, Nakhon Phanom 48000, Thailand*

^{||}*School of Physics, Suranaree University of Technology, Nakhon Ratchasima 30000,
Thailand*

[⊥]*Department of Applied Physics, Faculty of Sciences and Liberal Arts, Rajamangala
University of Technology Isan, Nakhon Ratchasima 30000, Thailand*

[#]*Interdisciplinary Center for Scientific Computing, Ruprecht-Karls University, Im
Neuenheimer Feld 205A, D-69120 Heidelberg, Germany*

[@]*MAX IV Laboratory, Lund University, P.O. Box 118, SE-22100 Lund, Sweden*

[△]*Fachbereich Physik, Freie Universität Berlin, Arnimallee 14, D-14195, Berlin, Germany*

E-mail: tsveta.miteva@upmc.fr; denis.ceolin@synchrotron-soleil.fr




Figure 1: Radial density distributions of the singly-occupied natural orbital occupied by the excited electron corresponding to the $1s \rightarrow 4p$, $1s \rightarrow 3d$ and $1s \rightarrow 5p$ core excitations in K^+ (lower panel) and Cl^- (upper panel). The insets show the region of distances relevant for the overlap with the $1s$ core orbital whose radial density is shown as a grey shaded area.

References

- (1) Föhlisch, A.; Feulner, P.; Hennies, F.; Fink, A.; Menzel, D.; Sanchez-Portal, D.; Echenique, P. M.; Wurth, W. *Nature* **436**, 373.




Figure 2: Partial cross sections and charge transfer time extracted from Fig. ???. The blue and red curves are obtained by integrating the area of the $2p^{-2}$ and $2p^{-2}4p$ final states (1D state region only) at each photon energy step. From these curves we determine the charge transfer time τ_{CT} according to the formula $\tau_{CT} = \tau l/d$, with τ being the Cl 1s core-hole lifetime and l/d being the intensity ratio of the localized ($2p^{-2}4p$) and delocalized ($2p^{-2}$) states at a given excitation energy.¹ The green line defines the $Cl_{aq}^{-}(1s)$ ionization potential.