



# The all-seeing eye of resonant Auger electron spectroscopy: a study on aqueous KCl

Tsveta Miteva,<sup>\*,†</sup> Nikolai V. Kryzhevoi,<sup>‡</sup> Nicolas Sisourat,<sup>†</sup> Christophe Nicolas,<sup>¶</sup>  
Wandared Pokapanich,<sup>§</sup> Thanit Saisopa,<sup>||</sup> Prayoon Songsiriritthigul,<sup>||</sup> Yuttakarn  
Rattanachai,<sup>⊥</sup> Andreas Dreuw,<sup>#</sup> Jan Wenzel,<sup>#</sup> Jérôme Palaudoux,<sup>†</sup> Gunnar  
Öhrwall,<sup>@</sup> Ralph Püttner,<sup>△</sup> Lorenz S. Cederbaum,<sup>‡</sup> Jean-Pascal Rueff,<sup>†,¶</sup> and  
Denis Céolin<sup>\*,¶</sup>

<sup>†</sup>*Sorbonne Université, CNRS, Laboratoire de Chimie Physique Matière et Rayonnement,  
UMR 7614, F-75005 Paris, France*

<sup>‡</sup>*Theoretische Chemie, Physikalisch-Chemisches Institut, Universität Heidelberg, Im  
Neuenheimer Feld 229, D-69120 Heidelberg, Germany*

<sup>¶</sup>*Synchrotron SOLEIL, l'Orme des Merisiers, Saint-Aubin, F-91192 Gif-sur-Yvette Cedex,  
France*

<sup>§</sup>*Faculty of Science, Nakhon Phanom University, Nakhon Phanom 48000, Thailand*

<sup>||</sup>*School of Physics, Suranaree University of Technology, Nakhon Ratchasima 30000,  
Thailand*

<sup>⊥</sup>*Department of Applied Physics, Faculty of Sciences and Liberal Arts, Rajamangala  
University of Technology Isan, Nakhon Ratchasima 30000, Thailand*

<sup>#</sup>*Interdisciplinary Center for Scientific Computing, Ruprecht-Karls University, Im  
Neuenheimer Feld 205A, D-69120 Heidelberg, Germany*

<sup>@</sup>*MAX IV Laboratory, Lund University, P.O. Box 118, SE-22100 Lund, Sweden*


<sup>△</sup>*Fachbereich Physik, Freie Universität Berlin, Arnimallee 14, D-14195, Berlin, Germany*

E-mail: tsveta.miteva@upmc.fr; denis.ceolin@synchrotron-soleil.fr

## Abstract

X-ray absorption and Auger electron spectroscopies are powerful tools to probe the electronic structure and immediate surroundings of ions in solution. In this work we use a combination of these methods together with *ab initio* calculations to study the electronic structure and decay in aqueous KCl at the K-edges of  $\text{K}^+$  and  $\text{Cl}^-$ . Although the two ions are isoelectronic, their Auger electron spectra exhibit notably different features. This is due to the fact that the energetic order of the  $1s^{-1}3d$  and  $1s^{-1}4p$  core excited states is inverted in  $\text{K}^+$  with respect to  $\text{Cl}^-$ , and moreover the two states lie energetically close in the bare  $\text{K}^+$  ion leading to their mixing in the presence of the solvent. As a result, the dipole forbidden  $1s^{-1}3d$  state is populated upon K-shell excitation of aqueous  $\text{K}^+$ . The resonant Auger decay of this state results in a separate feature in the Auger electron spectrum of  $\text{K}^+$  which is absent in the spectrum of  $\text{Cl}^-$ . The results of this work represent a pioneering study of the decay processes initiated by photoabsorption in the tender x-ray regime close to threshold in liquids and are thus of importance in unveiling the mechanisms of radiation damage in biologically relevant systems.

## Graphical TOC Entry



## Keywords

Solvated ions, resonant Auger spectroscopy, X-ray absorption spectroscopy

X-ray absorption (XAS) and Auger electron spectroscopies (AES) are powerful tools to study the electronic structure and the nearest environment of atoms and molecules in gas, liquid and solid phase. Understanding how atoms or molecules respond to irradiation with x-rays gives insight into the structure of solutions (see Ref.<sup>1</sup> and references therein), and the mechanism of radiation damage<sup>2-4</sup>. Upon absorption of an x-ray photon, either core excited or core ionized states of a specific atom are populated depending on the photon energy. The relaxation of these highly energetic states involves an ultrafast cascade of intraatomic processes, such as radiative and Auger decays. Furthermore, if the initially excited or ionized species is embedded in an environment, interatomic processes such as charge and energy transfer<sup>4-8</sup> are possible.

It has been well understood in atoms and molecules in gas phase that the course of a decay cascade depends on the character of the initially populated states<sup>9-16</sup>. In a solution, the electronic decay processes initiated by x-ray photoabsorption are different compared to those in rare gas clusters due to the shorter distances and stronger interatomic interactions. In particular, the solvent molecules have two effects – first, they strongly affect the excited<sup>17</sup> or ionized states of the ion and second, they can participate in the decay processes, leading to the population of charge-separated final states, and ionization of the surrounding environment<sup>4-8</sup>. Moreover, the process of delocalization of the initially excited electron also occurs in aqueous solutions<sup>18,19</sup>. In the case of pure water, the delocalization of the O1s excited electron takes place on a femtosecond to sub-femtosecond time scale depending on the photon energy, thus being commensurate with the lifetime of the O1s core hole which is  $\sim 3.6$  fs<sup>18</sup>. The O K-edge is located in the soft x-ray range of photon energies,  $\sim 533$  eV in water vapor<sup>20</sup>. Going higher in photon energy, to the tender and hard x-ray regimes, the lifetimes of the core ionized or core excited states become even shorter, on the order of 1 fs. And thus, it is even more imperative to reveal whether the delocalization of the core excited electron occurs within the lifetime of the core hole.

The aim of this work is to elucidate the nature of the states populated upon x-ray

irradiation of solvated ions in the tender x-ray regime, and furthermore, to understand whether the process of delocalization influences the resonant Auger decay. To this end, we used Auger electron spectroscopy together with x-ray absorption spectroscopy in the tender x-ray regime to study aqueous potassium chloride at the K-edges of both  $\text{K}^+$  and  $\text{Cl}^-$ . In particular, we demonstrate experimentally that at photon energies below the K-edges of the two ions, core excited states are populated. These states undergo resonant Auger decay within less than 1 fs. In both ions, there is a competition between resonant Auger decay and delocalization of the excited electron. Using the core-hole clock method we show that in  $\text{K}^+$  delocalization at the pre-edge is slower than the Auger process. However, in the case of  $\text{Cl}^-$ , the rate of delocalization is of the same order as that of the resonant Auger process due to the energetic proximity of the core excited state to the ionization threshold. Moreover, we observe that although the  $\text{K}^+$  and  $\text{Cl}^-$  ions are isoelectronic, they have different fingerprints in the resonant Auger spectra. With the aid of high-level *ab initio* calculations of the initial and final states of the resonant Auger process of both the bare ions and their microsolvated clusters, we demonstrate that these differences result from different electronic structures of the two ions, thus confirming that the combination of XAS and AES techniques is a sensitive probe of the electronic structure of solutions.

For the present experiment we used the newly operational microjet setup that was specifically designed for the HAXPES station of the GALAXIES beamline<sup>21,22</sup>. A differentially-pumped tube in which the microjet head is inserted, is mounted on a 3-axes motorized manipulator in front of the spectrometer lens. Two holes of 2 mm diameter allow the photons to go in and out. At the end of the tube and in front of the lens, a 500  $\mu\text{m}$  diameter hole skimmer allows the electrons created at the interaction point to go in the direction of the spectrometer. The microjet head is mostly composed of a 30  $\mu\text{m}$  diameter vertical glass capillary facing a temperature-controlled catcher in CuBe having a 300  $\mu\text{m}$  hole, and a camera. Piezo motors allow their precise alignment relative to each other and to the photon beam. The catcher is placed at a distance of about 5 mm from the capillary and is permanently




Figure 1: Schematic representation of the resonant (left) and normal (right) Auger processes of the isoelectronic  $\text{K}^+$  and  $\text{Cl}^-$  ions. The charges of the final Auger states in the two ions are also shown.

pumped in order to extract the liquid. For the present experiment, a 0.5M KCl aqueous solution is injected in the capillary by a high performance liquid chromatography (HPLC) pump with a constant flux of 1.6 ml/min. The alignment of the setup is performed on the KCl aqueous solution by measuring the water O1s x-ray photoelectron peak intensity and by optimizing the liquid vs gas phase ratio. The pressure in the main chamber is kept below the  $10^{-5}$  mbar range whereas it is kept at about  $10^{-4}$  mbar in the differentially-pumped tube when the HPLC pump is ON. Our equipment is an updated version of the equipment used in Ref.<sup>23</sup> The aqueous potassium chloride solution was prepared by mixing >99% KCl salt with deionized water. Filtering and degazing procedures were systematically performed before injecting the solution. The spectrometer resolution of about 0.6 eV was achieved with the 500 eV pass energy and 0.5 mm slits. The photon energy resolution achieved at 2.8 keV and 3.6 keV was about 0.3 eV and 0.4 eV, respectively. The experimental 2D maps representing the evolution of the KLL Auger spectra in the vicinity of the  $\text{Cl}^-$  and  $\text{K}^+$  K-edges, as a function of the photon energy, are shown in Figs. 2 and 3, respectively. The aqueous  $\text{K}^+$  and  $\text{Cl}^-$  1s ionization potentials were measured at  $h\nu = 5$  keV and calibrated on the

liquid contribution of the O1s XPS spectrum.<sup>24</sup> The maps were also calibrated using the O1s photoelectron line of liquid water but at photon energies close to the potassium and chloride 1s ionization thresholds.

The theoretical X-ray absorption spectra were computed for the hexa-coordinated clusters of both ions,  $\text{K}^+(\text{H}_2\text{O})_6$  and  $\text{Cl}^-(\text{H}_2\text{O})_6$ , which can be considered as representatives of the complete first solvation shell of the two ions<sup>25–27</sup>. The two structures shown in Fig. 4 were optimized at the DFT level of theory using the B3LYP functional and the 6-311++G(2d,2p) basis set<sup>28,29</sup>. The geometry optimization was performed with the Gaussian 09 package<sup>30</sup>. In order to obtain a realistic structure for  $\text{K}^+$  corresponding to the bulk solution, we carried out constrained geometry optimization by choosing the equilibrium gas-phase geometries<sup>31,32</sup> belonging to the  $D_3$  point group and then increasing the angle  $\theta$  between the K-O bond and the  $C_3$  axis to  $55^\circ$ . This angle was chosen such that the O-K-O angles are around the maxima of the angular distributions obtained from quantum mechanics/molecular mechanics simulations in Ref.<sup>27</sup>. Moreover, we fixed the K-O distance to 2.840 Å, such that it corresponds to the distances from other theoretical and experimental works<sup>25–27</sup>.

The energies and transition moments of the core excited states of the bare ions and micro-solvated clusters were computed with the algebraic diagrammatic construction method for the polarization propagator<sup>33</sup> within the core-valence separation approximation<sup>34–36</sup> (CVS-ADC(2)x) as implemented in the Q-Chem package<sup>37–40</sup>. In the case of  $\text{Cl}^-$  the 6-311++G(3df,3pd) basis set<sup>28,41</sup> (excluding f functions) was used on all atoms, whereas in the case of  $\text{K}^+$  we used the 6-311+G(2d,p) basis set<sup>28,29</sup> on all atoms, and two additional sets of s, p and d diffuse functions were added on K. The use of a smaller basis set in the case of K is due to the higher number of atomic orbitals compared to the case of Cl, and therefore, prohibitively high cost of the CVS-ADC(2)x computation. In our calculations the core space comprises the 1s orbital of  $\text{K}^+$  or  $\text{Cl}^-$ , whereas the remaining occupied orbitals are included in the valence space. For the calculations of the XAS spectra we used the  $C_2$  point group in the case of  $\text{K}^+(\text{H}_2\text{O})_6$  and  $\text{Cl}^-(\text{H}_2\text{O})_6$ . To account for the lifetime broadening due to the Auger decay

of the core excited states, we convolved the theoretical spectra with a Lorentzian function of FWHM 0.74 eV and 0.62 eV in the case of  $\text{K}^+$  and  $\text{Cl}^-$ , respectively<sup>42</sup>. Additionally, we convolved the theoretical spectra with a Gaussian profile to also account for the experimental resolution (see Fig. 4). We analyzed the core excited states by expanding the natural orbitals occupied by the excited electron (singly occupied natural orbitals, SONOs)  $\psi_i$  of the microsolvated clusters in the basis of SONOs of the bare  $\text{K}^+$  or  $\text{Cl}^-$  ions,  $\chi_{nl}$

$$\psi_i = \sum_{nl} a_{nl}^i \chi_{nl} \quad (1)$$

where  $n$  and  $l$  stand for the principal and orbital quantum numbers as described in Ref.<sup>17</sup>. The expansion coefficients  $a_{nl}^i$  show the degree of delocalization of the excited electron and the mixing of the core excited states in the ligand field created by the surrounding water molecules (see Fig. 4).


The final states following KLL resonant Auger decay of  $\text{K}^+(\text{H}_2\text{O})_6$  and  $\text{Cl}^-(\text{H}_2\text{O})_6$  were computed at the Configuration Interaction Singles (CIS) level using the Graphical Unitary Group Approach (GUGA) as implemented in the GAMESS-US package<sup>43–45</sup>. In order to account for the relaxation effects upon core ionization, we employed a restricted open-shell Hartree-Fock reference wave function with a hole in the 2s orbital of both  $\text{K}^+$  and  $\text{Cl}^-$ . We used the 6-311++G(2d,2p) basis set<sup>28,29,41</sup> on all atoms. Additionally, the basis set was augmented with two sets of s, p, d diffuse functions in the case of  $\text{K}^+$ , and three sets of s, p, d diffuse functions in the case of  $\text{Cl}^-$ . The larger basis set employed in the case of Cl was necessary in order to ensure the convergence of the excited states. The active space comprises the 2s and 2p orbitals of K/Cl with occupancy fixed to 6 and all virtual orbitals with occupancy fixed to 1. The remaining doubly occupied orbitals were frozen in the calculation.<sup>46</sup>



# Results and discussion

## Normal Auger decay

The  $KL_{2,3}L_{2,3}$  normal Auger decay following K-shell ionization of aqueous  $K^+$  and  $Cl^-$  can be written as follows



It populates the  $2p^{-2}(^3P, ^1D, ^1S)$  final states. The  $^3P$  final states have a very low intensity since the corresponding transitions are forbidden from angular momentum and parity conservation rules. In the case of  $K_{aq}^+$  the maxima of the  $^1S$  and  $^1D$   $KL_{2,3}L_{2,3}$  Auger lines at photon energy  $h\nu = 3616.0$  eV are located at 2958.0 eV and 2968.5 eV kinetic energy, respectively (see Fig. 2). For  $Cl_{aq}^-$ , the lines corresponding to the  $Cl^+ 2p^{-2}(^1S)$  and ( $^1D$ ) states are located at 2373.1 eV and 2382.0 eV kinetic energy for photon energy  $h\nu = 2830.0$  eV (see Fig. 3).

The  $KL_{2,3}L_{2,3}$  normal Auger lines may disperse with photon energy close to threshold due to energy exchange between the photoelectron and Auger electron called post-collision interaction (PCI). As a result of this interaction, first, an Auger peak becomes asymmetric with a high kinetic energy tail, and second, its intensity maximum is shifted to higher kinetic energies close to threshold<sup>47,48</sup>. These effects are seen in our spectra – the asymmetric tail of the peaks resulting from PCI is clearly shown on the Auger spectra at photon energies 3616 eV in the case of  $K^+$ , and 2830 eV in the case of  $Cl^-$ .

Further, we compare the positions of the normal KLL Auger lines of both  $Cl_{aq}^-$  and  $K_{aq}^+$  close to threshold with those recorded far from threshold, at photon energies  $h\nu = 5$  keV (see Ref.<sup>8</sup> for details). In the latter case, the maxima of the  $^1D$  and  $^1S$  states appear at 2381.0 eV and 2372.0 eV for  $Cl_{aq}^-$ , and 2967.4 eV and 2957.0 eV for  $K_{aq}^+$ , respectively. The




Figure 2: 2D map showing the kinetic energy of the electrons emitted in  $KL_{2,3}L_{2,3}$  Auger decay vs the photon energy in the vicinity of the K-edge of aqueous  $K^+$ . The black curve on the left represents the experimental partial electron yield spectrum of  $K^+$  obtained after integrating over the kinetic energies of the Auger electrons in the energy range presented on the figure. The upper panel shows two spectra at photon energies 3610.7 eV, and 3616 eV below and above the ionization potential at 3611.9 eV, respectively. The vertical bars in the resonant Auger spectrum measured at 3610.7 eV indicate the energy positions of the theoretical final  $2p^{-2} 3d$  (blue) and  $2p^{-2} 4p$  (red) resonant Auger states of  $K^+(H_2O)_6$  of doublet spin multiplicity. The features A, B and C are discussed in the text.

maxima were found to be shifted by  $\sim 1$  eV towards lower kinetic energies as compared to the spectra measured close to threshold. The magnitude of the shift is constant in the photon energy range of  $\sim 8$  eV above threshold (see Figs. 2 and 3) and similar for the two ions. A possible explanation of the shift observed in our experiment is given in Ref.,<sup>49</sup> which focuses on the Auger decay of large Kr clusters close to the  $3d_{5/2}$  ionization threshold. The observed  $4p^{-2}(^1S, ^1D, ^3P)$  Auger peaks were found to be shifted by 0.7 eV to higher kinetic energies compared to the positions of the peaks far above threshold. Moreover, the shift did not vary




Figure 3: 2D map showing the kinetic energy of the electrons emitted in  $KL_{2,3}L_{2,3}$  Auger decay vs the photon energy in the vicinity of the K-edge of aqueous  $Cl^-$ . The black curve on the left represents the experimental partial electron yield spectrum of  $Cl^-$  obtained after integrating over the kinetic energies of the Auger electrons in the energy range presented on the figure. The upper panel shows two spectra at photon energies 2825.2 eV, and 2830.0 eV below and above the ionization potential at 2825.4 eV, respectively. The vertical bars in the resonant Auger spectrum at 2825.2 eV indicate the energy positions of the theoretical  $2p^{-2}4p$  states of  $Cl^-(H_2O)_6$  of doublet spin multiplicity.


with the photon energy close to threshold. Consequently, it was proposed that these features originate from a process of internal ionization, i.e. excitation of the photoelectron into the conduction band, followed by normal Auger decay. The observed shift was explained as resulting from the PCI-like interaction between the electron excited in the conduction band and the Auger electron. Further investigations are, however, planned in the case of liquid samples.

Finally, the normal Auger  $^1D$  main line of  $K^+$  differs from that of  $Cl^-$  by the presence of a large shoulder (feature A on Fig. 2) on the low kinetic energy side at about 2965 eV kinetic energy. This shoulder is attributed to electron transfer from the solvent water molecules to

the unoccupied 3d orbitals of  $K^{+8}$ . In the case of  $Cl^{-}$ , there is no experimental evidence of such intense electron transfer processes.

## Resonant Auger decay

The  $KL_{2,3}L_{2,3}$  Auger decay following resonant K-shell excitation of solvated  $K^{+}$  and  $Cl^{-}$  can be written as follows



where  $V$  and  $V'$  denote the unoccupied orbitals in the presence of the  $1s^{-1}$  and  $2p^{-2}$  core holes, respectively. The character of these states is discussed below.

The pre-edge regions of the x-ray absorption spectra of  $K^{+}$  and  $Cl^{-}$  shown in the left panels of Figs. 2 and 3 do not exhibit any high intensity peaks owing to the lifetime broadening and energetic proximity of the core excited states to the ionization threshold. Consequently, solely from these absorption spectra, one cannot conclude whether there are core excited states in the pre-edge structure. Instead these states have to be identified by their resonant Auger features which differ from the normal Auger features of core ionized states. Thus, for  $Cl^{-}$ , the lowest core excited state is located at 2825.2 eV, which agrees very well with the position of the  $Cl^{-} 1s \rightarrow 4p$  excitation determined from Cl K-edge XAS experiments in hydrated metal dichlorides,  $MgCl_2 \cdot 6H_2O$  and  $SrCl_2 \cdot 6H_2O$ <sup>50</sup>, and tetrahedral  $MCl_4^{-}$  complexes (where  $M = Cu^{II}, Ni^{II}, Co^{II}, Fe^{II}$  and  $Fe^{III}$ )<sup>51</sup>. In the case of  $K^{+}$ , there are two dispersive features with maxima at photon energies of 3611.2 eV (B) and 3611.6 eV (C), respectively (Fig. 2). The positions of these two core excited states are close to the energy of the  $1s^{-1}4p$  excitation in bare  $K^{+}$  at 3610.7 eV<sup>52</sup>.

The resonant Auger features produced in the decay of these core excited states are quite different for  $Cl^{-}$  and  $K^{+}$ . In the 2D map of  $Cl^{-}$  shown in Fig. 3 there are two dispersive

features, indicated with diagonal dashed lines, each on the high kinetic energy side of the  $^1S$  and  $^1D$  main lines. The maxima of these features are at 2825.2 eV photon energy and 2374.6 and 2383.4 eV kinetic energy, respectively. In the case of  $K^+$  (see Fig. 2), the dispersive line related to the  $^1S$  main line cannot be clearly identified due to the presence of strong background. Instead two dispersive features related to the  $^1D$  main line are observed (features B and C indicated with diagonal dashed lines on Fig. 2). Feature B exhibits a maximum at  $h\nu = 3611.2$  eV and 2969.2 eV kinetic energy. The additional feature C appears as a separate island away from the main lines on the 2D map of  $K^+$ . It is located at  $h\nu = 3611.6$  eV and 2978.1 eV kinetic energy, thus it is separated by approximately 400 meV in photon energy and 8.3 eV in kinetic energy from the maximum of feature B.

In order to rationalize the pre-edge region of the experimental XAS spectra and the differences in the AES spectra of  $K^+_{aq}$  and  $Cl^-_{aq}$ , we computed the lowest core excited states of the bare  $K^+$  and  $Cl^-$  ions and their hexa-coordinated clusters. The theoretical XAS spectra are presented in Fig. 4. In the bare ions (lowermost panels of Fig. 4), the lowest energy peak corresponds to the dipole allowed  $1s^{-1}4p$  state. The next dipole allowed state,  $1s^{-1}5p$ , is located 4.3 eV and 10.8 eV higher in the cases of  $K^+$  and  $Cl^-$ , respectively. Together with the two dipole allowed transitions, we show the dipole forbidden  $1s^{-1}3d$  states of the bare ions as blue crosses at photon energies 3611.2 eV in the case of  $K^+$  and 2831.7 eV in the case of  $Cl^-$ , respectively. It is noteworthy that the positions of the  $1s^{-1}4p$  and  $1s^{-1}3d$  states are inverted in  $K^+$  and  $Cl^-$ . In the case of  $Cl^-$  the  $1s^{-1}4p$  excitation has lower energy and the  $1s^{-1}3d$  excitation is close to the  $1s^{-1}5p$  state. On the contrary, in  $K^+$  the  $1s^{-1}3d$  excitation has lower energy and lies below the  $1s^{-1}4p$  state. We note in passing that the intensity of the  $Cl^-(1s^{-1}4p)$  state is lower than that of the  $Cl^-(1s^{-1}5p)$  state contrary to what is observed in  $K^+$ . This difference can be explained by the lower electron density of the 4p compared to the 5p electron in the region close to the core hole which thus results in the lower oscillator strength of the  $1s^{-1}4p$  compared to the  $1s^{-1}5p$  transition in  $Cl^-$  (see Fig. 1 in SI).




Figure 4: XAS spectra of the lowest K-shell resonant transitions in the bare  $\text{K}^+$  (lower left panel) and  $\text{Cl}^-$  (lower right panel) ions and their 6-coordinated clusters (upper left panel,  $\text{K}^+(\text{H}_2\text{O})_6$ , and upper right panel,  $\text{Cl}^-(\text{H}_2\text{O})_6$ ). The theoretical stick spectra are convoluted with a Lorentzian profile of FWHM 0.74 eV for  $\text{K}^+$  and 0.62 eV for  $\text{Cl}^-$  (dashed line)<sup>8</sup> and a Voigt profile to account for both the lifetime broadening and the experimental resolution (full line). The stick spectrum corresponds to the projections  $|a_{nl}^i|^2$  of the SONOs corresponding to the core excited states of the 6-coordinated clusters on the basis of SONOs corresponding to the  $1s^{-1}3d$ ,  $1s^{-1}4p$ , and  $1s^{-1}5p$  states in the bare  $\text{K}^+$  and  $\text{Cl}^-$  ions (Eq. 1). The grey sticks in the stick spectrum correspond to contributions from higher-lying atomic core excitations or from excitations to the solvent molecules. The theoretical XAS spectra of both  $\text{K}^+$  and  $\text{Cl}^-$  were shifted to higher photon energies such that the excitation energies of the lowest core excited states correspond to the experimentally determined energies – 3610.7 eV in the case of  $\text{K}^+$ , and 2825.2 eV in the case of  $\text{Cl}^-$ . The experimental ionization thresholds are depicted as grey boxes starting at photon energies of 3611.9 eV ( $\text{K}^+_{\text{aq}}$ ) and 2825.4 eV ( $\text{Cl}^-_{\text{aq}}$ ).

The water molecules in the first solvation shell have several effects on the core excited states. First, upon addition of water molecules, the degeneracy of the  $1s^{-1}4p$  state is lifted and the intensity of the resulting states in the cluster drops. Moreover, the character of these states changes as can be seen on the upper panels of Fig. 4. They are no longer of pure atomic character but they rather interact with states of the neighboring water molecules (shown as grey bars in Fig. 4) and with other closely lying states of the bare ion, such as the dipole allowed  $1s^{-1}5p$  and dipole forbidden  $1s^{-1}3d$  state. Thus, the latter also acquire intensity in the cluster due to mixing with the dipole allowed states in the ligand field of the

solvent. A similar effect was observed in the XAS spectra of microsolvated clusters of  $\text{Na}^+$  and  $\text{Mg}^{2+}$ <sup>17</sup>.

Further we assume that only the lowest peak in the theoretical XAS spectra is populated in the experiment for two reasons. First, due to the lifetime broadening, it spreads over approximately 2 eV which coincides with the width of the pre-edge structure in the experimental XAS spectra. Second, the splitting between the first core excited state and the ionization threshold in the experiment is 1.2 eV for  $\text{K}^+$  and 0.2 eV for  $\text{Cl}^-$ , and thus it is smaller than the splitting between the first and second peak in the theoretical spectra (1.5 eV for  $\text{K}^+$  and  $\sim 3$  eV for  $\text{Cl}^-$ , Fig. 4). In the 6-coordinated cluster (Fig. 4 upper left panel), which represents the complete first solvation shell around  $\text{K}^+$ , the lowest peak in the spectrum contains three states. The lowest and highest lying states are split by approximately 0.5 eV and they have mixed 4p and 5p character. The low intensity state in between these two states has a predominantly  $1s^{-1}3d$  character. Since the dispersive feature B appears at lower excitation energies compared to the feature C, we assume that it is produced in the resonant Auger decay of the lowest core excited states of  $\text{K}^+$ , which are predominantly of  $1s^{-1}4p$  character. Moreover, we can attribute the feature C to the resonant Auger decay of the low intensity dipole forbidden  $1s^{-1}3d$  state. Thus, we explain both the energy splitting of  $\sim 400$  meV photon energy of the two features, and the fact that island C has lower intensity than feature B (Fig. 2).

In the hexa-coordinated cluster of  $\text{Cl}^-$ , the solvent molecules have little influence on the position and character of the first state. Due to the large energy separation from other  $\text{Cl}^-$  states it has mainly  $\text{Cl}^- 1s^{-1}4p$  character with some admixture of states of the nearest water molecules. We therefore attribute the two dispersive features associated with the  $^1\text{S}$  and  $^1\text{D}$  main lines on the 2D map of  $\text{Cl}^-$  to the resonant Auger decay of this core excited state involving mostly the 4p orbitals of chloride.

To fully characterize the dispersive features on the experimental 2D maps, we also computed the lowest  $\text{K}^{2+}[2p^{-2}\text{nl}](\text{H}_2\text{O})_6$  and  $\text{Cl}^0[2p^{-2}\text{nl}](\text{H}_2\text{O})_6$  states of the hexa-coordinated

clusters corresponding to the lowest final spectator resonant Auger states. The energy positions of these lines are shown as bars on the experimental resonant Auger spectra in the upper panels of Figs. 2 and 3. In both cases, we shifted the lowest  $2p^{-2}(^1D)4p$  states such that the kinetic energies coincide with the maxima of the dispersive features on the high kinetic energy part of the  $^1D$  main line. Note that we only indicate resonant Auger transitions to the spin allowed final states of doublet multiplicity. Since the initial core excited states are populated by photon excitation, only doublet states are efficiently populated in the resonant Auger decay. The energetic positions of the  $2p^{-2}nl$  states of  $K^+(H_2O)_6$  and  $Cl^-(H_2O)_6$  are substantially different, reflecting the fact that the 3d unoccupied orbitals of  $K^+$  are lower than the 4p orbitals. This is the opposite of what is observed in  $Cl^-$ , where the  $2p^{-2}3d$  states (not shown) are located on the lower kinetic energy side of the respective  $2p^{-2}4p$  states.

As mentioned above, we attribute the island B to the decay of the lowest lying core excited state, which has predominantly  $1s^{-1}4p$  character. Supposing that this state undergoes mostly pure spectator resonant Auger decay, which is the case of the  $1s^{-1}4p$  state in the isoelectronic Ar atom<sup>53</sup>, then the lowest states of  $2p^{-2}4p$  character are populated resulting in resonant Auger lines located between 2969.0 and 2970.5 eV kinetic energy. As can be seen from the Auger electron spectrum at  $h\nu = 3610.7$  eV (upper panel of Fig. 2), the lowest  $2p^{-2}4p$  states of  $K^+(H_2O)_6$  starting at electron energy of 2969 eV are separated by  $\sim 5$  and 12-13 eV from the two groups of  $2p^{-2}3d$  states located at higher electron kinetic energies. Thus, the group of  $2p^{-2}3d$  states at  $\sim 2975$  eV lies closer to the position of island C. Consequently, we attribute this dispersive feature as originating from the resonant Auger decay of the  $1s^{-1}3d$  excitation to the group of  $2p^{-2}3d$  states. The splitting between the  $2p^{-2}4p$  and  $2p^{-2}3d$  states in our calculation is smaller than the splitting between the islands B and C. This difference may be due to the fact that we do not account for the effect of distant solvent shells in our calculation. Concerning the higher lying group of  $2p^{-2}3d$  states at kinetic energies between 2982 and 2983 eV, we conclude that these states are not populated via the Auger process



since no additional experimental features are observed.

The energy difference between the spectral features A and C provides another argument in favor of the  $2p^{-2}3d$  character of the feature C. The feature A originates from electron transfer processes from water molecules (W) to the doubly core ionized potassium ion and has the configuration  $K^{2+}(2p^{-2}3d)W^{-1}$ . The lowest ionization potential of liquid water is about 11.16 eV<sup>54</sup> which fits well with the observed A-C splitting. Based on the above energetic arguments we attribute the island C as originating from resonant Auger decay to the  $K^{2+} 2p^{-2}3d$  final state.

In the computed  $Cl^0[2p^{-2}nl](H_2O)_6$  spectrum there are two groups of states split by about 7 eV (see upper panel of Fig. 3). The lower kinetic energy group corresponds to the  $2p^{-2}(^1S)4p$  final states, whereas the higher kinetic energy group corresponds to the  $2p^{-2}(^1D)4p$  final states. The splitting between these two groups is in good agreement with the experimental splitting between the dispersive features on the high kinetic energy sides of the  $^1S$  and  $^1D$  main peaks. Consequently, we attribute these dispersive features as resulting from the resonant Auger decay of the  $1s^{-1}4p$  core excited state of  $Cl^-_{aq}$  to the  $2p^{-2}(^1S)4p$  and  $2p^{-2}(^1D)4p$  final states. Similar dispersive features originating from the decay of the Cl ( $1s^{-1}4p$ ) state were observed on the 2D map of chloromethane  $CH_3Cl$  recorded in the vicinity of the Cl K-edge in gas phase.<sup>55</sup> In this case, however, additional lower lying core excited states are observed. They result from excitation to the LUMO of  $CH_3Cl$ , which is a linear combination of the C 2p and Cl 3p atomic orbitals. Since the 3p shell is fully occupied in  $Cl^-$ , such core excited states are not observed in our experiment. We note that neither  $2p^{-2}3d$  nor charge transfer states are found in the  $Cl^-_{aq}$  case.

In summary, we studied the electronic structure of aqueous solution of KCl at the K-edges of both K and Cl using a combination of x-ray absorption and Auger electron spectroscopy in the tender x-ray regime, and *ab initio* calculations. The Auger electron spectra of both ions as a function of photon energy exhibit features of normal as well as resonant Auger processes. The latter process proceeds differently for aqueous  $K^+$  and  $Cl^-$  due to the population of the

dipole forbidden  $K^+ 1s^{-1}3d$  state in a solution. The spectator Auger decay of this state produces an additional dispersive feature which is manifested as a separate peak in the Auger electron spectrum. In the case of  $Cl^-$  only fingerprints of the population and Auger decay of the dipole allowed  $1s^{-1}4p$  excitation are observed in the spectrum.

Our work shows that the combination of x-ray absorption and resonant Auger spectroscopy is a sensitive probe of the electronic structure of solvated ions. The reported results are an important first step in the study of the chains of relaxation steps triggered by photoabsorption in the tender x-ray regime. The Auger processes considered here are inevitably followed by multiple intra- and interatomic electronic decays, such as interatomic Coulombic decay (ICD) and electron-transfer mediated decay (ETMD). As a result of the latter processes, genotoxic free radicals and slow electrons are formed in the vicinity of the metal center. The magnitude of the damage inflicted upon the environment and the energies of the emitted electrons depend on the initial Auger step, and can therefore be controlled by tuning the energy of the radiation. Consequently, the results of this work can have implications in understanding radiation chemistry and radiation damage in biologically relevant systems.

## Acknowledgement

We thank Prof. Nobuhiro Kosugi and Dr. Matjaž Žitnik for the fruitful discussions. Experiments were performed at the GALAXIES beamline, SOLEIL Synchrotron, France (Proposal No. 20140160). The authors are grateful to the SOLEIL staff for assistance during the beamtime. This project has received funding from the Research Executive Agency (REA) under the European Union’s Horizon 2020 research and innovation programme Grant agreement No. 705515. Campus France and the PHC SIAM exchange program are acknowledged for financial support (project No. 38282QB). L. S. Cederbaum and N. V. Kryzhevoi acknowledge the financial support of the European Research Council (ERC) (Advanced Investigator Grant No. 692657) and the Deutsche Forschungsgemeinschaft (DFG research unit 1789).

## Supporting Information Available

- suppinfo.pdf: contains 1) the radial density distributions of the core excited states of the bare ions; 2) partial cross sections and charge transfer time extracted from the experimental 2D map near the Cl 1s edge.

## References

- (1) Smith, J. W.; Saykally, R. J. Soft X-ray Absorption Spectroscopy of Liquids and Solutions. *Chem. Rev.* **2017**, *117*, 13909–13934, PMID: 29125751.
- (2) O'Neill, P.; Stevens, D. L.; Garman, E. F. Physical and chemical considerations of damage induced in protein crystals by synchrotron radiation: a radiation chemical perspective. *J. Synchrotron Radiat.* **2002**, *9*, 329–332.
- (3) Carugo, O.; Carugo, K. D. When X-rays modify the protein structure: radiation damage at work. *Trends Biochem. Sci.* **2005**, *30*, 213–219.
- (4) Stumpf, V.; Gokhberg, K.; Cederbaum, L. S. The role of metal ions in X-ray-induced photochemistry. *Nat. Chem.* **2016**, *8*, 237–241.
- (5) Pokapanich, W.; Bergersen, H.; Bradeanu, I. L.; Marinho, R. R. T.; Lindblad, A.; Legendre, S.; Rosso, A.; Svensson, S.; Björneholm, O.; Tchapyguine, M. et al. Auger Electron Spectroscopy as a Probe of the Solution of Aqueous Ions. *J. Am. Chem. Soc.* **2009**, *131*, 7264–7271.
- (6) Pokapanich, W.; Kryzhevoi, N. V.; Ottosson, N.; Svensson, S.; Cederbaum, L. S.; Öhrwall, G.; Björneholm, O. Ionic-charge dependence of the intermolecular Coulombic decay time-scale for aqueous ions probed by the core-hole clock. *J. Am. Chem. Soc.* **2011**, *133*, 13430.

- (7) Unger, I.; Seidel, R.; Thürmer, S.; Pohl, M. N.; Aziz, E. F.; Cederbaum, L. S.; Muchová, E.; Slavíček, P.; Winter, B.; Kryzhevoi, N. V. Observation of electron-transfer-mediated decay in aqueous solution. *Nat. Chem.* **2017**, *9*, 708.
- (8) Céolin, D.; Kryzhevoi, N. V.; Nicolas, C.; Pokapanich, W.; Choksakulporn, S.; Songsiriritthigul, P.; Saisopa, T.; Rattanachai, Y.; Utsumi, Y.; Palaudoux, J. et al. Ultrafast Charge Transfer Processes Accompanying *KLL* Auger Decay in Aqueous KCl Solution. *Phys. Rev. Lett.* **2017**, *119*, 263003.
- (9) Stoychev, S. D.; Kuleff, A. I.; Tarantelli, F.; Cederbaum, L. S. On the interatomic electronic processes following Auger decay in neon dimer. *J. Chem. Phys.* **2008**, *129*, 074307.
- (10) Demekhin, P. V.; Scheit, S.; Stoychev, S. D.; Cederbaum, L. S. Dynamics of interatomic Coulombic decay in a Ne dimer following the  $K-L_1L_{2,3}(^1P)$  Auger transition in the Ne atom. *Phys. Rev. A* **2008**, *78*, 043421.
- (11) Demekhin, P. V.; Chiang, Y.-C.; Stoychev, S. D.; Kolorenč, P.; Scheit, S.; Kuleff, A. I.; Tarantelli, F.; Cederbaum, L. S. Interatomic Coulombic decay and its dynamics in NeAr following K-LL Auger transition in the Ne atom. *J. Chem. Phys.* **2009**, *131*, 104303.
- (12) Ouchi, T.; Sakai, K.; Fukuzawa, H.; Higuchi, I.; Demekhin, P. V.; Chiang, Y.-C.; Stoychev, S. D.; Kuleff, A. I.; Mazza, T.; Schöffler, M. et al. Interatomic Coulombic decay following Ne 1s Auger decay in NeAr. *Phys. Rev. A* **2011**, *83*, 053415.
- (13) Miteva, T.; Chiang, Y.-C.; Kolorenč, P.; Kuleff, A. I.; Cederbaum, L. S.; Gokhberg, K. The effect of the partner atom on the spectra of interatomic Coulombic decay triggered by resonant Auger processes. *J. Chem. Phys.* **2014**, *141*, 164303.
- (14) Travnikova, O.; Marchenko, T.; Goldsztejn, G.; Jänkälä, K.; Sisourat, N.; Carniato, S.; Guillemin, R.; Journal, L.; Céolin, D.; Püttner, R. et al. Hard-X-Ray-Induced Multistep Ultrafast Dissociation. *Phys. Rev. Lett.* **2016**, *116*, 213001.

- (15) Gokhberg, K.; Kolorenč, P.; Kuleff, A. I.; Cederbaum, L. S. Site- and energy-selective slow-electron production through intermolecular Coulombic decay. *Nature* **2014**, *505*, 661–663.
- (16) Trinter, F.; Schöffler, M. S.; Kim, H.-K.; Sturm, F. P.; Cole, K.; Neumann, N.; Vredenburg, A.; Williams, J.; Bocharova, I.; Guillemin, R. et al. Resonant Auger decay driving intermolecular Coulombic decay in molecular dimers. *Nature* **2014**, *505*, 664–666.
- (17) Miteva, T.; Wenzel, J.; Klaiman, S.; Dreuw, A.; Gokhberg, K. X-Ray absorption spectra of microsolvated metal cations. *Phys. Chem. Chem. Phys.* **2016**, *18*, 16671–16681.
- (18) Nordlund, D.; Ogasawara, H.; Bluhm, H.; Takahashi, O.; Odelius, M.; Nagasono, M.; Pettersson, L. G. M.; Nilsson, A. Probing the Electron Delocalization in Liquid Water and Ice at Attosecond Time Scales. *Phys. Rev. Lett.* **2007**, *99*, 217406.
- (19) Ottosson, N.; Odelius, M.; Spångberg, D.; Pokapanich, W.; Svanqvist, M.; Öhrwall, G.; Winter, B.; Björneholm, O. Cations Strongly Reduce Electron-Hopping Rates in Aqueous Solutions. *J. Am. Chem. Soc.* **2011**, *133*, 13489–13495.
- (20) Wagner, C. D.; Zatko, D. A.; Raymond, R. H. Use of the oxygen KLL Auger lines in identification of surface chemical states by electron spectroscopy for chemical analysis. *Anal. Chem.* **1980**, *52*, 1445–1451.
- (21) Céolin, D.; Ablett, J.; Prieur, D.; Moreno, T.; Rueff, J.-P.; Marchenko, T.; Journal, L.; Guillemin, R.; Pilette, B.; Marin, T. et al. Hard X-ray photoelectron spectroscopy on the {GALAXIES} beamline at the {SOLEIL} synchrotron. *J. Electron Spectrosc. Relat. Phenom.* **2013**, *190, Part B*, 188 – 192.
- (22) Rueff, J.-P.; Ablett, J. M.; Céolin, D.; Prieur, D.; Moreno, T.; Balédent, V.; Lassalle-Kaiser, B.; Rault, J. E.; Simon, M.; Shukla, A. The GALAXIES beamline at the SOLEIL synchrotron: inelastic X-ray scattering and photoelectron spectroscopy in the hard X-ray range. *J. Synchrotron Rad.* **2015**, *22*, 175–179.

- (23) Faubel, M.; Schlemmer, S.; Toennies, J. P. A molecular beam study of the evaporation of water from a liquid jet. *Z. Phys. D* **1988**, *10*, 269–277.
- (24) Winter, B.; Faubel, M. Photoemission from Liquid Aqueous Solutions. *Chem. Rev.* **2006**, *106*, 1176–1211, PMID: 16608177.
- (25) Ohtaki, H.; Radnai, T. Structure and dynamics of hydrated ions. *Chem. Rev.* **1993**, *93*, 1157–1204.
- (26) Soper, A. K.; Weckström, K. Ion solvation and water structure in potassium halide aqueous solutions. *Biophys. Chem.* **2006**, *124*, 180 – 191.
- (27) Ma, H. Hydration structure of Na<sup>+</sup>, K<sup>+</sup>, F<sup>-</sup>, and Cl<sup>-</sup> in ambient and supercritical water: A quantum mechanics/molecular mechanics study. *Int. J. Quant. Chem.* **2014**, *114*, 1006–1011.
- (28) Krishnan, R.; Binkley, J. S.; Seeger, R.; Pople, J. A. Self-consistent molecular orbital methods. XX. A basis set for correlated wave functions. *J. Chem. Phys.* **1980**, *72*, 650–654.
- (29) Blaudeau, J.-P.; McGrath, M. P.; Curtiss, L. A.; Radom, L. Extension of Gaussian-2 (G2) theory to molecules containing third-row atoms K and Ca. *J. Chem. Phys.* **1997**, *107*, 5016–5021.
- (30) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A. et al. Gaussian 09 Revision D.01. Gaussian Inc. Wallingford CT 2009.
- (31) Lee, H. M.; Kim, J.; Lee, S.; Mhin, B. J.; Kim, K. S. Aqua-potassium(I) complexes: Ab initio study. *J. Chem. Phys.* **1999**, *111*, 3995–4004.
- (32) Lee, H. M.; Kim, D.; Kim, K. S. Structures, spectra, and electronic properties of

- halide-water pentamers and hexamers,  $\text{X}\ddot{\text{A}}\ddot{\text{S}}(\text{H}_2\text{O})_{5,6}$  ( $\text{X}=\text{F}, \text{Cl}, \text{Br}, \text{I}$ ): Ab initio study. *J. Chem. Phys.* **2002**, *116*, 5509–5520.
- (33) Schirmer, J. Beyond the random-phase approximation: A new approximation scheme for the polarization propagator. *Phys. Rev. A* **1982**, *26*, 2395–2416.
- (34) Barth, A.; Schirmer, J. Theoretical core-level excitation spectra of  $\text{N}_2$  and  $\text{CO}$  by a new polarisation propagator method. *J. Phys. B At. Mol. Opt. Phys.* **1985**, *18*, 867.
- (35) Cederbaum, L. S.; Domcke, W.; Schirmer, J. Many-body theory of core holes. *Phys. Rev. A* **1980**, *22*, 206–222.
- (36) Barth, A.; Cederbaum, L. S. Many-body theory of core-valence excitations. *Phys. Rev. A* **1981**, *23*, 1038–1061.
- (37) Wenzel, J.; Wormit, M.; Dreuw, A. Calculating core-level excitations and x-ray absorption spectra of medium-sized closed-shell molecules with the algebraic-diagrammatic construction scheme for the polarization propagator. *J. Comp. Chem.* **2014**, *35*, 1900–1915.
- (38) Wenzel, J.; Wormit, M.; Dreuw, A. Calculating X-ray Absorption Spectra of Open-Shell Molecules with the Unrestricted Algebraic-Diagrammatic Construction Scheme for the Polarization Propagator. *J. Chem. Theory Comput.* **2014**, *10*, 4583–4598.
- (39) Wormit, M.; Rehn, D. R.; Harbach, P. H.; Wenzel, J.; Krauter, C. M.; Epifanovsky, E.; Dreuw, A. Investigating excited electronic states using the algebraic diagrammatic construction (ADC) approach of the polarisation propagator. *Mol. Phys.* **2014**, *112*, 774–784.
- (40) Shao, Y.; Gan, Z.; Epifanovsky, E.; Gilbert, A. T.; Wormit, M.; Kussmann, J.; Lange, A. W.; Behn, A.; Deng, J.; Feng, X. et al. Advances in molecular quantum

- chemistry contained in the Q-Chem 4 program package. *Mol. Phys.* **2015**, *113*, 184–215.
- (41) McLean, A. D.; Chandler, G. S. Contracted Gaussian basis sets for molecular calculations. I. Second row atoms,  $Z=11\text{--}18$ . *J. Chem. Phys.* **1980**, *72*, 5639–5648.
- (42) Krause, M. O.; Oliver, J. H. Natural widths of atomic K and L levels,  $K\alpha$  X-ray lines and several KLL Auger lines. *J. Phys. Chem. Ref. Data* **1979**, *8*, 329–338.
- (43) Brooks, B. R.; Laidig, W. D.; Saxe, P.; Handy, N. C.; Schaefer III, H. F. The Loop-Driven Graphical Unitary Group Approach: A Powerful Method for the Variational Description of Electron Correlation. *Phys. Scr.* **1980**, *21*, 312.
- (44) Brooks, B. R.; Schaefer, H. F. The graphical unitary group approach to the electron correlation problem. Methods and preliminary applications. *J. Chem. Phys.* **1979**, *70*, 5092–5106.
- (45) Schmidt, M. W.; Baldridge, K. K.; Boatz, J. A.; Elbert, S. T.; Gordon, M. S.; Jensen, J. H.; Koseki, S.; Matsunaga, N.; Nguyen, K. A.; Su, S. et al. General atomic and molecular electronic structure system. *J. Comp. Chem.* **1993**, *14*, 1347–1363.
- (46) Mosnier, J.-P.; Kennedy, E. T.; van Kampen, P.; Cubaynes, D.; Guilbaud, S.; Sisourat, N.; Puglisi, A.; Carniato, S.; Bizau, J.-M. Inner-shell photoexcitations as probes of the molecular ions  $\text{CH}^+$ ,  $\text{OH}^+$ , and  $\text{SiH}^+$ : Measurements and theory. *Phys. Rev. A* **2016**, *93*, 061401.
- (47) Russek, A.; Mehlhorn, W. Post-collision interaction and the Auger lineshape. *J. Phys. B At. Mol. Opt. Phys.* **1986**, *19*, 911.
- (48) Guillemin, R.; Sheinerman, S.; Püttner, R.; Marchenko, T.; Goldsztejn, G.; Journal, L.; Kushawaha, R. K.; Céolin, D.; Piancastelli, M. N.; Simon, M. Postcollision interaction



- effects in *KLL* Auger spectra following argon 1s photoionization. *Phys. Rev. A* **2015**, *92*, 012503.
- (49) Tchapyguine, M.; Kivimäki, A.; Peredkov, S.; Sorensen, S. L.; Öhrwall, G.; Schulz, J.; Lundwall, M.; Rander, T.; Lindblad, A.; Rosso, A. et al. Localized versus delocalized excitations just above the 3d threshold in krypton clusters studied by Auger electron spectroscopy. *J. Chem. Phys.* **2007**, *127*, 124314.
- (50) Sugiura, C. Influence of coordinating water on the chlorine K absorption spectra of hydrated metal dichlorides:  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$  and  $\text{SrCl}_2 \cdot 6\text{H}_2\text{O}$ . *J. Chem. Phys.* **1982**, *77*, 681–682.
- (51) Shadle, S. E.; Hedman, B.; Hodgson, K. O.; Solomon, E. I. Ligand K-edge x-ray absorption spectroscopic studies: metal-ligand covalency in a series of transition metal tetrachlorides. *J. Am. Chem. Soc.* **1995**, *117*, 2259–2272.
- (52) Hertlein, M. P.; Adaniya, H.; Amini, J.; Bressler, C.; Feinberg, B.; Kaiser, M.; Neumann, N.; Prior, M. H.; Belkacem, A. Inner-shell ionization of potassium atoms ionized by a femtosecond laser. *Phys. Rev. A* **2006**, *73*, 062715.
- (53) Céolin, D.; Marchenko, T.; Guillemin, R.; Journal, L.; Kushawaha, R. K.; Carniato, S.; Huttula, S.-M.; Rueff, J. P.; Armen, G. B.; Piancastelli, M. N. et al. Auger resonant-Raman study at the Ar K edge as probe of electronic-state-lifetime interferences. *Phys. Rev. A* **2015**, *91*, 022502.
- (54) Winter, B.; Weber, R.; Widdra, W.; Dittmar, M.; Faubel, M.; Hertel, I. V. Full Valence Band Photoemission from Liquid Water Using EUV Synchrotron Radiation. *J. Phys. Chem. A* **2004**, *108*, 2625–2632.
- (55) Goldsztejn, G.; Marchenko, T.; Püttner, R.; Journal, L.; Guillemin, R.; Carniato, S.; Selles, P.; Travnikova, O.; Céolin, D.; Lago, A. F. et al. Double-Core-Hole States in

Neon: Lifetime, Post-Collision Interaction, and Spectral Assignment. *Phys. Rev. Lett.* **2016**, *117*, 133001.