


练习 13.6

(a) (2)

(b) (4)

(c) (1) (d) (5)

练习 13.8



$$R = 10^3 \Omega$$

$$\omega = 10^3 \text{ rad/s} \quad R^2 - \omega^2 L^2 = 10^4$$

$$L = 10^{-3} \text{ H}$$

练习 13.11

$$(a) |H(j\omega)| = \frac{\frac{1}{j\omega C}}{R + \frac{1}{j\omega C}} = \frac{\omega C}{\sqrt{R^2 + \omega^2 C^2}} e^{-j\varphi}$$

$$\tan \varphi = \frac{R}{\omega C}$$

$$\angle H(j\omega) = \tan^{-1}\left(\frac{R}{\omega C}\right)$$

$$(b) |H(j\omega)| = \frac{j\omega L}{R + j\omega L} = \frac{\omega L}{\sqrt{R^2 + \omega^2 L^2}} e^{-j\varphi}$$

$$\angle H(j\omega) = \varphi = \tan^{-1}\left(-\frac{R}{\omega L}\right)$$

$$(c) |H(j\omega)| = \frac{R}{R + \frac{1}{j\omega C}} = \frac{R^2}{\sqrt{R^2 + \omega^2 C^2}}$$

$$\angle H(j\omega) = \tan^{-1}\left(-\frac{R}{\omega C}\right)$$

$$(d) |H(j\omega)| = \frac{R}{j\omega L + R} = \frac{R}{\sqrt{R^2 + \omega^2 L^2}} e^{-j\varphi}$$

$$\angle H(j\omega) = \tan^{-1}\left(\frac{\omega L}{R}\right)$$

练习 13.16

$$\frac{V_o(s)}{V_i(s)} = \frac{\frac{1}{j\omega C} \parallel (j\omega L + R_2)}{R_1 + [\frac{1}{j\omega C} \parallel (j\omega L + R_2)]} \cdot \frac{R_2}{j\omega L + R_2}$$

$$= \frac{R_2}{R_1 + R_2 + j\omega L + R_1 R_2 j\omega C - \omega^2 L C}$$

$$\frac{I_A(s)}{V_i(s)} = \frac{j\omega C \parallel (j\omega L + R_2)}{R_1 + [\frac{1}{j\omega C} \parallel (j\omega L + R_2)]}$$

$$= \frac{jR_2 \omega C - \omega^2 L}{R_1(1 - \omega^2 L C + R_2 j\omega C) + j\omega L + R_2}$$

问题 13.1

$$(1) Z_1 = \frac{R(j\omega C)}{R + j\omega C} = \frac{R - jR\omega C}{1 + \omega^2 C^2 R^2}$$

$$(2) Z_2 = \frac{1}{\frac{1}{R} + \frac{1}{j\omega L}} = \frac{\omega L^2 R + j\omega L R^2}{\omega^2 L^2 + R^2}$$

$$(3) Z_3 = \frac{1}{\frac{1}{j\omega C_1} + \frac{1}{\frac{1}{j\omega C_2} + R}} = \frac{1 + jR\omega C_2}{j(\omega C_1 + \omega C_2) - \omega^2 R C_1 C_2}$$

