

Homework (10)

1.

a) Solve the optimization problem

$$\min_{\mathbf{x}} f(x_1, x_2) := 2x_1 + 3x_2, \quad s.t. \quad \sqrt{x_1} + \sqrt{x_2} = 5,$$

using Lagrange multipliers.

- b) Visualize the contour lines of f as well as the set of feasible points, and mark the optimal solution \mathbf{x}^* .
- c) Find all its KKT points. Do they all correspond to local minima?
- d) Find all the saddle points of its Lagrangian function. Do they all correspond to local minima?

Homework (10)

2. With $f(\mathbf{x}) := x_1^2 + x_2^2$ for $\mathbf{x} \in \mathbb{R}^2$ consider


$$(P) \begin{cases} \min_{\mathbf{x}} f(\mathbf{x}) \\ -x_2 \leq 0 \\ x_1^3 - x_2 \leq 0 \\ x_1^3(x_2 - x_1^3) \leq 0 \end{cases}.$$

- a) Determine the linearizing cone, the tangent cone and the feasible direction cones at the (strict global) minimal point $\mathbf{x}_0 := (0, 0)^T$.
- b) Find all its KKT points. Do they all correspond to local minima?
- c) Find all the saddle points of its Lagrangian function. Do they all correspond to local minima?

Homework (10)

3. Determine a triangle with minimal area containing two disjoint disks with radius 1. Without loss of generalization, let $(0,0)$, $(x_1,0)$ and (x_2,x_3) with $x_1, x_3 \geq 0$ be the vertices of the triangle; (x_4, x_5) and (x_6, x_7) denote the centers of the disks.

- a) Formulate this problem as a minimization problem in terms of seven variables and nine constraints.
- b) $\mathbf{x}^* = (4 + 2\sqrt{2}, 2 + \sqrt{2}, 2 + \sqrt{2}, 1 + \sqrt{2}, 1, 3 + \sqrt{2}, 1)^T$ is a solution of this problem; calculate the corresponding Lagrange multipliers $\boldsymbol{\lambda}^*$, such that the KKT conditions are fulfilled.



Homework (10)

4. Find local extremizers (either minimizer or maximizer) for the following optimization problems:

$$\begin{aligned} 1) \quad & \min_{\mathbf{x}} \quad x_1^2 + 2x_1x_2 + 3x_2^2 + 4x_1 + 5x_2 + 6x_3 \\ & \text{s.t.} \quad x_1 + 2x_2 = 3 \\ & \quad \quad 4x_1 + 5x_3 = 6. \end{aligned}$$

$$\begin{aligned} 2) \quad & \max_{\mathbf{x}} \quad 4x_1 + x_2^2 \\ & \text{s.t.} \quad x_1^2 + x_2^2 = 9. \end{aligned}$$

$$\begin{aligned} 3) \quad & \min_{\mathbf{x}} \quad x_1x_2 \\ & \text{s.t.} \quad x_1^2 + 4x_2^2 = 1. \end{aligned}$$

Homework (10)

5. Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\mathbf{x}_0 \in \mathbb{R}^n$ be given, where $g(\mathbf{x}_0) \geq 0$. Consider the problem

$$\begin{array}{ll} \text{minimize} & \frac{1}{2} \|\mathbf{x} - \mathbf{x}_0\|^2 \\ \text{subject to} & g(\mathbf{x}) \leq 0. \end{array}$$

Suppose that \mathbf{x}^* is a solution to the problem and $g \in \mathcal{C}^1$. Use the KKT theorem to decide which of the following equations/inequalities hold:

1. $g(\mathbf{x}^*) \leq 0$.
2. $g(\mathbf{x}^*) = 0$.
3. $(\mathbf{x}^* - \mathbf{x}_0)^T \nabla g(\mathbf{x}^*) \leq 0$.
4. $(\mathbf{x}^* - \mathbf{x}_0)^T \nabla g(\mathbf{x}^*) = 0$.
5. $(\mathbf{x}^* - \mathbf{x}_0)^T \nabla g(\mathbf{x}^*) \geq 0$.

Homework (10)

6. Consider the problem with equality constraint

$$\begin{array}{ll}\text{minimize} & f(\mathbf{x}) \\ \text{subject to} & \mathbf{h}(\mathbf{x}) = \mathbf{0}\end{array}$$

We can convert the above into the equivalent optimization problem

$$\begin{array}{ll}\text{minimize} & f(\mathbf{x}) \\ \text{subject to} & \frac{1}{2} \|\mathbf{h}(\mathbf{x})\|^2 \leq 0.\end{array}$$

Write down the KKT condition for the equivalent problem (with inequality constraint) and explain why the KKT theorem cannot be applied in this case.