



# **Associations between bovine respiratory disease complex and the probability and latency of group-reared neonatal dairy calves to approach a novel object or stationary person**

Catie Cramer\* and Amy Stanton


University of Wisconsin- Madison  
Department of Dairy Science




Macrophages




Inflammatory  
Cytokines



(+)



Infectious Pathogens (-)



Anorexia, lethargy,  
decreased social and  
exploratory behavior



# Sickness Behavior

- Highly coordinated and evolved response
- Increase probability of survival
- Conserve energy for febrile response
- Nonspecific for illness
- Common across different species
- Interest in using this to detect disease



# Bovine Respiratory Disease (BRD)

- 12% calves affected on US dairies, 22.5% of deaths (NAHMS, USDA, 2010)
- Reduced growth (Virtala et al., 1996; Stanton et al., 2012)
- Increased likelihood of morbidity and age at first calving (Waltner-Toews, 1989; Stanton et al, 2012)
- Increased risk of not completing the first lactation (Bach, 2011)



# BRD Impacts Animal Welfare

- Evidence of pain and labored breathing (Theurer et al., 2013, White et al. 2013)
- Decrease welfare (Mellor and Stafford, 2004)





# Disease Detection in Group Housing

- Group housing increasing in popularity
  - Challenge for illness detection
  - Socially competitive environment
- Calf Health Scoring Chart (McGuirk, 2008)
  - Individual animal evaluation time consuming
- Automated feeder data (e.g. Borderas et al., 2009)
  - Expensive
- Screening tools needed
  - Low cost, valid, quick






# Approach Tests

- Test fearfulness and exploratory behavior in cattle (Arave et al., 1997; Lauber et al., 2005)
- Illness in mice decreased exploration of novel object (Haba et al., 2012)
- New opportunity for disease detection in calves?





# Objectives

- To determine **associations between BRD status** of group-housed calves and the **probability of approaching** a novel object (**OBJ**) and stationary person (**SP**)
- To determine **associations between BRD status** of group-housed calves and the **latency of approaching** a novel object (**OBJ**) and stationary person (**SP**)







# Hypotheses

- Calves with BRD will be less likely to approach compared to healthy calves
- Calves with BRD will have a longer latency to approach compared to healthy calves





# Materials and Methods

- Commercial dairy in Wisconsin, USA
- 79 calves in group housing
  - 8-10 calves per pen, 8 pens
- Research staff visited once per week
- Enrolled over 2 weeks
- Birth to 6 weeks of age
- *Ad lib* acidified milk
- Weaning began at 6 weeks





# Approach Tests

- Calves tested at beginning of farm visit
- 60 seconds to approach OBJ **AND** SP
  - All calves tested with both every week
- Tests recorded on video
  - Time of approach for each calf for 6 weeks









# Health Scores

- Performed after approach tests
- Umbilical palpation
- Diarrhea
- Respiratory disease (McGuirk, 2008)











### Calf Health Scoring Criteria





0	1	2	3
<b>Rectal temperature</b>			
100-100.9	101-101.9	102-102.9	≥103
<b>Cough</b>			
None	Induce single cough	Induced repeated coughs or occasional spontaneous cough	Repeated spontaneous coughs
<b>Nasal discharge</b>			
Normal serous discharge	Small amount of unilateral cloudy discharge	Bilateral, cloudy or excessive mucus discharge	Copious bilateral mucopurulent discharge



### Eye scores

Normal	Small amount of ocular discharge	Moderate amount of bilateral discharge	Heavy ocular discharge
			

### Ear scores

Normal	Ear flick or head shake	Slight unilateral droop	Head tilt or bilateral droop
			

# BRD Scoring Chart

0= Normal

3= Severely  
Abnormal


**Temp + Cough +  
Nasal + Eye or Ear  
= BRD Score**

**\*2 scores of 2+  
= BRD**

McGuirk, 2008




## Proportion of All Calves that Approached OBJ and SP and Prevalence of Diarrhea and BRD





## Proportion of Calves With and Without BRD that Approached





# BRD Status and Probability of Approach

- OBJ



- Calves **with BRD** were **0.5** (95% CL: 0.3 to 0.9) **times as likely** to approach OBJ compared to calves without BRD ( $P < 0.05$ )

- SP



- Calves **with BRD** were **0.5** (95% CL: 0.3 to 0.9) **times as likely** to approach SP compared to calves without BRD ( $P < 0.05$ )



# BRD Status and Latency to Approach

## •OBJ



- Calves **without BRD** were **0.6** (95% CL: 0.4 to 0.8) times **as likely** to approach OBJ **at any given time point** during the test, compared to calves with BRD ( $P < 0.01$ )

## •SP



- Calves **without BRD** were **0.6** (95% CL: 0.4 to 0.9) times **as likely** to approach SP **at any given time point** during the test compared to calves with BRD ( $P < 0.05$ )





# Summary

- Calves with BRD exhibit decreased exploratory behavior
- When calves with BRD approached, they did so more slowly compared to calves without BRD





# Screening Tool

- Sensitivity: 64% (OBJ) and 68% (SP)
  - Correctly identify 64 or 68% of calves with BRD
  - Miss about 30% of calves
- Specificity: 43% (OBJ) and 42% (SP)
  - Identify 42 or 43% of healthy calves
  - Rule out animals that are not sick



# Conclusion

- Approach tests may be useful in identifying calves with BRD in group housing
  - Low cost and quick
  - Should not be sole method
  - Require observation of calves
- Combine with other behavioral measures
- Screening tools are needed to detect disease early in group- housed calves to improve welfare



# Acknowledgements

- Farm Owners and Staff
- Dr. Amy Stanton
- Undergraduates: Liz Binversie, Alyssa Daniels, Jessi Dercks, Kaitlin Gelsinger, Lynzie Miller, Amy Moy, Sophia Wood, Aisha Hill
- Melissa Cornett and Tonia Peters







Questions?





# References

Arave, C. W., J. L. Albright, D. V. Armstrong, W. W. Foster, and L. L. Larson. 1992. Effects of isolation of calves on growth, behavior, and 1st lactation milk-yield of holstein cows. *J. Dairy Sci.* 75(12):3408-3415.

Borderas, T. F., A. M. B. d. Passille, and J. Rushen. 2009. Feeding behavior of calves fed small or large amounts of milk. *J. Dairy Sci.* 92(6).

Haba, R., N. Shintani, Y. Onaka, H. Wang, R. Takenaga, A. Hayata, A. Baba, and H. Hashimoto. 2012. Lipopolysaccharide affects exploratory behaviors toward novel objects by impairing cognition and/or motivation in mice: Possible role of activation of the central amygdala. *Behav. Brain Res.* 228(2):423-431.

Hart, B. L. 1988. Biological Basis of the Behavior of Sick Animals. *Neurosci. Biobehav. Rev.* 12(2):123-137.



## References cont.

Lauber, M. C. Y., P. H. Hemsworth, and J. L. Barnett. 2006. The effects of age and experience on behavioural development in dairy calves. *Appl. Anim. Behav. Sci.* 99(1-2):41-52.

McGuirk, S. M. 2008. Disease management of dairy calves and heifers. *Veterinary Clinics of North America, Food Animal Practice* 24(1).

Mellor, D. J. and K. J. Stafford. 2004. Animal welfare implications of neonatal mortality and morbidity in farm animals. *Vet. J.* 168(2):118-133.

Stanton, A. L., D. F. Kelton, S. J. LeBlanc, J. Wormuth, and K. E. Leslie. 2012. The effect of respiratory disease and a preventative antibiotic treatment on growth, survival, age at first calving, and milk production of dairy heifers. *J. Dairy Sci.* 95(9).

USDA. 2010. Heifer Calf Health and Management Practices on U.S. Dairy Operations, 2007. C. USDA:APHIS:VS, ed, Fort Collins, CO.


Virtala, A. M. K., G. D. Mechor, Y. T. Grohn, and H. N. Erb. 1996. The effect of calfhood diseases on growth of female dairy calves during the first 3 months of life in New York state. *J. Dairy Sci.* 79(6): 1040-1049.

Waltner-Toews, D., S. W. Martin, A. H. Meek. 1989. The effect of early calfhood health status on survivorship and age at first calving. *Canadian veterinary Reserves Journal* 50:314-417.



# Proportion of All Calves that Approached OBJ and SP During Four Different Time Points for All Weeks

Proportion of Calves that Approached





# Statistical Analysis

- Separate models for OBJ and SP
- Probability of Approach
  - Linear mixed model (PROC GLIMMIX in SAS)
- Latency of Approach
  - Cox proportional hazards regression (PROC PHREG in SAS)
- Models controlled for
  - Week (age of calf in weeks), pen, pen order, enrollment group, test order, and diarrhea status