Paper 2: Dual Impedance Networks and Metabolic Scaling
Vascular Architecture, Organ Health, Fractal Cost, and Kleiber's Law under the Minimum Reflection Principle

Hsi-Yu Huang[1, *]

[1] *Independent Researcher, Taipei, Taiwan*

(Dated: March 20, 2026)

We ask: given a blood vessel with impedance $Z_\mathrm{v} = 8\mu L/\pi r^4$ governed by vascular C2 remodeling, what architecture—and what metabolic scaling laws—must emerge?

Part A derives vascular architecture from the Minimum Reflection Principle (MRP). A blood vessel is a hydraulic transmission line. At every bifurcation, impedance mismatch produces a reflection Γ_v; C1 gives $\Gamma_\mathrm{v}^2 + T_\mathrm{v} = 1$. We derive Murray's Law ($r_p^3 = \sum r_i^3$) as the exact minimum of the vascular action $A_\mathrm{v} = \sum_j [Q_j^2 Z_j + \lambda V_j]$, with 100% numerical agreement for $n = 2, 3, 4$. Vascular C2 remodeling ($\Delta Z_\mathrm{v} = -\eta_v \Gamma_\mathrm{v} \tau_\mathrm{wall} \rho$) operates on the same principle as neural C2 but with $\eta_v \ll \eta_n$.

Part B derives the dual-network organ model. Every organ is served by two coincident networks: vascular (Γ_v) and neural (Γ_n). Organ health $H = T_\mathrm{n} \cdot T_\mathrm{v} = (1-\Gamma_\mathrm{n}^2)(1-\Gamma_\mathrm{v}^2)$ follows from multiplicative transmission. Four failure modes are classified by which network fails. The organ impedance hierarchy ranks ten organs by their resting Γ_v^2.

Part C derives metabolic scaling. A self-similar relay hierarchy with branching number n in d dimensions produces an irreducible action $A_\mathrm{cut} \propto M^{(d-1)/d}$ (for $d = 3$: $M^{2/3}$). The Murray bifurcation ratio $\beta = n^{1/d} = 2^{1/3}$ sets length and radius contraction across levels. Kleiber's Law ($B \propto M^{3/4}$) follows from capillary invariance. Dual-network packing yields a metabolic overhead $\Omega \approx 1.33$, with $\Omega_n \approx \beta$. Multi-scale verification confirms the irreducibility theorem across cardiovascular, musculoskeletal, and 29-tissue-atlas scales.

Part D demonstrates interface universality. The same $\Gamma = (Z_2 - Z_1)/(Z_2 + Z_1)$ governs seven physical domains across 14 orders of magnitude—from synapses to power grids. The Interface Isomorphism Proposition proves that Γ depends only on boundary impedance, not bulk properties.

Circularity disclosure. Internal experiments use simulation modules built from C1–C3; they demonstrate self-consistency, not empirical validity. External tests are reported in Papers 3–5 [3–5].

Part A: From Blood Vessel to Vascular Network

I. INTRODUCTION

A blood vessel is a tube carrying fluid under pressure. Its impedance—the ratio of pressure drop to volume flow—is a physical fact set by geometry and fluid properties.

Paper 1 [2] derives neural architecture and the mind from the Minimum Reflection Principle (MRP). Yet every organ is governed by a *second* impedance network: the vasculature. The pressure wave in an artery reflects at every bifurcation, stricture, or compliance mismatch, precisely as an electromagnetic wave reflects at a transmission-line discontinuity [7, 8].

This paper derives the vascular architecture from the same MRP, then combines both networks into the dual-network organ model, and finally shows that metabolic scaling laws (Murray, Kleiber) are necessary consequences of the same irreducible topology.

The derivation follows a strictly outward chain: single vessel (Sec. II) → Murray's Law (Sec. III) → vascular C2 (Sec. V) → dual-network organ model (Sec. VI) → fractal cost (Sec. IX) → Kleiber's Law (Sec. X) → interface universality (Sec. XIII).

II. VASCULAR IMPEDANCE PHYSICS

A. The blood vessel as transmission line

A cylindrical vessel of length L and radius r carrying a Newtonian fluid of viscosity μ has steady-state impedance given by the Poiseuille law:

$$Z_\mathrm{v} = \frac{\Delta P}{Q} = \frac{8\mu L}{\pi r^4}. \tag{1}$$

In the pulsatile regime, Womersley theory adds a frequency-dependent component:

$$Z_\mathrm{v}(\omega) = \frac{8\mu L}{\pi r^4} \cdot F(\alpha), \tag{2}$$

where $\alpha = r\sqrt{\omega \rho_f/\mu}$ is the Womersley number, $F(\alpha) \to 1$ for $\alpha \ll 1$ (Poiseuille limit) and $F(\alpha) \to j\alpha^2/8$ for $\alpha \gg 1$ (inertial limit) [7].

* llc.y.huangll@gmail.com

TABLE I. Electromagnetic–vascular transmission-line analogy.

EM quantity	Vascular	Unit
Voltage V	Pressure P	Pa
Current I	Flow Q	m^3/s
Impedance Z	$\Delta P/Q$	$\mathrm{Pa\,s/m}^3$
Γ	Γ_v	dimensionless
$T = 1 - \Gamma^2$	T_v	dimensionless

B. Three physical laws in the vascular domain

C1—Energy Conservation. $\Gamma_\mathrm{v}^2 + T_\mathrm{v} = 1$ at every vascular junction.

C2—Vascular Remodeling.

$$\Delta Z_\mathrm{v} = -\eta_v\, \Gamma_\mathrm{v}\, \tau_\mathrm{wall}\, \rho, \tag{3}$$

where τ_wall is wall shear stress (x_in), ρ is material supply (x_out), and η_v is the vascular remodeling rate.

C3—Impedance-Tagged Transport. Blood pressure waves carry source impedance as metadata.

C. Bifurcation reflection

At a symmetric bifurcation (n identical daughters), $Z_\mathrm{v}^\mathrm{down} = Z_\mathrm{v}^d/n$:

$$\Gamma_\mathrm{v} = \frac{Z_\mathrm{v}^d/n - Z_\mathrm{v}^p}{Z_\mathrm{v}^d/n + Z_\mathrm{v}^p}. \tag{4}$$

Pure impedance matching ($\Gamma_\mathrm{v} = 0$) gives $r_d = r_p/n^{1/4}$, which *disagrees* with Murray's Law. The resolution: MRP minimises the *total vascular action* including metabolic maintenance cost.

Remark 1 (Why impedance disparity is extreme (E1))**.** *Characteristic impedance Z depends on the physical properties of the medium—density, elastic modulus, membrane resistance—not on any single variable. In the human body, these properties span orders of magnitude:*

- Cortical neurons*: high membrane resistance ($R_m \sim 10^4\ \Omega\,\mathrm{cm}^2$), giving high characteristic impedance Z_n;*
- Skeletal muscle*: low membrane resistance, high density ($\rho \sim 1.06\ \mathrm{g/cm}^3$), giving low characteristic impedance $Z_m \ll Z_n$;*
- Cortical bone*: elastic modulus $E \sim 18$ GPa, two orders above soft tissue (~ 10 kPa).*

Direct connection between any two such media gives $\Gamma \to 1$ (total reflection). Multi-stage fractal matching—the vascular tree for hydraulic impedance, the neural relay hierarchy for electromagnetic impedance—is therefore not merely optimal but physically necessary*. Murray's Law (below) quantifies the exact geometry of the vascular solution.*

III. MURRAY'S LAW FROM THE VASCULAR ACTION PRINCIPLE

A. Vascular action functional

$$A_\mathrm{v} = \sum_j \left[Q_j^2\, Z_j + \lambda\, V_j\right], \tag{5}$$

where Q_j is flow, $Z_j = 8\mu L_j/\pi r_j^4$ is impedance, $V_j = \pi r_j^2 L_j$ is volume, and λ is the metabolic cost per unit volume.

B. Derivation for symmetric n-branching

Daughter contribution:

$$A_\mathrm{v}^d = \frac{Q^2\, 8\mu L_d}{n\,\pi\, r_d^4} + n\lambda\pi r_d^2 L_d. \tag{6}$$

Minimising $\partial A_\mathrm{v}^d/\partial r_d = 0$:

$$r_d^6 = \frac{16\, Q^2\, \mu}{n^2\, \lambda\, \pi^2}, \qquad r_p^6 = \frac{16\, Q^2\, \mu}{\lambda\, \pi^2}. \tag{7}$$

Taking the ratio:

Theorem 1 (Murray's Law from MRP (E0))**.** *For symmetric n-branching of a vessel with parent radius r_p, minimisation of the vascular action $A_v = \sum_j [Q_j^2 Z_j + \lambda V_j]$ yields*

$$\boxed{r_d = \frac{r_p}{n^{1/3}},} \tag{8}$$

equivalently $r_p^3 = \sum_{i=1}^n r_{d,i}^3$ (the cube law).

Proof. Setting $\partial A_\mathrm{v}^d/\partial r_d = 0$ in Eq. (6) gives $r_d^6 = 16Q^2\mu/(n^2\lambda\pi^2)$ and $r_p^6 = 16Q^2\mu/(\lambda\pi^2)$. The ratio $r_d^6/r_p^6 = 1/n^2$ gives $r_d/r_p = n^{-1/3}$. The 1/3 exponent—rather than 1/4 from pure impedance matching—arises from the balance of dissipation ($\propto r^{-4}$) and maintenance ($\propto r^{+2}$). □

Remark 2 (Physics, not optimisation (E1))**.** *The 1/3 exponent arises from the balance of dissipation ($\propto r^{-4}$) and maintenance ($\propto r^{+2}$). Pure impedance matching gives 1/4; MRP gives 1/3. Murray's Law is a physical law of impedance networks.*



FIG. 1. Murray's Law from the Vascular Action Principle. **Left**: Action $A_{\rm v}(r_d)$ vs. daughter radius for $n = 2$; minimum (star) at Murray prediction (dashed); pure impedance matching (dotted) overestimates. **Right**: Agreement for $n = 2, 3, 4$.



FIG. 2. Blood as crosstalk isolator. (a) In microwave circuits, ferrite absorber prevents reflection pulses from contaminating adjacent channels. (b) In biological tissue, blood with impedance $Z_{\rm blood}$ absorbs stray $\Gamma^2 P_{\rm in}$ pulses between neural interfaces, converting them to heat carried away by convective flow.

IV. MURRAY'S LAW: NUMERICAL VERIFICATION

For each branching number $n \in \{2, 3, 4\}$, the minimum of $A_{\rm v}^d(r_d)$ is located numerically and compared with Murray's prediction:

TABLE II. Murray's Law verification.

n	$r_d^{\rm MRP}$	$r_d^{\rm Murray}$	Agreement
2	$r_p/2^{1/3}$	$r_p/2^{1/3}$	exact
3	$r_p/3^{1/3}$	$r_p/3^{1/3}$	exact
4	$r_p/4^{1/3}$	$r_p/4^{1/3}$	exact

Agreement is exact (algebraic identity). The $\Gamma = 0$ solution gives $r_d = r_p/n^{1/4}$, overestimating by $n^{1/12}$. Full test: `tests/test_paper2_murray.py` (15 tests, 100% pass).

V. VASCULAR C2 REMODELING

Neural C2 operates with rate η_n (ms–hours); vascular C2 with η_v (days–weeks):

$$\Delta Z_{\rm v} = -\eta_v\, \Gamma_{\rm v}\, \tau_{\rm wall}\, \rho, \tag{9}$$

where $\tau_{\rm wall} = 4\mu Q/\pi r^3$.

C2 term	Neural	Vascular
Γ	$\Gamma_{\rm n}$	$\Gamma_{\rm v}$
$x_{\rm in}$	presynaptic activity	wall shear stress τ
$x_{\rm out}$	postsynaptic activity	material supply ρ
η	η_n (fast)	η_v (slow)

Key consequence: $\eta_v \ll \eta_n$ means vascular damage accumulates slowly (atherosclerosis over decades) and recovery is slow even with optimal conditions.

Part B: The Dual-Network Organ Model

VI. ORGAN HEALTH FROM MULTIPLICATIVE TRANSMISSION

A. Why two networks

Neural C2 requires material substrate (ρ) from vascular delivery. Vascular C2 requires neural control (autonomic signals regulate vascular tone). The two networks are *mutually prerequisite*.

Remark 3 (Blood as crosstalk isolator (E1))**.** *In microwave circuits, absorbing material is placed between adjacent channels to prevent reflection pulses in one channel from contaminating another (crosstalk isolation). Blood serves the same function: when a neural interface produces a reflection pulse $\Gamma_i^2 P_{in,i}$, part of the scattered energy propagates laterally. Blood, with its own characteristic impedance Z_{blood}, absorbs this stray energy and converts it to heat, which is carried away by convective flow (Eq. 11). This yields a third reason for the dual-network architecture beyond mutual prerequisiteness and thermal regulation:* impedance isolation between densely packed neural interfaces. *The blood–brain barrier is, in this view, an impedance wall that prevents vascular Z fluctuations from polluting neural Z; cerebrospinal fluid is an impedance-matched filler between brain and cranium. The brain's disproportionate share of cardiac output (∼15%) is driven not solely by oxygen demand but by the crosstalk-isolation requirements of its extremely dense relay architecture.*

B. Organ health

At each organ, the neural network transmits $T_{\rm n} = 1 - \Gamma_{\rm n}^2$ and the vascular network transmits $T_{\rm v} = 1 - \Gamma_{\rm v}^2$. Series connection:

$$\boxed{H = T_{\rm n} \cdot T_{\rm v} = (1 - \Gamma_{\rm n}^2)(1 - \Gamma_{\rm v}^2).} \tag{10}$$

$H = 1$: perfect transmission; $H = 0$: at least one network totally blocked.

C. Thermal bus

The vascular network carries heat:

$$\dot{Q}_{\text{organ}} = c_p\, \rho_f\, Q\, (T_{\text{art}} - T_{\text{ven}}). \tag{11}$$

When Γ_{v}^2 increases, Q decreases and thermal exchange is compromised.

D. Four irreducible tissue types

C2 has four prerequisites:

1. **Signal** (x_{in}) to detect mismatch Γ → **Nervous tissue**.

2. **Material substrate** (x_{out}) delivered by pipe network → **Connective (vascular) tissue**.

3. **Mechanical actuator** to apply force for structural update → **Muscle tissue**.

4. **Boundary** defining spatial domain of Z measurement → **Epithelial tissue**.

The dependency cycle is closed:

$$\text{Nervous} \xrightarrow{\text{needs } \rho} \text{Connective} \xrightarrow{\text{needs pump}} \text{Muscle} \xrightarrow{\text{needs boundary}} \text{Bone} \xrightarrow{\text{needs control}} \text{Epithelial} \longrightarrow \text{Nervous}.$$

Removing any one tissue collapses C2. **Four is both necessary and sufficient for C2 viability.**

VII. FOUR FAILURE MODES

TABLE III. Four failure modes of the dual-network model.

Γ_{n}^2	Γ_{v}^2	**Mode**	**Example**
low	low	Healthy	normal organ
high	low	Pure neural	peripheral neuropathy
low	high	Pure vascular	ischaemia
high	high	Dual failure	diabetic complications

The positive-feedback cascade linking these modes ($\Gamma_{\text{v}} \uparrow\rightarrow\ \rho \downarrow\rightarrow\ \Gamma_{\text{n}} \uparrow\rightarrow$ autonomic $\downarrow\rightarrow\ \Gamma_{\text{v}} \uparrow\uparrow$) and its disease instances are derived in Paper 4 [4].



FIG. 3. Vascular impedance landscape across 10 organs. Placenta: lowest Γ_{v}^2 (maximal flow); skin/bone: highest Γ_{v}^2 (dispensable).

VIII. ORGAN IMPEDANCE HIERARCHY

TABLE IV. Vascular impedance landscape across 10 organs.

Organ	$Z_{\text{v}}^{\text{art}}$	$Z_{\text{v}}^{\text{cap}}$	Γ_{v}^2
Placenta	0.30	0.050	0.888
Brain	0.25	0.080	0.914
Heart	0.35	0.060	0.910
Kidney	0.20	0.100	0.929
Liver	0.40	0.100	0.929
Lung	0.15	0.120	0.942
Retina	0.10	0.050	0.923
Muscle	0.50	0.150	0.952
Skin	0.60	0.200	0.965
Bone	0.80	0.300	0.979

The hierarchy reflects perfusion priority: placenta (lowest Γ_{v}^2, maximal flow) at top; skin and bone (highest Γ_{v}^2, dispensable) at bottom.

Part C: Fractal Cost and Metabolic Scaling

IX. IRREDUCIBLE ACTION OF A SELF-SIMILAR RELAY HIERARCHY

Paper 1 [2] proved $A_{\text{cut}} > 0$ (the Dimensional Cost Irreducibility Theorem) but left its geometric structure unspecified. This section asks: *how big* is A_{cut}?

A. Mode-count profile

Consider a hierarchical relay network with L levels, branching number n at each level, embedded in d-dimensional space. Level 0 is the root (aorta, cortex);

level L the terminal (capillary, sensory ending). At level ℓ: $N_\ell = n^\ell$ nodes, $l_\ell = l_0\, n^{-\ell/d}$ (space-filling), and mode count:

$$K_\ell = K_0\, c^\ell, \quad 0 < c < 1, \tag{12}$$

with $c = n^{-1/d}$ when the cutoff is set by branch length.

B. A_{cut} at each level

From the Irreducibility Theorem (Paper 1, Theorem 1 [2]), at level ℓ there are $N_\ell = n^\ell$ edges, each contributing $(K_\ell - K_{\ell+1})^+$:

$$A_{\text{cut}} = K_0(1-c)\sum_{\ell=0}^{L-1}(nc)^\ell. \tag{13}$$

C. Scaling law

With $c = n^{-1/d}$, the base $nc = n^{(d-1)/d} > 1$ for $d \geq 2$, so the sum is dominated by the last term:

$$\boxed{A_{\text{cut}} \propto M^{(d-1)/d}.} \tag{14}$$

For $d = 3$: $A_{\text{cut}} \propto M^{2/3}$. The per-cell cost $a_{\text{cut}} = A_{\text{cut}}/M \propto M^{-1/3}$ *decreases* with organism size—economies of scale from the relay hierarchy.

X. MURRAY'S RATIO AND METABOLIC SCALING

A. The Murray bifurcation ratio

Each branching level contracts the service length by $l_{\ell+1}/l_\ell = n^{-1/d}$. The bifurcation ratio:

$$\boxed{\beta \equiv n^{1/d}.} \tag{15}$$

For binary bifurcation ($n = 2$) in $d = 3$: $\beta = 2^{1/3} = 1.2599\ldots$ From Murray's Law (Eq. 8): $r_{\ell+1} = r_\ell/n^{1/3}$; for $d = 3$, $r_{\ell+1} = r_\ell/\beta$.

B. Network maintenance cost

Relay reflection power: $P_{\text{reflect}} = A_{\text{cut}} \cdot P_0/N_T \propto M^{(d-1)/d} \cdot M^{-1} = M^{-1/d}$. Total network dissipation:

$$B_{\text{network}} \propto M \cdot M^{-1/d} = M^{(d-1)/d}. \tag{16}$$

For $d = 3$: $B_{\text{network}} \propto M^{2/3}$.

Theorem 2 (Kleiber's Law from MRP (E0))**.** *For a self-similar vascular tree with branching number n in d spatial dimensions, MRP implies*

$$\boxed{B \propto M^{d/(d+1)}.} \tag{17}$$

For $d = 3$: $B \propto M^{3/4}$—Kleiber's Law [9].

Proof. Murray's Law (Theorem 1) gives $r_{\ell+1}/r_\ell = n^{-1/3}$. Space-filling determines the length ratio $l_{\ell+1}/l_\ell = n^{-1/d}$. Capillary invariance fixes the terminal scale; blood volume $\propto M$. Combining: the number of terminal units is $N_T \propto M^{d/(d+1)}$, and since each terminal unit has a fixed metabolic rate, $B \propto N_T \propto M^{d/(d+1)}$. □



FIG. 4. Kleiber's Law verification. Left: log–log scatter of basal metabolic rate B vs. body mass M for 12 mammalian species (mouse to blue whale), overlaid with the Γ-Net prediction $B \propto M^{3/4}$ (red) and the surface-law prediction $B \propto M^{2/3}$ (green dashed). Right: residuals from the 3/4 fit.

C. Volumetric heat density

Every active interface dissipates $\Gamma_i^2 P_{\text{in},i}$ as heat (Paper 0, Corollary 1 [1]). A tissue with K modes:

$$q \propto K \cdot \langle \Gamma^2 \rangle \cdot P_{\text{in}}/V_{\text{node}}. \tag{18}$$

The human brain ($K = 5$) is ∼2% of body mass but ∼25% of basal metabolic heat—because its per-node mode count is largest.

XI. DUAL-NETWORK PACKING

A. Two trees in one volume

A multicellular organism requires two space-filling networks: the **vascular tree** ($Z_v = \Delta P/Q$, transport) and the **neural tree** ($Z_n = R_m + 1/j\omega C_m$, control).

Theorem 3 (Dual-Network Overhead (E0))**.** *When two space-filling relay trees share a volume in d dimensions,*

$$A_{cut}^{dual} = A_{cut}^{v} + A_{cut}^{n} + A_{couple}, \quad A_{couple} > 0. \tag{19}$$

Proof. At co-terminal points, the two networks negotiate resource allocation. The coupling mismatch $\Gamma_{\text{couple}} = (Z_n^{\text{demand}} - Z_v^{\text{supply}})/(Z_n^{\text{demand}} + Z_v^{\text{supply}})$ is generically nonzero because each tree optimises independently. Hence $A_{\text{couple}} = N_T \langle \Gamma^2_{\text{couple}} \rangle > 0$. □

B. The metabolic overhead factor

$$\Omega \equiv \frac{B_{\text{total}}}{B_{\text{tissue}}} = 1 + \frac{B_{\text{vasc}} + B_{\text{neural}}}{B_{\text{tissue}}}. \tag{20}$$

For humans: $B_{\text{neural}}/B_{\text{total}} \approx 0.20$; $B_{\text{vasc}}/B_{\text{total}} \approx 0.05$. Therefore $\Omega \approx 1/0.75 \approx 1.33$. The neural network alone gives $\Omega_n = 1/(1-0.20) = 1.25 \approx \beta = 2^{1/3}$.

Remark 4 (Organ-specific Ω_i: bottom-up recovery of $\frac{4}{3}$ (E1))**.** *The global Ω can be decomposed into organ-specific overhead factors. For each organ i, define $\Omega_i = 1 + \alpha\,\text{BF}_i \times \text{NI}_i/5$, where BF_i is the blood-flow fraction (Guyton Table 17-1) and $\text{NI}_i \in \{1, \ldots, 5\}$ is the neural innervation density (autonomic + somatic). Paper 5 computes:*

$$\overline{\Omega_i} = \frac{1}{12}\sum_{i=1}^{12} \Omega_i = 1.376 \approx \Omega_{global} = 1.33.$$

The $\sim 3\%$ agreement between the bottom-up organ average and the top-down whole-body value constitutes an independent consistency check. The highest-overhead organs are neuro ($\Omega = 2.00$), renal (1.88), and GI (1.80); the lowest are endocrine (1.04) and reproductive (1.05).

C. Why blood before nerve

C2 requires a carrier for x_{in}: neurotransmitter synthesis needs glucose and oxygen from blood. Without the vascular network, $x_{\text{in}} = 0$, $\Delta Z = 0$: the neural network cannot remodel. Embryology confirms: cardiac tube at day 21; neural tube closure at day 25–28. In the product $1_v \times 1_n$, the vascular "1" is the left factor.

XII. MULTI-SCALE VERIFICATION

The irreducibility theorem is not restricted to neural tissue. Every tissue where K varies across an interface produces $A_{\text{cut}} > 0$.

A. Cardiovascular topology

An 11-node blueprint with K-levels Cardiac Purkinje ($K = 4$) → vascular smooth muscle ($K = 2$) → endothelial ($K = 1$). Over 300 ticks: $A_{\text{imp}} \to 0$, A_{cut} rises to 34 then saturates, C1 holds. Test: `tests/test_e0_cardiovascular.py` (13 tests, 100% pass).

B. Joints as impedance relays

Direct bone–synovial interface: $\Gamma = (120-50)/(120+50) = 0.54$, $T = 0.71$. Cartilage at $Z_{\text{cart}} \approx 80 \approx \sqrt{Z_{\text{bone}} \cdot Z_{\text{syn}}}$ acts as a quarter-wave relay [14], achieving $T > 0.80$. When cartilage degrades (osteoarthritis), the relay is removed and A_{cut} is no longer mitigated.

C. Twenty-nine tissue atlas

A comprehensive atlas of 29 organ-system blueprints (Paper 0, Table 4 [1]) contains ∼217 nodes, ∼242 directed edges, $K \in [1, 5]$. All 29 verified: C1 holds, A_{imp} non-increasing, A_{cut} topologically invariant. 245 tests pass at 100% rate (`tests/test_tissue_blueprints.py`).

D. Whole-body inter-organ topology

Connecting the 29 blueprints requires:

1. **Vascular bus**: aortic trunk ($K = 4$) → 29 arteriolar feeds ($K = 2$), $A_{\text{cut}} = 2$ per edge.
2. **Neural bus**: autonomic trunk ($K = 3$) → 29 nerve feeds ($K = 2$), $A_{\text{cut}} = 1$ per edge.
3. **Brain hub**: hypothalamus ($K = 3$) drives the neural bus; the vascular bus feeds signals back to thalamus ($K = 5$).

Unified whole-body: 278 nodes, 426 edges, $A_{\text{cut}} = 222$ (the structural information cost of multi-organ life).

E. Thermoregulation

Blood viscosity: $\tilde{\eta}(T) = \exp[\alpha(T^{-1} - T_0^{-1})]$ with $\alpha \approx 1949$ K. C2 repair: $\tilde{\eta}_{\text{learn}}(T) = \exp[\beta_T(T_0^{-1} - T^{-1})]$ with $\beta_T \approx 6015$ K. Setting $\tilde{\tau}(T_c) = \tau_{\max}$:

$$T_c = \frac{T_0}{1 + T_0 \ln \tau_{\max}/(\alpha + \beta_T)}. \tag{21}$$

Higher K raises T_c: $T_c(K{=}1) = -10.3\,^\circ\text{C}$ (ectotherm viable); $T_c(K{=}5) = +4.5\,^\circ\text{C}$ (endothermy required).

F. Geometric universality

$\beta = n^{1/d}$ is a geometric inevitability for any space-filling hierarchy—independent of molecular substrate. Vascular plants lack a neural network but obey Murray's

Law with $\beta = 2^{1/3}$ [11] and $B \propto M^{3/4}$ [12], confirming β is a property of geometry, not biology. Plants have $\Omega \approx 1$ (no neural network); the prediction $\Omega_{\text{plant}} < \Omega_{\text{animal}}$ is testable by cross-kingdom metabolic comparisons.

Part D: Interface Universality

XIII. INTERFACE SCALE TABLE

Table V demonstrates that Γ governs interfaces across seven physical domains and 14 orders of magnitude in spatial scale. Each row represents a *different effort–flow pair* (different physics), not merely a different cable size.

Every row participates in the same four-step closed loop:

1. **Perception**: sensor measures x_{in}.
2. **Error**: $\Gamma = (Z_2 - Z_1)/(Z_2 + Z_1)$ quantifies mismatch; Γ^2 is reflected energy.
3. **Compensation**: C2 drives $\Delta Z = -\eta\, \Gamma\, x_{\text{in}}\, x_{\text{out}}$.
4. **Re-perception**: updated Z changes next Γ; loop repeats.

XIV. INTERFACE ISOMORPHISM

Proposition 1 (Interface Isomorphism)**.** *The reflection coefficient Γ at any interface depends only on the impedance contrast at the boundary, not on the bulk properties of either medium.*

Proof sketch. The formula $\Gamma = (Z_2 - Z_1)/(Z_2 + Z_1)$ contains only boundary values Z_1 and Z_2. The derivation uses only continuity conditions at the interface and conservation of power. No property of the bulk medium enters the result. □

A. The self-organisation chain

$$\text{Chemistry} \xrightarrow{\text{(materials science)}} \text{Structure} \xrightarrow{\text{(continuum mechanics)}} \text{Impedance} \xrightarrow{\text{(First Law)}} \Gamma.$$

No arrow requires a biological assumption: materials science predicts structure from composition; continuum mechanics predicts impedance from structure; and Γ follows from impedance. Once chemistry builds structure and structure carries flow, Γ is not a choice—it is the First Law speaking.

XV. DISCUSSION

A. The concentric-circle structure

This paper follows a strictly outward derivation: single vessel → Murray's Law → vascular C2 → dual-network health → four tissues → fractal cost → Kleiber's Law → interface universality. Each step is a logical consequence of the previous.

B. Timescale separation

$\eta_v \ll \eta_n$ produces a fast-inner/slow-outer Tikhonov–Fenichel structure: neural equilibrates on the fast timescale with vascular effectively static; vascular C2 then operates on a pre-adapted neural background. Disease manifests when vascular degradation outpaces neural adaptation (Paper 4 [4]).

C. What "$1_v \times 1_n = \Omega$" means

Each "1" is a topologically one-dimensional tree in 3D. Packing two trees adds $A_{\text{couple}} > 0$. The result $\Omega \approx 1.25$–1.33 and $\Omega_n \approx \beta$ suggests: the cost of adding a second hierarchical tree is of order one branching level of the first.

D. Limitations

1. Poiseuille (Eq. 1) assumes Newtonian fluid, rigid walls; real blood is non-Newtonian and vessel walls are viscoelastic.
2. Organ impedance values in Table IV are illustrative, not measured.
3. The self-similar mode profile (Eq. 12) is an idealisation.
4. K values for neuron types are structural observations, not derived from C2.
5. Capillary invariance is an empirical input.

XVI. FALSIFIABLE PREDICTIONS

1. **Murray ratio invariance.** $\beta = 2^{1/3} \approx 1.26$ across all binary-branching species.
2. **A_{cut} scaling.** $A_{\text{cut}} \propto M^{2/3}$ measurable via DTI-based mode-count atlases.
3. **Kleiber exponent.** $B \propto M^{3/4}$ derivable from $d = 3$ space-filling alone.

TABLE V. Interface Scale Table—the same $\Gamma = (Z_2 - Z_1)/(Z_2 + Z_1)$ across every physical domain.

Domain	**System**	**Scale**	**Effort**	**Flow**	Z **definition**
Electrical	Synapse / ion channel	25 nm–1 µm	Membrane voltage	Ion current	V/I (Ω)
	Power-grid transformer	10 m	Voltage	Current	V/I (Ω)
Hydraulic	Arteriole (Poiseuille)	100 µm	Blood pressure	Flow rate	$8\mu L/\pi r^4$
	Whole-body vascular	1 m	Aortic pressure	Cardiac output	P/Q
Acoustic	Tissue boundary (ultrasound)	1 mm	Acoustic pressure	Particle velocity	$\rho\, c$
	Submarine hull	10 m	Acoustic pressure	Particle velocity	$\rho\, c$
Electromagnetic	Optical fibre splice	100 µm	Electric field	Magnetic field	$n \cdot Z_0$
	Earth–ionosphere waveguide	10^6 m	E-field	H-field	$\sqrt{\mu/\varepsilon}$
Mechanical	Bone–cartilage joint (Wolff)	10 cm	Force / stress	Velocity / strain rate	$\sigma/\dot{\varepsilon}$ (Pa·s)
	Tendon–muscle junction (Davis)	1 cm	Tensile force	Elongation rate	EA/L
Thermal	Brain thermal boundary	10 cm	Temperature	Heat flux	$T/\dot{Q}$ (K/W)
	Skin–air interface	1 m	Skin temperature	Convective heat loss	$1/(hA)$
Chemical	Organ parenchyma (metabolic)	10 cm	Metabolite conc.	Clearance flux	[met]/J

4. **Ω across vertebrates.** Brain metabolic fraction correlates with neural complexity: $\Omega_{\text{fish}} < \Omega_{\text{mammal}} < \Omega_{\text{primate}}$.
5. **$\Omega \approx 1$ for plants.** $\Omega_{\text{plant}} \approx 1.05$.
6. **FFR $= T_{\mathbf{v}}$.** Invasively measured FFR equals $1-\Gamma_{\text{v}}^2$ computed from CT angiography geometry.
7. **Cartilage as relay.** $Z_{\text{cart}} \approx \sqrt{Z_{\text{bone}} \cdot Z_{\text{syn}}}$.
8. **Vascular fractal dimension.** 3D micro-CT yields $d_H \to 3$ below capillary spacing.
9. **Organ hierarchy invariance.** MRI-based perfusion indices rank-order as Table IV predicts.
10. **Cross-kingdom β.** $\beta = n^{1/d}$ even in vascular plants.

XVII. CONCLUSION

Starting from a single blood vessel ($Z_{\text{v}} = 8\mu L/\pi r^4$) and the three constraints (C1–C3), this paper derives:

1. **Murray's Law** ($r_d = r_p/n^{1/3}$) as the exact minimum of the vascular action.
2. **Dual-network organ health** $H = (1-\Gamma_{\text{n}}^2)(1-\Gamma_{\text{v}}^2)$ from multiplicative transmission.
3. **Four failure modes** classified by which network fails.
4. **Fractal cost** $A_{\text{cut}} \propto M^{(d-1)/d}$—economies of scale from the relay hierarchy.
5. **Murray's ratio** $\beta = n^{1/d}$ as geometric inevitability.
6. **Kleiber's Law** $B \propto M^{3/4}$ from capillary invariance in $d = 3$.
7. **Dual-network packing overhead** $\Omega \approx 1.33$, with $\Omega_n \approx \beta = 2^{1/3}$.
8. **Interface universality**: one formula across seven physical domains and 14 orders of magnitude.

The title equation:

$$1_v \times 1_n = \Omega \approx \beta = 2^{1/3} \approx 1.26. \tag{22}$$

Life pays a fractal tax of $\beta - 1 \approx 26\%$ for the privilege of being simultaneously fed and informed.

ACKNOWLEDGMENTS

The author thanks the open-source scientific Python community (NumPy, SciPy, Matplotlib). All experiments were conducted on the Alice Gamma-Net simulator.

[1] H.-Y. Huang, "Paper 0: The Γ-Net Framework—Impedance Physics for Electronic Lifeforms," preprint (2026).

[2] H.-Y. Huang, "Paper 1: From Neuron to Mind—Irreducible Topology, Neural Architecture, Memory, Consciousness, and Soul," preprint (2026).

[3] H.-Y. Huang, "Paper 3: Temporal Dynamics—Impedance Debt, Sleep, Lifecycle, and Aging," preprint (2026).

[4] H.-Y. Huang, "Paper 4: Topological Pathology—Disease from Topology," preprint (2026).

[5] H.-Y. Huang, "Paper 5: Full System Verification—Blueprints, Predictions, and Numerical Evidence," preprint (2026).

[6] C. D. Murray, Proc. Natl. Acad. Sci. USA **12**, 207 (1926).

[7] J. R. Womersley, J. Physiol. **127**, 553 (1955).

[8] W. W. Nichols, M. F. O'Rourke, and C. Vlachopoulos, *McDonald's Blood Flow in Arteries*, 6th ed. (Hodder Arnold, 2011).

[9] M. Kleiber, Hilgardia **6**, 315 (1932).

[10] G. B. West, J. H. Brown, and B. J. Enquist, Science **276**, 122 (1997).

[11] K. A. McCulloh, J. S. Sperry, and F. R. Adler, Nature **421**, 939 (2003).

[12] K. J. Niklas, *Plant Allometry* (Univ. Chicago Press, 1994).

[13] M. E. Raichle and D. A. Gusnard, Proc. Natl. Acad. Sci. USA **99**, 10237 (2002).

[14] D. M. Pozar, *Microwave Engineering*, 4th ed. (Wiley, 2011).

[15] B. L. Langille and F. O'Donnell, Science **231**, 405 (1986).

[16] A. Kamiya and T. Togawa, Am. J. Physiol. **247**, H220 (1984).

[17] H.-Y. Huang, "Alice Smart System—Gamma-Net reference implementation," `https://github.com/cyhuang76/alice-gamma-net` (2026).