

Problem Set 7

Parvesh Adi Lachman

November 2023

1 Problem 1

Consider the algorithm below, which takes an $n \geq 0$ and finds its remainder when divided by $c \geq 1$

```
function REMAINDER( $n$ ):  
  if  $n \leq c - 1$  then  
    return  $n$   
  else  
    return REMAINDER( $n - c$ )  
  end if  
end function
```

Claim: Let $c \geq 1$. For any $n \geq 0$, $\text{remainder}(n) = n \bmod c$.

2 Problem 2

Claim: Let $n, c \geq 1$ and $c \leq n$. The number of simple paths of length c in the complete graph on n nodes is $\frac{n!}{(n-c-1)!}$ which is equal to $n(n-1) \cdots (n-c)$.

complete graph K_n : an undirected graph on n nodes with an edge between every pair of nodes.

simple path: a sequence of distinct nodes with edges between consecutive nodes in the sequence.

length of a path: the number of *edges* in the path (**not** number of nodes).

Proof: Let G be a complete graph on n nodes. Let v_1, v_2, \dots, v_n be the nodes of G . Let P be a simple path of length c in G . Then P is a sequence of c distinct nodes in G . Since P is a simple path, the nodes in P are distinct. Thus, P is a permutation of c distinct nodes in G . Since G has n nodes, there are n choices for the first node in P , $n-1$ choices for the second node in P , and so on. Thus, there are $n(n-1) \cdots (n-c)$ simple paths of length c in G .