

PART-TO-WHOLE ARRAY


$$A_{[S,x]}$$

$s_j \in \mathcal{S}$	x
x1111	100
x2222	200
x3333	300
x4444	400
x5555	500
x6666	600
x7777	700

total 2800

CROSSMAP

$$\mathcal{X} = (\mathcal{S}, \mathcal{T}, \mathcal{R}, \mathcal{W})$$



GRAPH
($\mathcal{S}, \mathcal{T}, \mathcal{R}, \mathcal{W}$)

	A1	B2	B3	C5	D6	D7
x1111	1	0	0	0	0	0
x2222	0	0.5	0.5	0	0	0
x3333	0	0	0	1	0	0
x4444	0	0	0	1	0	0
x5555	0	0	0	0	0.4	0.6
x6666	0	0	0	0	0.3	0.7
x7777	0	0	0	0	1	0

MATRIX
 C ($y = C^T x$)

s_j	t_k	w
x1111	A1	1
x2222	B2	0.5
x2222	B3	0.5
x3333	C5	1
x4444	C5	1
x5555	D6	0.4
x5555	D7	0.6
x6666	D6	0.3
x6666	D7	0.7
x7777	D6	1

LIST
 $E(s_j, t_k, w)$

PART-TO-WHOLE ARRAY

$$A'_{[\mathcal{T},y]}$$

$t_k \in \mathcal{T}$	y
A1	100
B2	100
B3	100
C5	700
D6	1080
D7	720

total 2800