

REVIEW ARTICLE

PCR inhibitors – occurrence, properties and removalC. Schrader¹, A. Schielke², L. Ellerbroek¹ and R. Johné¹¹ Food Hygiene and Safety Concepts, Federal Institute for Risk Assessment, Berlin, Germany² Department for Infectious Disease Epidemiology, Robert Koch Institute, Berlin, Germany**Keywords**

detection, environmental, food, polymerase chain reaction.

Correspondence

Reimar Johné, Federal Institute for Risk Assessment, Max-Dohrn-Strasse 8-10, 10589 Berlin, Germany. E-mail: reimar.johne@bfr.bund.de

2012/0617: received 5 April 2012, revised 14 June 2012 and accepted 27 June 2012

doi:10.1111/j.1365-2672.2012.05384.x

Summary

The polymerase chain reaction (PCR) is increasingly used as the standard method for detection and characterization of microorganisms and genetic markers in a variety of sample types. However, the method is prone to inhibiting substances, which may be present in the analysed sample and which may affect the sensitivity of the assay or even lead to false-negative results. The PCR inhibitors represent a diverse group of substances with different properties and mechanisms of action. Some of them are predominantly found in specific types of samples thus necessitating matrix-specific protocols for preparation of nucleic acids before PCR. A variety of protocols have been developed to remove the PCR inhibitors. This review focuses on the general properties of PCR inhibitors and their occurrence in specific matrices. Strategies for their removal from the sample and for quality control by assessing their influence on the individual PCR test are presented and discussed.

Background

In the last few decades, the polymerase chain reaction (PCR) has become one of the most powerful molecular biological tools. PCR may be used for diagnosis of infectious or hereditary diseases and for genetic analyses in a large variety of sample types. The PCR is a very rapid and sensitive method, in which genomic DNA is exponentially amplified by a DNA polymerase using specific primer molecules. Variations of the PCR comprise the reverse transcription (RT-) PCR using RNA as template, which is first transcribed into DNA by a reverse transcriptase, or real-time PCR, which uses fluorescent probes for the detection of the PCR product providing quantitative information. The PCR is an enzymatic reaction and therefore sensitive to inhibitors. The occurrence of such so-called PCR inhibitors, which comprise all substances that have a negative effect on the PCR, is a major drawback of the PCR. PCR inhibitors can originate from the sample or may be introduced during sample processing or nucleic acid extraction. The major consequence of a partly or total inhibition of the PCR is a decreased sensitivity or false-negative results, respectively.

In the following sections, the different classes of PCR inhibitors and their mechanisms of action are presented. The incidence of PCR inhibitors in different types of matrices such as clinical, food or environmental specimens will be reviewed. Finally, different possibilities to remove PCR inhibitors as well as strategies for quality controls of the PCR will be presented and discussed.

Substance classes of PCR inhibitors

PCR inhibitors are a very heterogeneous group of chemical substances. One certain matrix may contain many different inhibitory substances and the same inhibitors can be found in many different matrices. Organic as well as inorganic substances, which may be dissolved or solid, can appear as PCR inhibitors. Calcium ions are an example for inorganic substances with inhibitory effects on the PCR. However, most of the known inhibitors are organic compounds, for example, bile salts, urea, phenol, ethanol, polysaccharides, sodium dodecyl sulphate (SDS), humic acids, tannic acid, melanin as well as different proteins, such as collagen, myoglobin, haemoglobin, lactoferrin, immunoglobulin G (IgG) and proteinases (Rossen *et al.*

1992; Rådström *et al.* 2004). Besides the substance class, the concentration of the compound is important for its inhibitory effect. For a detailed review on the relation of the concentration of inhibitors (e.g. sodium chloride, magnesium chloride, sucrose, phenol, SDS, ethanol or cetrimonium bromide) and their inhibitory effects, see the article of Rossen *et al.* (1992).

PCR inhibitors can be found in a variety of biological materials (organs, blood, body fluids etc.), environmental samples (water, soil, air etc.) and food (meat, milk, fruits, vegetables, seafood etc.). In addition, inhibitory substances may also be unintentionally added during transport, sample processing (e.g. pre-concentration procedures) or nucleic acid extraction. Table 1 shows some examples of matrices and their typical inhibitors.

Mechanisms of action of PCR inhibitors

PCR inhibitors may interfere with different steps of a PCR analysis (Fig. 1). Generally, several PCR components, especially DNA, may adsorb to polymeric surfaces, for example, to the wall of vessels and reaction tubes, during sample processing, extraction or during PCR (Butot *et al.* 2007b; Fox *et al.* 2007; Gassilloud *et al.* 2007; Gonzalez *et al.* 2007). The efficacy of sample processing and nucleic acid extraction may be affected. Nucleases may degrade template RNA or DNA. Phenols may cross-link RNA under oxidizing conditions and thus hamper RNA isolation (Su and Gibor 1988; Wilkins and Smart 1996). In addition, the existence of polysaccharides may reduce the capacity to resuspend precipitated RNA (Wilkins and Smart 1996; Sipahioglu *et al.* 2006). Reverse transcription may be inhibited, for example, by direct interaction of the enzyme with melanin (Eckhart *et al.*

2000). The DNA used as template of the PCR can be modified or degraded by nucleases and other substances. Annealing of the primers to the DNA template may be disturbed by certain PCR inhibitors (Chandler *et al.* 1998; Abbaszadegan *et al.* 1999). As this effect is because of a competitive binding of the inhibitor to the template, appropriate primer design leading to higher melting points can overcome this problem (Huggett *et al.* 2008; Opel *et al.* 2010).

There are many PCR inhibitors that target the DNA polymerase directly or indirectly. Proteases or detergents present in the reaction can degrade this enzyme (Rossen *et al.* 1992; Powell *et al.* 1994). For example, urea (Saulnier and Andremonet 1992; Wilson 1997) and phenol (Katcher and Schwartz 1994) are known to degrade DNA polymerases. Calcium, collagen, haematin and tannic acid may inhibit polymerase activity (Opel *et al.* 2010). Melanin forms a reversible complex with the DNA polymerase (Eckhart *et al.* 2000) and polysaccharides may disturb the enzymatic process by mimicking the structure of nucleic acid (Peist *et al.* 2001). Humic acids interact with the template DNA and the polymerase thus preventing the enzymatic reaction even at low concentrations (Sutlović *et al.* 2005, 2008). Other substances react with cofactors of the polymerase. High concentrations of calcium may lead to a competitive binding by the DNA polymerase instead of magnesium and complexing agents, for example, tannic acid, deplete magnesium. In both cases, magnesium is no longer available as cofactor for the polymerase and its activity is decreased (Opel *et al.* 2010). For real-time PCR assays, the interference with the fluorescent probes or increased background fluorescence represents additional mechanisms of action for PCR inhibitors decreasing sensitivity. Examples of PCR

Table 1 Selected matrices and their identified polymerase chain reaction inhibitors

Matrix	Contained inhibitors	References
Clinical specimens (e.g. blood; muscle tissues)	Antiviral substances (e.g. acyclovir), Haemoglobin, Heparin, Hormones, IgG, Lactoferrin, Myoglobin	Al-Soud and Rådström (2001); Burkardt (2000); Rådström <i>et al.</i> (2004); Yedidag <i>et al.</i> (1996)
Stool	Complex polysaccharides, Bile salts, Lipids, Urate	Kreader (1996); Monteiro <i>et al.</i> (1997); Rådström <i>et al.</i> (2004); Chaturvedi <i>et al.</i> (2008)
Seafood, bivalves, oysters	Algae, Glycogen, Polysaccharides	Atmar <i>et al.</i> (1993, 1995); Richards (1999)
Berries	Phenols, Polysaccharides	Seeram <i>et al.</i> (2006); Wei <i>et al.</i> (2008)
Plants	Pectin, Polyphenols, Polysaccharides, Xylan	Demeke and Adams (1992); Henson and French (1993); John (1992); Sipahioglu <i>et al.</i> (2006); Su and Gibor (1988); Wan and Wilkins (1994); Wei <i>et al.</i> (2008); Wilkins and Smart 1996
Cheese, milk	Proteases (e.g. plasmin), Calcium ions,	Bickley <i>et al.</i> (1996); Powell <i>et al.</i> (1994); Rossen <i>et al.</i> (1992)
Water, environment	Debris, Fulmic acids, Humic acids, Humic material Metal ions, Polyphenol	Abbaszadegan <i>et al.</i> (1993); Ijzerman <i>et al.</i> (1997)
Palaeobiology, archaeology, forensic	Bone dust, Coprolite Peat extract, Clay-rich soil	Baar <i>et al.</i> (2011)




Figure 1 Schematic presentation of the attack points of polymerase chain reaction (PCR) inhibitors during sample preparation and PCR. The nucleic acids may interfere with surfaces of the vessels (1) or substances may react with nucleic acids (2) during sample processing and extraction. Other substances inhibit reverse transcription (3) and degrade or modify the template DNA (4). Annealing of primers to the template can be hampered (5) or the DNA polymerase is degraded, inhibited or altered (6). Finally, substances may interfere with binding of probes or with their fluorophores (7).

inhibitors and their proposed mechanisms of action are presented in Table 2. However, it has to be mentioned that for many inhibitory substances, the distinct mechanism of action is not known so far.

PCR inhibitors in clinical samples

In blood, serum or plasma samples, substances like IgG, haemoglobin and lactoferrin have been described as inhibitors of PCR (Al-Soud *et al.* 2000; Al-Soud and Rådström 2001). Anticoagulants, for example, heparin, may also inhibit the PCR (Costafreda *et al.* 2006). Many of these inhibitors are presumed to affect directly the RNA and not the enzymes of the reaction (Konet *et al.* 2000). In addition, hormones or antiviral substances like acyclovir can also affect amplification (Yedidag *et al.* 1996; Burkardt 2000).

The most critical component in urine samples is urea, which may lead to the degradation of the polymerase (Saulnier and Andreumont 1992; Wilson 1997). However, the effect is dependant on the concentration of urea in the sample and initiates with a concentration of about 50 mmol l⁻¹ (Khan *et al.* 1991). As the urine excretion in humans is – dependent on diet and age – in the range between 340 and 580 mmol urea per day, inhibitory concentrations are not unusual.

Stool and faecal samples contain highly variable components dependant on nutrition, gut flora, lifestyle and environment of the patient (Oikarinen *et al.* 2009). Inhibitors may include polysaccharides or chlorophyll originating from herbs and vegetables, bile salts, urea, glycolipids, haemoglobin and heparin (Lantz *et al.* 1997; Monteiro *et al.* 1997; Pontiroli *et al.* 2011).

The most important inhibitory factors present in bile samples are the bile acids and their corresponding salts. However, efficient amplification can be achieved using a polymerase which is not sensitive to these substances (Al-Soud *et al.* 2005). The same applies to the analysis of eye fluids or muscle tissue containing myoglobin (Wiedbrauk *et al.* 1995; Bélec *et al.* 1998).

PCR inhibitors in food and environmental samples

Many different PCR inhibitors have been identified in food. Fats, glycogen, polysaccharides, minerals as well as enzymes present in food may cause inhibition of the PCR (Powell *et al.* 1994; Richards 1999). In milk samples, PCR inhibition is mainly dependant on the concentration of calcium, whereas the fat content seems to have only minor influence on the amplification efficiency (Bickley *et al.* 1996). Additionally, Powell *et al.* (1994) identified plasmin, which degrades the *Taq* polymerase, as an inhibitor naturally occurring in milk.

In seafood, mainly polysaccharides seem to be responsible for PCR inhibition (Atmar *et al.* 1993, 1995). In addition, the glycogen content in the tissues of bivalve molluscs influences PCR efficiency (Enriquez *et al.* 1992; Richards 1999). Generally, the ability of bivalve molluscs to filter the water may lead to concentration of different inhibitory substances.

Plants carry many substances, such as polysaccharides, polyphenols, pectin and xylan, which may be co-extracted and thereafter hamper the PCR (Wei *et al.* 2008). Although for most of these polysaccharides (dextran, inulin, pectin or starch), no inhibitory effect was reported, some of them like dextran sulphate and ghatti gum affect the PCR efficiency. However, this effect was partly reversible by adding Tween 20, dimethyl sulphoxide or polyethylene glycol 400 (Demeke and Adams 1992). In another study, acidic polysaccharides (ghatti gum, xylan, dextran

Table 2 Examples of PCR inhibitors and their mechanisms of action

Inhibitor	Mechanism of action	References
Polyphenols	Co-precipitation with nucleic acid; reduction in the ability to resuspend precipitated RNA	John (1992), Sipahioglu <i>et al.</i> (2006), Su and Gibor (1988), Wan and Wilkins (1994) and Wilkins and Smart (1996)
Polysaccharides		
Bacterial cells	Degradation/sequestration of nucleic acids	Burkardt (2000), Katcher and Schwartz (1994), Peist <i>et al.</i> (2001), Rossen <i>et al.</i> (1992), Weyant <i>et al.</i> (1990) and Wilson (1997)
Cell debris		
Detergents		
PCR additives		
Proteins		
Polysaccharides		
Salts		
Solvents		
Polyphenols	Cross-linking with nucleic acids; change of chemical properties of nucleic acids	John (1992), Opel <i>et al.</i> (2010) and Wilkins and Smart (1996)
Polysaccharides		
Humic acids		
Collagen		
Melanin		
Humic acid		
Humic matter	Binding/adsorption to nucleic acid and enzymes	Abbaszadegan <i>et al.</i> (1993)
Haematin	Incomplete melting of DNA	Opel <i>et al.</i> (2010)
Indigo		
Metal ions	Reduction in specificity of primers	Abbaszadegan <i>et al.</i> (1993)
Detergents	Degradation of polymerases	Powell <i>et al.</i> (1994), Rossen <i>et al.</i> (1992), Saulnier and Andreumont (1992) and Wilson (1997)
Proteases		
Urea		
Calcium		
Collagen	Inhibition of DNA polymerase or reverse transcriptase activity	Al-Soud <i>et al.</i> (2000a), Al-Soud and Rådström (1998), Eckhart <i>et al.</i> (2000), Opel <i>et al.</i> (2010), Peist <i>et al.</i> (2001) and Wilkins and Smart (1996)
Haematin		
Herbal metabolites		
IgG		
Melanin		
Myoglobin		
Polysaccharides		
Sodium		
Tannic acid		
Polyphenols		
Tannic acid	Chelation of metal ions	Abbaszadegan <i>et al.</i> (1993) and Opel <i>et al.</i> (2010)
EDTA	Chelation of metal ions including Mg ⁺⁺	Rossen <i>et al.</i> (1992)
Calcium ions	Competition with co-factors of the polymerase	Bickley <i>et al.</i> (1996), Opel <i>et al.</i> (2010)
Antiviral substances (e.g. acyclovir)	Competition with nucleotides, inhibition of DNA elongation	Yedidag <i>et al.</i> (1996)
Exogenic DNA	Competition with template	Tamariz <i>et al.</i> (2006)

EDTA, ethylenediaminetetraacetic acid; PCR, polymerase chain reaction.

sulphate), polyphenols as well as inulin and pectin found in tea extracts were proven to inhibit PCR amplification (Peist *et al.* 2001). Berries are generally rich in phenols (e.g. anthocyanin, flavonol, ellagitannin, proanthocyanidin and phenolic acids) and polysaccharides (Seeram *et al.* 2006; Wei *et al.* 2008). Substances present in berries and tomatoes seem to especially inhibit real-time PCR assays using TaqMan probes, whereas conventional PCR assays are less affected (Love *et al.* 2008).

Environmental samples can be very diverse and derived from different compartments including soil, water or air.

This large variety leads to the presence of many different PCR inhibitors including those already described above. In addition, dead biomass and soil may contain humic and fulminic acids, which inhibit PCR even at low concentrations (Ijzerman *et al.* 1997). In sewage sludge, fats, proteins, polyphenols and heavy metals are found, and waste water contains polysaccharides, metal ions (e.g. iron and aluminium) and RNases – all of them are common PCR inhibitors of environmental samples (Shieh *et al.* 1995; Rock *et al.* 2010). Several different inhibitors have also been detected in animal feed (Löfström *et al.* 2004).

A large variety of water samples are frequently used for the detection of pathogens. To detect even low amounts of pathogens, large volumes of water are usually concentrated to very small volumes. However, this often results in concurrent concentration of the different inhibitors and increased interference with PCR (Abbaszadegan *et al.* 1993, 1999; Tsai *et al.* 1993; Jiang *et al.* 2005). Similar problems may occur during the collection of air samples, which is usually carried out by passing large air volumes through a variety of filters or other binding material. By this, inhibitory airborne components are often enriched leading to failure of PCR (Maher *et al.* 2001; Chen *et al.* 2010; Oppliger *et al.* 2011).

PCR inhibitors introduced during sample preparation

Inhibitors may be added to the sample during sample processing or during nucleic acid extraction. This includes powder from gloves (Demeke and Jenkins 2010), different salts (e.g. sodium chloride or potassium chloride), detergents or organic molecules [ethylenediaminetetraacetic acid (EDTA), sarkosyl, ethanol, isopropyl alcohol or phenol] (Weyant *et al.* 1990; Katcher and Schwartz 1994; Burkardt 2000; Peist *et al.* 2001; Demeke and Jenkins 2010). These substances may be necessary for efficient cell lysis or for the preparation of pure nucleic acids, but they may also cause PCR inhibition at certain concentrations. Ionic detergents (e.g. sodium deoxycholate, sarkosyl and SDS) are highly inhibitory for the PCR, whereas non-ionic detergents (e.g. Nonidet P-40, Tween 20, Triton X-100 and N-octyl glucoside) cause PCR inhibition only at relatively high concentrations (Weyant *et al.* 1990). EDTA is found in several elution buffers of purification kits for preservation of DNA, but at certain concentrations, it may deplete magnesium ions and thus inhibit DNA polymerase activity. Additives of the PCR mixture, such as dithiothreitol, dimethyl sulphoxide or mercaptoethanol, may also be inhibitory at certain concentrations. For some of the polymerases commonly used in PCR, the inhibitory concentrations of certain substances have been determined and can be retrieved from the manufacturer.

Contact between polymeric surfaces and UV-irradiated plastic tubes with PCR chemicals is reported to decrease the sensitivity of the PCR (Butot *et al.* 2007b; Fox *et al.* 2007; Gassilloud *et al.* 2007; Gonzalez *et al.* 2007), although Tamariz *et al.* (2006) did not find any influence of UV irradiation. However, this discrepancy might be explained by different dosages of UV light used, which have been shown to correlate with the inhibitory effect (Burgess and Hall 1999). Generally, the material of swabs or the composition of transport media may also influence the sensitivity of the PCR (Wadowsky *et al.* 1994).

General methods for removal of PCR inhibitors

Several methods for the removal of inhibitors or for the reduction of their effects have been proposed so far and some examples are shown in Table 3. Generally, the effects of the inhibitors may be reduced by selecting an appropriate method for sample processing and nucleic acid extraction, by the choice of a more robust DNA polymerase or by the use of specific PCR additives (Al-Soud and Rådström 2001). For example, guanidinium thiocyanate extraction may remove inhibitors from different sample matrices more efficient than other methods (Shieh *et al.* 1995; Hale *et al.* 1996). Other effective strategies include a phenol–chloroform extraction for the removal of inhibitory lipids or a treatment with activated carbon to eliminate inhibitory salts, such as urates (Wiedbrauk *et al.* 1995; Abolmaaty *et al.* 2007; Chaturvedi *et al.* 2008). These methods are reported to be more successful than gel filtration, treatment with proteinase K or heat treatment (Huppertz *et al.* 1993; Bergallo *et al.* 2006). Nevertheless, other reports show that the application of the phenol–chloroform extraction was not sufficient for complete removal of PCR inhibitors (Pachner and Delaney 1993).

Column chromatography using Sephacryl S-400, Sephadex G-200, Chelex, Sephadex or cetrimonium bromide can be used to remove salts and small proteins as well as polysaccharides from several sample matrices, for example, seminal fluid or stool (Da Silva *et al.* 1995; Schmidt *et al.* 1995; Hale *et al.* 1996; Croci *et al.* 2008). For example, cetrimonium bromide forms an insoluble complex with polysaccharides and denatured proteins and therefore efficiently removes them from the preparation (Alaeddini 2011). A repeated extraction using silica columns may also remove inhibitors (Kemp *et al.* 2006). In addition, cation exchange resins have been successfully used for the removal of PCR inhibitors (Jacobsen and Rasmussen 1992; Henson and French 1993). The use of magnetic silica beads for nucleic acid isolation has been repeatedly shown to efficiently remove a wide range of PCR inhibitors (Maher *et al.* 2001; Rutjes *et al.* 2005; Ngazoa *et al.* 2008; Sur *et al.* 2010).

Immunocapture methods are very efficient in removal of PCR inhibitors because the pathogen can be specifically separated from the sample matrix and the accordant inhibitors (Widjoatmodjo *et al.* 1992; Croci *et al.* 2008; Schrader *et al.* 2011). For example, antigen-capture PCR has been successfully used for the sensitive detection of hepatitis A virus in seafood (Arnal *et al.* 1999). However, because of the use of specific antibodies, this method is not applicable to highly variable pathogens, such as noroviruses (Atmar *et al.* 1995). In this case, human histo-blood group antigens or pig gastric mucin, which

Table 3 Strategies for the removal of PCR inhibitors from specific matrices

Matrix	Strategy for removal of inhibitors	References
Stool	Additional extraction steps Sedaphex G-200 chromatography Heat treatment before the PCR BSA or gp32 Selection of resistant polymerases Chloroform extraction Treatment with activated carbon Dilution of the sample	Al-Soud and Rådström (1998), Chaturvedi <i>et al.</i> (2008), Hale <i>et al.</i> (1996), Kreader (1996), Monteiro <i>et al.</i> (1997), Scipioni <i>et al.</i> (2008a,b) and Wilde <i>et al.</i> (1990)
Seafood, bivalves, oysters	PEG precipitation Cetyltrimethylammonium bromide treatment Proteinase K treatment Preparation of selected tissues (digestive gland) Activated carbon treatment	Abolmaaty <i>et al.</i> (2007), Atmar <i>et al.</i> (1993, 1995) and Jothikumar <i>et al.</i> (2005)
Plants	RNA precipitation Treatment with: Tween 20, DMSO, PEG 400, polyvinylpyrrolidone, β -mercaptoethanol, dithiothreitol dilution of nucleic acids cation exchanger Extraction with high concentration of borates Precipitation of polysaccharides proteinase K treatment Desiccation (65°C for 2 days)	Demeke and Adams (1992), John (1992), Henson and French (1993), Sipahioglu <i>et al.</i> (2006), Su and Gibor (1988), Wan and Wilkins (1994) and Wilkins and Smart (1996)
Berries	Chloroform/butanol extraction treatment with pectinases Use of conventional PCR instead of real-time PCR	Butot <i>et al.</i> (2007a), Dubois <i>et al.</i> (2002) and Love <i>et al.</i> (2008)
Cheese, meat	Selection of resistant polymerase	Al-Soud and Rådström (1998)
Cheese	Inactivation of proteases by hot NaOH extraction	Rossen <i>et al.</i> (1992)
Milk	Addition of BSA, protease inhibitors, magnesium ions Chelation of calcium ions	Bickley <i>et al.</i> (1996) and Powell <i>et al.</i> (1994)
Water, environmental samples	Combined treatment of Sedaphex G-100 and Chelex-100 Dialysis Extraction with solvents Ultrafiltration with positively charged membranes, UV irradiation Antigen-capture PCR	Abbaszadegan <i>et al.</i> (1993), Cannon and Vinjé (2008), Ijzerman <i>et al.</i> (1997) and Tamariz <i>et al.</i> (2006)
Airborne and environmental samples	Magnetic bead DNA capture method	Maher <i>et al.</i> (2001)

BSA, bovine serum albumin; PCR, polymerase chain reaction.

are proposed to be cellular receptors for noroviruses, have been successfully used for virus capturing and subsequent detection by PCR (Cannon and Vinjé 2008; Tian *et al.* 2008).

A more general and widely applied method is the dilution of the sample or the extracted nucleic acid, which will automatically result in a dilution of the PCR inhibitors (Widjoatmodjo *et al.* 1992; Monteiro *et al.* 1997; Eckhart *et al.* 2000; Scipioni *et al.* 2008a,b). However, the dilution clearly is accompanied by a decrease in sensitivity. An opponent strategy is the addition of substances to the PCR

mixture, which may include betaine, bovine serum albumin (BSA), dimethyl sulphoxide, formamide, glycerole, non-ionic detergents, polyethylene glycol, powdered milk, T4 bacteriophage gene 32 product (gp32) and proteinase inhibitors (Frackman *et al.* 1998; Al-Soud and Rådström 2000; Eckhart *et al.* 2000). Especially, BSA and gp32 are known to be effective against iron chloride, hemin, fulminic acid, humic acid, tannic acid, stool extracts and melanin (Al-Soud and Rådström 2000; Scipioni *et al.* 2008b; Opel *et al.* 2010). However, BSA is not effective against bile salts, bilirubin, EDTA, sodium chloride, SDS, Triton

X-100, calcium and collagen (Kreader 1996; Opel *et al.* 2010). Other described methods such as heating of the sample or DNase treatment were not always suitable for the removal of PCR inhibitors (Wilde *et al.* 1990).

The choice of the DNA polymerase and – if applicable – of the appropriate reverse transcription system is of great importance to prevent PCR inhibitory effects (Al-Soud and Rådström 1998; Löfström *et al.* 2004). Kermekchiev *et al.* (2009) showed that distinct mutations in the Taq DNA polymerase can overcome its inhibition by blood, plasma, haemoglobin, lactoferrin, serum IgG, soil extracts and humic acids. Baar *et al.* (2011) developed a chimeric polymerase consisting of parts from different polymerases of the genus *Thermus*, which shows a broad resistance against a variety of organic and inorganic inhibitors, as for example, humic acids, bone powder, fossilized excrements, coal tar or clay-rich earths. For real-time PCR using TaqMan probes, the 5' exonuclease activity of the Taq polymerase is essential. In an experiment with 15 different Taq polymerases, considerable differences in the intensity of the fluorescence in different sample types were obvious (Kreuzer *et al.* 2000).

Manufacturer of commercially available kits for nucleic acid purification and PCR use a variety of the above-mentioned strategies for removal of PCR inhibitors and increasing robustness of PCR enzymes. Several studies investigated the performance of such kits (e.g. Ribao *et al.* 2004; Levesque-Sergerie *et al.* 2007; Demeke and Jenkins 2010); however, the efficiency is strongly dependent on the used matrix and no general recommendations can be retrieved from those studies.

Methods for removal of specific PCR inhibitors

Several methods have been developed for removal of specific classes of inhibitors. Polysaccharides, which are mainly found in seafood or berries, may hamper the resuspension of precipitated nucleic acids (Atmar *et al.* 1993, 1995; Butot *et al.* 2007a). Either the precipitation of the polysaccharides before RNA isolation or application of an RNA isolation method without any polysaccharide contamination is used to avoid these negative effects (Fang *et al.* 1992; Wilkins and Smart 1996). This includes treatment with Tween 20, DMSO, polyethylene glycol or activated carbon (Demeke and Adams 1992; Abolmaaty *et al.* 2007). Pectinase treatment has been successfully used for the analysis of berries (Butot *et al.* 2007a).

Phenols can directly interact with RNA. Removal of phenols can be achieved by precipitation using polyvinylpyrrolidone (John 1992; Wilkins and Smart 1996). High concentrations of borates protect the RNA from interaction with polyphenols (Wan and Wilkins 1994; Wilkins

and Smart 1996). Sipahioglu *et al.* (2006) removed these inhibitors from leaves by drying at 65°C for 2 days and conserving them at 4°C under hermetic conditions.

Humic and fulminic acids, which are often present in dead biomass, soil and water samples may be removed by dialysis, liquid–liquid extraction, flocculation using polyvalent cations, gel extraction, column-based methods and ultrafiltration, the latter one being most successful (Tsai and Olson 1992; Abbaszadegan *et al.* 1993; Tsai *et al.* 1993; Ijzerman *et al.* 1997; Braid *et al.* 2003). Queiroz *et al.* (2001) used electropositive filters for the analysis of sewage and water samples, which preferentially bind microorganisms and avoid co-purification of the inhibitory substances.

For the specific removal of other substances, only a few methods have been published. Urea present in urine samples may be effectively removed by dialysis or ultrafiltration (Khan *et al.* 1991). Proteases, which might occur in milk, may be eliminated by the addition of protease inhibitors or BSA (Powell *et al.* 1994). The inhibitory effects caused by calcium ions may be compensated by the addition of magnesium ions. Another possibility is the use of different chelating agents, which catch the calcium ions, resulting in increased amplification rates (Bickley *et al.* 1996).

Assessing PCR inhibitions by control reactions

Despite extensive efforts to eliminate inhibitors from the sample, inhibitory substances which can affect PCR may still be present. Different analytical methods can be used to determine the presence of inorganic or organic compounds, for example, the amount of dissolved organic carbon indicative for humic acids (Chen *et al.* 2003; Rock *et al.* 2010). However, for most of the inhibitory substances, no practicable tools for their analysis are available.

To determine the inhibitory effect of all substances present in a nucleic acid preparation, it has been suggested to carry out PCR control reactions. This is especially necessary to exclude false-negative results, which may result from complete inhibition of the PCR, even in the presence of the target sequence (Parshionikar *et al.* 2004). Such controls generally consist of defined amounts of nucleic acids or microorganisms, which are added to the sample and analysed parallel to the target sequence. By comparison of the detected amount of the control with that originally added, the performance of the assay including the presence of PCR inhibitors can be assessed. Different types of controls can be distinguished according to the controlled steps (process control, amplification control) and the implementation of the control reaction (internal or external control).

A process control is added at the starting point of sample analysis, for example, before nucleic acid extraction, and passes therefore all preparation steps. In contrast, an amplification control is added to the nucleic acid extracted from the sample thus controlling only the performance of the PCR itself. An internal control is analysed in the same tube as the target, whereas the external control is analysed in a separate aliquot of the sample (Nolan *et al.* 2006).

The controls can be divided into competitive and non-competitive amplification controls. Competitive controls are amplified using the same primers and the same conditions but can be distinguished from the target by product length or sequence (Hoorfar *et al.* 2004; Villanova *et al.* 2007). One limiting factor of this application may be the concurrent use of primers, which may lead to a general decrease in PCR sensitivity (Hoorfar *et al.* 2004). In contrast, non-competitive controls composed of unrelated sequences are amplified by different primers as the target sequence and thus can also be used universally (Dingle *et al.* 2004; Hoorfar *et al.* 2004; Dreier *et al.* 2005; Villanova *et al.* 2007). A possible disadvantage of non-competitive controls is the use of separate primers, which may behave different than those used for the target sequence (Villanova *et al.* 2007). Generally, by using internal controls in multiplex PCR assays, the co-amplification of the control may inhibit the amplification of the target sequence, which might be solved by limiting the concentration of the control primers and control RNA (Hofmann 2003; Gall *et al.* 2007).

One special kind of non-competitive control is the endogenous control, which uses housekeeping genes that directly originate from the sample (Hoffmann *et al.* 2005). Endogenous controls act as process controls, which control the whole sample processing procedure. However, a high concentration of the endogenous nucleic acids may hamper the amplification of the target sequence (Hoffmann *et al.* 2005). Externally added process controls are used mostly when endogenous nucleic acids are not present due to the sample type (e.g. environmental samples) or sample preparation method (e.g. using virus purification). One possibility of such controls is the so-called armored RNA, which is packaged into phage proteins and thus resistant to RNases (Pasloske *et al.* 1998; Drosten *et al.* 2001; Hietala and Crossley 2006). Another strategy uses bacteriophages or animal viruses added to the sample. For example, bacteriophage MS2 and mengoviruses, feline calicivirus and enterovirus have been successfully used as process controls in food analysis (Dreier *et al.* 2005; Costafreda *et al.* 2006; Da Silva *et al.* 2007; Rolfe *et al.* 2007; Comelli *et al.* 2008; Lowther *et al.* 2008; Mattison *et al.* 2009; Blaise-Boisseau *et al.* 2010).

Conclusions

In conclusion, PCR inhibitors are a heterogeneous class of substances that act at different steps of the diagnostic procedure. They are present in a large variety of sample types and may lead to decreased PCR sensitivity or even false-negative PCR results. Several strategies have been developed to remove PCR inhibitors during sample preparation. However, as at least in complex matrices, it cannot be guaranteed that the preparations are free of PCR inhibitors, all reactions should be analysed for the presence of inhibitory effects. The development of standardized controls has therefore been recommended for substantiated evaluation of diagnostic PCR results and for comparison of the efficiencies of different PCR protocols (Hoorfar *et al.* 2003, 2004; Mattison *et al.* 2009; Lees and CEN WG6 TAG4 2010). On the basis of the results of such efficiency-controlled assays, it might be necessary to modify the steps of sample analysis to remove specific PCR inhibitors or to use a PCR system, which is less sensitive to the inhibitory substances. This might be carried out by the application of general methods for removal of PCR inhibitors, for example, by changing the commercially available kits for nucleic acid preparation and PCR analysis. However, additional steps for removal of specific inhibitors may be necessary. The overview presented here might provide a collection of published methods, which can be selected and tested based on the sample type. In other cases, the different mechanisms of action of PCR inhibitors described here might be a starting point for the development of new methods for their removal or inactivation.

Conflict of interest

There is no conflict of interests to declare.

References

- Abbaszadegan, M., Huber, M.S., Gerba, C.P. and Pepper, I.L. (1993) Detection of enteroviruses in groundwater with the polymerase chain reaction. *Appl Environ Microbiol* **59**, 1318–1324.
- Abbaszadegan, M., Stewart, P. and LeChevallier, M. (1999) A strategy for detection of viruses in groundwater by PCR. *Appl Environ Microbiol* **65**, 444–449.
- Abolmaaty, A., Gu, W., Witkowski, R. and Levin, R.E. (2007) The use of activated charcoal for the removal of PCR inhibitors from oyster samples. *J Microbiol Methods* **68**, 349–352.
- Alaeddini, R. (2011) Forensic implications of PCR inhibition – a review. *Forensic Sci Int Genet* **6**, 297–305.
- Al-Soud, W.A. and Rådström, P. (1998) Capacity of nine thermostable DNA polymerases To mediate DNA

- amplification in the presence of PCR-inhibiting samples. *Appl Environ Microbiol* **64**, 3748–3753.
- Al-Soud, W.A. and Rådström, P. (2000) Effects of amplification facilitators on diagnostic PCR in the presence of blood, feces, and meat. *J Clin Microbiol* **38**, 4463–4470.
- Al-Soud, W.A. and Rådström, P. (2001) Purification and characterization of PCR-inhibitory components in blood cells. *J Clin Microbiol* **39**, 485–493.
- Al-Soud, W.A., Jonsson, L.J. and Rådström, P. (2000) Identification and characterization of immunoglobulin G in blood as a major inhibitor of diagnostic PCR. *J Clin Microbiol* **38**, 345–350.
- Al-Soud, W.A., Ouis, I.S., Li, D.Q., Ljungh, S. and Wadström, T. (2005) Characterization of the PCR inhibitory effect of bile to optimize real-time PCR detection of *Helicobacter* species. *FEMS Immunol Med Microbiol* **44**, 177–182.
- Arnal, C., Ferre-Aubineau, V., Besse, B., Mignotte, B., Schwartzbrod, L. and Billaudel, S. (1999) Comparison of seven RNA extraction methods on stool and shellfish samples prior to hepatitis A virus amplification. *J Virol Methods* **77**, 17–26.
- Atmar, R.L., Metcalf, T.G., Neill, F.H. and Estes, M.K. (1993) Detection of enteric viruses in oysters by using the polymerase chain reaction. *Appl Environ Microbiol* **59**, 631–635.
- Atmar, R.L., Neill, F.H., Romalde, J.L., Le Guyader, F., Woodley, C.M., Metcalf, T.G. and Estes, M.K. (1995) Detection of Norwalk virus and hepatitis A virus in shellfish tissues with the PCR. *Appl Environ Microbiol* **61**, 3014–3018.
- Baar, C., d'Abbadie, M., Vaisman, A., Arana, M.E., Hofreiter, M., Woodgate, R., Kunkel, T.A. and Holliger, P. (2011) Molecular breeding of polymerases for resistance to environmental inhibitors. *Nucleic Acids Res* **39**, e51.
- Bélec, L., Authier, J., Eliezer-Vanerot, M.C., Piedouillet, C., Mohamed, A.S. and Gherardi, R.K. (1998) Myoglobin as a polymerase chain reaction (PCR) inhibitor: a limitation for PCR from skeletal muscle tissue avoided by the use of *Thermus thermophilus* polymerase. *Muscle Nerve* **21**, 1064–1067.
- Bergallo, M., Costa, C., Gribaudo, G., Tarallo, S., Baro, S., Negro, P.A. and Cavallo, R. (2006) Evaluation of six methods for extraction and purification of viral DNA from urine and serum samples. *New Microbiol* **29**, 111–119.
- Bickley, J., Short, J.K., McDowell, D.G. and Parkes, H.C. (1996) Polymerase chain reaction (PCR) detection of *Listeria monocytogenes* in diluted milk and reversal of PCR inhibition caused by calcium ions. *Lett Appl Microbiol* **22**, 153–158.
- Blaise-Boisseau, S., Hennechart-Collette, C., Guillier, L. and Perelle, S. (2010) Duplex real-time qRT-PCR for the detection of hepatitis A virus in water and raspberries using the MS2 bacteriophage as a process control. *J Virol Methods* **166**, 48–53.
- Braid, M.D., Daniels, L.M. and Kitts, C.L. (2003) Removal of PCR inhibitors from soil DNA by chemical flocculation. *J Microbiol Methods* **52**, 389–393.
- Burgess, L.C. and Hall, J.O. (1999) UV light irradiation of plastic reaction tubes inhibits PCR. *Biotechniques* **27**, 252, 254–254, 256.
- Burkardt, H.J. (2000) Standardization and quality control of PCR analyses. *Clin Chem Lab Med* **38**, 87–91.
- Butot, S., Putallaz, T. and Sanchez, G. (2007a) Procedure for rapid concentration and detection of enteric viruses from berries and vegetables. *Appl Environ Microbiol* **73**, 186–192.
- Butot, S., Putallaz, T., Croquet, C., Lamothe, G., Meyer, R., Joosten, H. and Sanchez, G. (2007b) Attachment of enteric viruses to bottles. *Appl Environ Microbiol* **73**, 5104–5110.
- Cannon, J.L. and Vinjé, J. (2008) Histo-blood group antigen assay for detecting noroviruses in water. *Appl Environ Microbiol* **74**, 6818–6819.
- Chandler, D.P., Wagnon, C.A. and Bolton, H. (1998) Reverse transcriptase (RT) inhibition of PCR at low concentrations of template and its implications for quantitative RT-PCR. *Appl Environ Microbiol* **64**, 669–677.
- Chaturvedi, U., Tiwari, A.K., Ratta, B., Ravindra, P.V., Rajawat, Y.S., Palia, S.K. and Rai, A. (2008) Detection of canine adenoviral infections in urine and faeces by the polymerase chain reaction. *J Virol Methods* **149**, 260–263.
- Chen, W., Westerhoff, P., Leenheer, J.A. and Booksh, K. (2003) Fluorescence excitation-emission matrix regional integration to quantify spectra for dissolved organic matter. *Environ Sci Technol* **37**, 5701–5710.
- Chen, P.S., Tsai, F.T., Lin, C.K., Yang, C.Y., Chan, C.C., Young, C.Y. and Lee, C.H. (2010) Ambient influenza and avian influenza virus during dust storm days and background days. *Environ Health Perspect* **118**, 1211–1216.
- Comelli, H.L., Rimstad, E., Larsen, S. and Myrmel, M. (2008) Detection of norovirus genotype I.3b and II.4 in bioaccumulated blue mussels using different virus recovery methods. *Int J Food Microbiol* **127**, 53–59.
- Costafreda, M.I., Bosch, A. and Pintó, R.M. (2006) Development, evaluation, and standardization of a real-time TaqMan reverse transcription-PCR assay for quantification of hepatitis A virus in clinical and shellfish samples. *Appl Environ Microbiol* **6**, 3846–3855.
- Croci, L., Dubois, E., Cook, N., de Medici, D., Schultz, A., China, B., Rutjes, S., Hoorfar, J. et al. (2008) Current methods for extraction and concentration of enteric viruses from fresh fruit and vegetables: towards international standards. *Food Anal Methods* **1**, 73–84.
- Da Silva, N., Zardoya, R., Santurde, G., Solana, A. and Castro, J.M. (1995) Rapid and sensitive detection of the bovine viral diarrhoea virus genome in semen. *J Virol Methods* **55**, 209–218.

- Da Silva, A.K., Le Saux, J.C., Parnaudeau, S., Pommepuy, M., Elimelech, M. and Le Guyader, F.S. (2007) Evaluation of removal of noroviruses during wastewater treatment, using real-time reverse transcription-PCR: different behaviors of genogroups I and II. *Appl Environ Microbiol* **73**, 7891–7897.
- Demeke, T. and Adams, R.P. (1992) The effects of plant polysaccharides and buffer additives on PCR. *Biotechniques* **12**, 332–334.
- Demeke, T. and Jenkins, G.R. (2010) Influence of DNA extraction methods, PCR inhibitors and quantification methods on real-time PCR assay of biotechnology-derived traits. *Anal Bioanal Chem* **396**, 1977–1990.
- Dingle, K.E., Crook, D. and Jeffery, K. (2004) Stable and noncompetitive RNA internal control for routine clinical diagnostic reverse transcription-PCR. *J Clin Microbiol* **42**, 1003–1011.
- Dreier, J., Stormer, M. and Kleesiek, K. (2005) Use of bacteriophage MS2 as an internal control in viral reverse transcription-PCR assays. *J Clin Microbiol* **43**, 4551–4557.
- Drosten, C., Seifried, E. and Roth, W.K. (2001) TaqMan 5'-nuclease human immunodeficiency virus type 1 PCR assay with phage-packaged competitive internal control for high-throughput blood donor screening. *J Clin Microbiol* **39**, 4302–4308.
- Dubois, E., Agier, C., Traore, O., Hennechart, C., Merle, G., Cruciere, C. and Laveran, H. (2002) Modified concentration method for the detection of enteric viruses on fruits and vegetables by reverse transcriptase-polymerase chain reaction or cell culture. *J Food Prot* **65**, 1962–1969.
- Eckhart, L., Bach, J., Ban, J. and Tschachler, E. (2000) Melanin binds reversibly to thermostable DNA polymerase and inhibits its activity. *Biochem Biophys Res Commun* **271**, 726–730.
- Enriquez, R., Frosner, G.G., Hochstein-Mintzel, V., Riedemann, S. and Reinhardt, G. (1992) Accumulation and persistence of hepatitis A virus in mussels. *J Med Virol* **37**, 174–179.
- Fang, G., Hammar, S. and Grumet, R. (1992) A quick and inexpensive method for removing polysaccharides from plant genomic DNA. *Biotechniques* **13**, 52–54, 56.
- Fox, D.H., Huang, C.K., Du, J., Chang, T.Y. and Pan, Q. (2007) Profound inhibition of the PCR step of CF V3 multiplex PCR/OLA assay by the use of UV-irradiated plastic reaction tubes. *Diagn Mol Pathol* **16**, 121–123.
- Frackman, S., Kobs, G., Simpson, D. and Storts, D. (1998) Beatine and DMSO: enhancing agents for PCR. *Promega Notes* **65**, 27–30.
- Gall, A., Hoffmann, B., Teifke, J.P., Lange, B. and Schirrmeier, H. (2007) Persistence of viral RNA in rabbits which overcome an experimental RHDV infection detected by a highly sensitive multiplex real-time RT-PCR. *Vet Microbiol* **120**, 17–32.
- Gassilloud, B., Huguet, L., Maul, A. and Gantzer, C. (2007) Development of a viral concentration method for bottled water stored in hydrophobic support. *J Virol Methods* **142**, 98–104.
- Gonzalez, A., Grimes, R., Walsh, E.J., Dalton, T. and Davies, M. (2007) Interaction of quantitative PCR components with polymeric surfaces. *Biomed Microdevices* **9**, 261–266.
- Hale, A.D., Green, J. and Brown, D.W. (1996) Comparison of four RNA extraction methods for the detection of small round structured viruses in faecal specimens. *J Virol Methods* **57**, 195–201.
- Henson, J.M. and French, R. (1993) The polymerase chain reaction and plant disease diagnosis. *Annu Rev Phytopathol* **31**, 81–109.
- Hietala, S.K. and Crossley, B.M. (2006) Armored RNA as virus surrogate in a real-time reverse transcriptase PCR assay proficiency panel. *J Clin Microbiol* **44**, 67–70.
- Hoffmann, B., Beer, M., Schelp, C., Schirrmeier, H. and Depner, K. (2005) Validation of a real-time RT-PCR assay for sensitive and specific detection of classical swine fever. *J Virol Methods* **130**, 36–44.
- Hofmann, M.A. (2003) Construction of an infectious chimeric classical swine fever virus containing the 5'UTR of bovine viral diarrhoea virus, and its application as a universal internal positive control in real-time RT-PCR. *J Virol Methods* **114**, 77–90.
- Hoorfar, J., Cook, N., Malorny, B., Wagner, M., De Medici, D., Abdulmawjood, A. and Fach, P. (2003) Diagnostic PCR: making internal amplification control mandatory. *Lett Appl Microbiol* **38**, 79–80.
- Hoorfar, J., Malorny, B., Abdulmawjood, A., Cook, N., Wagner, M. and Fach, P. (2004) Practical considerations in design of internal amplification controls for diagnostic PCR assays. *J Clin Microbiol* **42**, 1863–1868.
- Huggett, J.F., Novak, T., Garson, J.A., Green, C., Morris-Jones, S.D., Miller, R.F. and Zumla, A. (2008) Differential susceptibility of PCR reactions to inhibitors: an important and unrecognised phenomenon. *BMC Res Notes* **1**, 70.
- Huppertz, H.I., Schmidt, H. and Karch, H. (1993) Detection of *Borrelia burgdorferi* by nested polymerase chain reaction in cerebrospinal fluid and urine of children with neuroborreliosis. *Eur J Pediatr* **152**, 414–417.
- Ijzerman, M.M., Dahling, D.R. and Fout, G.S. (1997) A method to remove environmental inhibitors prior to the detection of waterborne enteric viruses by reverse transcription-polymerase chain reaction. *J Virol Methods* **63**, 145–153.
- Jacobsen, C.S. and Rasmussen, O.F. (1992) Development and application of a new method to extract bacterial DNA from soil based on separation of bacteria from soil with cation-exchange resin. *Appl Environ Microbiol* **58**, 2458–2462.
- Jiang, J., Alderisio, K.A., Singh, A. and Xiao, L. (2005) Development of procedures for direct extraction of

- Cryptosporidium* DNA from water concentrates and for relief of PCR inhibitors. *Appl Environ Microbiol* **71**, 1135–1141.
- John, M.E. (1992) An efficient method for isolation of RNA and DNA from plants containing polyphenolics. *Nucleic Acids Res* **20**, 2381.
- Jothikumar, N., Lowther, J.A., Henshilwood, K., Lees, D.N., Hill, V.R. and Vinje, J. (2005) Rapid and sensitive detection of noroviruses by using TaqMan-based one-step reverse transcription-PCR assays and application to naturally contaminated shellfish samples. *Appl Environ Microbiol* **71**, 1870–1875.
- Katcher, H.L. and Schwartz, I. (1994) A distinctive property of Tth DNA polymerase: enzymatic amplification in the presence of phenol. *Biotechniques* **16**, 84–92.
- Kemp, B.M., Monroe, C. and Smith, D.G. (2006) Repeat silica extraction: a simple technique for the removal of PCR inhibitors from DNA extracts. *J Archaeol Sci* **33**, 1680–1689.
- Kermekchiev, M.B., Kirilova, L.I., Vail, E.E. and Barnes, W.M. (2009) Mutants of Taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. *Nucleic Acids Res* **37**, e40.
- Khan, G., Kangro, H.O., Coates, P.J. and Heath, R.B. (1991) Inhibitory effects of urine on the polymerase chain reaction for cytomegalovirus DNA. *J Clin Pathol* **44**, 360–365.
- Konet, D.S., Mezencio, J.M., Babcock, G. and Brown, F. (2000) Inhibitors of RT-PCR in serum. *J Virol Methods* **84**, 95–98.
- Kreider, C.A. (1996) Relief of amplification inhibition in PCR with bovine serum albumin or T4 gene 32 protein. *Appl Environ Microbiol* **62**, 1102–1106.
- Kreuzer, K.A., Bohm, A., Lass, U., Peters, U.R. and Schmidt, C.A. (2000) Influence of DNA polymerase on quantitative PCR results using TaqManTM probe format in the LightCyclerTM instrument. *Mol Cell Probes* **14**, 57–60.
- Lantz, P.-G., Matsson, M., Wadström, T. and Rådström, P. (1997) Removal of PCR inhibitors from human faecal samples through the use of an aqueous two-phase system for sample preparation prior to PCR. *J Microbiol Methods* **28**, 159–167.
- Lees, D. and CEN WG6 TAG4 (2010) International standardisation of a method for detection of human pathogenic viruses in Molluscan shellfish. *Food Environ Virol* **2**, 146–155.
- Levesque-Sergerie, J.P., Duquette, M., Thibault, C., Delbecchi, L. and Bissonnette, N. (2007) Detection limits of several commercial reverse transcriptase enzymes: impact on the low- and high-abundance transcript levels assessed by quantitative RT-PCR. *BMC Mol Biol* **8**, 93.
- Löfström, C., Knutsson, R., Axelsson, C.E. and Radström, P. (2004) Rapid and specific detection of *Salmonella* spp. in animal feed samples by PCR after culture enrichment. *Appl Environ Microbiol* **70**, 69–75.
- Love, D.C., Casteel, M.J., Meschke, J.S. and Sobsey, M.D. (2008) Methods for recovery of hepatitis A virus (HAV) and other viruses from processed foods and detection of HAV by nested RT-PCR and TaqMan RT-PCR. *Int J Food Microbiol* **126**, 221–226.
- Lowther, J.A., Henshilwood, K. and Lees, D.N. (2008) Determination of norovirus contamination in oysters from two commercial harvesting areas over an extended period, using semiquantitative real-time reverse transcription PCR. *J Food Prot* **71**, 1427–1433.
- Maher, N., Dillon, H.K., Vermund, S.H. and Unnasch, T.R. (2001) Magnetic bead capture eliminates PCR inhibitors in samples collected from the airborne environment, permitting detection of *Pneumocystis carinii* DNA. *Appl Environ Microbiol* **67**, 449–452.
- Mattison, K., Brassard, J., Gagne, M.J., Ward, P., Houde, A., Lessard, L., Simard, C., Shukla, A. et al. (2009) The feline calicivirus as a sample process control for the detection of food and waterborne RNA viruses. *Int J Food Microbiol* **132**, 73–77.
- Monteiro, L., Bonnemaïson, D., Vekris, A., Petry, K.G., Bonnet, J., Vidal, R., Cabrita, J. and Megraud, F. (1997) Complex polysaccharides as PCR inhibitors in feces: *Helicobacter pylori* model. *J Clin Microbiol* **35**, 995–998.
- Ngazoa, E.S., Fliss, I. and Jean, J. (2008) Quantitative study of persistence of human norovirus genome in water using TaqMan real-time RT-PCR. *J Appl Microbiol* **104**, 707–715.
- Nolan, T., Hands, R.E., Ogunkolade, W. and Bustin, S.A. (2006) SPUD: a quantitative PCR assay for the detection of inhibitors in nucleic acid preparations. *Anal Biochem* **351**, 308–310.
- Oikarinen, S., Tauriainen, S., Viskari, H., Simell, O., Knip, M., Virtanen, S. and Hyöty, H. (2009) PCR inhibition in stool samples in relation to age of infants. *J Clin Virol* **44**, 211–214.
- Opel, K.L., Chung, D. and McCord, B.R. (2010) A study of PCR inhibition mechanisms using real time PCR. *J Forensic Sci* **55**, 25–33.
- Oppliger, A., Masclaux, F.G. and Niculita-Hirzel, H. (2011) Assessment of airborne microorganisms by real-time PCR: optimistic findings and research challenges. *Front Biosci (Schol Ed)* **3**, 445–453.
- Pachner, A.R. and Delaney, E. (1993) The polymerase chain reaction in the diagnosis of Lyme neuroborreliosis. *Ann Neurol* **34**, 544–550.
- Parshionkar, S.U., Cashdollar, J. and Fout, G.S. (2004) Development of homologous viral internal controls for use in RT-PCR assays of waterborne enteric viruses. *J Virol Methods* **121**, 39–48.
- Pasloske, B.L., WalkerPeach, C.R., Obermoeller, R.D., Winkler, M. and DuBois, D.B. (1998) Armored RNA technology for production of ribonuclease-resistant viral RNA controls and standards. *J Clin Microbiol* **36**, 3590–3594.

- Peist, R., Honsel, D., Twieling, G. and Löffert, D. (2001) PCR inhibitors in plant DNA preparations. *QIAGEN News* **3**, 7–9.
- Pontirol, A., Travis, E.R., Sweeney, F.P., Porter, D., Gaze, W.H., Mason, S., Hibberd, V., Holden, J. et al. (2011) Pathogen quantitation in complex matrices: a multi-operator comparison of DNA extraction methods with a novel assessment of PCR inhibition. *PLoS ONE* **6**, e17916.
- Powell, H.A., Gooding, C.M., Garrett, S.D., Lund, B.M. and McKee, R.A. (1994) Protease inhibition of the detection of *Listeria monocytogenes* in milk using the polymerase chain reaction. *Lett Appl Microbiol* **18**, 59–61.
- Queiroz, A.P., Santos, F.M., Sassaroli, A., Harsi, C.M., Monezi, T.A. and Mehnert, D.U. (2001) Electropositive filter membrane as an alternative for the elimination of PCR inhibitors from sewage and water samples. *Appl Environ Microbiol* **67**, 4614–4618.
- Rådström, P., Knutsson, R., Wolffs, P., Lovenklev, M. and Löfström, C. (2004) Pre-PCR processing: strategies to generate PCR-compatible samples. *Mol Biotechnol* **26**, 133–146.
- Ribao, C., Torrado, I., Vilarino, M.L. and Romalde, J.L. (2004) Assessment of different commercial RNA-extraction and RT-PCR kits for detection of hepatitis A virus in mussel tissues. *J Virol Methods* **115**, 177–182.
- Richards, G.P. (1999) Limitations of molecular biological techniques for assessing the virological safety of foods. *J Food Prot* **62**, 691–697.
- Rock, C., Alum, A. and Abbaszadegan, M. (2010) PCR inhibitor levels in concentrates of biosolid samples predicted by a new method based on excitation-emission matrix spectroscopy. *Appl Environ Microbiol* **76**, 8102–8109.
- Rolfe, K.J., Parmar, S., Mururi, D., Wreghitt, T.G., Jalal, H., Zhang, H. and Curran, M.D. (2007) An internally controlled, one-step, real-time RT-PCR assay for norovirus detection and genotyping. *J Clin Virol* **39**, 318–321.
- Rossen, L., Nørskov, P., Holmström, K. and Rasmussen, O.F. (1992) Inhibition of PCR by components of food samples, microbial diagnostic assays and DNA-extraction solutions. *Int J Food Microbiol* **17**, 37–45.
- Rutjes, S.A., Italiaander, R., van den Berg, H.H., Lodder, W.J. and de Roda Husman, A.M. (2005) Isolation and detection of enterovirus RNA from large-volume water samples by using the NucliSens miniMAG system and real-time nucleic acid sequence-based amplification. *Appl Environ Microbiol* **71**, 3734–3740.
- Saulnier, P. and Andreumont, A. (1992) Detection of genes in feces by booster polymerase chain reaction. *J Clin Microbiol* **30**, 2080–2083.
- Schmidt, B.L., Aberer, E., Stockenhuber, C., Klade, H., Breier, F. and Luger, A. (1995) Detection of *Borrelia burgdorferi* DNA by polymerase chain reaction in the urine and breast milk of patients with Lyme borreliosis. *Diagn Microbiol Infect Dis* **21**, 121–128.
- Schrader, C., Johne, R., Schielke, A. and Ellerbroek, L. (2011) Food associated viruses and their detection – a review. *J Food Saf Food Qual* **62**, 36–51.
- Scipioni, A., Mauroy, A., Ziant, D., Saegerman, C. and Thiry, E. (2008a) A SYBR Green RT-PCR assay in single tube to detect human and bovine noroviruses and control for inhibition. *Virol J* **5**, 94.
- Scipioni, A., Bourgot, I., Mauroy, A., Ziant, D., Saegerman, C., Daube, G. and Thiry, E. (2008b) Detection and quantification of human and bovine noroviruses by a TaqMan RT-PCR assay with a control for inhibition. *Mol Cell Probes* **22**, 215–222.
- Seeram, N.P., Adams, L.S., Zhang, Y., Lee, R., Sand, D., Scheuller, H.S. and Heber, D. (2006) Blackberry, black raspberry, blueberry, cranberry, red raspberry, and strawberry extracts inhibit growth and stimulate apoptosis of human cancer cells in vitro. *J Agric Food Chem* **54**, 9329–9339.
- Shieh, Y.S., Wait, D., Tai, L. and Sobsey, M.D. (1995) Methods to remove inhibitors in sewage and other fecal wastes for enterovirus detection by the polymerase chain reaction. *J Virol Methods* **54**, 51–66.
- Sipahioglu, H.M., Usta, M. and Ocak, M. (2006) Use of dried high-phenolic laden host leaves for virus and viroid preservation and detection by PCR methods. *J Virol Methods* **137**, 120–124.
- Su, X. and Gibor, A. (1988) A method for RNA isolation from marine macro-algae. *Anal Biochem* **174**, 650–657.
- Sur, K., McFall, S.M., Yeh, E.T., Jangam, S.R., Hayden, M. A., Stroupe, S.D. and Kelso, D.M. (2010) Immiscible phase nucleic acid purification eliminates PCR inhibitors with a single pass of paramagnetic particles through a hydrophobic liquid. *J Mol Diagn* **12**, 620–628.
- Sutlović, D., Definis, G.M., Andelinovic, S., Gugic, D. and Primorac, D. (2005) Taq polymerase reverses inhibition of quantitative real time polymerase chain reaction by humic acid. *Croat Med J* **46**, 556–562.
- Sutlović, D., Gamulin, S., Definis-Gojanovic, M., Gugic, D. and Andjelinovic, S. (2008) Interaction of humic acids with human DNA: proposed mechanisms and kinetics. *Electrophoresis* **29**, 1467–1472.
- Tamariz, J., Voynarovska, K., Prinz, M. and Caragine, T. (2006) The application of ultraviolet irradiation to exogenous sources of DNA in plasticware and water for the amplification of low copy number DNA. *J Forensic Sci* **51**, 790–794.
- Tian, P., Engelbrektson, A. and Mandrell, R. (2008) Two-log increase in sensitivity for detection of norovirus in complex samples by concentration with porcine gastric mucin conjugated to magnetic beads. *Appl Environ Microbiol* **74**, 4271–4276.
- Tsai, Y.L. and Olson, B.H. (1992) Rapid method for separation of bacterial DNA from humic substances in sediments for polymerase chain reaction. *Appl Environ Microbiol* **58**, 2292–2295.

- Tsai, Y.L., Sobsey, M.D., Sangermano, L.R. and Palmer, C.J. (1993) Simple method of concentrating enteroviruses and hepatitis A virus from sewage and ocean water for rapid detection by reverse transcriptase-polymerase chain reaction. *Appl Environ Microbiol* **59**, 3488–3491.
- Villanova, G.V., Gardiol, D., Taborda, M.A., Reggiardo, V., Tanno, H., Rivadeneira, E.D., Perez, G.R. and Giri, A.A. (2007) Strategic approach to produce low-cost, efficient, and stable competitive internal controls for detection of RNA viruses by use of reverse transcription-PCR. *J Clin Microbiol* **45**, 3555–3563.
- Wadowsky, R.M., Laus, S., Libert, T., States, S.J. and Ehrlich, G.D. (1994) Inhibition of PCR-based assay for *Bordetella pertussis* by using calcium alginate fiber and aluminum shaft components of a nasopharyngeal swab. *J Clin Microbiol* **32**, 1054–1057.
- Wan, C.Y. and Wilkins, T.A. (1994) A modified hot borate method significantly enhances the yield of high-quality RNA from cotton (*Gossypium hirsutum* L.). *Anal Biochem* **223**, 7–12.
- Wei, T., Lu, G. and Clover, G. (2008) Novel approaches to mitigate primer interaction and eliminate inhibitors in multiplex PCR, demonstrated using an assay for detection of three strawberry viruses. *J Virol Methods* **151**, 132–139.
- Weyant, R.S., Edmonds, P. and Swaminathan, B. (1990) Effect of ionic and nonionic detergents on the *Taq* polymerase. *Biotechniques* **9**, 308–309.
- Widjoatmodjo, M.N., Fluit, A.C., Torensma, R., Verdonk, G. P. and Verhoef, J. (1992) The magnetic immuno polymerase chain reaction assay for direct detection of salmonellae in fecal samples. *J Clin Microbiol* **30**, 3195–3199.
- Wiedbrauk, D.L., Werner, J.C. and Drevon, A.M. (1995) Inhibition of PCR by aqueous and vitreous fluids. *J Clin Microbiol* **33**, 2643–2646.
- Wilde, J., Eiden, J. and Yolken, R. (1990) Removal of inhibitory substances from human fecal specimens for detection of group A rotaviruses by reverse transcriptase and polymerase chain reactions. *J Clin Microbiol* **28**, 1300–1307.
- Wilkins, T.A. and Smart, L.B. (1996) Isolation of RNA from plant tissue. In *A Laboratory Guide to RNA: Isolation, Analysis, and Synthesis* ed. Krieg, P.A. pp. 21–42. New York: Wiley-Liss, Inc.
- Wilson, I.G. (1997) Inhibition and facilitation of nucleic acid amplification. *Appl Environ Microbiol* **63**, 3741–3751.
- Yedidag, E.N., Koffron, A.J., Mueller, K.H., Kaplan, B., Kaufman, D.B., Fryer, J.P., Stuart, F.P. and Abecassis, M. (1996) Acyclovir triphosphate inhibits the diagnostic polymerase chain reaction for cytomegalovirus. *Transplantation* **62**, 238–242.