

(6 problems, 80 minutes, 100 points)

1. (30 points) Evaluate the following definite and indefinite integrals. Please show your work clearly, and simplify when possible.

(a) $\int_{-1}^2 \frac{x^5 dx}{\sqrt{x^3 + 17}};$

(b) $\int \frac{e^{\sin(\pi x)} \cos(\pi x) dx}{1 + e^{2 \sin(\pi x)}}.$

① Let $u = x^3 + 17$ $du = 3x^2 dx$

Change bounds: $u = (-1)^3 + 17 = 16$
 $u = (2^3) + 17 = 25$

$\int_{16}^{25} \frac{1}{3} \frac{x^3}{\sqrt{u}} du = \int_{16}^{25} \frac{1}{3} \frac{u-17}{\sqrt{u}} du$

$= \int_{16}^{25} \frac{1}{3} (u^{1/2} - 17u^{-1/2}) du$

$= \frac{2}{9} u^{3/2} - \frac{34}{3} u^{1/2} \Big|_{16}^{25}$

$= \frac{2}{9} 5^3 - \frac{34}{3} (5) - \left(\frac{2}{9} 4^3 - \frac{34}{3} (4) \right)$

$= \boxed{\frac{20}{9}}$

② $u = \sin(\pi x)$
 $du = \cos(\pi x) \cdot \pi dx$

$= \int \frac{1}{\pi} \frac{e^u}{1 + e^{2u}} du$

$v = e^u$

$dv = e^u du$

$= \int \frac{1}{\pi} \frac{1}{1+v^2} dv$

$= \frac{1}{\pi} \arctan(v) + C$

$= \boxed{\frac{1}{\pi} \arctan(e^{\sin(\pi x)}) + C}$

(CONTINUE TO NEXT PAGE)

2. (14 points) In this problem use the trapezoid rule (which is the same as the average of the left and right Riemann sums). Suppose you have a table of velocities in m/sec of a fast bicycle A and a slower one B. At time $t = 0$, A passes B. The table has the velocities v_0^A, \dots, v_4^A of A at 15-sec intervals for a minute, and similarly for B. Using the trapezoid rule, write an expression in terms of the v -values for the distance in meters that A is ahead of B after the minute is over.

t	$v_A(t)$	$v_B(t)$
0	v_0^A	v_0^B
15	v_1^A	v_1^B
30	v_2^A	v_2^B
45	v_3^A	v_3^B
60	v_4^A	v_4^B

The distance A over B = initial distance + $\int_0^{60} v_A(t) - v_B(t) dt$
 bicycle A is ahead of bicycle B
 (This is 0 in the problem)

Right Riemann Sum


Trapezoid = $\frac{1}{2} \left[15 \left[(v_1^A - v_1^B) + (v_2^A - v_2^B) + (v_3^A - v_3^B) + (v_4^A - v_4^B) \right] \right.$
 $\left. + 15 \left[(v_0^A - v_0^B) + (v_1^A - v_1^B) + (v_2^A - v_2^B) + (v_3^A - v_3^B) \right] \right]$
 left Riemann Sum

(CONTINUE TO NEXT PAGE)

3. (10 points) The ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where $a > b$, has major axis $2a$ ("longest diameter"), has minor axis $2b$ ("shortest diameter"), and has area πab . Suppose that the ellipse

$$\frac{x^2}{49} + \frac{y^2}{9} = 1$$

is being covered by a big rectangle whose lower horizontal side is being pulled downward at 0.3 units/sec. At the instant when the lower side of the rectangle crosses the x -axis, what is the rate at which the area of the ellipse is being covered?



$$\int_{3-0.3t}^3 2\sqrt{49(1-y^2/9)} dy$$

$$\Rightarrow - \int_3^{3-0.3t} 2\sqrt{49(1-y^2/9)} dy$$

By the FTC,

$$\frac{d}{dt} \int_3^{3-0.3t} 2\sqrt{49(1-y^2/9)} dy = -2\sqrt{49(1-\frac{(3-0.3t)^2}{9})} \cdot (-0.3)$$

The instant when the rectangle hits $y=0$, is precisely at

$$3-0.3t=0 \Rightarrow 3=\frac{3}{10}t \Rightarrow t=10$$

\therefore The rate at which the area of the ellipse is being covered is

$$-2\sqrt{49(1-\frac{(3-0.3(10))^2}{9})}(-0.3) = 0.6\sqrt{49} = \boxed{4.2 \frac{m^2}{sec}}$$

(CONTINUE TO NEXT PAGE)



4. (24 points) In the xy -plane draw the circle $x^2 + y^2 = 100$ and the line segment joining the points $(0, 10)$ and $(6, 8)$. Let R be the region in the circle that's above the line segment. Write each of the following in terms of definite integrals. Do **NOT** evaluate the integrals.

- (a) The volume using the washer method when R is revolved around the x -axis.
 (b) The volume using the washer method when R is revolved around the y -axis.
 (c) The volume using the shell method when R is revolved around the line $x = -2$.

Note


$$6^2 + 8^2 = 100$$

so the pt $(6, 8)$ lies on the circle




$$\begin{aligned} \text{Outer radius} &= \sqrt{100 - y^2} \\ \text{Inner radius} &= 30 - 3y \end{aligned}$$

$$\text{Volume} = \pi \int_8^{10} (\sqrt{100 - y^2})^2 - (30 - 3y)^2 dy$$



$$\begin{aligned} \text{Outer radius} &= \sqrt{100 - x^2} \\ \text{Inner radius} &= -\frac{1}{3}x + 10 \end{aligned}$$

$$\text{Volume} = \pi \int_0^6 (\sqrt{100 - x^2})^2 - (-\frac{1}{3}x + 10)^2 dx$$




$$\begin{aligned} \text{Height} &= \sqrt{100 - x^2} - (-\frac{1}{3}x + 10) \\ \text{Radius} &= x + 2 \end{aligned}$$

(CONTINUE TO NEXT PAGE)

$$\text{Volume} = 2\pi \int_0^6 (x+2) [\sqrt{100 - x^2} - (-\frac{1}{3}x + 10)] dx$$

5. (12 points) A solid is formed as follows. Its base is the region in the xy -plane bounded by the curve $y = x^{-3/2}$, the x -axis, the line $x = 1$, and the line $x = 3$. Its cross-section by a plane perpendicular to the x -axis at x is a square resting on the xy -plane whose side is the line joining $(x, 0)$ to $(x, x^{-3/2})$. Find the volume of the solid. Please show your work clearly.



$$\begin{aligned}
 \text{Volume} &= \int_1^3 \text{Area of Square} \, dx = \int_1^3 (x^{-3/2})^2 \, dx \\
 &= \int_1^3 x^{-3} \, dx \\
 &= \left. \frac{-1}{2} x^{-2} \right|_1^3 = \frac{-1}{2} \left[\frac{1}{9} - 1 \right] \\
 &= \frac{8}{18} = \boxed{\frac{4}{9}}
 \end{aligned}$$

enterline(CONTINUE TO NEXT PAGE)

6. (10 points) During a certain journey of 200 miles the fuel efficiency of your car at a given point depended on several factors — road conditions, weather, traffic, etc. Let $f(x)$ denote the fuel efficiency, measured in miles per gallon, at a distance x from the start of the journey. Let

$$I = \int_0^{200} \frac{dx}{f(x)}.$$

What is the practical meaning of I , and what are the units of I ?

$$\frac{1}{f(x)} \text{ has units } \frac{1}{\frac{\text{miles}}{\text{gal}}} = \frac{\text{gal}}{\text{miles}}$$

Since $dx = \text{miles}$, then

$$I = \frac{\text{gal.}}{\text{miles}} \cdot \text{miles} = \text{gallons}$$

The practical meaning of I is how much fuel it took you to drive 200 miles.