

Attention-Deficit Hyperactivity Disorder (ADHD): Does New Research Support Old Concepts?

Lydia Mary Furman

J Child Neurol 2008 23: 775

DOI: 10.1177/0883073808318059

The online version of this article can be found at:

<http://jcn.sagepub.com/content/23/7/775>

Published by:



<http://www.sagepublications.com>

Additional services and information for *Journal of Child Neurology* can be found at:

Email Alerts: <http://jcn.sagepub.com/cgi/alerts>

Subscriptions: <http://jcn.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

Citations: <http://jcn.sagepub.com/content/23/7/775.refs.html>

>> [Version of Record](#) - Jul 24, 2008

[What is This?](#)

Attention-Deficit Hyperactivity Disorder (ADHD): Does New Research Support Old Concepts?

Lydia Mary Furman, MD

Objective: To examine the evidence for and against the classification of attention-deficit hyperactivity disorder (ADHD) as a valid disease entity, as defined by the *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition (DSM-IV), criteria. **Data Sources:** Sources included but were not limited to published literature on ADHD accessed via PubMed (<http://www.ncbi.nlm.nih.gov/PubMed/>). **Study Selection:** Peer-reviewed research, review articles, consensus statements, "white papers," and proceedings of professional meetings were used. **Data Extraction:** Focused on evidence base and scientific validity of conclusions. **Data Synthesis:** Evidence for a genetic or

neuroanatomic cause of ADHD is insufficient. Experimental work shows that executive function deficits do not explain ADHD. The psychometric properties of widely used ADHD rating scales do not meet standards expected for disease identification. **Conclusions:** ADHD is unlikely to exist as an identifiable disease. Inattention, hyperactivity, and impulsivity are symptoms of many underlying treatable medical, emotional, and psychosocial conditions affecting children.

Keywords: ADHD; attention-deficit hyperactivity disorder; stimulant

Attention-deficit hyperactivity disorder (ADHD) is now practically a household word. Estimates suggest that anywhere from 2.4 to 4.4 million children aged 4 to 17 years in the United States may meet published criteria for ADHD, yet this disease has no established biological cause or objective diagnostic test.¹⁻³ Critics have challenged the validity of the diagnosis.⁴⁻⁸ Roughly 1 in 25 children in the United States has been prescribed stimulant medications, with a steadily increasing price tag; in 2003, \$2.4 billion dollars were spent on ADHD drugs in the United States.⁹ The crux of this problem has been highlighted by publication of the 3-year outcomes of the National Institutes of Mental Health sponsored Multimodal Treatment Study of Children with Attention Deficit Hyperactivity Disorder, which enrolled 579 ADHD-diagnosed children.¹⁰⁻¹⁵ Although neither blinded nor placebo controlled, the study reported at 36 months that all treatment groups *with and without stimulant medication* were improved from baseline and did not differ

significantly on any measure of outcome, and that children receiving (as compared with not receiving) stimulant medications showed significant symptom deterioration from 24 to 36 months and higher delinquency ratings at 24 and 36 months.¹⁰⁻¹⁵ These unexpected outcomes are not easily explained by the orthodox paradigm of ADHD as a neurobehavioral disease best treated with stimulants. The results give pause to thoughtful clinicians and suggest the need for more rigorous and transparent diagnostic and therapeutic methods.

Evidence for an Etiology of ADHD

A *genetic cause* for ADHD is postulated. A prominent ADHD researcher states, "Family studies have identified a 2- to 8-fold increase in the risk for ADHD in parents and siblings of children with ADHD. Various twin and adoption studies have also highlighted the highly genetic nature of ADHD."¹⁶ Jay Joseph has elegantly unraveled the evidence behind the rhetoric and points out that not only does "familial" not distinguish between "nature" (genes) and "nurture" (environmental factors), but the logic behind family, twin, and adoption studies is fundamentally flawed.¹⁷⁻²⁰ Most ADHD reviews confuse "familial" with "genetic," and trivialize the potential role of nongenetic factors with short lists such as "food additives/diet, lead contamination, cigarette and alcohol exposure, maternal smoking during pregnancy, and low birth

From the Division of General Academic Pediatrics, Rainbow Babies and Children's Hospital, Cleveland, Ohio.

The author reports no financial disclosure or conflict of interest.

Address correspondence to: Lydia Mary Furman, MD, Division of General Academic Pediatrics, Room 784, MS 6019, Rainbow Babies and Children's Hospital, 11100 Euclid Avenue, Cleveland, OH 44106; e-mail: Lydia.Furman@uhhospitals.org.

Furman LM. Attention-deficit hyperactivity disorder (ADHD): does new research support old concepts? *J Child Neurol*. 2008;23:775-784.

weight.”¹⁶ In fact, nongenetic familial factors are powerful and difficult to measure and include the effects of culture, religion, learning, parenting, and socioeconomic adversity.

Twin studies that purport to demonstrate a genetic cause for ADHD are scientifically untenable. Studies are based on the “equal environment assumption,” which assumes that, contrary to evidence, (a) the only difference between monozygotic and dizygotic twins is greater genetic similarity between the former than the latter and (b) monozygotic and dizygotic twin pairs share the same behavior-influencing environment. An extended literature documents that monozygotic as compared with dizygotic twins do not experience equal environments; rather, monozygotic twins share greater psychological and physical closeness, are treated more similarly by their parents, and more report “extreme closeness” (65% vs 19%) and being “inseparable as children” (73% vs 19%) than dizygotic twins.^{21,22} No studies report on ADHD-diagnosed twins who are reared apart. Greater concordance between monozygotic than dizygotic twins for the diagnosis of ADHD does not distinguish between the roles of genes and family factors or lead to the conclusion that ADHD is genetic.

Methodologic problems with adoption studies have been elucidated, again by Joseph.¹⁷⁻²⁰ Researchers propose that the effects of genes and environment can be differentiated by comparing children with (and without) ADHD who are living with their own parents versus with adoptive parents. However, it is well appreciated that parents seeking to adopt are screened for mental health problems whereas biologic parents are not, and most children in adoptive care have endured traumatic experiences or emotional upset. Thus, neither adoptive and biological parents nor their children are comparable with regard to nongenetic factors. No adoption study uses blinded diagnosticians to identify children with ADHD, and no study has assessed the biological relatives of adoptees. These scientific limitations preclude meaningful contribution to an understanding of ADHD.

Psychiatric geneticists initially embraced genome scans and candidate gene studies as ways to identify a genetic cause of ADHD. It is clear that genome-wide scans have yielded negative results.²³ Candidate molecular gene research has focused primarily on neurotransmitters affected by ADHD medications. This literature is largely inaccessible to those not schooled in epidemiologic analysis of molecular genetic studies, creating reliance on reviews that are excessively enthusiastic and frankly misleading, describing results as having “produced substantial evidence implicating several genes in the etiology [of ADHD].”²³ Such a summary sidesteps several scientific issues.²⁴⁻²⁷ A methodologic white paper on gene–disease associations cautions, “Regarding the strength of association, many of the genetic variants so far identified as influencing susceptibility to common diseases are associated with a low relative and absolute risk. Therefore exclusion of noncausal

explanations for association is crucial.”^{25(p307)} This task has not been undertaken for ADHD.

Waldman and Gizer²⁶ outline areas for future research, illuminating difficulties with current studies. Individual study replicability has been poor, with both positive and negative results for each candidate gene. The most obvious explanation is small sample size with low statistical power. Multiple studies have sought to replicate findings of an association of the dopamine D4 and D5 receptor genes (DRD4 and DRD5), the dopamine transporter gene (DAT), the dopamine β -hydroxylase gene (DBH), the serotonin transporter and receptor 1B genes (5-HTT and HTR1B), and the synaptosomal-associated protein 25 gene (SNAP25) with ADHD. Meta-analyses have produced mixed results, for example, regarding whether or not the dopamine transporter gene DAT is related to ADHD.^{28,29} The validity of meta-analysis for observational molecular genetics studies is debated, because the technique combines results of studies that may be disparate in design and subject ascertainment and exclusion.²⁶ Sample pooling is preferable because it combines data from individuals. However, family-based studies show pooled odds ratios that are barely significant (eg, dopamine D4 receptor gene 1.16 [1.03-1.13] and the dopamine transporter gene 1.13 [1.03-1.24]), whereas case–control studies, which show slightly higher odds ratios, are weakened by inherent susceptibility to the effects of population stratification.^{23,30}

It is difficult to extrapolate a causal role for single markers in a candidate gene to ADHD as a whole, because individual genes are unlikely to act directly and independently of surrounding genes. It is also difficult to link candidate genes to specific measurable qualities of ADHD. This search for an “endophenotype” (a simplified phenotype of a more complex disorder that is closer to the biological cause or candidate gene) has not yielded an objective measure or mechanism.³¹ Finally, studies of candidate genes have ignored the role of individual characteristics such as age, gender, age of onset, and gene–environment interactions, which are known to affect rates of ADHD diagnosis and have played a key role in other medical genetics stories (eg, breast cancer and BRCA1).²⁶ Swanson et al note,

The existing studies of genetic and environmental factors do not meet the standards for modern molecular genetic studies . . . [which] emphasize that gene–gene and gene–environment interactions are likely to be present and require large sample sizes to detect and describe.^{32(p53)}

Given the extraordinary variability of children labeled as ADHD, many of whose symptoms change over time, and the lack of any measurable endophenotype, the likelihood that any of the candidate molecular genes has a causal role in ADHD is as small as the very low pooled odds ratios would indicate, and it is misleading and inaccurate to state that any substantive “genetic cause” of ADHD has been identified.³³

The evidence for a neuroanatomic cause/locus for ADHD has been similarly misrepresented. Studies of neuroanatomy are compromised by small numbers; use of “controls” with other neurologic diagnoses; inclusion of subjects who are not medication naive; failure to account for confounders, including prenatal and perinatal factors and other psychiatric diagnoses; and use of a cross-sectional rather than longitudinal approach (which is critical because myelination continues into early adulthood).³⁴ Leo and Cohen³⁵ have provided a critical review of studies on the neuroanatomy of ADHD, and they conclude that imaging of the neural circuitry of learning is at best a scientific goal for the future, not a current accomplishment. Executive function, deficits of which are proposed by Barkley as central to ADHD, is localized to the prefrontal cortex and frontostriatal regions, yet the most careful and methodologically acceptable study of children with and without ADHD did not find a significant difference in these regions after controlling for brain volume; rather, a difference in cerebellar volume was noted.^{36,37} Other studies of overall and region-specific cerebellar volumes showed inconsistent rather than confirmatory results.³⁸ It is highly likely that complex brain functions recruit neural networks throughout the cortex and are not mapped identically in all individuals.³⁹⁻⁴¹

The original unifying *neuropsychological theory* postulated for ADHD was a deficiency in executive function, specifically in inhibitory control.³⁶ A recent study comparing 75 boys with “severe” ADHD with 70 matched controls, however, found no group differences on executive function inhibitory control tests.⁴² Meta-analyses have demonstrated that executive function dysfunction is “neither necessary nor sufficient to have a causal role in ADHD.”^{43(p1336)} No single executive function task defines all children with ADHD or ADHD subtype. Meta-analyses show low to moderate effect sizes across studies ($d = 0.4-0.7$; $d = 0.6-0.8$) and overlapping performance distributions, with some ADHD children testing as normal on all tests.^{43,44} One meta-analysis found that only 16% to 51% of children with ADHD were impaired on any 1 of 5 typical tests of executive function and that 21% of children with ADHD, compared with 53% of controls, failed none of multiple tests of executive function.⁴⁴ Another meta-analysis of 83 studies found significant differences between groups on all 13 executive function tasks, but significant group differences were not found in 35% of comparisons (59 of 168).⁴³ “Clearly the strong predictions of the executive function deficit hypothesis of ADHD are not supported. The cognitive literature is thus incompatible with the assumption of pathophysiological homogeneity—of a single core deficit [for ADHD].”^{39(p119)}

It is not surprising that more children diagnosed with ADHD than controls show impairment on tests of executive function. Because an ADHD diagnosis requires impairment in 2 settings, and in most cases 1 setting is school, children who have academic and educational problems are being compared with those who do not. It is also not surprising

that children with academic problems have nonidentical results on neurocognitive testing.

Other neuropsychological theories of ADHD include executive function “subset” theories (working memory manipulation problems, “hot” [affect] driven rather than “cool” [abstract] executive function deficits), the theory of optimal stimulation/arousal deficit, reward response deficit theories, and the cognitive-energetic model.^{39,45-50} A recent scholarly attempt to clarify the relationship between language disorders, spatial working memory manipulation deficits, and ADHD is confounded by significant group differences in intelligence quotients, by absence of significant group differences in ADHD Conners’s Rating Scale scores, and by use of outcome measures (tests of verbal memory) that are also components of tests that measure intelligence quotient.⁴⁹ Almost all samples of children with ADHD as compared to controls have significantly lower mean full-scale intelligence quotient (mean differences of 10 points are typical) and significantly higher rates of reading and language disorder; whether these differences should be statistically controlled for is debated.^{36,49} Additionally, neurocognitive testing studies typically use small sample sizes and exclude children with intelligence quotients less than 80 and children with psychiatric diagnoses, limiting the ability to generalize results.

Neuropsychological research on ADHD has largely ignored environmental and personality factors, which have been important to understanding of other childhood pathologies. Test results are analyzed without consideration for emotional factors. Investigators use reward points for task success as a proxy for affect, motivation, and task saliency.⁵¹ In “How emotions inform judgement and regulate thought,” Clore and Huntsinger⁵² review the importance of affect (feelings) to cognition and cognitive testing. Test anxiety, self-esteem, and interest in task on the day of testing are also not evaluated and are likely to differentially affect performance and response variability.⁵³

Other Causes of ADHD Symptoms

Factors associated with the diagnosis of ADHD include lower socioeconomic level; family conflict; punitive parenting methods; parental affective disorders, such as maternal depression and anxiety; parental substance abuse; parental disruptive, antisocial, and criminal behavior; increased and early TV viewing; and intellectual disability.⁵⁴⁻⁶⁵ In a prospective longitudinal study of a national sample of Canadian children aged 2 to 7 years that controlled for relevant confounders, statistically significant predictors of high and persistent hyperactivity included male sex, maternal depression, hostile parenting, and maternal prenatal smoking.⁶⁵ Enduring associations between environmental, psychosocial, and parenting influences and the diagnosis of ADHD do not prove causality, but should not be ignored.

The very common behaviors described in the *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition (DSM-IV), criteria for ADHD can be conceptualized as *symptoms* of childhood mental, emotional, and cognitive problems rather than as a neurobehavioral disease. This approach explains the remarkable heterogeneity of children whose behavior is diagnosed as ADHD. Children who meet criteria for ADHD may have one or more urgent and treatable conditions, including

1. Ongoing or past physical or sexual abuse, ranging from punitive parenting to explicit abuse;
2. Mental retardation, cognitive limitations, or one or more specific learning disabilities;
3. Hypervigilance and apprehension because of an unsafe or unpredictable environment at home or school;
4. Psychiatric diagnoses, including depression and anxiety;
5. Emotional and developmental difficulties because of conflicts or life stresses that would benefit from psychotherapy; and
6. A mismatch between parental or educator expectations and child behavior and performance—this may be due to developmentally inappropriate or child-inappropriate expectations.

Children with apparently “pure” attentional problems whose grades improve with stimulant treatment may have short-term improvement without long-term benefit, and the child’s apparent improvement (or worsening) can mask need for treatment of an underlying problem. A mainstream ADHD researcher noted that the “costs of [ADHD] treatment reflect dramatic underutilization of psychosocial treatments” and recommended the use of behavioral treatments, including educational interventions and parent education before medication.⁶⁶ In a list of 10 “Limitations of Pharmacological Interventions When Used Alone” included are “removes incentives for parents and teachers/schools to work on other treatments,” “does not affect several important variables (eg, academic achievement, concurrent family problems, peer relationships),” and “uniform lack of evidence for beneficial long-term effects.”⁶⁶

The Parent Trap

Clinicians may be reluctant to explore causes for symptoms diagnosed as ADHD. Without obvious psychosocial stressors, with an intelligence quotient above 80 and parents who are involved and caring (and often professionals), it may seem untenable that a child’s behaviors could be anything but a “brain disease.” It is falsely assumed that the only other possibility is that the child’s difficulties are the parents’ fault.

This unhelpful and fallacious dichotomous approach prevents children from obtaining needed treatment. It

also fails to engage the parents in the supportive role to which they are optimally suited; they know the child best and are best positioned to alter home care in ways that can help the child. The simplistic concept of a “neurotransmitter deficiency” is well outdated, and even generalized “behavior treatments” that may not address the individual child’s needs show some efficacy in decreasing symptoms.^{67,68} Therapy that is tailored to the individual child’s difficulties and includes parental support has not received sufficient attention.

Comorbidities or Etiologies?

The range of “comorbidities” described in ADHD includes oppositional defiant disorder (30% to 60%); conduct disorder (9% to 19%); mood disorders, including depression (18% to 60%); anxiety disorders (23% to 34%); and learning disorders (10% to 90%).^{10,69-76} The significant degree of overlap between ADHD and oppositional defiant disorder and conduct disorder suggests that the same children are being identified: “Almost all children younger than 12 years of age who meet criteria for ODD [oppositional defiant disorder] or CD [conduct disorder] will almost always meet criteria for ADHD.”^{76(p526)} Depression and anxiety may be difficult to identify in children, both have documented strong continuity in adulthood, and neither condition is treated with stimulant medications.⁷⁷⁻⁸⁵ In fact, children with major affective disorders including depression may have dysphoria and worsening with stimulants, whereas children with anxiety plus ADHD diagnosis are much less likely to have response to stimulants.^{11,81,82,86-88} An increasing number of neuropsychologists are exploring the overlap between learning disabilities and ADHD.^{48,49} Although stimulants cause motoric slowing and improved attention and performance on rote tasks for both control and ADHD-diagnosed individuals, they do not improve reading ability, knowledge acquisition, or knowledge application.⁸⁹⁻⁹⁶ Because early and intensive treatment of learning disabilities improves educational outcomes, the use of stimulant medications as first-line treatment for academic failure is worrisome.⁹⁷⁻⁹⁹ “Comorbidities” mimic ADHD, may be difficult to diagnose accurately, and have poor outcomes without appropriate therapy. It is likely that “comorbidities” produce symptoms mistakenly diagnosed as ADHD.

The Rating Scales— Searching for Diagnostic Validity

Children aged 6 years and older are diagnosed with ADHD if they are identified as having 6 or more hyperactive/impulsive or inattentive behaviors from a checklist in the DSM-IV and the behaviors occur “to an extent that is disruptive and inappropriate for developmental level,”

cause “clinically significant impairment” in 2 settings, and were present for more than 6 months with onset before age 7 years.¹⁰⁰ In practice, many more children of younger ages are identified, and many clinicians identify children who do not strictly meet the criteria. However, the validity of this behavior-driven diagnostic method is debated. Horwitz and Wakefield¹⁰¹ examine this clinical problem as it applies to the *DSM-IV* diagnosis of depression.¹⁰¹ In his review of their book, McHugh notes that this approach “may obscure important distinctions between a patient with a mental disorder and a person who is responding naturally to life circumstances.” This concern applies equally to the diagnosis of ADHD.¹⁰² A highly reliable diagnostic method may repeatedly identify the same individual(s), yet does not create a meaningful or coherent diagnostic category.

Snyder et al¹⁰³ reviewed the accuracy of ADHD rating scales using American Academy of Neurology evidence standards. Eleven scales, including the widely used Conners’s Rating Scale–Revised, were eliminated because of major design flaws such as use of the same sample for development and validation of the tool (Conners’s Rating Scale–Revised), use of the evaluated scale as the clinical standard, and convergent validity with no prior validation of either scale. Thirteen scales could be evaluated, none met all standards, and a pooled mean accuracy of 69% ($\pm 7\%$) for 9 studies with 2228 participants was reported. Not evaluated were the Vanderbilt ADHD rating scales, which have been widely promoted by the American Academy of Pediatrics in their ADHD Toolkit.¹⁰⁴ The forms were partially funded by McNeil Consumer and Specialty Pharmaceuticals, whose parent pharmaceutical company manufactures Concerta, a sustained release stimulant medication for ADHD.¹⁰⁵

Investigators developing the Vanderbilt ADHD Parent Rating Scale recruited parents from an urban K–12 school with 6171 students and screened with teacher ratings that identified 1536 (23%) as having ADHD.^{106,107} Teachers had no information on longevity of symptoms, 2 setting behaviors, or age of onset of symptoms, as required by *DSM-IV* for diagnosis. Parents of these 1536 students received anonymous letters requesting participation, of whom 288 (18.7%) responded and 241 (15.7%) completed 3 interviews that were compared with teacher evaluations. The responding parents differed from nonresponders in that significantly more of their children were receiving stimulant medications. During the 2-year study children were not interviewed, observed, or directly assessed at any time. Investigators used the ADHD section of the Diagnostic Interview Schedule for Children (DISC-IV) to validate the Vanderbilt ADHD Parent Rating Scale, even though “no formal validity testing has been performed” for this section of the DISC-IV.¹⁰⁸ They found good concurrent validity (0.79)—not unexpectedly because the same 18 questions are used on both scales—but very low interrater reliability between teachers and parents and low test–retest reliability.

The investigators concluded that the Vanderbilt ADHD Parent Rating Scale has “good psychometric properties in a high risk population.” Alternately, using Snyder’s evaluative caveats, one would conclude that the Vanderbilt ADHD Parent Rating Scale does not have good psychometric properties because (a) it underwent testing with a small pre-screened population whose diagnosis was not defined by *DSM-IV* criteria or interview, and (b) convergent validity was demonstrated with the DISC-IV without prior validation of either scale.

Development of the Vanderbilt ADHD Diagnostic Teacher Rating Scale involved teachers of 8257 children (year 1) and 4323 children (year 2) in a suburban school system (6.9% African American, 17.2% low income).^{109–111} Internal consistency for the scale was high, but the authors report,

It was not possible to obtain a precise measure of the presence of AD/HD by which to compare the scale to a “gold standard” . . . The teachers’ responses to the question about the child having been diagnosed with AD/HD provided an approximation . . . One limitation of this study is that all information was obtained from teacher reports, which does not allow for an independent determination of the diagnoses.^{110(pp149-150)}

Thus, validation of the Vanderbilt ADHD Diagnostic Teacher Rating Scale was compromised by use of a population that did not meet standardized *DSM-IV* criteria for ADHD, which is demographically at lower risk for ADHD than the general population, and by use of the same informants for both the clinical diagnosis and rating scale development.

The problem of parent–teacher rating scale discrepancy is not easily dismissed because *DSM-IV* requires impairment in 2 settings. In evaluating the Vanderbilt forms, Leslie et al¹¹² reported a discrepancy rate such that 40% of children no longer met criteria for diagnosis with both raters. Wolraich et al¹⁰⁷ found teacher–parent concordance so low that diagnostic rates decreased among all subtypes of ADHD, leading the authors to suggest that criteria be relaxed. Other studies show similar results.^{113–115} Knowing whether a behavior is “inappropriate for developmental level” requires knowledge of “appropriate” for developmental level, and persistent discrepancies may reflect the work of untrained informants (parents and teachers) using subjective criteria. Scales do not include objective measures or child interview and do not adjust for any child characteristic (eg, age, intelligence quotient, gender). Because a child’s behavior depends on setting-specific structure, expectations, and adult guidance, a home–school discrepancy is not surprising.

Rating scales are called “objectives” because quantitative information is generated. Missing from the evaluation of most children labeled as ADHD are “projectives,” or tests that use a free response and require interpretation. Examples include the Children’s Apperception Test, which presents a

child with 10 pictures to describe. Projectives evaluate personality structure and dynamics and assess coping and defense mechanisms and can identify possible underlying reasons for attentional or behavioral problems, such as anxiety, depression, or areas of emotional conflict that can respond to appropriate therapy.

Medications

Few clinicians who attempt to treat ADHD do so without prescribing stimulant medications. Both older short-acting preparations and newer more expensive sustained release preparations of mixed amphetamine salts and methylphenidate are marketed, in addition to the non-stimulant atomoxetine. No published evidence exists that demonstrates better comparative effectiveness of any of these pharmacologic treatments for ADHD.³ Choice of medication is anecdotal and physician driven.

Side effects of treatment are infrequently studied prospectively. Rates range from 5% to 25% for anorexia, listlessness, nausea, somnolence, and mood swings, with at least 1 side effect reported in 22% of children in a large study.¹¹⁶ Side effects of higher morbidity and lower frequency include tics, hallucinations, and bizarre behavior. Few studies examine the opinions of the child who is receiving the medication, but when opinion is sought, 12.7% to 18.8% of children reported that they feel worse on stimulants.¹¹⁷

Few ADHD studies were actually designed to monitor growth, and long-term studies that continue through the pubertal growth spurt to adult height have not been performed. Clinicians face a significant challenge interpreting results because most studies are performed by researchers funded by pharmaceutical firms that manufacture the studied medications, and results revealing growth decrements are summarized with reassuring language that minimizes findings.^{118,119} For example, growth decrements in a prospective study of 0.23 cm and 1.23 kg after 21 months in osmotic-release methylphenidate treated as compared with untreated children are described as "clinically insignificant."¹²⁰ The Multimodal Treatment Study of Children with Attention Deficit Hyperactivity Disorder documented average growth of 2.0 cm and 2.7 kg less in newly medicated as compared with never medicated children at 36 months, without evidence of growth rebound.¹²¹ This information was obtained prospectively and appears solid. Longer-term studies also show measurable effects of stimulants on height and weight gain.¹²²

Whether the reported rate of sudden death in children receiving amphetamine and methylphenidate products (0.3-0.2/100 000 patient years) actually differs from baseline rates of sudden death in childhood is not known because of limitations in the reliability of spontaneous report data.^{3,123} Elevations of blood pressure (3.6 mm Hg systolic, 2.5 mm Hg diastolic) and resting heart rate (3.4

beats per minute) during 2 years of stimulant treatment are described as "insignificant"; however, because long-term studies have not been performed, and prehypertension is causally linked to cardiovascular mortality, this conclusion may be premature.¹²⁴⁻¹²⁶

Conflict of Interest

Pharmaceutical firms that manufacture ADHD medications play a large role in promotion of ADHD as a disease requiring medication. Pharmaceutical firms fund informational brochures from professional organizations such as the American Academy of Pediatrics, funded the development of the Vanderbilt ADHD rating scales in the American Academy of Pediatrics "ADHD Toolkit," contribute substantively to parent support groups, including Child and Adults with Attention-Deficit/Hyperactivity Disorder (26.11% of its \$1 169 000 budget for 2006-2007), and fund continuing medical education, speakers, and physician researchers.^{104,105,127-129} Over half of the principal investigators participating in the National Institutes of Mental Health Multimodal Treatment Study of Children with Attention Deficit Hyperactivity Disorder reported multiple financial disclosures due to pharmaceutical funding received as speakers, consultants, and researchers.¹³⁻¹⁵ Thirteen of the 21 individuals who created the *DSM-IV* criteria for diagnosis of ADHD have financial ties to manufacturers of ADHD medications.^{130,131}

There is evidence from the fields of economics and psychology that self-interest bias "does not appear to be strategic or deliberate,"^{132(p110)} yet is pervasive, and "It takes extraordinarily little to bias an individual's interpretation and processing of information."^{133(p42)} There is a growing awareness that acceptance of gifts from pharmaceutical firms is unethical, that pharmaceutical sponsorship of "research" may contribute significantly to publication bias, and that physicians who accept fees to perform as consultants may not be serving the best interests of their patients.¹³⁴⁻¹⁴¹ Physicians should consider the possibility that extensive pharmaceutical funding has essentially controlled research and treatment for children with symptoms of hyperactivity and inattention and has driven an explosion in the number of children who must be diagnosed as ADHD.

Summary

It is critical for unbiased clinicians to consider the possibility that ADHD is not a neurobehavioral disease but rather a constellation of symptoms that require attention (no pun intended). Children and adolescents manifesting hyperactivity, impulsivity, or inattention may have treatable medical, emotional, or psychosocial conditions that require urgent care. Stimulant medications are costly, have no proven

long-term efficacy, and may mask problems without benefiting the child. Parents, educators, and physicians are urged to reevaluate their thinking on this controversial topic.

References

- Stein RK. Measurement of ADHD outcomes: implications for future research. *Ambul Pediatr*. 2007;7:140-143.
- Froehlich TE, Lanphear BP, Epstein JN, Barbaresi WJ, Katusic SK, Kahn RS. Prevalence, recognition and treatment of attention-deficit/hyperactivity disorder in a national sample of US children. *Arch Pediatr Adolesc Med*. 2007;161:857-864.
- McDonagh MS, Peterson K, Dana T, Thakurta S. Drug class review on pharmacologic treatments for ADHD. Final Report. Oregon Evidence-based Practice Center, Oregon Health and Science University. <http://www.ohsu.edu/drugeffectiveness/reports/documents/ADHD%20Final%20Report%20Update%20.pdf>. Published December 2007. Accessed January 12, 2008.
- Breggin PR. Questioning the treatment for ADHD. *Science*. 2001;291:595.
- McGuiness D. Attention deficit disorder: the emperor's clothes, animal "pharm," and other fiction. In: Fisher S, Greenburg RP, eds. *The Limits of Biological Treatments for Psychological Distress*. Hillsdale, NJ: Lawrence Erlbaum; 1989:151-187.
- Timimi S. *Naughty Boys: Anti-Social Behaviour, ADHD and the Role of Culture*. Basingstoke, UK: Palgrave Macmillan; 2005.
- Carey WB. Is ADHD a valid disorder? In: Jensen PS, Cooper JR, eds. *Attention Deficit Hyperactivity Disorder: State of the Science. Best Practices*. Kingston, NJ: Civic Research Institute; 2002.
- Furman L. What is attention-deficit/hyperactivity disorder (ADHD)? *J Child Neurol*. 2005;20:994-1003.
- Scheffler RM, Hinshaw SP, Modrek S, Levine P. The global market for ADHD medications. *Health Aff*. 2007;26:450-457.
- A 14-month randomized clinical trial of treatment strategies for attention-deficit/hyperactivity disorder. The MTA Cooperative Group. Multimodal Treatment Study of Children with ADHD. *Arch Gen Psychiatry*. 1999;56:1073-1086.
- MTA Cooperative Group. National Institute of Mental Health Multimodal Treatment Study of ADHD follow-up: changes in effectiveness and growth after the end of treatment. *Pediatrics*. 2004;113:762-769.
- MTA Cooperative Group. National Institute of Mental Health Multimodal Treatment Study of ADHD follow-up: 24-month outcomes of treatment strategies for attention-deficit/hyperactivity disorder. *Pediatrics*. 2004;113:754-761.
- Swanson JM, Hinshaw SP, Arnold LA, et al. Secondary evaluations of MTA 36-month outcomes: propensity score and growth mixture model analyses. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1003-1014.
- Molina B, Flory K, Hinshaw SP, et al. Delinquent behavior and emerging substance use in the MTA at 36 months: prevalence, course, and treatment effects. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1028-1040.
- Jensen PS, Arnold LE, Swanson JM, et al. 3-Year follow-up of the NIMH MTA study. *J Am Acad Child Adolesc Psychiatry*. 2007;46:989-1002.
- Banerjee TD, Middleton F, Faraone SV. Environmental risk factors for attention-deficit hyperactivity disorder. *Acta Paediatr*. 2007;96:1269-1274.
- Joseph J. Not in their genes: a critical view of the genetics of attention-deficit hyperactivity disorder. *Dev Rev*. 2000;20:539-567.
- Joseph J. Problems in psychiatric genetic research: A reply to Faraone and Biederman. *Dev Rev*. 2000;20:582-593.
- Joseph J. Adoption study of ADHD. *J Am Acad Child Adolesc Psychiatry*. 2002;41:1389-1391.
- Joseph J. *The Missing Gene: Psychiatry, Heredity, and the Fruitless Search for Genes*. New York, NY: Algora; 2006.
- Scarr S, Carter-Saltzman L. Twin method: defense of a critical assumption. *Behav Genet*. 1979;9:527-542.
- Kringle E. *Heredity and Environment in the Functional Psychoses: An Epidemiological-Clinical Study*. Oslo, Norway: Univeritetsforlaget; 1967.
- Faraone SV, Khan SA. Candidate gene studies of attention-deficit/hyperactivity disorder. *J Clin Psychiatry*. 2006;67(suppl 8): 13-20.
- Elia J, Devoto M. ADHD genetics: 2007 update. *Curr Psychiatry Rep*. 2007;9:434-439.
- Little J, Bradley L, Bray MS, et al. Reporting, appraising, and integrating data on genotype prevalence and gene-disease associations. *Am J Epidemiol*. 2002;156:300-310.
- Waldman ID, Gizer IR. The genetics of attention deficit hyperactivity disorder. *Clin Psychol Rev*. 2006;26:396-432.
- Risch N, Teng J. The relative power of family-based and case-control designs for linkage disequilibrium studies of complex human diseases I. DNA pooling. *Genome Res*. 1998;8:1273-1288.
- Li D, Sham PC, Owen MJ, He L. Meta-analysis shows significant association between dopamine system genes and attention deficit hyperactivity disorder (ADHD). *Hum Mol Genet*. 2006;15:2276-2284.
- Yang B, Chan RC, Jing J, Li T, Sham P, Chen RY. A meta-analysis of association studies between the 10-repeat allele of a VNTR polymorphism in the 3'-UTR of dopamine transporter gene and attention deficit hyperactivity disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2007;144:541-550.
- Faraone SV, Khan SA. Candidate gene studies of attention deficit/hyperactivity disorder. *J Clin Psychiatry*. 2006;67(suppl 8): 13-20.
- Doyle A, Willcutt EG, Seidman LJ, et al. Attention-deficit/hyperactivity disorder endophenotypes. *Biol Psychiatry*. 2005;57: 1324-1335.
- Swanson JM, Kinsbourne M, Nigg J, et al. Etiologic subtypes of attention-deficit/hyperactivity disorder: brain imaging, molecular genetic and environmental factors and the dopamine hypothesis. *Neuropsychol Rev*. 2007;17:39-59.
- Lahey BB, Pelham WE, Loney J, Lee SS, Willcutt E. Instability of the DSM-IV subtypes of ADHD from preschool through elementary school. *Arch Gen Psychiatry*. 2005;62:896-902.
- Thatcher RW. Maturation of the human frontal lobes: physiological evidence for staging. *Dev Neuropsychol*. 1991;7: 397-419.
- Leo J, Cohen D. Broken brains or flawed studies? A critical review of ADHD neuroimaging research. *J Mind Behav*. 2003; 24:29-56.
- Barkley RA. Attention-deficit/hyperactivity disorder, self-regulation, and time: toward a more comprehensive theory. *J Dev Behav Pediatr*. 1997;18:271-279.
- Castellanos FX, Lee PP, Sharp W, et al. Developmental trajectories of brain volume abnormalities in children and adolescents with attention-deficit/hyperactivity disorder. *JAMA*. 2002; 288:1740-1748.

38. Seidman LJ, Valera EM, Makris N. Advancing the neuroscience of ADHD, structural brain imaging of attention-deficit/hyperactivity disorder. *Biol Psychiatry*. 2005;57:1263-1272.
39. Castellanos FX, Sonuga-Barke EJS, Milham MP, Tannock R. Characterizing cognition in ADHD: beyond executive dysfunction. *Trends Cogn Sci*. 2006;10:117-123.
40. Sanai N, Mirzadeh Z, Berger MS. Functional outcome after language mapping for glioma resection. *N Engl J Med*. 2008;358:18-27.
41. Snyder SH. Seeking god in the brain. *N Engl J Med*. 2008;358:6-7.
42. Rhodes SM, Coghill DR, Matthews K. Neuropsychological functioning in stimulant-naïve boys with hyperkinetic disorder. *Psychol Med*. 2005;35:1109-1120.
43. Wilcutt EG. Validity of the executive function theory of attention deficit/hyperactivity disorder: a meta-analytic review. *Biol Psychiatry*. 2005;57:1336-1446.
44. Nigg JT, Wilcutt EG, Doyle AE, Sonuga-Barke EJS. Causal heterogeneity in attention deficit/hyperactivity disorder: do we need neuropsychological subtypes? *Biol Psychiatry*. 2005;57:1224-1230.
45. Nigg JT. Neuropsychologic theory and findings in attention-deficit/hyperactivity disorder: the state of the field and salient challenges for the coming decade. *Biol Psychiatry*. 2005;57:1424-1435.
46. Solanto MV, Abikoff H, Sonuga-Barke E, et al. The ecological validity of delay aversion and response inhibition as measures of impulsivity in AD/HD: A supplement to the NIMH multimodal treatment study of AD/HD. *J Abnorm Child Psychol*. 2001;29:215-228.
47. Zentall S, Zentall T. Optimal stimulation: a model of disordered activity and performance in normal and deviant children. *Psychol Bull*. 1983;94:446-471.
48. Denckla MB. A theory and model of executive function: a neuropsychological perspective. In: Lyon GA, Krasnegor NA, eds. *Attention, Memory and Executive Function*. Baltimore, MD: Paul H. Brooks; 1983:263-278.
49. Martinussen R, Tannock R. Working memory impairments in children with attention-deficit hyperactivity disorder with and without comorbid language learning disorders. *J Clin Exp Neuropsychol*. 2006;28:1073-1094.
50. Sonuga-Barke EJS. Causal models of attention-deficit/hyperactivity disorder: from common simple deficits to multiple developmental pathways. *Biol Psychiatry*. 2005;57:1231-1238.
51. Wodka EL, Mahone EM, Blankner JG, et al. Evidence that response inhibition is a primary deficit in ADHD. *J Clin Exp Neuropsychol*. 2007;29:345-356.
52. Clore GL, Huntsinger JR. How emotions inform judgement and regulate thought. *Trends Cogn Sci*. 2007;11:393-399.
53. Volkow ND, Wang GJ, Fowler JS, et al. Evidence that methylphenidate enhances the saliency of a mathematical task by increasing dopamine in the human brain. *Am J Psychiatry*. 2004;161:1173-1180.
54. Biederman J, Milberger S, Faraone SV, et al. Family-environment risk factors for attention deficit hyperactivity disorder: a test of Rutter's indicators of adversity. *Arch Gen Psychiatry*. 1995;52:464-470.
55. Biederman J, Faraone S, Milberger S, et al. Predictors of persistence and remission of ADHD into adolescence: results from a four-year prospective follow-up study. *J Am Acad Child Adolesc Psychiatry*. 1996;35:343-351.
56. Chronis AM, Lahey BB, Pelham WE, Kipp HL, Baumann BL, Lee SS. Psychopathology and substance abuse in parents of young children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry*. 2003;42:1424-1432.
57. Chronis AM, Lahey BB, Pelham WE, et al. Maternal depression and early positive parenting predict future conduct problems in young children with attention-deficit/hyperactivity disorder. *Dev Psychol*. 2007;43:70-82.
58. Connor DF, Edwards G, Fletcher KE, Baird J, Barkley RA, Steingard RJ. Correlates of comorbid psychopathology in children with ADHD. *J Am Acad Child Adolesc Psychiatry*. 2003;42:193-200.
59. Elgar FJ, Curtis LJ, McGrath PJ, Waschbusch DA, Stewart SH. Antecedent-consequence conditions in maternal mood and child adjustment: a four-year cross-lagged study. *J Clin Child Adolesc Psychol*. 2003;42:362-374.
60. Fisher M. Parenting stress and the child with attention-deficit/hyperactivity disorder. *J Clin Child Psychol*. 1990;19:337-346.
61. Johnson JG, Cohen P, Kasen S, Brooke JS. Extensive television viewing and the development of attention and learning difficulties during adolescence. *Arch Pediatr Adolesc Med*. 2007;161:480-486.
62. Johnston C, Mash EJ. Families of children with attention-deficit/hyperactivity disorder: review and recommendations for future research. *Clin Child Fam Psychol Rev*. 2001;4:183-207.
63. Siminoff E, Pickles A, Wood N, Gringras P, Chadwick O. ADHD symptoms in children with mild intellectual disability. *J Am Acad Child Adolesc Psychiatry*. 2007;46:591-600.
64. Antshel KM, Phillips MH, Gordon M, Barkley R, Faraone SV. Is ADHD a valid disorder in children with intellectual delays? *Clin Psychol Rev*. 2006;26:555-572.
65. Romano E, Tremblay RE, Farhat A, Côté S. Development and prediction of hyperactive symptoms from 2 to 7 years in a population-based sample. *Pediatrics*. 2006;117:2101-2109.
66. Pelham W. What should we treat and how should we treat it? Paper presented at: The Spectrum of Developmental Disabilities XXIX: ADHD—Beyond the Basics; March 28, 2007; Baltimore MD.
67. Piffner LJ, Yee Mikami A, Huang-Pollack C, Easterlin B, Zalecki C, McBurnett K. A randomized, controlled trial of integrated home-school behavioral treatment for ADHD, predominantly inattentive type. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1041-1050.
68. Van den Hoofdakker B, van den Veen-Mulders L, Stema S, Emmelkamp P, Minderaa RB, Nauta M. Effectiveness of behavioral parent training for children with ADHD in routine clinical practice: a randomized clinical study. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1263-1271.
69. Pliszka SR. Psychiatric comorbidities in children with attention deficit hyperactivity disorder: implications for management. *Paediatr Drugs*. 2003;5:741-50.
70. Dilsaver SC, Henderson-Fuller S, Akiskal HS. Occult mood disorders in 104 consecutively presenting children referred for the treatment of attention-deficit/hyperactivity disorder in a community mental health clinic. *J Clin Psychiatry*. 2003;64:1170-1176.
71. Brown TE, ed. *Attention-Deficit Disorders and Comorbidities in Children, Adolescents, and Adults*. Washington, DC: American Psychiatric Press; 2000.

72. Butler SF, Arredondo DE, McCloskey V. Affective comorbidity in children and adolescents with attention deficit hyperactivity disorder. *Ann Clin Psychiatry*. 1995;7:51-55.
73. Wolraich ML, Hannah JN, Baumgaertel A, Brown J. Comparison of diagnostic criteria for attention deficit/hyperactivity disorder in a county-wide sample. *J Am Acad Child Adolesc Psychiatry*. 1996;35:319-324.
74. Wilens TE, Biederman J, Brown S, et al. Psychiatric comorbidity and functioning in clinically referred preschool children and school-age youth with ADHD. *J Am Acad Child Adolesc Psychiatry*. 2002;41:262-268.
75. Semrud-Clikeman M, Biederman J, Sprich-Buckminster S, Lehman BK, Faraone SV, Norman D. Comorbidity between ADDH and learning disability: a review and report in a clinically referred sample *J Am Acad Child Adolesc Psychiatry*. 1992;31:439-448.
76. Pliszka SR. Patterns of psychiatric comorbidity with attention deficit/hyperactivity disorder. *Child Adolesc Psychiatry Clin N Am*. 2000;9:525-540.
77. DelBello MP, Soutullo CA, Hendricks W, Niemeier RT, McElroy SL, Strakowski SM. Prior stimulant treatment in adolescents with bipolar disorder: association with age at onset. *Bipolar Disord*. 2001;3:53-57.
78. Furman E. What is depression in childhood? *Child Anal*. 1992;4:101-123.
79. Bhatia SK, Bhatia SC. Childhood and adolescent depression. *Am Fam Physician*. 2007;75:73-80.
80. Pine DS, Cohen E, Cohen P, Brook J. Adolescent depressive symptoms as predictors of adult depression: moodiness or mood disorder? *Am J Psychiatry*. 1999;156:133-135.
81. Coyle JT. Depression and bipolar support alliance consensus statement on the unmet needs in diagnosis and treatment of mood disorders in children and adolescents. *J Am Acad Child Adolesc Psychiatry*. 2003;42:1494-1503.
82. Compton SN. Cognitive-behavioral psychotherapy for anxiety and depressive disorders in children and adolescents: an evidence-based medicine review *J Am Acad Child Adolesc Psychiatry*. 2004;43:930-951.
83. Pine DS, Cohen P, Brook J. Adolescent fears as predictors of depression. *Biol Psychiatry*. 2001;50:721-724.
84. Ryan ND. Diagnosing pediatric depression. *Biol Psychiatry*. 2001;49:1050-1054.
85. Lewinsohn PM, Rohde P, Seeley JR, Klein DN, Gotlib IH. Natural course of adolescent major depressive disorder in a community sample: predictors of recurrence in young adults. *Am J Psychiatry*. 2002;157:1584-1591.
86. Pliszka SR. Effect of anxiety on cognition, behavior, and stimulant response in ADHD. *J Am Acad Child Adolesc Psychiatry*. 1989;28:882-887.
87. Goetz H, Back-Bennet O, Zelnik N. Differential stimulant response on attention in children with comorbid anxiety and oppositional defiant disorder. *J Child Neurol*. 2007;22:538-542.
88. Bedard AC, Tannock R. Anxiety, methylphenidate response, and working memory in children with ADHD. *J Atten Disord*. 2008;11:546-557.
89. Barkley RA, Cunningham CE. Do stimulant drugs improve the performance of hyperkinetic children? *Clin Pediatr*. 1978;17:85-92.
90. Douglas VI, Barr RG, O'Neill ME, Britton BG. Short term effects of methylphenidate on the cognitive, learning and academic performance of children with attention deficit disorder in the laboratory and the classroom. *J Child Psychol Psychiatry*. 1986;27:191-211.
91. Bernstein GA, Carroll ME, Crosby RD, Perwein AR, Go FS, Benowitz NL. Caffeine effects on learning, performance and anxiety in normal school-age children. *J Am Acad Child Adolesc Psychiatry*. 1994;33:407-415.
92. Rappaport JL, Buchsbaum MS, Zahn TP, Weingartner H, Ludlow C, Mikkelsen EJ. Dextroamphetamine: cognitive and behavioral effects in normal prepubertal boys. *Science*. 1978;199:560-563.
93. Rapport MD, Denney C, DuPaul GJ, Gardner MJ. Attention deficit disorder and methylphenidate: normalization rates, clinical effectiveness and response prediction in 76 children. *J Am Acad Child Adolesc Psychiatry*. 1994;33:882-893.
94. Rappaport JL, Buchsbaum MS, Weingartner H, Zahn TP, Ludlow C, Mikkelsen EJ. Dextroamphetamine: its cognitive and behavioral effects in normal and hyperactive boys and men. *Arch Gen Psychiatry*. 1980;37:933-943.
95. Rie HE, Rie ED, Stewart S, Ambuel JP. Effects of ritalin on underachieving children: a replication. *Am J Orthopsychiatry*. 1976;46:313-322.
96. Rhodes SM, Coghill DR, Matthews K. Acute neuropsychological effects of methylphenidate in stimulant drug-naïve boys with ADHD II—broader executive and non-executive domains. *J Child Psychol Psychiatry*. 2006;47:1184-1194.
97. Grizenko N, Bhat M, Schwartz G, Ter-Stepanian M, Joobar R. Efficacy of methylphenidate in children with attention-deficit hyperactivity disorder and learning disabilities: a randomized crossover trial. *J Psychiatry Neurosci*. 2006;31:46-51.
98. Foorman BR, Breier JI, Fletcher JM. Interventions aimed at improving reading success: an evidence-based approach. *Dev Neuropsychol*. 2003;24:613-639.
99. O'Connor RE, Harty KR, Fulmer D. Tiers of intervention in kindergarten through third grade. *J Learn Disabil*. 2005;38:532-538.
100. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. Washington, DC: American Psychological Association; 1994.
101. Horwitz AV, Wakefield JC. *The Loss of Sadness. How Psychiatry Transformed Normal Sorrow into Depressive Disorder*. New York, NY: Oxford University Press; 2007.
102. McHugh PR. Book review: the loss of sadness. *N Eng J Med*. 2007;357:947.
103. Snyder SM, Hall JR, Cornwell SL, Quintana H. Review of clinical validation of ADHD behavior rating scales. *Psychol Rep*. 2006;99:363-378.
104. American Academy of Pediatrics Arizona Chapter. ADHD Toolkit. <http://www.azaap.org/for-health-care-providers/adhd.aspx>. Accessed January 26, 2008.
105. Van Landeghem K, Hess C. Children's mental health: an overview and key considerations for health system stakeholders. NICHM Issue Paper. National Institute for Health Care Management Foundation. <http://www.nihcm.org/pdf/CMHReport-FINAL.pdf>. Published February 2005. Accessed January 26, 2008.
106. Wolraich ML, Lambert W, Doffing MA, Bickman L, Simmons T, Worley K. Psychometric properties of the Vanderbilt ADHD diagnostic parent rating scale in a referred population. *J Pediatr Psychol*. 2003;28:559-567.
107. Wolraich ML, Lambert EW, Bickman L, Simmons T, Doffing MA, Worley KA. Assessing the impact of parent and teacher agreement on diagnosing attention-deficit hyperactivity disorder. *J Dev Behav Pediatr*. 2004;25:41-47.
108. Shaffer D, Fisher P, Lucas CP, Dulcan MK, Schwab-Stone ME. The NIMH Diagnostic Interview Schedule for Children Version IV (DISC-IV): description, differences from previous

- versions, and reliability of some common diagnoses. *J Am Acad Child Adolesc Psychiatry*. 2000;39:28-38.
109. Wolraich ML, Hannah JN, Baumgaertel A, Brown J. Comparison of diagnostic criteria for attention deficit/hyperactivity disorder in a county-wide sample. *J Am Acad Child Adolesc Psychiatry*. 1996;35:319-324.
 110. Wolraich ML, Feurer I, Hannah JN, Pinnock TY, Baumgaertel A. Obtaining systematic teacher reports of disruptive behavior disorders utilizing DSM-IV. *J Abnorm Child Psychol*. 1998;26:141-152.
 111. Wolraich ML, Hannah JN, Baumgaertel A, Feurer ID. Examination of DSM-IV criteria for attention deficit/hyperactivity disorder in a county-wide sample. *J Dev Behav Pediatr*. 1998;19:162-168.
 112. Leslie LK, Weckerly J, Plemmons D, Landsverk J, Eastman S. Implementing the American Academy of Pediatrics attention-deficit/hyperactivity disorder diagnostic guidelines in primary care settings. *Pediatrics*. 2004;114:129-140.
 113. Lahey BB, McBurnett K, Piacentini JC. Agreement of parent and teacher rating scales with comprehensive clinical assessment of attention deficit disorder with hyperactivity. *J Psychopathol Behav Assess*. 1987;9:429-439.
 114. Mitsis EM, McKay KE, Schulz KP, Newcorn JH, Halperin JM. Parent-teacher concordance for DSM-IV attention-deficit/hyperactivity disorder in a clinic-referred sample. *J Am Acad Child Adolesc Psychiatry*. 2000;39:308-313.
 115. Tripp G, Schaughency EA, Clarke B. Parent and teacher rating scales in the evaluation of attention-deficit hyperactivity disorder: contribution to diagnosis and differential diagnosis in clinically referred children. *J Dev Behav Pediatr*. 2006;27:209-218.
 116. Barbaresi WJ, Katusic SK, Colligan RC, Weaver AL, Leibson CL, Jacobsen SJ. Long-term stimulant medication treatment of attention-deficit/hyperactivity disorder: results from a population-based study. *J Dev Behav Pediatr*. 2006;27:1-10.
 117. Klimkeit E, Graham C, Lee P, Morling M, Russo D, Tonge B. Children should be seen and heard: self-report of feelings and behaviours in primary-school-age children with ADHD. *J Atten Disord*. 2006;10:181-191.
 118. Faraone S, Giefer E. Long-term effects of methylphenidate transdermal delivery system treatment of ADHD on growth. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1138-1147.
 119. Faraone SV, Biederman J, Monuteaux M, Spencer T. Long-term effects of extended-release mixed amphetamine salts treatment of attention-deficit/hyperactivity disorder on growth. *J Child Adolesc Psychopharmacol*. 2005;15:191-202.
 120. Spencer TJ, Faraone SV, Biederman J, et al. Does prolonged therapy with a long-acting stimulant suppress growth in children with ADHD? *J Am Acad Child Adolesc Psychiatry*. 2006;45:527-537.
 121. Swanson JM, Elliott GR, Greenhill LL, et al. Effects of stimulant medication on growth rates across 3 years in the MTA follow-up. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1015-1027.
 122. Charach A, Figueroa M, Chen S, Ickowicz A, Schachar R. Stimulant treatment over 5 years: effects on growth. *J Am Acad Child Adolesc Psychiatry*. 2006;45:415-421.
 123. Villalba M, Racoosin J. Follow up of AERS search identifying cases of sudden death occurring with drugs used for the treatment of attention deficit hyperactivity disorder (ADHD). Office of Drug Safety. http://www.fda.gov/OHRMS/DOCKETS/AC/06/briefing/2006-4210b_07_01_safetyreview.pdf. Published February 28, 2006. Accessed February 15, 2008.
 124. Findling RL, Biederman J, Wilens TE, et al; for the SLI381.301 and .302 Study Groups. Short- and long-term cardiovascular effects of mixed amphetamine salts extended release in children. *J Pediatr*. 2005;147:348-354.
 125. Israeli E, Schochat T, Korzets Z, Tekes-Manova D, Bernheim J, Golan E. Prehypertension and obesity in adolescents: a population study. *Am J Hypertens*. 2006;19:708-712.
 126. Julius S. Blood pressure lowering only or more? Has the jury reached its verdict? *Am J Cardiol*. 2007;100:32J-37J.
 127. Assessing the safety of ADHD medications. CME-Certified Monograph. Medical Crossfire/Liberty Communications Network. Vol. 7. <http://www.medicalcrossfire.com/ADHD.pdf>. Published October 2006. Accessed February 15, 2008.
 128. American Academy of Pediatrics. *What is ADHD Anyway? Questions from Teens* [Trifold brochure]. Elk Grove Village, IL: American Academy of Pediatrics; 2005.
 129. Child and Adults with attention-deficit/hyperactivity disorder income and expenditures 2006-2007. <http://www.chadd.org/AM/Template.cfm?Section=Reports1&Template=/CM/ContentDisplay.cfm&ContentID=5148>. Accessed February 11, 2008.
 130. Cosgrove L, Krinsky S, Vijayaraghavan M, Schneider L. Financial ties between DSM-IV panel members and the pharmaceutical industry. *Psychother Psychosom*. 2006;75:154-160.
 131. Okie S. ADHD in adults. *N Engl J Med*. 2006;354:2637-2644.
 132. Babcock L, Loewenstein G. Explaining bargaining impasse: the role of self-serving biases. *J Econ Perspect*. 1997;11:109-126.
 133. Katz D, Caplan AL, Merz JF. All gifts large and small: toward an understanding of the ethics of pharmaceutical industry gift-giving. *Am J Bioeth*. 2003;3:39-46.
 134. Tungaraza T, Poole R. Influence of drug company authorship and sponsorship on drug trial outcomes. *Br J Psychiatry*. 2007;191:82-83.
 135. Perlis RH, Perlis CS, Wu Y, Hwang C, Joseph M, Nierenberg AA. Industry sponsorship and financial conflict of interest in the reporting of clinical trials in psychiatry. *Am J Psychiatry*. 2005;162:1957-1960.
 136. Lesser LI, Ebbeling CB, Goozner M, Wypij D, Ludwig DS. Relationship between funding source and conclusion among nutrition-related scientific articles. *PLoS Med*. 2007;4:e5.
 137. Procyshyn RM, Chau A, Fortin P, Jenkins W. Prevalence and outcomes of pharmaceutical industry-sponsored clinical trials involving clozapine, risperidone, or olanzapine. *Can J Psychiatry*. 2004;49:601-606.
 138. Turner EH, Matthews AM, Lindartos E, Tell RA, Rosenthal R. Selective publication of antidepressant trials and its influence on apparent efficacy. *N Engl J Med*. 2008;358:252-260.
 139. Lexchin J. Interactions between physicians and the pharmaceutical industry: what does the literature say? *CMAJ*. 1993;149:1401-1407.
 140. Brennan TA, Rothman DJ, Blank L, et al. Health industry practices that create conflicts of interest: a policy proposal for academic medical centers. *JAMA*. 2006;295:429-433.
 141. Campbell EG, Weissman JS, Ehringhaus S, et al. Institutional academic-industry relationships. *JAMA*. 2007;298:1779-1786.