

Standard errors entered as standard deviations: four analyses of electronic versus face-to-face cognitive behavioural therapy in Luo et al. (2020), its addendum, corrigendum and letter of concern

Constantin Yves Plessen^{1*}

¹ Psychotherapy and Psychotherapy Research (CBT), Center for Psychotherapy, University of Graz, Graz, Austria

* constantin.plessen@uni-graz.at



Abstract

All four published estimates recompute from their inputs; only the letter's inputs are standard deviations. 2388 specifications give g from -0.04 to 0.36 ; no interval favours eCBT.

Copyright attribution to authors.

This work is a submission to Journal of Robustness Reports.

License information to appear upon publication.

Publication information to appear upon publication.

Received Date

Accepted Date

Published Date

1 Target article

C. Luo, N. Sanger, N. Singhal, K. Patrick, I. Shams, H. Shahid, P. Hoang, J. Schmidt, J. Lee, S. Haber, M. Puckering, N. Buchanan *et al.*, *A comparison of electronically-delivered and face to face cognitive behavioural therapies in depressive disorders: A systematic review and meta-analysis*, *EClinicalMedicine* **24**, 100442 (2020), doi:[10.1016/j.eclinm.2020.100442](https://doi.org/10.1016/j.eclinm.2020.100442)

1 Goal

The target pooled 14 trials of electronic (eCBT) against face-to-face cognitive behavioural therapy for depression: -1.73 [95% CI $-2.72, -0.74$], $I^2 = 98\%$, read as eCBT more effective [1]. An addendum reinterpreted it as less change for eCBT [2]; a corrigendum gave -0.92 [$-1.76, -0.09$] and declared its conclusions unchanged [3], although the original abstract still claims the opposite; a letter of concern re-extracted the trials and found 0.20 [$-0.05, 0.44$] [4] (positive favours face-to-face CBT throughout; the letter prints the opposite sign). I asked whether each estimate follows from its document's inputs, and how robust the effect is.

2 Methods

I recomputed each estimate from its document's inputs; reproduction means agreement at printed precision. I re-extracted all trials from the primary reports and cross-checked against both published extractions. Hedges' g was pooled with metafor [5] in a metaMultiverse [6] specification curve [7, 8] over six factors: data source (corrigendum, letter, mine), effect-size metric [9, 10], instrument, inclusion set, estimator and the handling of the change-score-only trial [11].

3 Results

Kriston et al. identified each arm's entered "SD" as the standard error (SE) of the mean change [4]. I confirm it in 27 of 28 arms; RevMan's formula on these entries returns -1.73 , -0.79 and -0.92 exactly, with I^2 of 97 to 98%. The letter's estimate reproduces exactly from its supplement.

Two findings are not in the letter of concern. First, Wright 2005 [11] reports baseline and change scores only. The original correctly derived the post-treatment mean as baseline minus endpoint change; the corrigendum "corrected" it to the week-4 completer change score, wrong column and time point, which the letter did not report. Second, the inconsistency for which the original downgraded its evidence is an artefact of the SEs: with SDs, I^2 is 68% (Figure 1A), 0% without the MoodGYM trials. Across 2388 specifications (Figure 1B) over three data sources, g spans -0.04 to 0.36 (median 0.17 , IQR 0.13 to 0.21); only 12 intervals exclude zero, all favouring face-to-face CBT. Data source shifts the median by at most 0.03 : with SDs, the corrigendum's table gives 0.10 $[-0.13, 0.33]$, my re-extraction 0.15 $[-0.09, 0.39]$, or 0.20 $[-0.04, 0.45]$ under the letter's specification. Inclusion is the largest source of spread: without the MoodGYM trials ($n = 19, 44$) the median is 0.05 ; no other factor shifts it by more than 0.12 . No specification comes within 0.5 of the published -0.92 .

4 Conclusion

All four published estimates recompute exactly from their printed inputs, but the original, addendum and corrigendum are obtained only by entering standard errors as standard deviations; with SDs, three independent extractions and every defensible analysis give estimates not comparable in magnitude to the published ones. This reanalysis undercuts the target's claim that eCBT is more effective and the corrigendum's claim of unchanged conclusions; it corroborates the letter. As in the letter, every quantity changes once SDs replace SEs: estimates, intervals, heterogeneity. The interval includes zero in 99.5% of specifications: the evidence is compatible with small differences in either direction; the slight advantage for face-to-face CBT rests on two small MoodGYM trials.

Acknowledgments and Disclosures

Reproducibility For each of the four documents: *Original* [1]: the printed -1.73 $[-2.72, -0.74]$ is recovered exactly from the forest-plot inputs with DerSimonian-Laird and RevMan's variance, but those inputs are standard errors, so the recovered value is not a standardised mean difference. *Addendum* [2]: its re-reading of -1.73 is the same number, and its -0.79 $[-1.60, 0.02]$ with Choi 2014 removed recovers to within rounding, with the same caveat; its

A. Input: the corrigendum's Table 1, the quantity entered as "SD", and the effect sizes it implies

Δ = mean change (pre minus post); SEM = $\sqrt{SD_{pre}^2/n + SD_{post}^2/n}$, the value entered as each arm's SD (purple);

SD_{Δ} = SD of the change score, $r = 0.6$. Published = Δ / SEM as in the corrected forest plot, sign reversed. Wright 2005 as in the corrigendum.

Study	n (e / f)	eCBT			Face-to-face CBT			Hedges' g (SE), positive favours face-to-face		
		Δ	SEM	SD_{Δ}	Δ	SEM	SD_{Δ}	Published	Change SMD	Post SMD
Andersson 2013	33 / 36	10.0	1.9	7.9	7.0	1.6	6.4	-1.70 (0.28)	-0.41 (0.24)	-0.39 (0.24)
Choi 2014	56 / 63	9.9	1.3	6.3	13.7	1.3	6.3	2.97 (0.27)	0.60 (0.19)	-0.05 (0.18)
Glueckauf 2012	7 / 7	8.0	3.5	7.8	2.8	3.5	6.1	-1.76 (0.66)	-0.69 (0.55)	-0.95 (0.56)
Himelhoch 2013	16 / 18	6.3	2.5	6.9	8.7	2.1	5.8	1.02 (0.37)	0.37 (0.35)	0.05 (0.34)
Kalapatapu 2014	50 / 53	9.4	1.5	7.4	10.0	1.1	5.8	0.45 (0.20)	0.09 (0.20)	0.12 (0.20)
Kay-Lambkin 2009	32 / 35	11.5	2.8	10.1	18.3	2.4	9.1	2.59 (0.34)	0.70 (0.25)	0.04 (0.24)
Luxton 2016	62 / 59	13.0	2.2	10.9	18.0	2.2	10.7	2.26 (0.23)	0.45 (0.18)	0.15 (0.18)
Mohr 2012	163 / 162	9.2	0.7	6.0	10.3	0.8	6.7	1.46 (0.13)	0.17 (0.11)	0.13 (0.11)
Nelson 2003	14 / 14	7.7	2.9	8.0	1.9	3.9	9.5	-1.64 (0.45)	-0.63 (0.39)	-0.54 (0.38)
Poppelars 2016	51 / 50	4.7	2.4	11.0	4.0	2.4	10.9	-0.29 (0.20)	-0.06 (0.20)	-0.11 (0.20)
Sethi 2010	9 / 10	0.7	3.4	7.5	11.8	1.9	4.1	3.91 (0.85)	1.79 (0.54)	2.22 (0.58)
Sethi 2013	23 / 21	5.5	1.2	4.0	13.6	1.6	5.5	6.37 (0.77)	1.67 (0.35)	1.38 (0.34)
Wagner 2014	32 / 30	10.6	2.1	8.0	11.1	2.1	7.4	0.24 (0.26)	0.07 (0.25)	0.01 (0.25)
Wright 2005	15 / 15	20.4	3.0	7.3	14.7	2.8	7.0	-1.95 (0.45)	-0.78 (0.38)	0.12 (0.37)
Pooled (DL) [95% CI]								0.92 [0.09, 1.76]	0.23 [-0.05, 0.51]	0.10 [-0.13, 0.33]
I^2								97%	78%	68%

B. Output: every reasonable analysis of the same 14 trials

2388 distinct specifications, sorted by estimate. Pooled Hedges' g with 95% CI. Positive favours face-to-face CBT.

Dotted lines: the published pooled estimates, each placed as its own paper interpreted it; all divided mean differences by standard errors instead of SDs.




Figure 1: Input and output of the reanalysis. **A:** the corrigendum's Table 1 per study. Δ is the mean change (pre minus post); $SEM = \sqrt{SD_{pre}^2/n + SD_{post}^2/n}$ is what the authors entered as each arm's "SD" (purple); SD_{Δ} is the SD of the change score at pre-post $r = 0.6$, the value Luxton 2016 reports. Effect sizes are Hedges' g (SE), positive favouring face-to-face CBT: published (Δ/SEM , sign reversed from the printed plot), change-score SMD, post-test SMD; pooled rows are DerSimonian-Laird. **B:** pooled g with 95% CI for 2388 distinct specifications sorted by estimate, coloured by data source (named in the same colours in the first dashboard block), with the published estimates placed as each paper read them (dotted) and each specification's factor levels below. Change-score SMD uses $r \in \{0.3, 0.6\}$ (the letter's value; Luxton's), standardised mean change $r \in \{0.3, 0.5, 0.7, 0.9\}$. Levels implied by other rows are omitted. Full table and source comparison: online supplement.

post-intervention analysis (0.21 [−0.05, 0.46]) could not be reproduced: the corrigendum’s table gives 0.10 [−0.13, 0.33] (DerSimonian-Laird, RevMan variance) and the pre-correction values in the corrigendum’s footnotes give 0.14 [−0.13, 0.40], or 0.19 [−0.09, 0.46] without Choi 2014. *Corrigendum* [3]: −0.92 [−1.76, −0.09] recovers exactly from its forest-plot values, which disagree with its own Table 1 in four cells and whose Wright 2005 entries are week-4 change scores; the computation is correct but the inputs are standard errors. *Letter of concern* [4]: the printed −0.20 [−0.44, 0.05], $Q = 41.88$, $I^2 = 69\%$, reproduces exactly from its supplementary table with DerSimonian-Laird, change-score SMD at $r = 0.3$ and RevMan’s small-sample correction; all 14 per-study values match at two decimals, and the pooled estimate is stable under REML (Hartung-Knapp interval −0.13 to 0.53, positive favouring face-to-face). No authors’ reply to the letter was found.

Code and Data Availability All data (published tables, screenshots of the primary-study tables, the reconciled extraction with per-value provenance), numbered R scripts, outputs and an extended version of this report are in the FAIR [12] repository at <https://doi.org/10.5281/zenodo.23162504> (GitHub: <https://github.com/cyplessen/ecbt-fcbt-reanalysis>). Effect sizes were computed with metafor [5]; the specification curve used metaMultiverse [6], whose pre-post effect-size support (version 0.3.1) was developed for this reanalysis; the project-level implementation is included in the repository. The letter’s supplementary workbook is transcribed verbatim in data/.

Author Contributions C.Y.P.: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Visualization, Writing - original draft, Writing - review & editing.

Funding The author received no financial support for the research, authorship, and/or publication of this article.

Conflicts of Interest The author maintains the metaMultiverse R package used in the analysis [6], and published two blog posts on the target article in 2020, since removed, which served as extraction records for this work. No other competing interests.

References

- [1] C. Luo, N. Sanger, N. Singhal, K. Patrick, I. Shams, H. Shahid, P. Hoang, J. Schmidt, J. Lee, S. Haber, M. Puckering, N. Buchanan *et al.*, *A comparison of electronically-delivered and face to face cognitive behavioural therapies in depressive disorders: A systematic review and meta-analysis*, *EClinicalMedicine* **24**, 100442 (2020), doi:10.1016/j.eclinm.2020.100442.
- [2] C. Luo, N. Sanger, N. Singhal, K. Patrick, I. Shams, H. Shahid, P. Hoang, J. Schmidt, J. Lee, S. Haber, M. Puckering, N. Buchanan *et al.*, *Addendum to ‘A comparison of electronically-delivered and face to face cognitive behavioural therapies in depressive disorders: A systematic review and meta-analysis’*, *EClinicalMedicine* **27**, 100580 (2020), doi:10.1016/j.eclinm.2020.100580.
- [3] N. Sanger, Z. Samaan and L. Thabane, *Corrigendum to ‘A comparison of electronically-delivered and face to face cognitive behavioural therapies in depressive disorders: A systematic review and meta-analysis’*, *eClinicalMedicine* **41**, 101182 (2021), doi:10.1016/j.eclinm.2021.101182.

- 95 [4] L. Kriston, S. Liebherz and M. Köhnen, *Concerns about ‘A comparison of electronically-*
96 *delivered and face to face cognitive behavioural therapies in depressive disorders:*
97 *A systematic review and meta-analysis’*, *eClinicalMedicine* **54**, 101763 (2022),
98 doi:[10.1016/j.eclinm.2022.101763](https://doi.org/10.1016/j.eclinm.2022.101763).
- 99 [5] W. Viechtbauer, *Conducting meta-analyses in R with the metafor package*, *Journal of*
100 *Statistical Software* **36**(3), 1 (2010), doi:[10.18637/jss.v036.i03](https://doi.org/10.18637/jss.v036.i03).
- 101 [6] C. Y. Plessen, *metaMultiverse: Principled Multiverse Meta-Analysis in R*,
102 doi:[10.5281/zenodo.22706036](https://doi.org/10.5281/zenodo.22706036), R package version 0.3.1 (2026).
- 103 [7] U. Simonsohn, J. P. Simmons and L. D. Nelson, *Specification curve analysis*, *Nature*
104 *Human Behaviour* **4**, 1208 (2020), doi:[10.1038/s41562-020-0912-z](https://doi.org/10.1038/s41562-020-0912-z).
- 105 [8] C. Y. Plessen, E. Karyotaki, C. Miguel, M. Ciharova and P. Cuijpers, *Exploring the efficacy*
106 *of psychotherapies for depression: a multiverse meta-analysis*, *BMJ Mental Health* **26**(1),
107 e300626 (2023), doi:[10.1136/bmjment-2022-300626](https://doi.org/10.1136/bmjment-2022-300626).
- 108 [9] B. J. Becker, *Synthesizing standardized mean-change measures*, *British Journal*
109 *of Mathematical and Statistical Psychology* **41**, 257 (1988), doi:[10.1111/j.2044-](https://doi.org/10.1111/j.2044-8317.1988.tb00901.x)
110 [8317.1988.tb00901.x](https://doi.org/10.1111/j.2044-8317.1988.tb00901.x).
- 111 [10] S. B. Morris, *Estimating effect sizes from pretest-posttest-control group designs*, *Organiza-*
112 *tional Research Methods* **11**, 364 (2008), doi:[10.1177/1094428106291059](https://doi.org/10.1177/1094428106291059).
- 113 [11] J. H. Wright, A. S. Wright, A. M. Albano, M. R. Basco, L. J. Goldsmith, T. Raffield
114 and M. W. Otto, *Computer-assisted cognitive therapy for depression: Maintaining effi-*
115 *cacy while reducing therapist time*, *American Journal of Psychiatry* **162**, 1158 (2005),
116 doi:[10.1176/appi.ajp.162.6.1158](https://doi.org/10.1176/appi.ajp.162.6.1158).
- 117 [12] M. D. Wilkinson, M. Dumontier, I. J. Aalbersberg *et al.*, *The FAIR guiding principles*
118 *for scientific data management and stewardship*, *Scientific Data* **3**, 160018 (2016),
119 doi:[10.1038/sdata.2016.18](https://doi.org/10.1038/sdata.2016.18).