

Revisiting the Approximate Carathéodory Problem via the Frank-Wolfe Algorithm

Cyrille W. Combettes

Joint work with Sebastian Pokutta



Conference on Data Science and Optimization
Fields Institute, Toronto, Canada

November 20th, 2019

Table of contents

- 1 Introduction
- 2 The Frank-Wolfe algorithm
- 3 Sparsity bounds via Frank-Wolfe
- 4 The Fully-Corrective Frank-Wolfe algorithm

Theorem (Carathéodory [1907])


Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}




Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}




Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}




Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}
- Can we reduce $n + 1$ when we can afford an ϵ -approximation?




Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}
- Can we reduce $n + 1$ when we can afford an ϵ -approximation?




Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}
- Can we reduce $n + 1$ when we can afford an ϵ -approximation?




Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}
- Can we reduce $n + 1$ when we can afford an ϵ -approximation?




Introduction

Theorem (Carathéodory [1907])

Every point in the convex hull of a set $\mathcal{V} \subset \mathbb{R}^n$ is the convex combination of at most $n + 1$ points in \mathcal{V} .

- In \mathbb{R}^2 , every point in $\text{conv}(\mathcal{V})$ is the convex combination of at most 3 points in \mathcal{V}
- Can we reduce $n + 1$ when we can afford an ϵ -approximation?
- Define the *sparsity* of x as the minimum number of vertices necessary to form x as a convex combination



The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

- Applications in game theory, combinatorial optimization, and machine learning

The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

- Applications in game theory, combinatorial optimization, and machine learning

Theorem

Let $p \geq 2$. Then there exists $x \in \text{conv}(\mathcal{V})$ with sparsity $\mathcal{O}(pD_p^2/\epsilon^2)$ satisfying $\|x - x^\|_p \leq \epsilon$, where $D_p = \sup_{v,w \in \mathcal{V}} \|w - v\|_p$.*

The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

- Applications in game theory, combinatorial optimization, and machine learning

Theorem

Let $p \geq 2$. Then there exists $x \in \text{conv}(\mathcal{V})$ with sparsity $\mathcal{O}(pD_p^2/\epsilon^2)$ satisfying $\|x - x^\|_p \leq \epsilon$, where $D_p = \sup_{v,w \in \mathcal{V}} \|w - v\|_p$.*

- This result is independent of the space dimension n

The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

- Applications in game theory, combinatorial optimization, and machine learning

Theorem

Let $p \geq 2$. Then there exists $x \in \text{conv}(\mathcal{V})$ with sparsity $\mathcal{O}(pD_p^2/\epsilon^2)$ satisfying $\|x - x^*\|_p \leq \epsilon$, where $D_p = \sup_{v,w \in \mathcal{V}} \|w - v\|_p$.

- This result is independent of the space dimension n
- The bound is tight [Mirrokni et al., 2017]

The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

- Applications in game theory, combinatorial optimization, and machine learning

Theorem

Let $p \geq 2$. Then there exists $x \in \text{conv}(\mathcal{V})$ with sparsity $\mathcal{O}(pD_p^2/\epsilon^2)$ satisfying $\|x - x^\|_p \leq \epsilon$, where $D_p = \sup_{v,w \in \mathcal{V}} \|w - v\|_p$.*

- This result is independent of the space dimension n
- The bound is tight [Mirrokni et al., 2017]
- Probabilistic proofs by Pisier [1981] and Barman [2015]

The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

- Applications in game theory, combinatorial optimization, and machine learning

Theorem

Let $p \geq 2$. Then there exists $x \in \text{conv}(\mathcal{V})$ with sparsity $\mathcal{O}(pD_p^2/\epsilon^2)$ satisfying $\|x - x^*\|_p \leq \epsilon$, where $D_p = \sup_{v,w \in \mathcal{V}} \|w - v\|_p$.

- This result is independent of the space dimension n
- The bound is tight [Mirrokni et al., 2017]
- Probabilistic proofs by Pisier [1981] and Barman [2015]
- Deterministic proof by Mirrokni et al. [2017] using mirror descent [Nemirovsky and Yudin, 1983] on the dual problem

The approximate Carathéodory problem

Problem

Find $x \in \text{conv}(\mathcal{V})$ with high sparsity satisfying $\|x - x^*\|_p \leq \epsilon$.

- Applications in game theory, combinatorial optimization, and machine learning

Theorem

Let $p \geq 2$. Then there exists $x \in \text{conv}(\mathcal{V})$ with sparsity $\mathcal{O}(pD_p^2/\epsilon^2)$ satisfying $\|x - x^\|_p \leq \epsilon$, where $D_p = \sup_{v,w \in \mathcal{V}} \|w - v\|_p$.*

- This result is independent of the space dimension n
- The bound is tight [Mirrokni et al., 2017]
- Probabilistic proofs by Pisier [1981] and Barman [2015]
- Deterministic proof by Mirrokni et al. [2017] using mirror descent [Nemirovsky and Yudin, 1983] on the dual problem
- Can we solve $\min_{x \in \text{conv}(\mathcal{V})} \|x - x^*\|_p$ by sequentially picking up vertices?


The Frank-Wolfe algorithm

Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

$$f(x) = \|x - x^*\|_2^2$$




The Frank-Wolfe algorithm

Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

$$f(x) = \|x - x^*\|_2^2$$




The Frank-Wolfe algorithm

Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t (v_t - x_t)$
 - 5: **end for**
-

$$f(x) = \|x - x^*\|_2^2$$




The Frank-Wolfe algorithm

Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

$$f(x) = \|x - x^*\|_2^2$$




The Frank-Wolfe algorithm

Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

$$f(x) = \|x - x^*\|_2^2$$




The Frank-Wolfe algorithm

Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

$$f(x) = \|x - x^*\|_2^2$$




The Frank-Wolfe algorithm

Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

$$f(x) = \|x - x^*\|_2^2$$



The Frank-Wolfe algorithm


Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

- FW minimizes f over $\text{conv}(\mathcal{V})$ by sequentially picking up vertices

$$f(x) = \|x - x^*\|_2^2$$



The Frank-Wolfe algorithm


Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

- FW minimizes f over $\text{conv}(\mathcal{V})$ by sequentially picking up vertices
- The final iterate x_T has sparsity at most $T + 1$

$$f(x) = \|x - x^*\|_2^2$$



The Frank-Wolfe algorithm


Frank & Wolfe [1956], Levitin & Polyak [1966]

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

- FW minimizes f over $\text{conv}(\mathcal{V})$ by sequentially picking up vertices
- The final iterate x_T has sparsity at most $T + 1$
- This is like a greedy method for the approximate Carathéodory problem!

$$f(x) = \|x - x^*\|_2^2$$



The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$

The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$
- Then x_T has sparsity at most $T + 1$ and satisfies $\|x_T - x^*\|_p \leq \epsilon$

The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$
- Then x_T has sparsity at most $T + 1$ and satisfies $\|x_T - x^*\|_p \leq \epsilon$
- Classical convergence results of FW and variants require smoothness and/or strong convexity of f w.r.t. a norm $\|\cdot\|$

The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$
- Then x_T has sparsity at most $T + 1$ and satisfies $\|x_T - x^*\|_p \leq \epsilon$
- Classical convergence results of FW and variants require **smoothness** and/or strong convexity of f w.r.t. a norm $\|\cdot\|$

$$f(y) - f(x) - \langle \nabla f(x), y - x \rangle \leq \frac{L}{2} \|y - x\|^2$$

The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$
- Then x_T has sparsity at most $T + 1$ and satisfies $\|x_T - x^*\|_p \leq \epsilon$
- Classical convergence results of FW and variants require **smoothness** and/or **strong convexity** of f w.r.t. a norm $\|\cdot\|$

$$\frac{S}{2} \|y - x\|^2 \leq f(y) - f(x) - \langle \nabla f(x), y - x \rangle \leq \frac{L}{2} \|y - x\|^2$$

The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$
- Then x_T has sparsity at most $T + 1$ and satisfies $\|x_T - x^*\|_p \leq \epsilon$
- Classical convergence results of FW and variants require **smoothness** and/or **strong convexity** of f w.r.t. a norm $\|\cdot\|$

$$\frac{S}{2} \|y - x\|^2 \leq f(y) - f(x) - \langle \nabla f(x), y - x \rangle \leq \frac{L}{2} \|y - x\|^2$$

- For $p \geq 2$, $f(x) = \|x - x^*\|_p^2$ is smooth w.r.t. $\|\cdot\|_p$ but it is strongly convex only when $p = 2$

The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$
- Then x_T has sparsity at most $T + 1$ and satisfies $\|x_T - x^*\|_p \leq \epsilon$
- Classical convergence results of FW and variants require **smoothness** and/or **strong convexity** of f w.r.t. a norm $\|\cdot\|$

$$\frac{S}{2} \|y - x\|^2 \leq f(y) - f(x) - \langle \nabla f(x), y - x \rangle \leq \frac{L}{2} \|y - x\|^2$$

- For $p \geq 2$, $f(x) = \|x - x^*\|_p^2$ is smooth w.r.t. $\|\cdot\|_p$ but it is strongly convex only when $p = 2$
- Replace strong convexity with a weaker condition: the PL inequality w.r.t. $\|\cdot\|$ [Polyak, 1963, Łojasiewicz, 1963]

$$f(x) - \min_{\mathbb{R}^n} f \leq \frac{1}{2\mu} \|\nabla f(x)\|_*^2$$

The approximate Carathéodory problem via FW

- Apply FW to $f(x) = \|x - x^*\|_p^2$ and count the number of iterations T to achieve ϵ^2 -convergence: $\|x_T - x^*\|_p^2 \leq \epsilon^2$
- Then x_T has sparsity at most $T + 1$ and satisfies $\|x_T - x^*\|_p \leq \epsilon$
- Classical convergence results of FW and variants require **smoothness** and/or **strong convexity** of f w.r.t. a norm $\|\cdot\|$

$$\frac{S}{2} \|y - x\|^2 \leq f(y) - f(x) - \langle \nabla f(x), y - x \rangle \leq \frac{L}{2} \|y - x\|^2$$

- For $p \geq 2$, $f(x) = \|x - x^*\|_p^2$ is smooth w.r.t. $\|\cdot\|_p$ but it is strongly convex only when $p = 2$
- Replace strong convexity with a weaker condition: the PL inequality w.r.t. $\|\cdot\|$ [Polyak, 1963, Łojasiewicz, 1963]

$$f(x) - \min_{\mathbb{R}^n} f \leq \frac{1}{2\mu} \|\nabla f(x)\|_*^2$$

- For $p \geq 2$, $f(x) = \|x - x^*\|_p^2$ is $2(p-1)$ -smooth and 2-PL w.r.t. $\|\cdot\|_p$

Sparsity bounds via FW convergence rates

Levitin & Polyak [1966], Guélat & Marcotte [1986], Jaggi [2013], Garber & Hazan [2015]

- $p \geq 2$
- $\mathcal{C} \subset \mathbb{R}^n$ be a compact convex set
- $\mathcal{V} \subseteq \partial\mathcal{C}$ be the compact set of interest (e.g., $\mathcal{C} = \text{conv}(\mathcal{V})$)
- We want a sparse approximate convex decomposition of $x^* \in \text{conv}(\mathcal{V})$

Sparsity bounds via FW convergence rates

Levitin & Polyak [1966], Guélat & Marcotte [1986], Jaggi [2013], Garber & Hazan [2015]

- $p \geq 2$
- $\mathcal{C} \subset \mathbb{R}^n$ be a compact convex set
- $\mathcal{V} \subseteq \partial\mathcal{C}$ be the compact set of interest (e.g., $\mathcal{C} = \text{conv}(\mathcal{V})$)
- We want a sparse approximate convex decomposition of $x^* \in \text{conv}(\mathcal{V})$

Assumptions	FW rate	Sparsity bound
–	$\frac{4(p-1)D_p^2}{t+2}$	$\frac{4(p-1)D_p^2}{\epsilon^2} = \mathcal{O}\left(\frac{pD_p^2}{\epsilon^2}\right)$
\mathcal{C} is S_p -strongly convex	$\frac{\max\{9(p-1)D_p^2, 1152(p-1)^2/S_p^2\}}{(t+2)^2}$	$\mathcal{O}\left(\frac{\sqrt{p}D_p + p/S_p}{\epsilon}\right)$
$x^* \in \text{relint}_p(\mathcal{C})$ with radius r_p	$\left(1 - \frac{1}{p-1} \frac{r_p^2}{D_p^2}\right)^t \epsilon_0$	$\mathcal{O}\left(\frac{pD_p^2}{r_p^2} \ln\left(\frac{1}{\epsilon}\right)\right)$

Sparsity bounds via FW convergence rates

Levitin & Polyak [1966], Guélat & Marcotte [1986], Jaggi [2013], Garber & Hazan [2015]

- $p \geq 2$
- $\mathcal{C} \subset \mathbb{R}^n$ be a compact convex set
- $\mathcal{V} \subseteq \partial\mathcal{C}$ be the compact set of interest (e.g., $\mathcal{C} = \text{conv}(\mathcal{V})$)
- We want a sparse approximate convex decomposition of $x^* \in \text{conv}(\mathcal{V})$

Assumptions	FW rate	Sparsity bound
–	$\frac{4(p-1)D_p^2}{t+2}$	$\frac{4(p-1)D_p^2}{\epsilon^2} = \mathcal{O}\left(\frac{pD_p^2}{\epsilon^2}\right)$
\mathcal{C} is S_p -strongly convex	$\frac{\max\{9(p-1)D_p^2, 1152(p-1)^2/S_p^2\}}{(t+2)^2}$	$\mathcal{O}\left(\frac{\sqrt{p}D_p + p/S_p}{\epsilon}\right)$
$x^* \in \text{relint}_p(\mathcal{C})$ with radius r_p	$\left(1 - \frac{1}{p-1} \frac{r_p^2}{D_p^2}\right)^t \epsilon_0$	$\mathcal{O}\left(\frac{pD_p^2}{r_p^2} \ln\left(\frac{1}{\epsilon}\right)\right)$

- FW adapts to the geometry of the problem to yield higher sparsity

Fully-Corrective Frank-Wolfe algorithm

Algorithm Frank-Wolfe (FW)

- 1: $x_0 \in \mathcal{V}$
 - 2: **for** $t = 0$ **to** $T - 1$ **do**
 - 3: $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$
 - 4: $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$
 - 5: **end for**
-

Fully-Corrective Frank-Wolfe algorithm

Algorithm Frank-Wolfe (FW)

```
1:  $x_0 \in \mathcal{V}$ 
2: for  $t = 0$  to  $T - 1$  do
3:    $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$ 
4:    $x_{t+1} \leftarrow x_t + \gamma_t (v_t - x_t)$ 
5: end for
```

- Selected vertices in FW may be redundant, can we fix this?

Fully-Corrective Frank-Wolfe algorithm

Algorithm Frank-Wolfe (FW)

```
1:  $x_0 \in \mathcal{V}$ 
2: for  $t = 0$  to  $T - 1$  do
3:    $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$ 
4:    $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$ 
5: end for
```

- Selected vertices in FW may be redundant, can we fix this?

Algorithm Fully-Corrective Frank-Wolfe (FCFW)

```
1:  $x_0 \in \mathcal{V}$ 
2:  $\mathcal{S}_0 \leftarrow \{x_0\}$ 
3: for  $t = 0$  to  $T - 1$  do
4:    $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$ 
5:    $\mathcal{S}_{t+1} \leftarrow \mathcal{S}_t \cup \{v_t\}$ 
6:    $x_{t+1} \leftarrow \arg \min_{\text{conv}(\mathcal{S}_{t+1})} f$ 
7: end for
```

Fully-Corrective Frank-Wolfe algorithm

Algorithm Frank-Wolfe (FW)

```
1:  $x_0 \in \mathcal{V}$ 
2: for  $t = 0$  to  $T - 1$  do
3:    $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$ 
4:    $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$ 
5: end for
```

- Selected vertices in FW may be redundant, can we fix this?

Algorithm Fully-Corrective Frank-Wolfe (FCFW)

```
1:  $x_0 \in \mathcal{V}$ 
2:  $\mathcal{S}_0 \leftarrow \{x_0\}$ 
3: for  $t = 0$  to  $T - 1$  do
4:    $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$ 
5:    $\mathcal{S}_{t+1} \leftarrow \mathcal{S}_t \cup \{v_t\}$ 
6:    $x_{t+1} \leftarrow \arg \min_{\text{conv}(\mathcal{S}_{t+1})} f$ 
7: end for
```

Fully-Corrective Frank-Wolfe algorithm

Algorithm Frank-Wolfe (FW)

```
1:  $x_0 \in \mathcal{V}$ 
2: for  $t = 0$  to  $T - 1$  do
3:    $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$ 
4:    $x_{t+1} \leftarrow x_t + \gamma_t(v_t - x_t)$ 
5: end for
```


- Selected vertices in FW may be redundant, can we fix this?

Algorithm Fully-Corrective Frank-Wolfe (FCFW)

```
1:  $x_0 \in \mathcal{V}$ 
2:  $\mathcal{S}_0 \leftarrow \{x_0\}$ 
3: for  $t = 0$  to  $T - 1$  do
4:    $v_t \leftarrow \arg \min_{v \in \mathcal{V}} \langle \nabla f(x_t), v \rangle$ 
5:    $\mathcal{S}_{t+1} \leftarrow \mathcal{S}_t \cup \{v_t\}$ 
6:    $x_{t+1} \leftarrow \arg \min_{\text{conv}(\mathcal{S}_{t+1})} f$ 
7: end for
```


Fully-Corrective Frank-Wolfe algorithm

- We randomly generated 1000 vertices and $x^* \in \text{conv}(\mathcal{V})$
- Here arbitrarily chose $p = 4$




Fully-Corrective Frank-Wolfe algorithm

- Here x^* is generated by a convex combination of only 50 vertices



Fully-Corrective Frank-Wolfe algorithm


- Here x^* is generated by a convex combination of only 50 vertices



- FCFW obtains an *exact* convex decomposition of x^* once it picks up all its vertices


Fully-Corrective Frank-Wolfe algorithm

- FCFW matches the theoretical lower bound!



Fully-Corrective Frank-Wolfe algorithm

- FCFW matches the theoretical lower bound!



- Can we derive a precise convergence rate for FCFW?

Thank you!

<https://arxiv.org/pdf/1911.04415.pdf>

References

- S. Barman. Approximating Nash equilibria and dense bipartite subgraphs via an approximate version of Carathéodory's theorem. *STOC*, 2015.
- C. Carathéodory. Über den Variabilitätsbereich der Koeffizienten von Potenzreihen, die gegebene Werte nicht annehmen. *Mathematische Annalen*, 1907.
- M. Frank and P. Wolfe. An algorithm for quadratic programming. *Naval Research Logistics Quarterly*, 1956.
- D. Garber and E. Hazan. Faster rates for the Frank-Wolfe method over strongly-convex sets. *ICML*, 2015.
- E. S. Levitin and B. T. Polyak. Constrained minimization methods. *USSR Computational Mathematics and Mathematical Physics*, 1966.
- S. Łojasiewicz. Une propriété topologique des sous-ensembles analytiques réels. *Colloques Internationaux du CNRS*, 1963.
- V. Mirrokni, R. Paes Leme, A. Vladu, and S. C.-W. Wong. Tight bounds for approximate Carathéodory and beyond. *ICML*, 2017.
- A. S. Nemirovsky and D. B. Yudin. *Problem complexity and method efficiency in optimization*. Wiley, 1983.
- G. Pisier. Remarques sur un résultat non publié de B. Maurey. *Séminaire d'Analyse Fonctionnelle*, École Polytechnique, 1981.
- B. T. Polyak. Gradient methods for the minimisation of functionals. *USSR Computational Mathematics and Mathematical Physics*, 1963.