

[Skip to main content](#)

Advertisement

Search 

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

Menu

- [Explore journals](#)
- [Get published](#)
- [About BMC](#)
- [Login](#)

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

[Manage Settings](#)[✓ Accept All Cookies](#)

- [Articles](#)
- [Submission Guidelines](#)
- Research
- Open Access
- [Published: 28 January 2015](#)

Increasing the inspiratory time and I:E ratio during mechanical ventilation aggravates ventilator-induced lung injury in mice

- [Holger C Müller-Redetzky](#)¹✉,
- [Matthias Felten](#)¹,
- [Katharina Hellwig](#)¹,
- [Sandra-Maria Wienhold](#)¹,
- [Jan Naujoks](#)¹,
- [Bastian Opitz](#)¹,
- [Olivia Kershaw](#)²,
- [Achim D Gruber](#)²,
- [Norbert Suttorp](#)¹ &
- [\[...\]](#)
- [Martin Witzenrath](#)¹
- [- Show fewer authors](#)

Critical Care volume 19, Article number: 23 (2015) [Cite this article](#)

- 23k Accesses
- 11 Citations

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

Lung-protective ventilation reduced acute respiratory distress syndrome (ARDS) mortality. To minimize ventilator-induced lung injury (VILI), tidal volume is limited, high plateau pressures are avoided, and positive end-expiratory pressure (PEEP) is applied. However, the impact of specific ventilatory patterns on VILI is not well defined. Increasing inspiratory time and thereby the inspiratory/expiratory ratio (I:E ratio) may improve oxygenation, but may also be harmful as the absolute stress and strain over time increase. We thus hypothesized that increasing inspiratory time and I:E ratio aggravates VILI.

Methods

VILI was induced in mice by high tidal-volume ventilation (HV_T 34 ml/kg). Low tidal-volume ventilation (LV_T 9 ml/kg) was used in control groups. PEEP was set to 2 cm H_2O , FiO_2 was 0.5 in all groups. HV_T and LV_T mice were ventilated with either I:E of 1:2 (LV_T 1:2, HV_T 1:2) or 1:1 (LV_T 1:1, HV_T 1:1) for 4 hours or until an alternative end point, defined as mean arterial blood pressure below 40 mm Hg. Dynamic hyperinflation due to the increased I:E ratio was excluded in a separate group of animals. Survival, lung compliance, oxygenation, pulmonary permeability, markers of pulmonary and systemic inflammation (leukocyte differentiation in lung and blood, analyses of pulmonary interleukin-6, interleukin-1 β , keratinocyte-derived chemokine, monocyte chemoattractant protein-1), and histopathologic pulmonary changes were analyzed.

Results

LV_T 1:2 or LV_T 1:1 did not result in VILI, and all individuals survived the ventilation period. HV_T 1:2 decreased lung compliance, increased pulmonary neutrophils and cytokine expression, and evoked marked histologic signs of lung injury. All animals survived. HV_T 1:1 caused further significant worsening of oxygenation, compliance and increased pulmonary proinflammatory cytokine expression, and pulmonary and blood neutrophils. In the HV_T 1:1 group, significant mortality during mechanical ventilation was observed.

Conclusion

According to the “baby lung” concept, mechanical ventilation-associated stress and strain in overinflated regions of ARDS lungs was simulated by using high tidal-volume ventilation. Increase of inspiratory time and I:E ratio significantly aggravated VILI in mice, suggesting an impact of a “stress/strain \times time product” for the pathogenesis of VILI. Thus increasing the inspiratory time and I:E ratio should be critically considered.

Introduction

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

Thus, a certain safety threshold for VILI does not seem to exist, and any effort to minimize VILI further might be of relevance, particularly for the most severely ill ARDS patients [5].

Of note, little is known regarding the impact of ventilator adjustments on VILI. The absolute inspiratory lung strain, which is defined as the end-inspiratory transpulmonary pressure, and the absolute lung strain, defined as V_T/FRC , are central drivers of VILI [5]. We hypothesized that, in addition, the duration of lung stress and strain is relevant, proposing a Time \times Stress/strain product that affects VILI.

Increasing the inspiration-to- expiration ratio (I:E) and thereby the inspiratory time (t_i) of the respiratory cycle can improve oxygenation. The two main underlying mechanisms are probably prolonged gas exchange during inspiration in lung areas that do not take part in gas exchange during expiration, and recruitment of lung tissue due to increased intrinsic PEEP generated by dynamic hyperinflation [6,7]. It is tempting to use this intervention to improve oxygenation at bedside as, despite all efforts made to stabilize an appropriate residual volume by titrating PEEP, almost always, recruitable lung regions remain. Conversely, increasing I:E will result in an increased Time \times Stress/strain product that might aggravate VILI. A previous experimental study [8] and a recently published review of numerous animal models of VILI underscore this hypothesis [9].

In this study, we therefore investigated the impact of I:E on VILI in an experimental VILI mouse model and found that an increased I:E ratio significantly aggravates VILI in mice, suggesting the relevance of a role of a Stress/strain \times Time in the pathogenesis of VILI.

Material and methods

Ethics statement

All animal experiments were approved by institutional (Charité-Universitätsmedizin Berlin) and governmental (Landesamt für Gesundheit und Soziales Berlin; G 0130/12) authorities.

Mice

Female C57BL/6 mice (8 to 10 weeks; 18 to 20 g; Charles River, Sulzfeld, Germany) were used.

Mechanical ventilation

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ Accept All Cookies

A urinary catheter was inserted. Mean arterial blood pressure, heart rate, peripheral oxygen saturation (MouseOx; Starr Life Science Corp., Pittsburgh, PA, USA) and urine output were measured. Mice were ventilated by using a special rodent ventilation system, which continuously recorded airway pressure, respiratory rate, and tidal volume (flexiVent; Scireq, Montreal, QC, Canada). After preparation, a recruitment maneuver was performed (increasing of the airway pressure to 30 cmH₂O), and mice were ventilated for 4 hours with the following ventilator settings:


Low tidal-volume (LV_T) groups

Mice were ventilated with a tidal volume of 9 ml/kg, respiratory rate of 160 per minute, and I:E ratio of either 1:2 or 1:1 (LV_T 1:2; LV_T 1:1). A deep inspiration (30 cmH₂O for 1 second), was performed every 10 minutes, in addition to the applied positive end-expiratory pressure (PEEP) to avoid atelectasis. Notably, this protocol does not cause measurable lung injury in mice [10].

High tidal-volume (HV_T) groups

Mice were ventilated with a tidal volume of 34 ml/kg, respiratory rate of 70 per minute, and I:E ratio of 1:2 or 1:1, respectively (HV_T 1:2; HV_T 1:1).

In all I:E 1:1 groups, the inspiratory time was prolonged by adding an inspiratory hold after completion of lung inflation, thereby leaving pressure and flow acceleration during inspiration identical between the corresponding LV_T and HV_T 1:2 groups, as schematically illustrated in Figure 1.



We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

between the corresponding LV_T and HV_T 1:2 groups.

[Full size image](#) >

To generate baseline values at the beginning of mechanical ventilation, a group of mice referred to as nonventilated control (ctr) was anesthetized and operated as outlined earlier. After an identical recruitment maneuver, ctr mice were ventilated for 5 minutes with adjustments identical to those of the LV_T 1:2 mice. After measurement of baseline lung functions and hemodynamics, the experiment was terminated.

PEEP of 2 cmH₂O and an FiO₂ of 0.5 were applied throughout the experiments in all LV_T and HV_T groups. After 235 minutes of mechanical ventilation (MV), the I:E ratio was switched to 1:2 in all ventilated groups, and the inspired oxygen fraction was increased to an FiO₂ of 1.0. After 240 minutes of MV, mice were killed by rapid exsanguination via the carotid artery catheter.

An alternative end point was defined as decrease of mean arterial blood pressure below 40 mm Hg, as this safely predicts death in this model. The I:E ratio was then switched to 1:2, and the inspired oxygen fraction was increased to an FiO₂ of 1.0. After a further 5 minutes of MV, mice were killed by exsanguination via the carotid artery catheter.

To exclude dynamic hyperinflation in the HV_T 1:1 group, an additional set of animals was ventilated according to the HV_T 1:2 ventilation pattern for 30 minutes, and then the I:E ratio was increased to 1:1 for 30 minutes. This procedure was repeated. Mean airway pressure and dynamic compliance were recorded.

Lung function

After the initial recruitment maneuver, dynamic elastance, resistance, and compliance were measured by using a forced oscillation technique. Measurements were repeated every 10 minutes throughout the experiment. In addition, static-compliance values were determined after exsanguination.

Blood gas analyses

Blood samples were analyzed for p_aO₂ with a blood-gas analyzer (ABL-800; Radiometer, Copenhagen, Denmark). The P/F ratio was calculated as P/F = p_aO₂/FiO₂. Oxygenation Index was calculated as OI = mean airway pressure × FiO₂/p_aO₂.

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

Lungs were flushed. RNA was extracted with TRIzol (Ambion; Life Technologies, Darmstadt, Germany) treatment of lung homogenates and reverse-transcribed by using a high-capacity reverse transcription kit (Applied Biosystems, Life Technologies, Darmstadt, Germany). Quantitative PCR (qRT-PCR) was performed on ABI 7300 by using TaqMan gene-expression assays (Applied Biosystems). The PCR conditions included initial denaturation in one cycle of 2 minutes at 50°C and 10 minutes at 95°C, followed by 40 cycles of 15 seconds at 95°C, 20 seconds at 60°C, and 1 minute at 72°C. The input was normalized to the average expression of GAPDH. Primer and probe sequences are provided in Table S1 in Additional file [1](#).

Leukocytes in BAL fluid and blood

Leukocytes in BALF were differentiated with flow cytometry, according to their side-scatter/forward-scatter characteristics, and CD45, Gr-1, and F4-80 expression (FACSCalibur; BD, Heidelberg, Germany). Blood leukocytes were quantified and differentiated with flow cytometry by using TruCount-Tubes according to cellular side-scatter/forward-scatter characteristics and CD45, Gr-1, and CD3 expression.

Quantification of cytokines

Cytokines were quantified from total protein of flushed and homogenized left lungs and from plasma samples by using the multiplex cytokine assay technique (BioRad, Hercules, CA, USA).

Histopathology

Lung samples were fixed in 4% formaldehyde solution and routinely embedded in paraffin. The 5- μ m-thick sections were cut, dewaxed, and stained with hematoxylin and eosin (H&E) or periodic acid-Schiff (PAS). Histopathology was performed by two European College of Veterinary Pathologists (ECVP) board-certified pathologists, who were blinded to the study groups.

Data analyses

Data are expressed as box-and-whisker plots, or columns (mean \pm SEM). For comparison between groups, a Mann–Whitney *U* test was performed. *P* values <0.05 were considered statistically significant. For survival analyses, a log rank test was applied.

Results

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

[Manage Settings](#)

[✓ Accept All Cookies](#)

Increasing the inspiratory time and I:E ratio during MV increased mortality in VILI

All mice of the low tidal-volume groups ventilated with an I:E ratio of either 1:2 or 1:1 (LV_T 1:2; LV_T 1:1) and of the high tidal-volume group ventilated with an I:E ratio of 1:2 survived the procedures. Increasing the I:E ratio in the HV_T group to 1:1 resulted in premature termination of the experiment in 13 of 14 mice because of dropping of mean arterial blood pressure below 40 mm Hg (alternative end point), corresponding to a 92.1% mortality in the HV_T 1:1 group (Figure 2).

Figure 2

Increasing the inspiratory time and I:E ratio during MV increased mortality in VILI. Mice were mechanically ventilated for 4 hours with either low tidal volume (LV_T 9 ml/kg) or high tidal volume (HV_T 34 ml/kg) and an inspiratory/expiratory ratio of 1:2 or 1:1, respectively. If the mean arterial pressure decreased below 40 mm Hg, the experiment was prematurely terminated, as this predicts death with certainty in this model. $n = 13$ -14 each group; *** $P < 0.001$.

[Full size image](#) >

Increasing the inspiratory time and I:E ratio increased lung injury

Lungs from ctr, LV_T 1:2, and LV_T 1:1 showed no macroscopic or histologic signs of lung injury. HV_T 1:2 ventilated mice exhibited significant histopathologic signs of lung injury, whereas only distinct signs of injury were seen macroscopically. HV_T 1:1 led to dramatic macroscopic and histopathologic lesions. In both HV_T groups but not in the controls, the lung architecture was compromised by severe alveolar collapse and emphysema. Histopathology revealed severe perivascular edema, damage of the alveolar walls with desquamation of alveolar epithelial cells type I and formation of hyaline membranes, increasing numbers of intraalveolar cells (neutrophils and macrophages) and occasional necrosis of bronchiolar epithelium. Severe lung lesions were observed histologically on HE-stained tissues, with no differences seen between both groups (Figure 3 and Figure S2 in Additional file 3).

Figure 3

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

Periodic acid-Schiff (PAS) reaction clearly visualized hyaline membranes diffusely distributed throughout the lung parenchyma, indicative of marked damage of the alveolar membrane. However, in HV_T 1:2 lungs, hyaline membranes appeared only occasionally as continuous thin layers on the alveolar surface, while lungs of the HV_T 1:1 group had thicker hyaline membranes, which commonly completely covered the surface of dilated alveoli (Figure 4). Because pulmonary vascular leakage is a hallmark of ARDS and VILI, we quantified lung permeability by measuring the albumin concentration in bronchoalveolar lavage fluid (BALF) and plasma and by calculating the BALF/plasma albumin ratio. Compared with ctr mice, LV_T 1:2 and LV_T 1:1 did not result in increased permeability. In contrast, HV_T 1:2 mice showed a trend toward increased permeability compared with ctr and LV_T groups, whereas HV_T 1:1 evoked a dramatic increase in pulmonary vascular permeability (Figure 5).

Figure 4

Increasing the inspiratory time and I:E ratio during MV increased histopathologic signs of lung injury. Mice were mechanically ventilated for 4 hours with either low tidal volume (LV_T 9 ml/kg) or high tidal volume (HV_T 34 ml/kg) and an inspiratory/expiratory ratio of 1:2 or 1:1, respectively. An alternative end point was defined as decreasing of mean arterial blood pressure below 40 mm Hg, which predicts death with certainty in this model. Controls (ctr) were subjected to LV_T 1:2 ventilation only during operation and were killed before the 4-hour ventilation protocol started. Histopathology of lungs from ctr, HV_T 1:2, and HV_T 1:1 groups, stained with the periodic acid-Schiff (PAS) reaction, are shown: In contrast to the ctr group, both ventilated groups had damage of the alveolar walls with septal thickening, necrosis, and desquamation of alveolar epithelial cells type I, formation of hyaline membranes (red arrows), and increased numbers of intraalveolar cells (predominantly neutrophils and macrophages, black arrows). PAS reaction highlighted the more severe and more continuous as well as thicker hyaline membranes along the alveolar surfaces of lungs from the HV_T 1:1 group. Top panel: magnification × 200; Bottom panel: magnification × 400. Representative images from each group (*n* = 4 each) are shown.

[Full size image](#) >

Figure 5

Increasing the inspiratory time and I:E ratio during MV increased pulmonary permeability in VILI. Mice were mechanically ventilated for 4 hours with either low tidal volume (LV_T 9 ml/kg) or high tidal volume (HV_T 34 ml/kg) and an inspiratory/expiratory ratio of 1:2 or 1:1, respectively. An alternative end point was defined as decreasing of mean arterial blood pressure below 40 mm Hg, which predicts

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

Peripheral oxygen saturation was measured continuously throughout the experiment. Partial pressure of oxygen in arterial blood and mean airway pressure were measured at the end of the experiment, and P/F ratio as well as oxygenation index (OI) were calculated. Whereas LV_T 1:2, LV_T 1:1, and HV_T 1:2 groups showed stable oxygenation regarding SpO₂ and P/F throughout the experiment (Figure 6A,B), the oxygenation index implied a reduced oxygenation capacity in HV_T 1:2 mice compared to ctr and LV_T groups (Figure 6C). HV_T 1:1 resulted in severe impairment of oxygenation compared with ctr, LV_T, and HV_T 1:2 groups (Figure 6A to C).

Figure 6

Increasing the inspiratory time and I:E ratio reduced oxygenation capacity in VILI. Mice were mechanically ventilated for 4 hours with either low tidal volume (LV_T 9 ml/kg) or high tidal volume (HV_T 34 ml/kg) and an inspiratory/expiratory ratio of 1:2 or 1:1, respectively. An alternative end point was defined as decreasing of mean arterial blood pressure below 40 mm Hg, which predicts death with certainty in this model. Controls (ctr) were subjected to LV_T 1:2 ventilation only during operation and were killed before the 4-hour ventilation protocol started. **(A)** Pulse oximetry revealed stable oxygen saturation in LV_T 1:2, LV_T 1:1, and HV_T 1:2 groups, whereas the HV_T 1:1 ventilated mice developed a decrease of oxygen saturation during the 4-hour ventilation period. End-point measurements of arterial partial pressure of oxygen were performed, and the P/F ratio **(B)**, and the oxygenation index were calculated **(C)**. $n = 13$ to 14 in each group; *** $P < 0.001$.

[Full size image](#) >

Increasing the inspiratory time and I:E ratio deteriorated lung function in VILI

Dynamic elastance was quantified every 10 minutes. While both LV_T groups showed stable elastance during the experiment, a slight increase over time in the HV_T 1:2 group, and a strong increase in the HV_T 1:1 group were observed (Figure 7A). Dynamic and static compliance at the respective end points of the experiment showed impaired compliance in the HV_T 1:2 compared with LV_T 1:2 mice. HV_T 1:1 led to a dramatic decrease in lung compliance (Figure 7B,C).

Figure 7

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

Increasing the inspiratory time and I:E ratio: impact on hemodynamics and markers of tissue perfusion

HV_T animals were ventilated with an alternating I:E ratio (1:2 versus 1:1) in 30-minute intervals for 120 minutes, and mean arterial blood pressure was measured. Changing of the I:E ratio had no impact on mean arterial blood pressure (see Figure S3A in Additional file 4). In animals ventilated for 4 hours (LV_T 1:2, LV_T 1:1, HV_T 1:2, and HV_T 1:1), cumulative urine output and blood lactate levels at the respective experimental end points were quantified. No difference in urine output between the groups was evident, whereas HV_T 1:1 revealed slightly higher lactate levels than the HV_T 1:2 group (see Figure S3 B, C in Additional file 4).

Increasing the inspiratory time and I:E ratio exacerbated the inflammatory response in VILI

We measured transcription of the proinflammatory cytokines IL-1 β , IL-6, KC, and MCP-1 (Figure 8A) and protein concentrations of IL-1 β , IL-6, KC, and MCP-1 in lung homogenates (Figure 8B). HV_T 1:2 increased IL-1 β , IL-6, KC, and MCP-1 mRNA expression compared with ctr and LV_T mice. Expression of most of these proinflammatory mediators was further increased in the HV_T 1:1.

Figure 8

Increasing the inspiratory time and I:E ratio increased the production of proinflammatory cytokines in VILI. Mice were mechanically ventilated for 4 hours with either low tidal volume (LV_T 9 ml/kg) or high tidal volume (HV_T 34 ml/kg) and an inspiratory/expiratory ratio of 1:2 or 1:1, respectively. An alternative end point was defined as decreasing of mean arterial blood pressure below 40 mm Hg, which predicts death with certainty in this model. Controls (ctr) were subjected to LV_T 1:2 ventilation only during operation and were killed before the 4-hour ventilation protocol started. **(A)** mRNA levels of interleukin (IL)-1 β , IL-6, macrophage chemotactic protein (MCP)-1, and keratinocyte-derived cytokine (KC) were measured with quantitative reverse transcription polymerase chain reaction in lung homogenates and normalized to GAPDH levels. **(B)** Protein levels of IL-1 β , IL-6, MCP-1, and KC were determined in lung homogenates by multiplex ELISA technique. $n = 6$ to 8 each group; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

[Full size image](#) >

LV_T 1:2 and LV_T 1:1 resulted in a certain increment of BALF neutrophils compared with ctr mice. In line with the elevation of proinflammatory

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

Hg, which predicts death with certainty in this model. Controls (ctr) were subjected to LV_T 1:2 ventilation only during operation and were killed before the 4-hour ventilation protocol started. The fractions of neutrophils among leukocytes in the BALF (**A**) and in the blood (**B**) at the end point of the experiment are shown. $n = 5$ to 6 each group; $*P < 0.05$, $**P < 0.01$, $##P < 0.01$ versus all.

[Full size image](#) >

Furthermore, HV_T 1:1 exclusively resulted in an increased number of blood neutrophils, indicating a systemic inflammatory response (Figure 2B).

Discussion

We provide strong evidence that increasing the I:E ratio during mechanical ventilation can aggravate VILI, indicating that not only the absolute lung stress and strain but also the time in which the lung is exposed to stress and strain (the Time \times Stress and strain product) may affect the harm of mechanical ventilation.

VILI impairs survival of ARDS patients [2,3]. Besides cyclic opening and closing of lung during tidal ventilation, high airway pressures and high tidal volumes have been identified as main drivers of VILI. More precisely, not absolute airway pressure but the transpulmonary pressure termed lung stress, and not the absolute tidal volume, but its relation to the FRC, termed lung strain, are mechanical determinants of VILI [5]. Besides the amount of lung opening and closing that is correlated with ARDS mortality [11], the concept of intraparenchymal stress raisers during mechanical ventilation may have significant impact on the development of lung injury due to mechanical ventilation in the ARDS patient [5,12].

Recent clinical trials revealed that VILI is particularly relevant in patients with severe ARDS, and therefore, optimizing our ventilation strategies especially for those patients is desirable [13,14].

Lung stress and strain are not equally distributed throughout the respiratory cycle under MV, obviously being higher during inspiration than during expiration. The current study now provides evidence that not only absolute lung stress and strain but also increasing lung stress and strain in relation to the cycle time by prolonging the inspiratory time (t_i) and increasing the I:E ratio aggravate VILI. This is in line with the theory of weighted lung strain during MV by Carioni and colleagues [9].

Physical forces during mechanical ventilation are sensed by the lung and induce a biochemical response characterized by inflammation and endoepithelial permeability, referred to as biotrauma [1,15,16]. Therefore we assessed lung permeability, detailed lung histology and markers of pulmonary inflammation. Even after the short observational time of 4 hours, we detected a significant impact of the increased t_i and I:E ratio on

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

To test the hypothesis of this study, we implemented severely injurious ventilation in mice. The tidal volume of 34 ml/kg was extraordinarily high compared with the standard of lung-protective ventilation with 6 ml/kg in humans with ARDS. At first view, this might outrange the stress and strain applied during MV in ARDS patients. However, residual capacity in ARDS lungs is severely reduced, which is referred to as the “baby lung” of ARDS patients [24]. Notably, the sicker the patient and the lower the oxygenation capacity becomes, the greater is the reduction of the residual capacity and the intention to increase the relative portion of inspiratory time to improve oxygenation. As intensivists do not routinely quantify residual capacity at the bedside, we do not know how much lung strain is generated during MV, despite limiting the tidal volume to 6 ml/kg, especially in very severe ARDS. Further, ARDS is characterized by a high grade of tissue inhomogeneity, in which open, atelectatic, and collapsed but recruitable lung areas coexist, which locally results in lung stress exceeding the measured airway pressure by far [5].

Loss and inactivation of surfactant, a hallmark in ARDS, further aggravate local trauma [1]. Thus, applying high tidal volumes in healthy mouse lungs constitutes a reasonable experimental approach. Further, the currently used model of VILI meets the ATS criteria on lung injury in animals, including the evidence of inflammation, microscopic tissue injury, alteration of alveolar barrier function, and impaired oxygenation [25]. Vice versa, the observation that prolonging t_i in the LV_T groups did not increase detectable lung injury in healthy mice does not argue for the safety of an I:E ratio increase in lung-protective ventilation of ARDS patients.

In this study, it was highly important to control properly factors that might significantly bias the results. (a) Dynamic hyperinflation would increase intrinsic PEEP, which consecutively enhances residual volume and shifts tidal volume upward on the pressure/volume curve, eventually above the upper inflection point, resulting in augmented absolute lung stress and strain. Thus, we adjusted respiratory rates in both HV_T groups to 70 per minute to exclude dynamic hyperinflation in the HV_T I:E 1:1 group. (b) To keep dynamics of lung inflation identical between 1:1 and 1:2 groups, we prolonged t_i by adding an inspiratory hold. This excluded that a difference in pressure acceleration during inspiration or a difference of the total inflation time biased the results of the study. (c) Expiration was most probably similar in the HV_T 1:1 and HV_T 1:2, and in the LV_T 1:1 and LV_T 1:2 groups, respectively. Expiration is a passive process starting after the opening of the expiratory valve with end-inspiratory pressure being the driving force, which was similar in the respective groups. As dynamic hyperinflation could be excluded, exhalation was complete. (d) Respiratory rate, PEEP and FiO_2 were identical in the HV_T and the LV_T groups, respectively and anesthesia and operation procedures were identical in all groups.

Intrathoracic pressure directly affects cardiac function (for an excellent review, see [26,27]), and increased intrathoracic pressure due to increased I:E ratio may decrease cardiac output [28,29]. Reduction of venous return seems to be the central mechanism reducing cardiac output by high intrathoracic pressure, which can be ameliorated by sufficient fluid support. In our study, mice received liberal fluid support to minimize reduction of cardiac output. Blood pressure and urine output were not affected by increased I:E ratio. Nevertheless, lactate levels were slightly elevated in HV_T 1:1 compared with HV_T 1:2 mice. Thus a certain reduction of cardiac output cannot be excluded. However, circulatory failure and resultant shock as cause of the premature

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

Conclusion

The study design applied aimed to provide a proof of concept. The data show that beyond stress and strain, the time in which the lung is exposed to stress and strain (the Time \times Stress and strain product) has dramatic impact on VILI. Therefore, it seems reasonable to minimize the Time \times Strain product during MV. Particularly, increasing the I:E ratio should be critically revised in patients with ARDS.

Key messages

- Increasing inspiratory time und thereby the I:E ratio aggravates VILI.
- Beyond stress and strain, the time during which the lung is exposed to stress and strain (the Time \times Stress and strain product) has dramatic impact on VILI.

Abbreviations

ARDS:

Acute respiratory distress syndrome

BAL:

bronchoalveolar lavage

BALF:

bronchoalveolar lavage fluid

C_{dyn} :

dynamic compliance

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

[Manage Settings](#)

✓ Accept All Cookies

inspiratory fraction of oxygen

FRC:

functional residual capacity

HV_T:

high tidal volume

I:E ratio:

inspiratory-to-expiratory ratio

IL-1 β :

interleukin-1 beta

IL-6:

interleukin-6

KC:

keratinocyte-derived chemokine

LV_T:

low tidal volume

MCP-1:

monocyte chemoattractant protein-1

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](https://www.biomedcentral.com/privacy-policy).

› [Manage Settings](#)

✓ Accept All Cookies

p_aO_2 :

partial pressure of oxygen

PBS:

phosphate-buffered saline

PEEP:

positive end-expiratory pressure

$qRT-PCR$:

quantitative reverse transcription polymerase chain reaction

t_i :

inspiratory time

VILI:

ventilator-induced lung injury

V_T :

tidal volume

References

1. 1.

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ Accept All Cookies

Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. N Engl J Med. 1998;338:347–54.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

3. 3.

The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. N Engl J Med. 2000;342:1301–8.

- [Article](#)
- [Google Scholar](#)

4. 4.

Wolthuis EK, Vlaar APJ, Choi G, Roelofs JJTH, Juffermans NP, Schultz MJ. Mechanical ventilation using non-injurious ventilation settings causes lung injury in the absence of pre-existing lung injury in healthy mice. Crit Care. 2009;13:R1.

- [Article](#)
- [PubMed Central](#)
- [PubMed](#)
- [Google Scholar](#)

5. 5.

Gattinoni L, Carlesso E, Caironi P. Stress and strain within the lung. Curr Opin Crit Care. 2012;18:42–7.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

6. 6.

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

Lee SM, Kim WH, Ahn HJ, Kim JA, Yang MK, Lee CH, et al. The effects of prolonged inspiratory time during one-lung ventilation: a randomised controlled trial. *Anaesthesia*. 2013;68:908–16.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

8. 8.

Casetti AV, Bartlett RH, Hirschl RB. Increasing inspiratory time exacerbates ventilator-induced lung injury during high-pressure/high-volume mechanical ventilation. *Crit Care Med*. 2002;30:2295–9.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

9. 9.

Caironi P, Langer T, Carlesso E, Protti A, Gattinoni L. Time to generate ventilator-induced lung injury among mammals with healthy lungs: a unifying hypothesis. *Intensive Care Med*. 2011;37:1913–20.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

10. 10.

Reiss LK, Kowallik A, Uhlig S. Recurrent recruitment manoeuvres improve lung mechanics and minimize lung injury during mechanical ventilation of healthy mice. *PLoS One*. 2011;6:e24527.

- [Article](#)
- [PubMed Central](#)
- [CAS](#)
- [PubMed](#)

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

Mead J, Takishima T, Leith D. Stress distribution in lungs: a model of pulmonary elasticity. J Appl Physiol. 1970;28:596–608.

- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

13. 13.

Papazian L, Forel J-M, Gacouin A, Penot-Ragon C, Perrin G, Loundou A, et al. Neuromuscular blockers in early acute respiratory distress syndrome. N Engl J Med. 2010;363:1107–16.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

14. 14.

Guérin C, Reignier J, Richard J-C, Beuret P, Gacouin A, Boulain T, et al. Prone positioning in severe acute respiratory distress syndrome. N Engl J Med. 2013;368:2159–68.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

15. 15.

Han B. Ventilator-induced lung injury: role of protein-protein interaction in mechanosensation. Proc Am Thorac Soc. 2005;2:181–7.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

16. 16.

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

18. 18.

Müller-Redetzky HC, Suttrop N, Witzernath M. Dynamics of pulmonary endothelial barrier function in acute inflammation: mechanisms and therapeutic perspectives. Cell Tissue Res. 2014;355:657–73.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

19. 19.

Frank JA, Pittet J-F, Wray C, Matthay MA. Protection from experimental ventilator-induced acute lung injury by IL-1 receptor blockade. Thorax. 2008;63:147–53.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

20. 20.

Wolters PJ, Wray C, Sutherland RE, Kim SS, Koff J, Mao Y, et al. Neutrophil-derived IL-6 limits alveolar barrier disruption in experimental ventilator-induced lung injury. J Immunol. 2009;182:8056–62.

- [Article](#)
- [PubMed Central](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

21. 21.

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

[Manage Settings](#)[✓ Accept All Cookies](#)

Gurkan OU, He C, Zielinski R, Rabb H, King LS, Dodd-o JM, et al. Interleukin-6 mediates pulmonary vascular permeability in a two-hit model of ventilator-associated lung injury. *Exp Lung Res*. 2011;37:575–84.

- [Article](#)
- [PubMed Central](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

23. 23.

Meduri GU, Headley S, Kohler G, Stentz F, Tolley E, Umberger R, et al. Persistent elevation of inflammatory cytokines predicts a poor outcome in ARDS: plasma IL-1 beta and IL-6 levels are consistent and efficient predictors of outcome over time. *Chest*. 1995;107:1062–73.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

24. 24.

Gattinoni L, Pesenti A. The concept of “baby lung”. *Intensive Care Med*. 2005;31:776–84.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

25. 25.

Matute-Bello G, Downey G, Moore BB, Groshong SD, Matthay MA, Slutsky AS, et al. An official American Thoracic Society workshop report: features and measurements of experimental acute lung injury in animals. *Am J Respir Cell Mol Biol*. 2011;44:725–38.

- [Article](#)
- [CAS](#)
- [PubMed](#)

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

Feihl F, Broccard AF. Interactions between respiration and systemic hemodynamics, Part II: practical implications in critical care. *Intensive Care Med.* 2009;35:198–205.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

28. 28.

Chan K, Abraham E. Effects of inverse ratio ventilation on cardiorespiratory parameters in severe respiratory failure. *Chest.* 1992;102:1556–61.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

29. 29.

Meinhardt JP, Friess U, Bender HJ, Hirschl RB, Quintel M. Relationship among cardiac index, inspiration/expiration ratio, and perfluorocarbon dose during partial liquid ventilation in an oleic acid model of acute lung injury in sheep. *J Pediatr Surg.* 2005;40:1395–403.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

[Download references](#) ↓

Acknowledgements

We thank Marfa Polikarpova for skillful technical assistance. The work is part of the doctoral thesis of Matthias Felten.

Funding

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

- Holger C Müller-Redetzky
 - , Matthias Felten
 - , Katharina Hellwig
 - , Sandra-Maria Wienhold
 - , Jan Naujoks
 - , Bastian Opitz
 - , Norbert Suttorp
 - & Martin Witzernath
2. Department of Veterinary Pathology, Freie Universität Berlin, Berlin, Germany
- Olivia Kershaw
 - & Achim D Gruber

Authors

1. Holger C Müller-Redetzky

[View author publications](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

2. Matthias Felten

[View author publications](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

3. Katharina Hellwig

[View author publications](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

4. Sandra-Maria Wienhold

[View author publications](#)

You can also search for this author in

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

7. Olivia Kershaw

[View author publications](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

8. Achim D Gruber

[View author publications](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

9. Norbert Suttrop

[View author publications](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

10. Martin Witzernath

[View author publications](#)

You can also search for this author in

- [PubMed](#)
- [Google Scholar](#)

Corresponding author

Correspondence to [Holger C Müller-Redetzky](#).

Additional information

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

important intellectual content. NS was involved in the study design and participated in drafting the manuscript. All authors read and approved the final version of the manuscript.

Additional files

[Additional file 1: Table S1.](#)

Providing primer and probe sequences used for qRT-PCR.

[Additional file 2: Figure S1.](#)

Giving mean airway pressure and dynamic compliance measurements under alternating I:E ratios (1:2 and 1:1) during HV_T ventilation.

[Additional file 3: Figure S2.](#)

Providing HE images of all experimental groups.

[Additional file 4: Figure S3.](#)

Providing hemodynamic data.

Rights and permissions

[Reprints and Permissions](#)

About this article



We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

[Download citation](#) 

- Received: 06 October 2014
- Accepted: 20 January 2015
- Published: 28 January 2015
- DOI: <https://doi.org/10.1186/s13054-015-0759-2>

Share this article

Anyone you share the following link with will be able to read this content:

Get shareable link

Sorry, a shareable link is not currently available for this article.

Copy to clipboard

Provided by the Springer Nature SharedIt content-sharing initiative

Keywords

- Acute Respiratory Distress Syndrome
- Acute Respiratory Distress Syndrome Patient
- Recruitment Maneuver
- Inspiratory Time
- Dynamic Hyperinflation

[Download PDF](#) 

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

- [Discussion](#)
- [Conclusion](#)
- [Key messages](#)
- [Abbreviations](#)
- [References](#)
- [Acknowledgements](#)
- [Author information](#)
- [Additional information](#)
- [Additional files](#)
- [Rights and permissions](#)
- [About this article](#)

Advertisement

- **Figure 1**
 figure1


We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)


[View in article](#)[Full size image](#) >

- **Figure 4**




[View in article](#)[Full size image](#) >

- **Figure 5**




[View in article](#)[Full size image](#) >

- **Figure 6**




[View in article](#)[Full size image](#) >

- **Figure 7**




[View in article](#)[Full size image](#) >

- **Figure 8**



[View in article](#)[Full size image](#) >

- **Figure 9**



[View in article](#)[Full size image](#) >

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

> [Manage Settings](#)

✓ [Accept All Cookies](#)

2. Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, et al. Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome. N Engl J Med. 1998;338:347–54.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

3. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. N Engl J Med. 2000;342:1301–8.

- [Article](#)
- [Google Scholar](#)

4. Wolthuis EK, Vlaar APJ, Choi G, Roelofs JJTH, Juffermans NP, Schultz MJ. Mechanical ventilation using non-injurious ventilation settings causes lung injury in the absence of pre-existing lung injury in healthy mice. Crit Care. 2009;13:R1.

- [Article](#)
- [PubMed Central](#)
- [PubMed](#)
- [Google Scholar](#)

5. Gattinoni L, Carlesso E, Caironi P. Stress and strain

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

➤ [Manage Settings](#)

✓ Accept All Cookies

time and gas exchange. *Acta Anaesthesiol Scand*. 2013;57:613–22.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

7. Lee SM, Kim WH, Ahn HJ, Kim JA, Yang MK, Lee CH, et al. The effects of prolonged inspiratory time during one-lung ventilation: a randomised controlled trial. *Anaesthesia*. 2013;68:908–16.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

8. Casetti AV, Bartlett RH, Hirschl RB. Increasing inspiratory time exacerbates ventilator-induced lung injury during high-pressure/high-volume mechanical ventilation. *Crit Care Med*. 2002;30:2295–9.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

9. Caironi P, Langer T, Carlesso E, Protti A, Gattinoni L. Time to generate ventilator-induced lung injury among mammals with healthy lungs: a unifying hypothesis. *Intensive Care Med*. 2011;37:1913–20.

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

- [PubMed Central](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

11. Caironi P, Cressoni M, Chiumello D, Ranieri M, Quintel M, Russo SG, et al. Lung opening and closing during ventilation of acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2010;181:578–86.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

12. Mead J, Takishima T, Leith D. Stress distribution in lungs: a model of pulmonary elasticity. *J Appl Physiol*. 1970;28:596–608.

- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

13. Papazian L, Forel J-M, Gacouin A, Penot-Ragon C, Perrin G, Loundou A, et al. Neuromuscular blockers in early acute respiratory distress syndrome. *N Engl J Med*. 2010;363:1107–16.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

15. Han B. Ventilator-induced lung injury: role of protein-protein interaction in mechanosensation. Proc Am Thorac Soc. 2005;2:181–7.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

16. Tremblay LN, Slutsky AS. Ventilator-induced lung injury: from the bench to the bedside. Intensive Care Med. 2006;32:24–33.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

17. Kawano T, Mori S, Cybulsky M, Burger R, Ballin A, Cutz E, et al. Effect of granulocyte depletion in a ventilated surfactant-depleted lung. J Appl Physiol. 1987;62:27–33.

- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

18. Müller-Redetzky HC, Suttrop N, Witzernath M. Dynamics of pulmonary endothelial barrier function in acute inflammation: mechanisms and therapeutic perspectives. Cell Tissue Res. 2014;355:657–73.

- [Article](#)

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ Accept All Cookies

- [PubMed](#)
- [Google Scholar](#)

20. Wolters PJ, Wray C, Sutherland RE, Kim SS, Koff J, Mao Y, et al. Neutrophil-derived IL-6 limits alveolar barrier disruption in experimental ventilator-induced lung injury. *J Immunol.* 2009;182:8056–62.

- [Article](#)
- [PubMed Central](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

21. Goldman JL, Sammani S, Kempf C, Saadat L, Letsiou E, Wang T, et al. Pleiotropic effects of interleukin-6 in a “two-hit” murine model of acute respiratory distress syndrome. *Pulm Circ.* 2014;4:280–8.

- [Article](#)
- [PubMed Central](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

22. Gurkan OU, He C, Zielinski R, Rabb H, King LS, Dodd-o JM, et al. Interleukin-6 mediates pulmonary vascular permeability in a two-hit model of ventilator-associated lung injury. *Exp Lung Res.* 2011;37:575–84.

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ Accept All Cookies

consistent and efficient predictors of outcome over time. Chest. 1995;107:1062–73.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

24. Gattinoni L, Pesenti A. The concept of “baby lung”. Intensive Care Med. 2005;31:776–84.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)

25. Matute-Bello G, Downey G, Moore BB, Groshong SD, Matthay MA, Slutsky AS, et al. An official American Thoracic Society workshop report: features and measurements of experimental acute lung injury in animals. Am J Respir Cell Mol Biol. 2011;44:725–38.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

26. Feihl F, Broccard AF. Interactions between respiration and systemic hemodynamics, Part I: basic concepts. Intensive Care Med. 2009;35:45–54.

- [Article](#)

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ Accept All Cookies

- [Google Scholar](#)

28. Chan K, Abraham E. Effects of inverse ratio ventilation on cardiorespiratory parameters in severe respiratory failure. Chest. 1992;102:1556–61.

- [Article](#)
- [CAS](#)
- [PubMed](#)
- [Google Scholar](#)

29. Meinhardt JP, Friess U, Bender HJ, Hirschl RB, Quintel M. Relationship among cardiac index, inspiration/expiration ratio, and perfluorocarbon dose during partial liquid ventilation in an oleic acid model of acute lung injury in sheep. J Pediatr Surg. 2005;40:1395–403.

- [Article](#)
- [PubMed](#)
- [Google Scholar](#)



ISSN: 1364-8535

Contact us

- Submission enquiries: [Access here and click Contact Us](#)

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)

- [Press center](#)
- [Support and Contact](#)
- [Leave feedback](#)
- [Careers](#)

Follow BMC

- [BMC Twitter page](#) 
- [BMC Facebook page](#) 
- [BMC Weibo page](#) 

By using this website, you agree to our [Terms and Conditions](#), [Privacy statement](#) and [Cookies](#) policy. [Manage the cookies](#) we use in the preference centre.

SPRINGER NATURE

© 2020 BioMed Central Ltd unless otherwise stated. Part of [Springer Nature](#).

We use cookies to ensure the functionality of our website, to personalize content and advertising, to provide social media features, and to analyze our traffic. If you allow us to do so, we also inform our social media, advertising and analysis partners about your use of our website. You can decide for yourself which categories you want to deny or allow. Please note that based on your settings not all functionalities of the site are available. Further information can be found in our [privacy policy](#).

› [Manage Settings](#)

✓ [Accept All Cookies](#)