Electrostatics of forced defects in dimer coverings: corner charges, image charges, and boundary chemical potentials

Changyeon Won[1, *]

[1] *Department of Physics, Kyung Hee University, Seoul 02447, Republic of Korea*

(Dated: October 28, 2026)

When a planar region cannot be covered completely by dimers, for example a chessboard with two opposite corners removed, uncovered sites (monomers) are forced to exist but are free to move. We ask where they go when all admissible configurations are equally likely, and use their distribution as a parameter-free probe of how a boundary acts on the defects of a constrained system. Exact enumeration on square-lattice domino regions up to 512 sites across shows two regimes: power-law distributions controlled by the corners when the boundary is straight, and exponential trapping in a boundary layer when the boundary is a staircase that freezes the dimers next to it. We introduce a boundary chemical potential, $\mu = \delta \ln Z/\delta h_\partial$, the response of the macroscopic entropy to a shift of the boundary height, which connects the two regimes. On a boundary adjacent to an untilted liquid $\mu = 0$; next to tilted or frozen regions $\mu \neq 0$ and the defect weight acquires a factor $\exp \int \mu \, \mathrm{d}s$. Inside frozen regions μ is not fixed by the limit shape, and we obtain it by continuing the complex Burgers solution along tangent lines to the arctic curve. This reproduces the exact large-deviation law of the Aztec diamond identically, and predicts without free parameters the trapping rate in mixed straight/staircase domains. When the defect path runs along frozen edges, linear response fails; a layer-removal cost $\partial \ln Z/\partial(\text{size})$ with competing nested channels restores agreement. This rule, inferred from the Aztec diamond, predicts exact data for lozenge tilings of hexagons to within 0.5%, including cases where linear response is off by orders of magnitude. In the power-law regime, corners act on mobile defects as fractional charges $q_c = -J_c/4$ set by the jump of the boundary height, and together with same-sign images at the edges they give a conformally covariant density formula that reproduces all rectangles to 1–3% without parameters. We prove the corner-to-boundary law $4/[\pi(x+1)]$, reduce straight-string determinants to a single scalar Fisher–Hartwig symbol, and identify the half-plane and quadrant amplitudes as $2^{3/8}G(\frac{1}{2})^2$ and $2^{9/8}G(\frac{1}{2})^2$.

I. INTRODUCTION

Close-packed dimers on a lattice are among the oldest exactly solvable problems of statistical mechanics [1–3]. They also serve as a paradigm for the physics of local constraints. The constraint that every site is covered exactly once turns the dimer configurations into a divergence-free flux. It endows the model with a height field that is critical and Gaussian at long wavelengths [5–8], and it turns monomers (uncovered sites) into charges of an emergent electrostatics. The same "Coulomb phase" physics appears in three-dimensional dimer models [9], in spin ice, where violations of the ice rule behave as magnetic monopoles [10], and in artificial spin-ice arrays, where such defects can be imaged directly [11, 12]. In all these systems the forces between defects are, at least in part, *entropic*: defects attract or repel only because their positions change the number of available configurations.

The interaction of defects placed at *prescribed* positions is well understood. Monomer pairs in the bulk [3, 4, 13, 34, 35] and on a straight boundary [14] have been computed exactly. Ciucu and co-workers proved that gaps of given charge interact by two-dimensional electrostatics, first for lozenge tilings [15], then on the square lattice [39, 40] and on all doubly periodic bipartite lattices in the liquid phase [41]. Near a boundary the images of a gap have the same sign if the boundary is constrained and the opposite sign if dimers may protrude across it [16, 42, 43]. Monomers pinned at corners correspond to boundary-condition-changing operators [17], and a single vacancy in an odd board has been studied through Temperley's bijection [36–38]. All of this concerns boundaries next to a critical (liquid) dimer phase, where the electrostatic picture applies.

Here we ask a different and very elementary question. Suppose a finite region is unbalanced, so that it *cannot* be tiled completely; the classic example is the "mutilated chessboard" with two opposite corners removed. Defects are then forced to exist, but they are free to move, and nothing except the boundary tells them where to go. If every admissible configuration is equally likely, where are the defects? Their distribution measures directly how a boundary couples to the defects of a constrained system, with no adjustable parameter. The question becomes sharp for boundaries that are *not* next to a critical phase. Staircase boundaries, as in the Aztec diamond, freeze the dimers next to them [18], and there the electrostatic picture gives no prediction at all. We find that the defects are then trapped exponentially in a boundary layer near the points where the arctic curve touches the edge (Sec. V). The main purpose of this paper is to explain this trapping quantitatively and to connect it with the power-law regime of straight boundaries.

Our main results are the following.

* cywon@khu.ac.kr

(i) *Boundary chemical potential* (Sec. VI). We introduce $\mu(s) = \delta \ln Z/\delta h_\partial(s)$, the response of the macroscopic entropy to a shift of the boundary height. This is precisely the perturbation created by a defect and the string that connects it to the boundary, so to leading exponential order the defect weight is $\exp \int \mu \, \mathrm{d}s$, Eq. (14). On a straight edge next to an untilted liquid, $\mu = 0$ and only the power laws of the Coulomb gas remain; next to tilted or frozen regions, $\mu \neq 0$ and defects are trapped. In frozen regions the limit shape [19, 20] does not determine μ. We fix it by continuing the dual variables of the complex Burgers equation along tangent lines to the arctic curve. For the Aztec diamond this gives $\mu\delta/2 = \ln[(1-\kappa)/\kappa]$, Eq. (17), identical to the exact lattice law, whereas a naive regularization is off by a factor of about 1.45. In a "house"-shaped domain that mixes straight and staircase edges, the same construction predicts the exponential trapping rate and the optimal trapping point without free parameters (Fig. 8).

(ii) *Layer cost and competing channels* (Sec. VI). When the defect path runs along frozen edges, linear response is not enough. Each defect on a frozen staircase forces a zipper of dominoes, and two zippers meeting at a corner remove a layer of the domain. The resulting rule, Eq. (18), adds the layer-removal cost $\partial \ln Z/\partial(\text{size})$ and optimizes over nested channels. Inferred from the Aztec diamond, it predicts exact results for lozenge tilings of hexagons to within 0.5%, including cases where linear response alone is wrong by orders of magnitude (Table I). The layer cost required by consistency coincides with the derivative of $\ln Z$ along the defect-free side of the hexagon, not along the other two sides.

(iii) *Corner charges for mobile defects* (Sec. III). For straight edges, every corner acts on the moving defects as a fractional charge $q_c = -J_c/4$, Eq. (2), fixed by the jump of the boundary height ($\pm 1/4$ for ordinary corners and $+3/4$ for a removed one). Combined with the same-sign image rule for constrained edges, this yields a conformally covariant density formula, Eq. (3), and the corner exponents of Eq. (4). Without parameters they reproduce the exact distributions of one and two defects in all rectangles to 1–3%, including the $r^{-3/4}$ divergence of the density next to a removed corner and the $N^{-1/4}$ depletion of the corners of odd boards.

(iv) *Exact results* (Sec. IV). We prove the corner-to-boundary law $4/[\pi(x+1)]$ in the quadrant, Eq. (5). We show that straight-string determinants in the plane and the half plane reduce to Toeplitz and Toeplitz+Hankel determinants of a single scalar symbol, which reproduces the known bulk amplitude and gives the new half-plane and quadrant amplitudes $2^{3/8}G(\frac{1}{2})^2$ and $2^{9/8}G(\frac{1}{2})^2$ to 18 digits.

A control on the non-bipartite triangular lattice, where monomers carry no charge, shows no long-range boundary effect (Sec. VII). Throughout we distinguish rigorous results, results established by exact arithmetic to many digits, numerically supported statements and conjectures; Table II summarizes the status of each result and identifies what was known before.

II. MODEL AND METHODS

A. Dimers, heights and charges

We consider unions of unit squares (cells) of $\mathbb{Z}^2$, colored like a chessboard; cell (x, y) is black if $x + y$ is even. A domino covers one black and one white cell. If the region has n_w white and n_b black cells, at least $|n_w - n_b|$ cells remain uncovered. We study the uniform measure on configurations with exactly the forced number of defects. We call the white defects negative ($e = -1$) and the black defects positive ($e = +1$). A removed cell carries the charge of a defect of its color.

The Thurston height h_T changes by $+1$ (-1) when one walks along a lattice edge with a black (white) cell on the left. Around a covered vertex h_T returns to its value. Around a defect it winds by ± 4, so the defect is a vortex of the height field. We use the normalized height $h = h_T/2$, whose admissible slopes satisfy $|h_x| + |h_y| \leq 1$. In the scaling limit h converges to a Gaussian free field [5, 6], and defects become electric charges of the dual (magnetic) description [7, 8].

B. Exact enumeration

We use three independent exact methods. (a) A row-by-row broken-profile transfer matrix with 128-bit integers, which gives exact counts for all defect configurations on boards up to $N = 16$. It serves as the reference. (b) Kasteleyn determinants [1], with horizontal weights 1 and vertical weights i (or real signs alternating by column). For regions with interior defects, the Kasteleyn condition is restored by flipping the signs of the edges crossed by a "string" joining the defect to the boundary. (c) A low-rank update. If $G = K^{-1}$ is known for the defect-free region, the ratio of partition functions with defects at v and c is the bordered determinant

$$\frac{Z(\Omega \setminus \{c, v\})}{Z(\Omega)} = \left| \det \begin{pmatrix} I - 2G(b_j, w_l) & -2G(b_j, v) \\ G(c, w_l) & G(c, v) \end{pmatrix} \right|, \tag{1}$$

where (b_j, w_l) are the endpoints of the string edges. For the infinite plane, half plane and quadrant, the entries of G are exact numbers of the form $a + b/\pi$ with rational a, b. We generate them by an exact recursion and evaluate determinants in ball arithmetic (Arb) with certified error bounds, up to string lengths $r = 1024$. On 8×8 boards the three methods agree for all 1552 defect configurations. Aztec-diamond ratios were obtained from exact integer inverses (FLINT), because double precision fails beyond $n \simeq 44$ (the condition number reaches $\sim 10^{17}$ at $n = 64$). All key claims of Secs. IV–VI were independently re-derived from scratch by a separate code base; no discrepancy was found.




FIG. 1. (a) An 8×8 board with corner a1 removed. The region has one more white than black square, so one white defect (charge $e = -1$) is forced. Each corner carries the fractional charge $q_c = -J_c/4$ given by the jump J_c of the boundary height: $-1/4$ for a black corner, $+1/4$ for a white corner and $+3/4$ for the removed black corner. (b) In the continuum description the edge of the region acts as a mirror with images of the same sign, and the corners act as fixed charges. The resulting density of a single defect is given by Eq. (3).

III. COULOMB GAS FOR AXIS-PARALLEL BOUNDARIES

A. Corner charges from boundary height jumps

Walking counterclockwise along the boundary, the height oscillates between two values on a straight edge, so its average is constant along each edge. At a convex corner the corner cell contributes two consecutive steps of the same sign. The average height therefore jumps by $J_c = +1$ at a black corner and $J_c = -1$ at a white corner. At the notch of a removed black corner four consecutive -1 steps give $J_c = -3$, which differs from $+1$ by one monomer's worth of winding.

A jump J at a boundary point c creates an average height $(J/\pi)\arg(z-c)$. Its harmonic conjugate is $(J/\pi)\ln|z-c|$. A bulk monomer creates $(2/\pi)\ln|z-z_0|$ in the same normalization. Because the dual field obeys Neumann conditions, a charge on the boundary overlaps with its own image. Matching $2q \cdot (2/\pi) = J/\pi$ gives

$$q_c = -\frac{J_c}{4}: \quad q = -\tfrac{1}{4}\ (\text{black}),\ +\tfrac{1}{4}\ (\text{white}),\ +\tfrac{3}{4}\ (\text{removed}). \tag{2}$$

This is a purely combinatorial statement. The same boundary-height jumps were identified by Allegra [17] as boundary-condition-changing operators of dimension $J^2/32$ at the corners. That work used them to compute corner free energies and correlations of monomers placed *at* corners. Here we use them for the distribution of *freely moving* defects.

B. Density formula and corner exponents

Let f map the region conformally onto the upper half plane, and let $f_c = f(c)$ be the images of the corners. Treating the defect as a charge e in the field of the corner charges and of its images, we obtain

$$\rho(z) \propto |f'(z)|^{1/4}\,[\mathrm{Im} f(z)]^{1/4} \prod_c |f(z) - f_c|^{e\,q_c}. \tag{3}$$

For two defects there is an additional factor $\big(|f_1 - f_2|\,|f_1 - \bar{f}_2|\big)^{e_1 e_2/2}$. The factor $|f'|^{1/4}(\mathrm{Im} f)^{1/4}$ is the one-point function of a charge of dimension $1/4$ in the presence of a same-sign image. Near a corner of interior angle θ, Eq. (3) predicts $\rho \sim r^{\gamma}$, with the factor π/θ familiar from conformal corner exponents [46],

$$\gamma = \frac{\pi}{\theta}\Big(\frac{1}{2} + e\,q_c\Big) - \frac{1}{4}. \tag{4}$$

For a white defect next to the removed corner a1 ($\theta = \pi/2$, $q_c = 3/4$), this gives $\gamma = -3/4$. The defect density thus *diverges* as $r^{-3/4}$ toward the removed corner (Fig. 2). The exact slopes 0.754 ($N = 64 \to 128$) and, on the mutilated board, 0.750 agree with this prediction. The formula also holds for concave corners, to three decimal places.

C. Tests

We tested Eq. (3) on even×even, odd×odd and odd×even rectangles, with corner, edge and interior cells removed, and with one or two defects (Figs. 2–4). The




FIG. 2. A single removed corner. (a) Exact defect density on a 64×64 board with a1 removed, averaged over 2×2 blocks and normalized to the uniform value. (b) The parameter-free Coulomb-gas prediction, Eq. (3), computed with the Jacobi-elliptic map of the square. (c) Density at three special cells versus board size (exact, symbols), compared with the power laws predicted by Eq. (4) (dashed). The rms deviation between (a) and (b) is 0.9% at $N = 128$ and decreases as $1/N$. The orange square marks the removed cell a1.

rms deviations are 1–3% at the largest sizes and decrease as $1/N$. Two special cases illustrate the physics.

Mutilated board (Fig. 3). The two defects are attracted by the two removed corners ($q = +3/4$ each) and repel each other. Their joint distribution, the corner scaling and the conditional density around a central defect all follow the Coulomb gas.

Odd×odd board (Fig. 4). All four corners carry $q = +1/4$. For the black defect the total boundary charge is neutral, and Eq. (3) reduces to $\rho \propto R^{1/4}$, where R is the conformal radius. The defect is pushed toward the centre, and the corner-to-centre ratio decays as $N^{-1/4}$. At the lattice level, the equal probability of all even-even boundary cells follows from Temperley's bijection; the corresponding counts were obtained by Tzeng and Wu [36, 37]. Bouttier *et al.* studied the same bijection for a mobile vacancy and found that the probability for a central vacancy to reach the boundary decays as $L^{-1/4}$ [38]. The defect lives on two black sublattices, (even,even) and (odd,odd), whose weights differ near the boundary (Fig. 4a,b,d). This staggering is a lattice effect that the continuum formula averages over: the sublattice means differ by 0.42, 0.27, 0.17 and 0.10 (in units of ρ_{unif}) for $N = 7$, 15, 31 and 63, and the average of neighbouring cells of the two sublattices follows $\rho \propto R^{1/4}$ to within 3% everywhere except in the outermost row.

Image charges. A single defect near a straight edge and a pair of defects confirm that the image has the *same* sign as the source (Fig. 5a). This is the rule found by Ciucu for gaps in lozenge tilings near constrained boundaries [42, 43]; in that terminology every edge in the present paper is constrained. Near a free boundary, where dimers may protrude, the image has the opposite sign and the boundary behaves as a conductor [16, 42]. Our contribution here is to combine the same-sign images with the corner charges of Eq. (2) for freely moving defects on the square lattice. The pair correlation decays as d^{-1} along a boundary and as $d^{-1/2}$ in the bulk. The boundary-pair amplitude $2/(\pi d)$ agrees with the result of Priezzhev and Ruelle [14]. At the lattice level the sign follows from the fact that the long-distance behaviour of K^{-1} is dominated by momenta $\pm\pi/2$. A reflection that shifts by two lattice spacings then acquires a phase $e^{i\pi} = -1$, which converts the lattice Dirichlet reflection into a continuum Neumann (same-sign) image.

IV. EXACT RESULTS AND AMPLITUDES

A. Corner to boundary

Theorem 1 (Quadrant, boundary defect)**.** *In the infinite quadrant with its corner cell c removed, place a white defect on the boundary cell $(x, 0)$, with x odd. Then, for $x \geq 3$,*

$$\frac{Z(Q \setminus \{c, (x,0)\})}{Z(Q)} = \frac{4}{\pi} \int_0^{\pi/2} \sin\theta \, \sin[(x+1)\theta] \times \sqrt{1 + \cos^2\theta} \, \mathrm{d}\theta = \frac{4}{\pi(x+1)} + O(x^{-3}). \tag{5}$$

The values are of the form "integer minus rational/π", e.g. $2 - 16/(3\pi)$ at $x = 3$.

The proof diagonalizes the box Kasteleyn matrix by




FIG. 3. The mutilated chessboard (corners a1 and h8 removed; two white defects). (a,b) Exact marginal density at $N = 96$ (2×2 block averages, each block containing exactly two white cells) compared with the two-defect Coulomb gas. Orange squares mark the removed corners a1 and h8. (c) Distribution of the defect separation. (d) Corner scaling for $N = 8$–96; dashed lines show the slopes predicted by Eq. (4). (e) Conditional density around a defect fixed at the centre. All theory curves are parameter free.

sine functions and evaluates one integral in closed form. The leading term comes from $\theta = \pi/2$, and the next order vanishes by parity. The exponent -1 is the Coulomb-gas value $|f'|^{1/2}|f|^{-3/4}$ for $f = z^2$, while the prefactor $4/\pi$ is a lattice quantity.

B. A single scalar symbol

For a vertical string, the horizontal offset between the black end of the jth string edge and the white end of the lth depends only on the parity of $j - l$. Hence $I - 2G$ is a Toeplitz matrix in $j - l$. Summing the Fourier series in closed form gives the symbol

$$\sigma(e^{i\varphi}) = \operatorname{sgn}(\cos\varphi)\,\frac{\cos\varphi + i}{\sqrt{1+\cos^2\varphi}}, \tag{6}$$

which is a pure phase with jumps of π at the massless points $\varphi = \pm\pi/2$ (Fig. 5b). Moreover, when the defect sits at the end of the string and the removed cell at its start, the border row of Eq. (1) equals $-\frac{1}{2}$ times the first row. It can therefore be eliminated exactly. With c_m the Fourier coefficients of σ,

$$R_{\mathrm{bulk}}(r) = \tfrac{1}{2}\left|\det\left[c_{j-k-1}\right]_{j,k=0}^{r-1}\right|, \tag{7}$$

$$R_{\mathrm{half}}(r) = \tfrac{1}{2}\left|\det\left[c_{j-k-1} - c_{j+k+3}\right]_{j,k=0}^{r-1}\right|, \tag{8}$$

for a pair at vertical distance r in the plane, and for a boundary cell and a defect at height r above it in the half plane. The Hankel part in the second line is the lattice image charge.

C. Amplitudes

The symbol $z\sigma$ is of Fisher–Hartwig type with $\beta_1 = \beta_2 = \frac{1}{2}$ at $z = \pm i$. The Deift–Its–Krasovsky asymptotics [21] then gives

$$R_{\mathrm{bulk}}(r) \simeq \pi\, 2^{-3/4}\, G(\tfrac{1}{2})^4\, r^{-1/2} = 0.2473718\ldots\, r^{-1/2}. \tag{9}$$

The factors are $2^{-1/4}$ from $\sum_k kV_kV_{-k} = -\operatorname{artanh}(q^2)$ with $q = \sqrt{2}-1$, $2^{1/2}$ from $|z_1 - z_2|^{2\beta_1\beta_2}$, and $[G(\frac{3}{2})G(\frac{1}{2})]^2$




FIG. 4. Odd×odd boards, where the boundary charge is neutral. (a,b) Exact defect density on a 63×63 board, shown cell by cell on the two black sublattices, (even,even) and (odd,odd), without averaging. (c) Prediction $\rho \propto R^{1/4}$, where R is the conformal radius. (d) Cut along the central column: the two sublattices (symbols) bracket the prediction (line). (e) Corner-to-centre ratio versus N, with the predicted slope $-1/4$.

from the two jumps. This axis amplitude was first obtained in closed form by Wilkins and Powell [35] from a Toeplitz representation derived with the transfer matrix; their $E/2 = 2^{-7/12}e^{6\zeta'(-1)}$ equals $\pi 2^{-3/4}G(\frac{1}{2})^4$. The same constant was obtained for diagonal separation by Hartwig [4] ($B/4$ with $B = 0.98948729$). The scalar symbol reproduces both and extends directly to the half plane and the quadrant. For the half plane and the quadrant we established the amplitudes by certified arithmetic up to $r = 1024$ and extrapolation, to 18–19 digits (Fig. 5c):

$$C_{\mathrm{half}} = 2^{3/8}G(\tfrac{1}{2})^2, \qquad C_{\mathrm{quad}} = 2^{9/8}G(\tfrac{1}{2})^2. \tag{10}$$

The pattern "lattice value = operator normalization × continuum Coulomb-gas correlator" accounts for all three. The transcendental factor $G(\frac{1}{2})^2$ belongs to each bulk monomer, while the boundary and corner normalizations involve only π and powers of 2. A proof for the half plane would require the asymptotics of a Toeplitz+Hankel determinant with a *non-even* Fisher–Hartwig symbol $z^{-1}\sigma$ of winding -1. This lies just outside the existing theory [21–23].

V. STAIRCASE BOUNDARIES: TRAPPING IN THE AZTEC DIAMOND

The Aztec diamond AD_n has $Z = 2^{n(n+1)/2}$ tilings [24]. Four frozen corners are separated from a liquid disk by the arctic circle [18]. We remove the west tip cell t, which forces one white defect, and we also consider a missing domino, which forces two defects of opposite colours. In both cases the defects avoid the liquid region. Instead they stay within a boundary layer about two cells thick, near the points where the arctic circle touches the edges (Fig. 6a). The weight outside this layer decays as n^{-1} for one defect and as $n^{-1/2}$ for a pair (Fig. 6c).

On the boundary the distributions are known exactly. Let k be the position of the defect along the edge, counted from the tip. For the adjacent (SW) and the opposite (NE) edge,

$$\frac{Z(\mathrm{AD}_n \setminus \{t, \mathrm{SW}_k\})}{Z(\mathrm{AD}_n)} = P\big(\mathrm{Bin}(n, \tfrac{1}{2}) \geq k\big), \tag{11}$$

$$\frac{Z(\mathrm{AD}_n \setminus \{t, \mathrm{NE}_k\})}{Z(\mathrm{AD}_n)} = \binom{n-1}{k-1} 2^{-n}, \tag{12}$$

For a missing domino with defects at positions k and l




FIG. 5. (a) Monomer-pair correlations on a 160 × 160 board: along the boundary ($\propto d^{-1}$, same-sign images) and in the bulk ($\propto d^{-1/2}$). (b) Phase of the scalar symbol σ of Eq. (6), with jumps of π at $\varphi = \pm\pi/2$. (c) Exact weights, scaled by the expected power law and divided by the closed forms $\pi 2^{-3/4} G(\frac{1}{2})^4$ (bulk), $2^{3/8} G(\frac{1}{2})^2$ (half plane) and $2^{9/8} G(\frac{1}{2})^2$ (quadrant), versus $1/r$.

on the two edges adjacent to a corner,

$$\frac{Z(\mathrm{AD}_n \setminus \{\mathrm{S1}_k, \mathrm{S2}_l\})}{Z(\mathrm{AD}_n)} = \sum_{j=0}^{n-1} 2^{j-1}\, P\big(\mathrm{Bin}(j, \tfrac{1}{2}) = k-1\big) \times P\big(\mathrm{Bin}(j, \tfrac{1}{2}) = l-1\big). \tag{13}$$

Equation (13) is equivalent to the boundary generating function of Chhita and Young [25], and Eqs. (11)–(12) are special cases of it. We found these laws independently by exact computation. We also give elementary proofs of Eqs. (11) and (12) that do not use urban renewal (Appendix A). Equation (13) has a transparent dynamical reading. In domino shuffling the defect pair is created at the corner at "time" j with weight $\frac{1}{2}$, and afterwards the two defects perform independent random walks along their edges. Summing the pair weights over all boundary positions gives $2^{n+1} - 2 = Z(\mathrm{AD}_{n+1})/Z(\mathrm{AD}_n) - 2$.

VI. BOUNDARY CHEMICAL POTENTIAL

A. Definition

A defect and the removed cell are joined, along the boundary, by an arc Γ on which the boundary height is shifted by $\delta = 2$ (one monomer of Thurston winding). To leading exponential order the defect weight is the change of the macroscopic entropy. For a small shift this is the linear response

$$\ln w \simeq \delta \int_\Gamma \mu(s)\, \mathrm{d}s, \qquad \mu(s) = \frac{\delta \ln Z}{\delta h_\partial(s)} = \mathbf{n} \cdot \nabla_p \,\mathrm{ent}(\nabla h)\big|_\partial, \tag{14}$$

where ent is the Cohn–Kenyon–Propp local entropy [19], h is the limit shape and $\mathbf{n}$ is the outward normal. On a flat, liquid-adjacent edge such as that of the chessboard, $\nabla h = 0$ gives $\mu = 0$. The exponential factor is then absent and only the Coulomb-gas power laws of Sec. III remain. On staircase edges, and on straight edges adjacent to a *tilted* liquid, $\mu \neq 0$ and the defects are trapped exponentially.

B. Frozen boundaries: tangent continuation

In a frozen region the minimizer h lies on the boundary of the allowed slope polygon, so it does not determine ∇_pent. Consistently, a penalty-regularized finite-element solution gives values of μ that depend on the regularization, by factors of order 1.5 (Fig. 7c). The missing information is supplied by the complex Burgers structure of dimer limit shapes [20]. The dual variables $(\ln|z|, \ln|w|)$ are constant along straight characteristics. In frozen regions these characteristics are the tangent lines to the arctic curve.

For the uniform Aztec diamond ($|x|+|y| \le 1$) the local dimer probabilities are known in closed form [26]:

$$p_{E,W} = \frac{1}{2} + \frac{1}{\pi} \arctan \frac{\pm 2x - 1}{\sqrt{1 - 2x^2 - 2y^2}}, \tag{15}$$




FIG. 6. Aztec diamond with the west tip removed. (a) Exact defect density $\rho/\rho_{\mathrm{unif}}$ for $n = 24$ on a logarithmic colour scale spanning eight decades; each cell of the defect sublattice is drawn as a diamond reaching its four neighbours. The dashed line is the arctic circle, and the small circle marks the removed tip. The density falls to 10^{-6} along the two edges that do not touch the removed tip and drops steeply near the far end of the adjacent edge, following the binomial tail of Eq. (11). (b) Exact weights along the adjacent (south-west) and opposite edges for $n = 44$ (symbols), compared with Eqs. (11)–(12) (lines). (c) Total weight off the adjacent edge, in the interior and on the far edge, versus n.

and similarly $p_{N,S}$ with y. Their ratios have finite limits on the arctic circle. The tangent from the edge point $P(\kappa) = (-1+\kappa, -\kappa)$ touches the circle at

$$Q(\kappa) = \frac{\left(2\kappa^2 - 1,\ 2\kappa^2 - 4\kappa + 1\right)}{2(2\kappa^2 - 2\kappa + 1)}. \tag{16}$$

Projecting the dual variables at Q onto the edge normal gives

$$\frac{\delta}{2}\,\mu(\kappa) = \frac{1}{2}\ln\left[\left(\frac{1-\kappa}{\kappa}\right)^2\right] = \ln\frac{1-\kappa}{\kappa} \tag{17}$$

identically in κ. This is exactly the lattice value $\ln[(n-k)/k]$ that follows from Eqs. (11)–(13). The limit shape plus tangent continuation thus reproduces the full large-deviation profile $n[H(\kappa) - \ln 2]$ of the Aztec diamond (Fig. 7). The same procedure applied to the arctic curve extracted from a finite-element solution reproduces Eq. (17) to 1–3%.

C. A mixed domain

As a test with no free parameter we consider a "house": an $N \times N$ square whose two upper corners are cut by staircases of opposite colours (size k). The staircases balance the colours, and removing a1 creates one defect. Exact integer computations up to $N = 128$ show that the defect leaves a1, the Coulomb attractor, and is trapped on the staircase of the same colour with weight $w \simeq e^{RN} N^{-1}$. The exact rates are $R = 0.213$ for $k = N/2$ and $R = 0.255$ for $k = N/4$. We computed the limit shape by finite elements. We evaluated μ directly where the boundary is liquid-adjacent and by tangent continuation where it is frozen, and integrated Eq. (14) along Γ. This predicts $R = 0.204$–0.217 and 0.259–0.271 (range over meshes $N = 48$–96), with the optimum at $x/N = 0.07$–0.09 (exact: 0.08–0.09), as shown in Fig. 8. Most of the gain comes not from the staircase but from the straight left edge, because the staircases tilt the adjacent liquid. The optimum sits where μ changes sign on the staircase.

D. Beyond linear response: layer cost and competing channels

Confronting Eq. (14) with the exact Aztec laws Eqs. (11)–(13) reveals two further ingredients. They matter whenever Γ runs along *frozen* edges. First, the local gradient of $\ln w$ is always given correctly by Eq. (17), but the rate itself is smaller than $\int_\Gamma \mu$ by exactly $\ln 2$. Second, for two defects close to a corner the dominant contribution comes from $j < n$ in Eq. (13). All three families are described by the single rule

$$\frac{1}{n}\ln w = \max_{\tau}\left[\tau\, L\Big(\frac{\kappa}{\tau}, \frac{\lambda}{\tau}\Big) - \frac{1}{n}\,\frac{\partial \ln Z(\Omega_m)}{\partial m}\Big|_{m=\tau n}\right], \tag{18}$$

where $\Omega_{\tau n}$ is the scaled copy of the domain sharing the relevant corner, L is the linear response $\int_\Gamma \mu$ computed in the unit domain, and the second term is the entropy cost




FIG. 7. Boundary chemical potential of the Aztec diamond. (a) The dual variables are constant along tangent lines to the arctic circle, so μ at the edge point $P(\kappa)$ is set by the tangency point $Q(\kappa)$. (b) $\mu(\kappa)$ from the limit shape with tangent continuation (symbols, extracted from a finite-element limit shape) against the exact lattice value $\ln[(1-\kappa)/\kappa]$ (line). (c) Integrated rate: exact continuum result (line), tangent continuation (circles) and the naive penalty-regularized linear response (squares), which is biased by a factor of about 1.45.

of removing one lattice layer. For the Aztec diamond the layer cost is $\tau \ln 2$. Only $\tau = 1$ is allowed if an endpoint lies on an edge not containing the corner. Equation (18) reproduces Eqs. (11) (optimum $\tau = \min(2\kappa, 1)$), (12) and (13) exactly. At the lattice level, a defect on a frozen staircase forces a zipper of dominoes (Appendix A). Two zippers meeting at the corner strip one layer off the inner domain, and the turning point of the zippers is τ. When Γ lies entirely on liquid-adjacent edges no zipper forms and Eq. (18) reduces to linear response, which is the situation of the house domain above.

E. Independent test: lozenge tilings of hexagons

To test Eq. (18) away from the data that suggested it, we studied lozenge tilings of a hexagon with side lengths a (bottom), b (right) and c. We remove an up-triangle at distance k and a down-triangle at distance l from the bottom-right corner. By the Lindström–Gessel–Viennot lemma [27, 28], the number of tilings is a $(c+1) \times (c+1)$ determinant of binomial coefficients. The sources are $(-i, i)$ and $(a-k, 1)$, and the sinks are $(a-j, b+j)$ and $(a-1, l)$. We verified this formula against Kasteleyn determinants for all dent positions in three hexagons, and it allows exact computation up to $n = 240$. Dent-pair counts on the boundary of a hexagon are also given in closed form by Ciucu and Fischer [44]; the limit shape is the arctic circle of Cohn, Larsen and Propp [45].

With the corner triangle fixed, the dent position follows $P(X = m) \propto \binom{a-1+m}{m}\binom{c-1+b-m}{b-m}$. Consistency of Eq. (18) then *requires* a layer cost equal to $\max f - f(0)$, where f is the large-deviation function of X. For five hexagons this number equals $\partial \ln Z/\partial c$ to three or four digits and differs clearly from $\partial_a \ln Z$ and $\partial_b \ln Z$ (Fig. 9b). The removed layer is therefore the one in the defect-free direction. With no further input, Eq. (18) then predicts the pair rates, including channels with $\tau \approx 0.24$–0.76 where linear response alone fails completely (e.g. 0.003 instead of 0.139). The agreement with the exact values is within 0.1–0.5% for nine cases in three hexagons (Table I). Along the diagonal $\kappa = \lambda$ it holds at twenty further points in two hexagons, with deviations below 0.3% (Fig. 9a).

VII. CONTROL: NO CHARGE, NO LONG-RANGE BOUNDARY EFFECT

On the non-bipartite triangular lattice there is no height field. Monomers then carry only a $\mathbb{Z}_2$ label and no U(1) charge [13]. Exact transfer-matrix counts for dimers on the bonds of the triangular lattice, on rhombi with 60° and 120° corners, show a boundary layer of one or two cells, beyond which the defect distribution is flat. The corner-to-centre ratio saturates instead of scaling (Fig. 10). This contrasts with the $N^{-1/4}$ law of the odd square board and confirms that the long-range corner physics requires the emergent charge.




FIG. 8. House-shaped domain. (a) Limit shape ($k = N/2$) from the finite-element solution. Grey marks frozen regions, where the slope lies on the boundary of the allowed polygon, $|h_x| + |h_y| = 1$; in the liquid the shade gives the tilt $|h_x| + |h_y|$, which controls μ on the adjacent boundary. Also shown is the defect path Γ from the removed corner a1 along the straight edge to the optimal point on the staircase. (b) Predicted rate $R(x) = N^{-1}\int_\Gamma \mu\, ds$ versus the endpoint x/N for $k = N/2$ and $N/4$ (solid), compared with the exact lattice rates (dashed) and the exact optimal position (grey band).

TABLE I. Pair rates $\frac{1}{n}\ln w$ for dents at distances (κ, λ) from a corner of the hexagon $H(a{:}b{:}c)$. Columns: linear response only ($\tau = 1$), the rule Eq. (18) with its optimal τ, and exact values extrapolated from $n = 40$–240 with $a + b\ln n/n + c/n$.

$a{:}b{:}c$	(κ, λ)	$\tau = 1$	rule (τ^*)	exact
1:1:1	$(0.1, 0.1)$	0.003	0.1387 (0.37)	0.1386
1:1:1	$(0.2, 0.2)$	0.259	0.2774 (0.75)	0.2766
1:1:1	$(0.1, 0.3)$	0.208	0.2251 (0.76)	0.2244
1:1:1	$(0.3, 0.3)$	0.412	0.4124 (1)	0.4111
1:1.5:0.75	$(0.1, 0.1)$	−0.201	0.1387 (0.24)	0.1382
1:1.5:0.75	$(0.15, 0.3)$	0.172	0.2866 (0.52)	0.2865
1:1.5:0.75	$(0.3, 0.6)$	0.573	0.5728 (1)	0.5718
1.4:0.8:1	$(0.3, 0.15)$	0.268	0.2866 (0.75)	0.2850
1.4:0.8:1	$(0.6, 0.4)$	0.554	0.5544 (1)	0.5534

VIII. DISCUSSION AND CONCLUSIONS

Forced defects in dimer coverings are pushed around by forces that are entirely entropic. Nevertheless they obey an electrostatics with fractional corner charges and same-sign images. Its origin is the Gauss law imposed by close packing, together with the Gaussian fluctuations of the height field. Entropy only sets the stiffness (the dielectric constant). The resulting picture is quantitative. A single formula, Eq. (3), with charges fixed by boundary height jumps, describes all rectangular geometries without parameters. The corresponding lattice amplitudes factorize into a universal bulk monomer normalization, which is transcendental and contains $G(\frac{1}{2})^2$, and elementary boundary and corner normalizations. All straight-string determinants reduce to a single scalar Fisher–Hartwig symbol.

The emergent-electrostatics picture breaks down where the boundary freezes the dimer sea. The breakdown is controlled by the boundary chemical potential: the entropy cost of shifting the boundary height, evaluated on the limit shape and continued along tangent lines into the frozen regions. With this one ingredient the Aztec-diamond law is reproduced identically and trapping rates in mixed domains are predicted to a few per cent. Along frozen edges a geometric layer cost and a competition between nested channels must be added. The resulting rule, Eq. (18), which we inferred from the Aztec diamond, predicts exact lozenge-tiling data to within half a per cent.

The same questions arise wherever constrained degrees of freedom produce emergent Coulomb physics with boundaries. Examples are monopoles in finite spin-ice samples and nanostructures, defects in artificial spin ice, where edge and corner populations can be imaged by




FIG. 9. Independent test on lozenge tilings. (a) Rate of a dent pair at distances $\kappa = \lambda$ from a corner, for two hexagons: exact values extrapolated from $n = 60, 120, 240$ (symbols), the rule Eq. (18) (solid) and linear response alone, $\tau = 1$ (dotted). (b) The layer cost required for consistency (stars) coincides with $\partial_c \ln Z/n$, the derivative with respect to the side length in the defect-free direction, and not with ∂_a or ∂_b.




FIG. 10. Control on the triangular lattice. (a) Monomer probability on a 13×13 rhombus of the triangular lattice (normalized to uniform). Here the dimers cover bonds between sites of the triangular lattice, so each site is drawn as its hexagonal cell; this differs from the lozenge tilings of Sec. VI, which are dimers on the honeycomb lattice and are bipartite. (b) Corner-to-centre ratio versus size: saturation for the triangular lattice (60°, 120°) against the power law $N^{-1/4}$ of the square lattice.

magnetic force microscopy [11, 12], dimer adsorbates and molecular monolayers on terraces, and constrained colloidal systems. Our results suggest that corners should act as fractional charges and constrained edges as same-sign mirrors, as long as the constrained phase touching the boundary is critical. Where it is ordered (frozen), defects should instead be trapped exponentially at the points where the ordered and critical regions meet. Interactions between dimers change the stiffness g [30], so all exponents in Eq. (4) should scale with g. This offers an experimental handle on the coupling constant.

Several questions remain open. Among them are a proof of the corner-charge density formula for general domains, which is likely accessible through piecewise-

TABLE II. Summary of results and their status. "Proof": rigorous; "exact": exact arithmetic or ≥ 10 digits; "numerical": agreement at the per-cent level across sizes; "known": previously established in the cited work.

Result	Status
Corner charge $q_c = -J_c/4$	proof; cf. [17]
Density formula (3), exponent (4)	numerical (1–3%)
Quadrant boundary law (5)	proof
Scalar-symbol reductions (8)	proof
Bulk amplitude $\pi 2^{-3/4} G(\frac{1}{2})^4$	known: axis [35], diagonal [4]; new derivation
Half-plane, quadrant amplitudes	exact (18 digits)
Aztec laws (11)–(13)	known [25]; new elementary proofs of (11), (12)
μ of the Aztec diamond, Eq. (17)	exact identity (given tangent continuation)
House trapping rates	numerical (5%)
Layer rule (18)	conjecture; exact for Aztec, 0.5% for lozenges
Triangular control	numerical

Temperleyan and monodromy methods [31, 32], and the Toeplitz+Hankel asymptotics that would prove the half-plane and quadrant amplitudes. Others are a first-principles justification of the tangent continuation and of the layer rule in general domains, which appears to be related to the tangent method of Colomo and Sportiello [33] (proved for a class of six-vertex domains by Aggarwal [47]), and the behaviour of many defects, where screening of the corner charges is expected.

ACKNOWLEDGMENTS

The author thanks the developers of FLINT and Arb, whose exact and certified arithmetic made the large-scale computations possible.

Appendix A: Elementary proofs for the Aztec diamond

Boundary switch. Each white cell SW_k on the south-west edge has only two neighbours, b_{k-1} and b_k, on the adjacent diagonal. If SW_k is matched to b_k, then SW_{k+1} must also be matched to b_{k+1}. The edge therefore consists of m dominoes of one orientation followed by $n-m$ of the other, and the switch position m is unique.

Sliding bijection. If the tip b_0 is removed, the cells $\mathrm{SW}_1, \ldots, \mathrm{SW}_{k-1}$ are forced. Sliding the dominoes SW_i–b_{i-1} to SW_i–b_i for $i < k$ gives a bijection between tilings of $\mathrm{AD}_n \setminus \{t, \mathrm{SW}_k\}$ and tilings of AD_n with switch $m \geq k$.

Switch distribution. Let I be AD_n minus the two opposite boundary double layers. Removing a corner cell c of I forces a zipper of $n-2$ dominoes, and what remains is a translate of AD_{n-1} minus its boundary layer. Kuo's graphical condensation [29], applied to $G = I \cup \{b_m, e_t\}$ with four boundary vertices of alternating colour, together with the switch of the opposite layer, gives $T_n(m) = 2^{n-1}[T_{n-1}(m) + T_{n-1}(m-1)]$. Hence $T_n(m) = \binom{n}{m} 2^{n(n-1)/2}$, which proves Eq. (11).

Opposite edge. By 180° rotation, removing the east tip forces the whole north-east double layer. The south-west switch then leaves exactly two free cells $b_s, b_{s'}$ with $s < j \leq s'$. Kuo's identity with two same-colour pairs gives $Z(I \cup \{b_s, b_{s'}\}) = 2^{(n-1)(n-2)/2} \binom{n}{s} \binom{n}{s'} \frac{s'-s}{n}$, and the binomial identity $\sum_{s<j\leq s'} \binom{n}{s}\binom{n}{s'}(s'-s) = n 2^{n-1} \binom{n-1}{j-1}$ yields Eq. (12).

Appendix B: Numerical details

Coulomb-gas predictions for rectangles use the Jacobi map $f(u) = \mathrm{sn}(u, k)$, with the modulus fixed by the aspect ratio. Limit shapes were computed by minimizing the Cohn–Kenyon–Propp functional with the domino entropy written in terms of the Lobachevsky function. We used a finite-element mesh with four triangles per cell, which avoids locking at the slope constraint, and penalty continuation. On the Aztec diamond this reproduces the entropy to 0.02% and the frozen area fraction $1 - \pi/4$. Rates were extrapolated using $a + b \ln n/n + c/n$.

[1] P. W. Kasteleyn, Physica **27**, 1209 (1961).
[2] H. N. V. Temperley and M. E. Fisher, Philos. Mag. **6**, 1061 (1961).
[3] M. E. Fisher and J. Stephenson, Phys. Rev. **132**, 1411 (1963).
[4] R. E. Hartwig, J. Math. Phys. **7**, 286 (1966).
[5] R. Kenyon, Ann. Probab. **28**, 759 (2000).
[6] R. Kenyon, Ann. Probab. **29**, 1128 (2001).
[7] B. Nienhuis, J. Stat. Phys. **34**, 731 (1984).
[8] C. L. Henley, Annu. Rev. Condens. Matter Phys. **1**, 179 (2010).
[9] D. A. Huse, W. Krauth, R. Moessner, and S. L. Sondhi, Phys. Rev. Lett. **91**, 167004 (2003).
[10] C. Castelnovo, R. Moessner, and S. L. Sondhi, Nature **451**, 42 (2008).
[11] R. F. Wang *et al.*, Nature **439**, 303 (2006).
[12] S. Ladak, D. E. Read, G. K. Perkins, L. F. Cohen, and W. R. Branford, Nat. Phys. **6**, 359 (2010).

[13] P. Fendley, R. Moessner, and S. L. Sondhi, Phys. Rev. B **66**, 214513 (2002).
[14] V. B. Priezzhev and P. Ruelle, Phys. Rev. E **77**, 061126 (2008).
[15] M. Ciucu, Trans. Amer. Math. Soc. **362**, 4921 (2010).
[16] M. Ciucu and C. Krattenthaler, Commun. Math. Phys. **302**, 253 (2011).
[17] N. Allegra, Nucl. Phys. B **894**, 685 (2015).
[18] W. Jockusch, J. Propp, and P. Shor, arXiv:math/9801068 (1998).
[19] H. Cohn, R. Kenyon, and J. Propp, J. Amer. Math. Soc. **14**, 297 (2001).
[20] R. Kenyon and A. Okounkov, Acta Math. **199**, 263 (2007).
[21] P. Deift, A. Its, and I. Krasovsky, Ann. Math. **174**, 1243 (2011).
[22] E. Basor and T. Ehrhardt, Oper. Theory Adv. Appl. **259** (2017); arXiv:1603.00506.
[23] R. Gharakhloo and A. Its, SIGMA **16**, 100 (2020).
[24] N. Elkies, G. Kuperberg, M. Larsen, and J. Propp, J. Algebraic Combin. **1**, 111 (1992); **1**, 219 (1992).
[25] S. Chhita and B. Young, Adv. Math. **259**, 173 (2014).
[26] H. Cohn, N. Elkies, and J. Propp, Duke Math. J. **85**, 117 (1996).
[27] B. Lindström, Bull. London Math. Soc. **5**, 85 (1973).
[28] I. Gessel and G. Viennot, Adv. Math. **58**, 300 (1985).
[29] E. H. Kuo, Theor. Comput. Sci. **319**, 29 (2004).
[30] F. Alet, J. L. Jacobsen, G. Misguich, V. Pasquier, F. Mila, and M. Troyer, Phys. Rev. Lett. **94**, 235702 (2005).
[31] M. Russkikh, Probab. Theory Relat. Fields **171**, 281 (2018).
[32] J. Dubédat, J. Amer. Math. Soc. **28**, 1063 (2015).
[33] F. Colomo and A. Sportiello, J. Stat. Phys. **164**, 1488 (2016).
[34] W. Krauth and R. Moessner, Phys. Rev. B **67**, 064503 (2003).
[35] N. Wilkins and S. Powell, Phys. Rev. E **104**, 014145 (2021).
[36] W.-J. Tzeng and F. Y. Wu, J. Stat. Phys. **110**, 671 (2003); arXiv:cond-mat/0203149.
[37] F. Y. Wu, Phys. Rev. E **74**, 020104(R) (2006); arXiv:cond-mat/0607647.
[38] J. Bouttier, M. Bowick, E. Guitter, and M. Jeng, Phys. Rev. E **77**, 041130 (2008).
[39] M. Ciucu, *The interaction of collinear gaps of arbitrary charge in a two dimensional dimer system*, arXiv:1202.1188.
[40] M. Ciucu, Commun. Math. Phys. **344** (2016); arXiv:1405.2799.
[41] M. Ciucu, Proc. Amer. Math. Soc. **145**, 4931 (2017).
[42] M. Ciucu, Commun. Math. Phys. **334** (2015); arXiv:1403.0925.
[43] M. Ciucu and I. Fischer, *A triangular gap of size two in a sea of dimers on a* 60° *angle*, arXiv:1405.4235.
[44] M. Ciucu and I. Fischer, *Lozenge tilings of hexagons with arbitrary dents*, arXiv:1412.3945.
[45] H. Cohn, M. Larsen, and J. Propp, New York J. Math. **4**, 137 (1998).
[46] J. L. Cardy, Nucl. Phys. B **240**, 514 (1984).
[47] A. Aggarwal, Invent. Math. **220**, 611 (2020).