Deformable polyhedral spin bodies: multipole channel selection, entropic phase ladders, and non-Abelian orientational order

Changyeon Won[1, *]

[1] *Department of Physics, Kyung Hee University, Seoul 02447, Republic of Korea*

(Dated: November 3, 2026)

We study lattices of "spin bodies": clusters of n classical Heisenberg spins that are coupled antiferromagnetically and through higher-order exchange inside each cluster, and ferromagnetically through multi-order exchange between clusters that does not distinguish the spins of a cluster. Inside a cluster the antiferromagnetic exchange enforces a polyhedral arrangement (a regular tetrahedron for $n = 4$), so the low-energy object is an orientable polyhedron whose order-parameter space is $SO(3)/T$ with the non-Abelian fundamental group $2T$. Unlike rigid-body models, the polyhedra here are made of spins and can collapse. Decomposing all couplings into multipole channels l we prove that the zero-temperature state is uniform for any lattice, reduce the problem to n particles on a sphere, and obtain exact phase boundaries. Odd channels drive collapse into ferromagnetism and even channels into a collinear nematic, with the ferro–nematic boundary fixed by $z\, g_{\rm odd}(1) = f_{\rm odd}(1)$ for every n. A harmonic spin-wave plus configurational (spin-label) free energy then predicts, without fitting parameters, an entropy-driven shift of this boundary, a reentrant ferro-to-nematic transition, and a thin wedge of an equilibrium collinear ferrimagnet between them. Pre-registered Monte Carlo simulations in two and three dimensions confirm the predicted boundary slope $a = 0.392 \pm 0.010$ (theory 0.382–0.390), the reentrance temperature 0.2300 ± 0.0026 (theory 0.234 ± 0.008), and the ferrimagnetic wedge in both dimensions; including the anharmonic free energy by thermodynamic integration reproduces the wedge positions. Transitions with a cubic Landau invariant ($l = 2$) are first order, whereas the octupolar ($l = 3$) orientational transition of the tetrahedral phase is consistent with a continuous transition up to $L = 14$. For $n = 6$ the same free energy predicts a four-step ladder; slab coexistence confirms the first collinear step (the M4 phase) and places all boundaries within 0.02 of the prediction, but does not resolve the M2 phase. We also predict a degenerate prism–octahedron family selected by the ratio of cubic and quartic inter-cluster exchange.

I. INTRODUCTION

Orientational order of objects with polyhedral symmetry is a recurring theme in condensed matter. Tetrahedral and other polyhedral nematics have been proposed for bent-core liquid crystals [1, 2], the cyclic phase of spin-2 Bose–Einstein condensates carries a tetrahedral order parameter with non-Abelian vortices [3–5], and lattice gauge theories have mapped out the generic behavior of nematics with arbitrary point-group symmetry [6–8]. In magnetism, noncollinear antiferromagnets are naturally described by cluster multipoles [9], multipolar order is central to f-electron physics [10], and in breathing pyrochlores strongly bound tetrahedra are weakly coupled to each other [11, 12]. Molecular solids such as methane realize the quantum version of orientational order of tetrahedral rotors coupled by octupole interactions [13].

In most of these settings the polyhedron is either rigid by construction (gauge models, molecules) or is a single-site multipole. A different situation arises when the polyhedron is *made of spins*. Then the same exchange that makes the polyhedron also allows it to deform, merge spins, or collapse into a ferromagnet. Two questions follow: Which couplings between such deformable polyhedra stabilize orientational order, and which make the polyhedra collapse? And when competing collapsed states are degenerate in energy, what selects among them at finite temperature?

In this paper we answer both questions for a minimal model, which we call a lattice of spin bodies [Fig. 1]. Each body consists of n classical Heisenberg spins with antiferromagnetic (AF) bilinear and biquadratic exchange inside the body, and the bodies are coupled ferromagnetically by cubic and quartic exchange that treats the spins of a body as indistinguishable. Our main findings are:

(i) *Channel selection.* Writing every coupling in Legendre channels l, the zero-temperature problem reduces exactly to n particles on a sphere with a pair potential whose spectrum is $\kappa_l = f_l - z g_l$. Odd channels can only produce magnetization and even channels only nematic order. The ferromagnet–nematic boundary is $z g_{\rm odd}(1) = f_{\rm odd}(1)$ for every n and every truncation. In simulations, nematic transition temperatures are independent of the odd-channel coupling, and nematic bodies decouple completely when only odd channels are present.

(ii) *Entropic phase ladder.* Along the ferro–nematic line all collinear states have energies $-\delta M^2/2$ with $\delta = zG - J$, while the number of ways to choose which spins are reversed gives an entropy $\ln \binom{n}{k}$. The competition between a concave energy and a convex entropy produces a ladder of collinear phases. A harmonic spin-wave plus label free energy, with no adjustable parameters, predicts

* cywon@khu.ac.kr

the boundary slope, a reentrant transition, and a ferrimagnetic wedge; all three are confirmed quantitatively.

(iii) *Transition order from invariants.* The $l = 2$ nematic order parameter has a cubic invariant and its transitions are first order, while the $l = 3$ octupole of the tetrahedral phase has none. The tetrahedral orientational transition is consistent with continuous behavior up to $L = 14$, in contrast to the generically first-order transitions found for rigid polyhedral gauge models [8].

We also derive the defect algebra of the tetrahedral phase, and for $n = 6$ test a four-step entropic ladder (its first step is confirmed) and predict an exactly degenerate prism–octahedron family. All finite-temperature hypotheses were tested with decision rules fixed before the corresponding simulations; we report every verdict, including rules that failed by design, in Table IV.

The paper is organized as follows. Section II defines the model and the channel decomposition. Section III derives the zero-temperature phase diagram and exact boundaries. Section IV discusses symmetry, topology, and Landau invariants. Section V develops the entropic free energy. Section VI summarizes the Monte Carlo method and Sec. VII the results. Section VIII gives predictions for $n = 6$ and Sec. IX discusses implications.

II. MODEL AND MULTIPOLE CHANNELS

Each site a of a lattice with coordination z hosts a body of n unit vectors $\mathbf{S}_{a,i}$. The Hamiltonian is

$$H = \sum_a \sum_{i<j} \left[J x_{ij}^{aa} + K (x_{ij}^{aa})^2 \right] - \sum_{\langle ab \rangle} \sum_{i,j=1}^{n} \left[G (x_{ij}^{ab})^3 + H_4 (x_{ij}^{ab})^4 \right], \tag{1}$$

with $x_{ij}^{ab} = \mathbf{S}_{a,i} \cdot \mathbf{S}_{b,j}$ and $J = 1$. The inter-body sum runs over all n^2 pairs, so H is invariant under independent permutations S_n of the spins of each body. We use the square ($z = 4$) and simple cubic ($z = 6$) lattices and quote couplings as zG and zH_4.

Channel decomposition. Expanding a pair function in Legendre polynomials, $f(x) = \sum_l f_l P_l(x)$, and using the addition theorem gives, for a uniform state,

$$E_{\text{body}} = \frac{1}{2} \sum_l \kappa_l A_l + \text{const}, \tag{2}$$

$$A_l = \sum_{i,j} P_l(\mathbf{S}_i \cdot \mathbf{S}_j) = \frac{4\pi}{2l+1} |\mathbf{q}^{(l)}|^2,$$

where $q_m^{(l)} = \sum_i Y_{lm}(\mathbf{S}_i)$ is the body multipole and $\kappa_l = f_l - z g_l$. For Eq. (1),

$$\kappa_1 = J - \tfrac{3}{5} zG, \qquad \kappa_2 = \tfrac{2}{3} K - \tfrac{4}{7} zH_4,$$
$$\kappa_3 = -\tfrac{2}{5} zG, \qquad \kappa_4 = -\tfrac{8}{35} zH_4. \tag{3}$$

A monomial x^k has non-negative Legendre coefficients, so a ferromagnetic monomial coupling contributes $g_l \geq 0$ in every channel it contains. Note that $x^3 = (2P_3 + 3P_1)/5$: a "cubic" coupling also contains a dipole part.

Uniform-state theorem. If $g_l \geq 0$ for all l, the zero-temperature ground state is uniform on any lattice. On each bond, $-g_l \mathbf{q}_a \cdot \mathbf{q}_b \geq -\frac{g_l}{2}(|\mathbf{q}_a|^2 + |\mathbf{q}_b|^2)$, so the total energy is bounded below by a sum of single-body energies, and the bound is reached when all $\mathbf{q}_a$ coincide. The problem therefore reduces exactly to n particles on the unit sphere interacting through the pair potential

$$\Phi(x) = \sum_l \kappa_l P_l(x). \tag{4}$$

Positive κ_l spreads the particles (spherical designs [14]), and negative κ_l makes them merge, in direct analogy with the cluster criterion for bounded pair potentials [15]. Phases are therefore classified by the partition of n into coincident spins and the geometry of the resulting directions.

III. ZERO-TEMPERATURE PHASES AND EXACT BOUNDARIES

Figure 2 shows the ground-state phases for $n = 4$, obtained by minimizing Eq. (4) over all configurations, together with the analytic boundaries derived below. Five phases appear: the tetrahedron T ($M = 0$, a two-design), a ferrimagnet U (a coincident pair plus two single spins at 120°, $|M|/n = 1/4$ exactly), a bent pair of dimers D, the collinear nematic C ($\uparrow\uparrow\downarrow\downarrow$, $M = 0$), and the ferromagnet F. Note that partially collapsed phases (U, D) have no counterpart in rigid-polyhedron models.

Ferro–nematic theorem. The F and C states both reach the maximum $A_l = n^2$ in every even channel. In odd channels F has $A_l = n^2$ while C has $A_l = M^2 = 0$. Hence

$$E_F - E_C = \frac{n^2}{2} \sum_{l\,\text{odd}} \kappa_l \;\Rightarrow\; z\, g_{\text{odd}}(1) = f_{\text{odd}}(1), \tag{5}$$

with $h_{\text{odd}}(1) = [h(1) - h(-1)]/2$. For Eq. (1) this is $zG = J$, independent of K, H_4, and n. Odd channels can therefore only drive collapse into ferromagnetism, and even channels only into the collinear nematic.

Other boundaries. With the closed-form energies of Table I, T–F and T–C are

$$\kappa_1 + \kappa_2 = -\tfrac{4}{9}\kappa_3 - \tfrac{20}{27}\kappa_4, \qquad \kappa_2 = \tfrac{5}{9}\kappa_3 - \tfrac{20}{27}\kappa_4. \tag{6}$$

In physical units these read $(7/9)zG + (20/27)zH_4 = J + 2K/3$ and $zH_4 = 0.9K + 0.3zG$. The D phase is a pair of dimers at angle $c = \cos\theta$, with energy $4\Phi(1) + 4\Phi(c)$. It bends continuously out of C when $\Phi'(-1)$ changes sign,

$$\kappa_1 - 3\kappa_2 + 6\kappa_3 - 10\kappa_4 = 0 \;\Leftrightarrow\; zG = \frac{J - 2K + 4zH_4}{3}, \tag{7}$$




FIG. 1. Model. (a) A spin body of $n = 4$ spins. Intra-body AF exchange $Jx + Kx^2$ ($x = \mathbf{S}_i \cdot \mathbf{S}_j$) favors the regular tetrahedron. (b) Bodies on a lattice are coupled by $-\sum_{i,j}[Gx^3 + H_4x^4]$ between *all* pairs of spins of neighboring bodies, so the coupling cannot tell which spin is which. (c) Channel spectrum for $J = 1$, $K = 0.5$, $zG = 1$, $zH_4 = 0.5$: intra coefficients f_l, inter coefficients $-zg_l$, and their sum κ_l. Odd channels ($l = 1, 3$) carry magnetization; even channels ($l = 2, 4$) carry nematic order.

and meets F first-order at

$$\kappa_1 + \tfrac{3}{4}\kappa_2 + \tfrac{3}{8}\kappa_3 - \frac{9\kappa_2^2}{40\kappa_3} = 0 \quad (\kappa_4 = 0). \tag{8}$$

All boundaries agree with the numerical minimization to the grid resolution [Fig. 2] and with the low-temperature Monte Carlo phases.

Local stability. The Hessian of the tetrahedron splits under T_d into three rotational zero modes and E and T_2 modes with

$$\lambda_{T_2} = \tfrac{8}{3}(\kappa_1 + \kappa_2) - \tfrac{32}{3}\kappa_3 + \tfrac{160}{81}\kappa_4,$$
$$\lambda_E = 8\kappa_2 - \tfrac{40}{3}\kappa_3 - \tfrac{40}{9}\kappa_4. \tag{9}$$

For pure cubic inter-body coupling both eigenvalues *increase* with G, because the tetrahedron is a local maximum of the octupole A_3. The T phase therefore never becomes locally unstable, and its collapse is nucleation driven. The F state is locally stable when $\Phi'(1) = \sum_l \kappa_l\, l(l+1)/2 < 0$.

IV. SYMMETRY, TOPOLOGY, AND LANDAU INVARIANTS

Full symmetry and chirality. Every term of Eq. (1) depends only on dot products, so H is invariant under $O(3) = SO(3) \times \{1, -1\}$, where -1 is time reversal. All isotropy groups in Table I contain improper elements. For example, time reversal maps a tetrahedron onto its dual, which is a 90° rotation of the original. Hence $\pi_0 = 0$: there are no chiral domains and no vestigial chiral phase of the type found for chiral polyhedral nematics [6]. The allowed vestigial phases follow from the group chains $T_d \subset O_h$ (a cubatic phase with only $l = 4$ order) and $C_{\infty v} \subset D_{\infty h}$ (a nematic phase left behind by a melting ferromagnet).

Landau invariants. The number of rotational invariants of a multipole order parameter $\mathbf{q}^{(l)}$ follows from the character formula. Cubic invariants exist only for even l, because $(l\,l\,l; 0\,0\,0)$ vanishes for odd l (Table II). The nematic ($l = 2$) and cubatic ($l = 4$) transitions out of the isotropic phase are therefore first order in mean-field theory, whereas for the tetrahedral octupole ($l = 3$) Landau theory does *not* enforce a first-order transition [Fig. 3]. A first-order transition can still arise through the coupling $(3\,3\,4)$ to the induced hexadecapole, or through fluctuations. The two independent quartic invariants of $l = 3$ select between the two most symmetric octupole orbits, T_d (xyz) and D_{3h} ($\mathrm{Re}(x + iy)^3$). These are exactly the octupoles of the $n = 4$ tetrahedron and of the $n = 6$ prism (Sec. VIII), so both cluster sizes are two branches of a single octupolar Landau theory.

Defect algebra of the T phase. Lifting T to $SU(2)$ gives the binary tetrahedral group $2T$ of order 24 with seven conjugacy classes. Line defects (or point defects in 2D) are labeled by these classes. In an isotropic-stiffness approximation their energy scales with the square of the rotation angle [Fig. 4]. The 2π class and the $4\pi/3$ classes are unstable to splitting, so the elementary defects are the eight $2\pi/3$ elements. Non-commuting lines cannot cross freely [16, 17]. For example, the π lines about x and y leave a connecting rung with holonomy $[i, j] = -1$. On a periodic lattice the ordered product of plaquette holonomies equals the commutator of the two winding holonomies, so a single non-trivial defect is allowed on a torus. In the Abelian phases U, D ($\mathbb{Z}_4$) and C ($\mathbb{Z}_2$) lines




FIG. 2. Zero-temperature phases for $n = 4$. (a) Channel space with $\kappa_3 = -1$, $\kappa_4 = 0$, from direct minimization (colors) and exact boundaries (black): T–F, Eq. (6) (solid), D–F, Eq. (8) (dashed), and C–D, Eq. (7) (dotted). U′ is an umbrella variant of U. (b) Physical plane at $J = 1$, $K = 0.5$ with T–C (solid), T–F (dashed), and C–F (dotted) boundaries of Table I. (c) Sketches of the cluster states; $\times k$ marks k coincident spins.

TABLE I. Phases of the $n = 4$ model: isotropy group $H \subset O(3)$ (time reversal acts as inversion), fundamental group of the order-parameter space, non-zero multipoles, and zero-temperature energy per body for $L = 0$, $G_1 = 0$ in Eq. (1).

Phase	H	π_1	$\mathbf{q}^{(l)} \neq 0$	E_{body}
T	T_d	$2T$	$l = 3, 4$	$-2J + \frac{2}{3}K - z(\frac{16}{9}G + \frac{56}{27}H_4)$
U, D	C_{2v}	$\mathbb{Z}_4$	all	numerical
C	$D_{\infty h}$	$\mathbb{Z}_2$	$l = 2, 4$	$-2J + 6K - 8zH_4$
F	$C_{\infty v}$	0	all	$6J + 6K - 8z(G + H_4)$

cross freely.

Base space, target space, and dimensionality. Two spaces enter the coarse-grained description. The field is a map $\phi : \mathbb{R}^d \to \mathcal{M}$ from the lattice of bodies (the base space, $d = 2$ or 3) to the order-parameter manifold $\mathcal{M} = O(3)/H$, which is fixed by the cluster state. Defects of codimension k in the base space are classified by $\pi_{k-1}(\mathcal{M})$ and nonsingular textures by $\pi_d(\mathcal{M})$ [17]. The same phase therefore hosts different defects in 2D and in 3D (Table III). In 2D, the geometry of $\mathcal{M}$ decides what happens at $T > 0$. If $\mathcal{M}$ contained a flat $U(1)$ direction, spin waves would leave quasi-long-range order and vortex unbinding would produce a Kosterlitz–Thouless (KT) transition with a universal stiffness jump [23, 24]. Here the full symmetry is $O(3)$, with no $U(1)$ factor, so every $\mathcal{M}$ is a homogeneous space of $SO(3)$ with positive curvature. The corresponding 2D sigma models are asymptotically free [22]: correlations decay exponentially at any $T > 0$, with $\xi \propto \exp(c/T)$, consistent with the Mermin–Wagner theorem [25]. *No phase of the model has a true KT transition in 2D.* A finite π_1 still per-

TABLE II. Number of independent rotational invariants of $\mathbf{q}^{(l)}$.

l	1	2	3	4	6
cubic	0	1	0	1	1
quartic	1	1	2	2	3




FIG. 3. Landau free energies. (a) With a cubic invariant ($l = 2$), a second minimum appears at finite amplitude and the transition is first order. (b) Without one ($l = 3$), the minimum moves continuously away from zero. In the full theory the induced $l = 4$ component can still make the $l = 3$ transition weakly first order.




FIG. 4. Defect classes of the T phase, labeled by their rotation angle θ, with line energy $\propto \theta^2$ (isotropic-stiffness estimate). The box lists the energetically favored splittings; the C_2 splitting is marginal and depends on elastic anisotropy.

mits a defect-driven topological transition or crossover of the Kawamura–Miyashita type [26]. The only KT-like behavior is a crossover on the C→D line, where the director correlation length exceeds the system size and, within that length, $\mathbf{M}_\perp$ breaks the $U(1)$ of rotations about the director. In 3D continuous symmetries can break, the transition order follows the Landau counting above, and line defects (π_1) and, in the C and F phases, hedgehogs (π_2) appear.

Fusion rules in two dimensions. In 2D a point defect is labeled by the conjugacy class of its holonomy, because carrying it around other defects conjugates the holonomy. Two defects of classes a and b fuse into the class of $g_a g_b$ with $g_a \in a$, $g_b \in b$. For an Abelian π_1 this product is unique and defects behave as charges of a Coulomb gas, as in the $\mathbb{Z}_2$ (C) and $\mathbb{Z}_4$ (U, D) phases. For a non-Abelian π_1 the outcome depends on how the two defects were brought together relative to the others. If the relative holonomy is uniform over the class, as for paths randomized by thermal motion, the outcome c occurs with probability $p_{ab}^c = |\{g \in b : g_a g \in c\}|/|b|$. Figure 5 shows these statistical fusion rules for $2T$ and Dic_3, computed exactly from the groups. Three consequences follow. First, a defect and its antidefect annihilate only with probability $1/|C_a|$: 1/4 for the elementary $2\pi/3$ classes of the T phase and 1/6 for the π class, whose pair otherwise leaves a π defect (probability 2/3) or a 2π defect (1/6). Second, the same pair can fuse through channels of opposite sign of the interaction energy. With line energy $\propto \theta^2$, a channel is attractive when $\theta_c^2 < \theta_a^2 + \theta_b^2$. Two like $2\pi/3$ vortices (6^+6^+) fuse attractively into 6^- with probability 3/4 and repulsively into 3^+ with probability 1/4, whereas like vortices of an Abelian Coulomb gas always repel. Third, the central element -1 fuses deterministically with every class. The defect gas of the non-Abelian phases is therefore not a Coulomb gas: annihilation is suppressed, and a single unbinding temperature need not exist. This is one reason why the 2D defect-unbinding test (H2) could not be decided with a single-scale detector.

V. ENTROPIC SELECTION NEAR THE FERRO–NEMATIC LINE

At $zG = J$ the C and F states are degenerate [Eq. (5)], and so is every collinear state with k reversed spins and moment $M = n - 2k$. Near this line their energies are exactly

$$E_k - E_F = \tfrac{1}{2}x\,(n^2 - M^2), \qquad x = zG - J, \qquad (10)$$

which is concave in M. Two entropic contributions distinguish them. The first is the harmonic spin-wave

(a) T phase: $\pi_1 = 2T$ (24 elements)

	6^+	6^-	4	3^+	3^-	-1
6^+	6^- 3/4 3^+ 1/4	1 1/4 4 3/4	6^+ 1/2 3^- 1/2	4 3/4 -1 1/4	6^- 1/4 3^+ 3/4	3^-
6^-	1 1/4 4 3/4	6^+ 3/4 3^- 1/4	6^- 1/2 3^+ 1/2	6^+ 1/4 3^- 3/4	4 3/4 -1 1/4	3^+
4	6^+ 1/2 3^- 1/2	6^- 1/2 3^+ 1/2	1 1/6 4 2/3 -1 1/6	6^- 1/2 3^+ 1/2	6^+ 1/2 3^- 1/2	4
3^+	4 3/4 -1 1/4	6^+ 1/4 3^- 3/4	6^- 1/2 3^+ 1/2	6^+ 3/4 3^- 1/4	1 1/4 4 3/4	6^-
3^-	6^- 1/4 3^+ 3/4	4 3/4 -1 1/4	6^+ 1/2 3^- 1/2	1 1/4 4 3/4	6^- 3/4 3^+ 1/4	6^+
-1	3^-	3^+	4	6^-	6^+	1

(b) P/222 phases: $\pi_1 = \mathrm{Dic}_3$ (12)

	6	4_a	4_b	3	-1
6	1 1/2 3 1/2	4_b	4_a	6 1/2 -1 1/2	3
4_a	4_b	6 2/3 -1 1/3	1 1/3 3 2/3	4_a	4_b
4_b	4_a	1 1/3 3 2/3	6 2/3 -1 1/3	4_b	4_a
3	6 1/2 -1 1/2	4_a	4_b	1 1/2 3 1/2	6
-1	3	4_b	4_a	6	1

green: attractive channel; red: repulsive; framed: annihilation possible; shading: number of channels




FIG. 5. Fusion of point defects in two dimensions, computed from the groups (`scripts/fusion_rules.py`). (a) T phase, $\pi_1 = 2T$; classes are labeled by element order: $6^\pm$ ($\theta = 2\pi/3$), 4 (π), $3^\pm$ ($4\pi/3$), -1 (2π); the identity row and column are omitted. Row a fused with column b gives the listed classes with the listed probabilities for a uniform relative holonomy. Green: attractive channel ($\theta_c^2 < \theta_a^2 + \theta_b^2$); red: repulsive; framed cells: annihilation possible; shading: one, two, or three channels. (b) Prism and 222 phases, $\pi_1 = \mathrm{Dic}_3$. (c) Probability that a defect annihilates with its antidefect, one dot per class: 1 for the Abelian groups, $1/|C_a|$ otherwise.

free energy $\frac{T}{2}\sum_{\mathbf{q}} \ln\det{}'\mathcal{H}(\mathbf{q})$ of the uniform state, with global zero modes removed. The second is a configurational term $-T\ln\binom{n}{k}$, which counts the labelings of which spins are reversed. It is exact because the inter-body coupling fixes only M_a, not the labels, and it is convex in M. Defining $u = x/T$,

$$\frac{\Delta F_k}{T} = a_k u + \delta_k, \qquad a_k = \frac{n^2 - M^2}{2}, \tag{11}$$
$$\delta_k = \tfrac{1}{2}\ln\frac{\det{}'_k}{\det{}'_F} - \ln\binom{n}{k}.$$

The equilibrium state is the lower envelope of these straight lines [Fig. 6]. For $n = 4$ this envelope contains an intermediate collinear ferrimagnet M2 ($\uparrow\uparrow\uparrow\downarrow$) whenever $4\delta_{\mathrm{M2}} < 3\delta_C$, which holds. The predicted phase sequence on heating at fixed $zG > J$ is therefore F→M2→C: a *reentrant* nematic phase stabilized by entropy, with the ferromagnet at low temperature.

The theory gives three parameter-free predictions for the 2D model at $K = 0.5$, $zH_4 = 1.5$, which we tested: (1) The low-temperature slope of the boundary, $zG^*(T) = J + aT + bT^2$. Harmonic plus label theory gives $a = 0.382$ for a direct C|F boundary and $a = 0.390$ for the F|M2 boundary. The label term alone gives 0.224, and the harmonic term alone gives 0.158. (2) A reentrance temperature $T_{\mathrm{re}} \simeq 0.234 \pm 0.008$ at $zG = 1.1$, obtained from the low-temperature fit. (3) A wedge of M2 between $T_1 = 0.2431$ and $T_2 = 0.2646$ at $zG = 1.1$ (2D), and between 0.2440 and 0.2665 in 3D.

TABLE III. Order-parameter manifolds $\mathcal{M}$ and their consequences in two and three dimensions. Codimension-k defects are classified by $\pi_{k-1}(\mathcal{M})$: point defects in 2D and lines in 3D by π_1, hedgehogs in 3D by π_2; 2D textures by π_2. NA: non-Abelian. "Crossover": no long-range or quasi-long-range order at $T > 0$; $\xi \propto \exp(c/T)$.

Phase	$\mathcal{M}$	$\dim \mathcal{M}$	π_1	π_2	$d = 2$, $T > 0$	$d = 3$
T ($n = 4$)	$SO(3)/T$	3	$2T$	0	crossover; NA point vortices	order; NA lines
P, 222 ($n = 6$)	$SO(3)/D_3$	3	Dic_3	0	crossover; NA point vortices	order; NA lines
O ($n = 6$)	$SO(3)/O$	3	$2O$	0	crossover; NA point vortices	order; NA lines
U, D	$SO(3)/C_2$	3	$\mathbb{Z}_4$	0	crossover; $\mathbb{Z}_4$ vortices	order; $\mathbb{Z}_4$ lines
C	RP^2	2	$\mathbb{Z}_2$	$\mathbb{Z}$	crossover; half vortices, textures	order; half-vortex lines, hedgehogs
F, M2, M4	S^2	2	0	$\mathbb{Z}$	crossover; skyrmion-like textures	order; hedgehogs




FIG. 6. Entropic ladder, Eq. (11), for the 2D model at $zG = 1.1$. Thick segments mark the lower envelope (equilibrium phase); dotted lines are phase boundaries. (a) $n = 4$: F→M2→C, boundaries at $u = 0.411$ and 0.378. (b) $n = 6$: F→M4→M2→C at $u = 0.303$, 0.272, 0.261. Open circles on top: boundaries measured by slab coexistence, $u = (zG - J)/T$ ($n = 6$: M4 resolved, M2 not); crosses: thermodynamic-integration prediction.

Anharmonic correction. Higher orders enter through the measured anharmonic energy of each branch, $\delta e(T) = e(T) - e_0 - \frac{n_{\mathrm{dof}}}{2} T$. The Gibbs–Helmholtz relation gives

$$\delta f(T) = -T \int_0^T \frac{\delta e(T')}{T'^2}\, dT', \tag{12}$$

which reduces to $-cT^2$ when $\delta e = cT^2$ but needs no polynomial fit.

VI. MONTE CARLO METHOD AND DECISION RULES

We simulate Eq. (1) with single-spin Metropolis moves, rigid rotations of whole bodies, and merge/split moves that propose a spin near another spin of the same body. The merge/split proposal density is a von Mises–Fisher mixture, included exactly in the acceptance ratio and verified by mutation tests. Parallel tempering [18] is used across continuous and weakly first-order transitions. Strongly first-order points are bracketed by separate heating and cooling ladders, and their transition temperatures are fixed by direct slab coexistence [19]: two phases share an interface and the sign of the interface velocity is recorded. Equilibrium boundaries near the ferro–nematic line come from slab coexistence and from the Clausius–Clapeyron slope

$$\frac{dzG^*}{dT} = \frac{\Delta E/T}{\langle \partial_{zG} H \rangle_C - \langle \partial_{zG} H \rangle_F}, \tag{13}$$

measured in single-phase chains. Order parameters are the uniform multipole moments $m_l = |\sum_a \mathbf{q}_a^{(l)}|/(Nn)$, which are invariant under spin relabeling. First-order character is judged from energy histograms with an autocorrelation-corrected BIC [20], the Binder-type energy cumulant V, and the size dependence of the specific-heat maximum, $C_{\max} \propto N^a$. Defects are detected from plaquette holonomies in $2T$, obtained by Kabsch alignment [21] of each body to the tetrahedron, both on single bodies and on coarse-grained octupole moments. The stiffness is measured by parallel tempering from the sums of the first and second derivatives of the twisted inter-body bond energies (about the lab axes and about the instantaneous director), with block-bootstrap errors. Lattice sizes are $L = 16$–32 in 2D and $L = 8$–16 in 3D.

All hypotheses H1–H12 were stated, with numerical decision rules, before the corresponding production runs. Where a rule or an estimator later proved inadequate we kept the original verdict, dated the change in the repository, and report the sequence here. Three changes were made after data had been seen. (i) The slab zero-crossing estimator for the boundary slope was changed twice after the first temperature ($T = 0.10$) had been analyzed: from a global linear fit to separate fits on the two sides of the zero, then with seed resampling and a window-choice systematic. The $T = 0.10$ point is therefore not an independent test; the other four temperatures were analyzed after the estimator was fixed, and six alternative estimators change the slope by at most 0.008 (Sec. VII B). (ii) After the 2D wedge measurement, the primary zero-crossing error was changed from the bootstrap standard deviation, which was inflated by rare extrapolated resamples, to the half-width of the 16–84% interval; the change applies only to later measurements. (iii) For the 3D wedge, the pre-registered estimator reverted to first-sign-change interpolation in noisy bins and failed its rule. A seed-averaged estimator was then fixed, T_2 was remeasured with it, and the earlier T_1 data were recomputed; the latter is not an independent test. The failed verdicts remain in the record. The anharmonic correction was done first with a polynomial fit of the branch energies and then, with a rule fixed beforehand, by thermodynamic integration; both are reported. Table IV lists the outcomes.

TABLE IV. Pre-registered hypotheses and verdicts. "Rule" is the verdict of the rule fixed before the run; supplementary measurements are listed separately.

Hypothesis	Verdict (key numbers)
H1 octupolar T transition (3D)	consistent with continuous, $zG = 0.5, 1, 1.5$; $a(12\to14) = 0.14$
H2 defect unbinding (2D)	undecidable (detector resolution)
H3 nematic C, 3D	first order; $T_c = 0.356(2)$, latent heat 2.05(3)
H5 ferrimagnet U	supported (2D, 3D); $m_1 = 0.24$–0.25
H6 boundary slope	harmonic+label supported; $a = 0.392(10)$
H7 vestigial nematic	two separated transitions (4 points)
H12 C→D	consistent with continuous (4 points)
Reentrance F→C	rule: seen on heating only (rule design error); slab: $T_{\rm re} = 0.2300(26)$
M2 wedge 2D/3D	located; final rule (3D): confirmed (width 0.0185 vs $2\sigma = 0.0129$)
Anharmonic (TI)	wedge positions within 0.3–1.2σ (Table V); b at the 2σ edge
ladder $n = 6$ (2D)	M4 phase confirmed; M2 not resolved, final (gap 0.0098 vs $2\sigma = 0.014$)
stiffness T phase (2D)	asymptotic-freedom crossover supported (Fig. 10a)
stiffness C→D (2D)	KT-type crossover consistent; $T^* = 0.172$–0.175, steepening with L

VII. RESULTS

A. Channel selection at finite temperature

The theorem of Eq. (5) has a direct finite-temperature consequence: an order that lives only in even channels is insensitive to the odd-channel coupling G. Figure 7 shows three tests. First, along cut E ($K = -0.5$, $zH_4 = 0$) the inter-body coupling is purely odd, and collinear C bodies, which have $A_{\rm odd} = 0$, do not couple at all. In 3D at $zG = 0.5$ (C) the order parameter decays as $N^{-1/2}$ ($m_2\sqrt{N} = 0.94$ at both $L = 8$ and 10, for ordered and random starts), and the specific-heat maximum does not grow with size (4.84 and 4.77); there is no collective transition. D and F points with random starts did not order within the run length ($m_1 \approx 0$ at the lowest temperature, and five of the six χ_1 peaks sit at the ladder edge); as for the U phase, ordered starts are used. Second, along cut E′ ($K = -0.25$, $zH_4 = 0.25$) the nematic temperature ($T_2 = 0.240$–0.242 in 2D, 0.254 in 3D, for all sizes) is the same at $zG = 0.9$ and 1.0, while the magnetization of the D phase orders at $T_1 = 0.094$ and 0.18 in 2D (0.104 and 0.190 in 3D), respectively. Third, along cut D ($zH_4 = 1.5$), $T_2 = 0.618$–0.623 is again independent of zG. Because the D phase couples only through odd channels, its ordering temperature scales as $G\,M^2$, with $M^2 \propto zG - zG_c$ near the C–D boundary of Eq. (7). This estimate gives the correct trend in both cuts (ratio $T_1(zG{=}1)/T_1(zG{=}0.9)$: 2.8 predicted, 1.90 measured in 2D and 1.82 in 3D for E′; $T_1(0.8)/T_1(1.0)$: 0.32 predicted, 0.26(1) measured for E in 3D at both $L = 8$ and 10, where $T_1 = 0.0252(11)$ and 0.0974(29) at $L = 8$). In 3D, T_1 and T_2 remain separated as L grows (H7), giving a vestigial nematic between the D and isotropic phases. The C→D transition is consistent with being continuous (H12), as expected for a vector order parameter $\mathbf{M}_\perp$ without a cubic invariant.

B. Entropic selection, reentrance, and the M2 wedge

Figure 8 collects the results near the ferro–nematic line. Slab coexistence at $T = 0.10$–0.20 gives $a = 0.392 \pm 0.010$ and $b = 0.15 \pm 0.05$ (2D). This value excludes the label-only (0.224) and harmonic-only (0.158) predictions by more than 12σ and agrees with the harmonic plus label value. As the M2 wedge shows, the measured boundary is in fact the F|M2 line, whose prediction 0.390 coincides with an auxiliary rescan, 0.390 ± 0.007. Sensitivity tests over six zero-crossing estimators give $a = 0.384$–0.392, all with the same verdict. Figure 8 shows the individual zero crossings; the first $T = 0.175$




FIG. 7. Channel selection at finite temperature, from the individual runs. (a) Cut E′: the nematic temperature T_2 (red) does not depend on the odd-channel coupling zG, while the D-phase ordering temperature T_1 (green) does. Circles: 2D, $L = 16$ (open) and 24 (filled); squares: 3D, $L = 8$ (open) and 10 (filled); diamonds: 3D $L = 12$, where the ladder (0.8–1.2 T_1) does not reach T_2. Error bars are jackknife errors of the susceptibility peaks. Dashed (2D) and dotted (3D) lines: $T_1 \propto GM^2$ normalized at $zG = 1$. (b) Cut E, 3D, ordered start: T_1 of the D order at $L = 8$ (open squares) and 10 (filled) and the same scaling; the C state at $zG = 0.5$ shows no collective order (text). Cut D: T_2 from the specific-heat peak of heating and cooling ladders ($L = 16$, triangles) and an $L = 8$ trial (circles).

and 0.20 zeros lie at the edge of the scanned range and are replaced there by the values of a rescan with a larger range.

At $zG = 1.1$ the ferromagnet re-enters the nematic on heating. Because the boundary is strongly first order, heating ladders superheat F to $T = 0.316$ and cooling ladders never leave C. The pre-registered rule, which required both ladders, therefore could not be met. Direct coexistence gives the equilibrium $T_{\rm re} = 0.2300 \pm 0.0026$, inside the predicted window and consistent with the extrapolated fit 0.234 ± 0.008.

The M2 wedge was predicted before it was measured. In 2D, slab coexistence of F|M2 and M2|C gives $T_1 = 0.2338 \pm 0.0005$ and $T_2 = 0.2560 \pm 0.0019$. In 3D ($L = 12$), the final rule with a seed-averaged estimator gives $T_1 = 0.2356 \pm 0.0008$ and $T_2 = 0.2541 \pm 0.0064$, a width of 0.0185 that exceeds twice its uncertainty. An earlier 3D estimator failed on this criterion because it reverted to first-sign-change interpolation; that verdict is retained in the record.

Both positions lie about 0.01 below the harmonic prediction, with the same sign in 2D and 3D. Including the anharmonic free energy by thermodynamic integration, Eq. (12), of the measured branch energies moves all four wedge temperatures onto the measurements (Table V); the corrected values differ from the measurements by 0.3–1.2 combined standard deviations (by 2.0, 1.1, 0.5, and 0.3 times the measurement error alone). The curvature b is reproduced only at the 2σ edge (0.054(4) versus 0.15(5)); its prediction error is negligible, so this test is limited by the measurement. The integral uses branch energies from $T = 0.02$ to 0.26 on 16^2 (2D) and 8^3 (3D) lattices, which are smaller than the slabs. A polynomial fit of the same energies gives the same direction, but its prediction error (0.017–0.06) is dominated by the choice of fit range, which is why we use the integral.

TABLE V. Thermodynamic-integration (TI) prediction against slab measurement at $zG = 1.1$ (2D $L = 24$ slab, 3D $L = 12$ slab; branch chains 16^2 and 8^3). The prediction error is the propagated statistical error of the branch energies; Δ/σ divides the difference by the combined error. T_1: F→M2, T_2: M2→C; b is the curvature of the 2D boundary $zG^* = J + aT + bT^2$.

	harmonic	TI	measured	Δ/σ
T_1 (2D)	0.2431	0.2348(7)	0.2338(5)	1.2
T_2 (2D)	0.2646	0.2539(23)	0.2560(19)	0.7
T_1 (3D)	0.2440	0.2352(5)	0.2356(8)	0.4
T_2 (3D)	0.2665	0.2560(18)	0.2541(64)	0.3
b (2D)	0	0.054(4)	0.15(5)	1.9

C. Transition order

Figure 9 compares the effective exponent $a = d\ln C_{\max}/d\ln N$ across transitions. For the octupolar transition of the T phase (cut A), the exponents between the two largest sizes are 0.14(24) at $zG = 0.5$ ($L = 12 \to 14$), 0.22(18) at 1.0, and 0.17(11) at 1.5 (Fig. 9a), energy histograms are unimodal, and $|2/3 - V_{\min}|$ decreases with L. In 2D, the same exponent is consistent with zero




FIG. 8. Phases near the ferro–nematic line (2D, $K = 0.5$, $zH_4 = 1.5$). Thin colored lines: harmonic plus label predictions for F|M2 and M2|C. Black line and band: fit $zG^* = 1+aT+bT^2$; black symbols: the individual slab zero crossings (dots: input to the fit; open diamonds: rescan of $T = 0.175$ and 0.20 with a larger range, auxiliary). Gray lines: label-only and harmonic-only slopes. At $zG = 1.1$: wedge boundaries T_1 (F|M2, purple) and T_2 (M2|C, red) from slab coexistence in 2D (open circles) and 3D (squares) with their errors, and the thermodynamic-integration predictions (×).

(-0.14 for $L = 24 \to 32$ at $zG = 1$). We therefore classify this transition as consistent with continuous. At $zG = 0.5$ the global fit over $L = 8$–14 gives $a = 0.48$ (68% interval 0.42–0.53), larger than any standard 3D universality class, although the pairwise values fall from 0.49 and 0.67 for the smaller pairs to 0.14(24) for $12 \to 14$. This leaves room for weak first-order behavior with a correlation length exceeding L, or for crossover near a tricritical point; an exploratory point at $zG = 0.3$ did not support the latter. Transitions with an $l = 2$ invariant are clearly first order. In 3D at $zH_4 = 1$, branch energies and the coexistence window are independent of L, with latent heat 2.05 ± 0.03 per body, while the slab-coexistence T_c decreases from 0.3578(3) at $L = 12$ to 0.3554(2) at $L = 16$ (Fig. 9b). In 2D at $zH_4 = 2$ the specific-heat maximum grows with exponent 0.72(16) ($L = 16 \to 24$). The contrast between $l = 3$ and $l = 2$ follows the Landau counting of Table II. It also distinguishes deformable spin bodies from rigid polyhedral gauge models, for which first-order transitions are generic [8].

D. Partially collapsed ferrimagnet

The U phase, a dimer plus two single spins with $|M|/n = 1/4$, is found at finite temperature in both 2D and 3D from ordered starts (cut C: $K = 0$, $zH_4 = 0$, $zG = 1.0$ and 1.2). The measured magnetization $m_1 = 0.24$–0.25 matches the exact value, and the fraction of bodies in the {211} partition is 1.00 at low temperature. In 3D the upper end of the U interval (0.140/0.143 and 0.176/0.177 for $L = 8/10$) does not drift with L. U is therefore a genuine thermodynamic phase, not a finite-size effect.

E. Defects

In 2D the octupole correlation function decays algebraically over the accessible sizes. Because the $SO(3)/T$ sigma model is asymptotically free [22] and no phase of this $O(3)$ model can have quasi-long-range order in 2D (Sec. IV, Table III), we interpret this as a crossover with a correlation length that grows like $\exp(c/T)$, not as quasi-long-range order. Defects of the $2T$ classes are detected reliably at low temperature. Near the crossover, however, the frame of individual bodies becomes ill defined before defects proliferate, and a scale-ratio test of defect unbinding is undecidable at our resolution (H2). In 3D the frame and the defects disappear together at the first-order transitions, as expected.

The absence of a true KT transition can be tested directly through the stiffness $\Upsilon = L^{-d}[\langle \partial_\theta^2 H\rangle - \beta\langle(\partial_\theta H)^2\rangle]$, where θ is the twist per lattice spacing of all bodies about an axis (Fig. 10); the decision rules were fixed before the runs. In the T phase (cut A, $zG = 1$, $L = 16$, 24, 32) the axis-averaged Υ decreases with L at all five temperatures between 0.6 and $1.0\,T_{\rm peak}$, roughly linearly in $\ln L$ and without an L-independent plateau ($\Upsilon_{32}/\Upsilon_{16} - 1 = -0.35\%$ at $0.6\,T_{\rm peak}$ and -52% at $T_{\rm peak}$), the signature of an asymptotically free crossover rather than quasi-long-range order. As a post hoc observation that is not part of the rule, $(\Upsilon_{16} - \Upsilon_{32})/(T \ln 2)$ lies between 0.10 and 0.18 at the four temperatures from 0.6 to $0.9\,T_{\rm peak}$ (inset of Fig. 10a). Along the C→D line (cut E′, $zG = 1$) a π rotation about the director maps the D state onto its time-reversed state, so $\mathbf{M}_\perp$ carries only 2π vortices and the reference line is $2T/\pi$. The stiffness about the instantaneous director, $\Upsilon_\parallel$, is the same for all three sizes at low T and crosses the line $2T/\pi$ at $T^* = 0.1745$, 0.1727, 0.1720 for $L = 16$, 24, 32, with a crossing slope that steepens with L (13.4, 16.0, 18.7): a KT-type crossover, consistent with the director correlation length exceeding the system size, but not a transition. Above T^*, $\Upsilon_\parallel$ is slightly negative (-0.03), an artifact of using the instantaneous director that does not




FIG. 9. (a) Effective specific-heat exponent $a = d\ln C_{\max}/d\ln N$ for every pair of sizes $L_0 \to L_1$, with 16–84% errors from the block-jackknife errors of $C_{\max}$: the octupolar ($l = 3$) transition of the tetrahedral phase in 3D (filled circles) and 2D (open squares), and the 2D nematic ($l = 2$) transition at $zH_4 = 2$ (triangle). The band marks standard 3D continuous transitions and the dashed line first-order scaling. (b) T_c of the first-order 3D nematic transition ($zH_4 = 1$) from slab coexistence at two slab sizes; the latent heat is from window chains.

enter the rule. A bias of this size near T^* would move T^* by about 0.002 (0.03 divided by the crossing slope), comparable to the spread of $T^*(L)$ over the three sizes, so we do not interpret the L dependence of T^* quantitatively. One seed and three sizes were used, and at $L = 32$ the tempering ladder made only 6 round trips.

VIII. PREDICTIONS FOR SIX-SPIN BODIES

For $n = 6$ the intra-body exchange selects a two-design. Among six points these form a continuous family: two triangles at height $\pm 1/\sqrt{3}$, twisted by φ, interpolating between the prism P ($\varphi = 0$, D_{3h}) and the octahedron O ($\varphi = 60°$, O_h). Along this family

$$A_3 = \tfrac{10}{3}(1 + \cos 3\varphi), \qquad A_4 = \tfrac{119}{9} - \tfrac{70}{9}\cos 3\varphi, \quad (14)$$

so the inter-body energy is linear in $\cos 3\varphi$ with coefficient $\propto \frac{4}{3}G - \frac{16}{9}H_4$ [Fig. 11]. Cubic exchange selects the prism and quartic exchange selects the octahedron. On the line $zH_4 = \frac{3}{4}zG$ the entire family is degenerate. Because the octahedron has $A_3 = 0$, octahedral bodies couple only through $l = 4$. Odd couplings instead favor dimerization into a planar triangle of dimers ("222", $M = 0$, D_{3h}), whose octupole $A_3/n^2 = 0.625$ exceeds that of any six-point design. Its boundaries with C and F are $zH_4 = 0.8K + 0.4zG$ and $13.5zG + 11.25zH_4 = 18J + 9K$. The 222 phase shares the D_{3h} symmetry of the prism, so the two can be connected continuously by flattening, and its defects form the non-Abelian group Dic_3.

The entropic ladder of Eq. (11) for $n = 6$ contains all four collinear states F→M4→M2→C [Fig. 6(b)], with boundaries at $u = 0.303$, 0.272, and 0.261 (2D, $zG = 1.1$), that is $T = 0.330$, 0.367, and 0.383. The ground-state competitors of this family are the four collinear states for $K = 0.5$, $zH_4 = 1.5$ and $zG = 0.95$–1.2; non-collinear states ({321}, {222}, {2211}) lie at least 4.8 above. We tested the ladder by slab coexistence (2D, $L = 24$, three seeds, interface pairs M4|F, M2|M4, C|M2). The three boundaries are at $T = 0.3205(11)$, 0.3544(21), and 0.3641(45), i.e. 0.010, 0.013, and 0.019 below the harmonic values, the same sign as for $n = 4$ and growing with temperature. The M4 phase is resolved ($T_b - T_a = 0.034$ against $2\sigma = 0.005$); the M2 interval is not ($T_c - T_b = 0.0098$ against $2\sigma = 0.0099$). Two refinements, six seeds on a finer grid and runs three times longer, return the same boundary ($T_c = 0.364(11)$ and 0.364(7)) with larger, not smaller, errors, because the C|M2 interface moves at less than 3×10^{-4} per sweep and its velocity near the zero is lost in the seed-to-seed variation. The final pre-registered verdict is therefore *M4 confirmed, M2 not resolved* ($T_c - T_b = 0.0098$ against $2\sigma = 0.014$). Thermodynamic integration of the $n = 6$ branch energies (2D $L = 16$, $T = 0.02$–0.40), computed before the last remeasurement, predicts $T = 0.3204(11)$, 0.3521(22), and 0.3705(67), within 0.1, 0.8, and 0.8 combined standard deviations of the three measured boundaries, and an M2 width 0.018(8) against the measured 0.010. The M2 phase is thus predicted, but our slab method cannot resolve it. The concave-energy versus




FIG. 10. Stiffness of the 2D spin bodies, $L = 16$, 24, 32 (bands: $\pm 2\sigma$, 16–84% block-bootstrap half-widths). (a) T phase (cut A, $zG = 1$): axis-averaged Υ against T/T_{peak}; it decreases with L at all temperatures, by 0.35% at $0.6\,T_{\mathrm{peak}}$ to 52% at T_{peak} (grid lines: the five temperatures of the rule); inset: the small low-temperature decrease, $(\Upsilon_{16} - \Upsilon_L)/T$ for $L = 24$ and 32 ($\pm 1\sigma$; a post hoc view, not part of the rule). (b) C→D line (cut E′, $zG = 1$): stiffness about the instantaneous director, with the line $2T/\pi$ (dashed) and its crossings $T^*(L)$ (open circles).

convex-entropy mechanism generalizes to any n and produces $\lfloor n/2 \rfloor$ collinear steps when all of them lie on the lower envelope.

IX. DISCUSSION

What is new. Three features distinguish spin bodies from rigid polyhedra. First, couplings act channel by channel, and the parity of the channel decides whether a body can only magnetize or only align. This turns a many-parameter problem into a few exact inequalities [Eqs. (5)–(8)]. Second, collapse produces phases with no rigid analog: partially merged ferrimagnets (U, D) and collinear states whose relative weights are set by a counting entropy. The resulting ladder is a simple and quantitative example of order by disorder [27, 28] in which a discrete label entropy and the spin-wave entropy contribute comparably; both are needed to obtain the measured slope. Third, the transition order follows the Landau counting of invariants of the relevant multipole rather than a generic rule for polyhedral order.

Where the model meets experiment. The classical limit is natural when each "spin" is a mesoscopic moment, as in artificial spin systems and nanomagnet arrays [29, 30]. There, the dipolar interaction between elements contains exactly the multipole structure of Eq. (2), and clusters of elements can be engineered with strong internal coupling. The cluster-multipole description of noncollinear antiferromagnets [9] and breathing pyrochlores [11, 12] provides materials in which the channel selection rules may be tested directly. Many artificial spin systems are, however, planar because of shape anisotropy, and this changes the 2D physics qualitatively. Confining the spins to a plane reduces $O(3)$ to $O(2)$. With a higher-order intra-body term selecting the regular n-gon (a circular design), an indistinguishable inter-body coupling then gives $\mathcal{M} = U(1)/\mathbb{Z}_k$ with $k = n$, which is flat. A genuine KT transition follows, mediated by fractional vortices of winding $2\pi/k$. Balancing their energy $\pi\rho\, k^{-2} \ln(R/a)$ against the entropy $2T \ln(R/a)$ gives

$$T_{\mathrm{KT}} = \frac{\pi\rho}{2k^2}, \tag{15}$$

where ρ is the orientational stiffness of the bodies. Because only the lowest circular harmonic compatible with the polygon couples neighboring bodies, ρ is itself fixed by the channel decomposition. The chain polygon symmetry → coupling channel → stiffness → T_{KT} is a direct prediction for planar arrays of nanomagnet clusters.

Limitations. All analytic predictions rest on the uniform-state theorem (exact at $T = 0$) and on low-temperature expansions. Size dependence was checked with two or three sizes, 3D wedge positions with one slab size, and the $n = 6$ ladder with one size and three seeds; the anharmonic integral uses chains smaller than the slabs; the stiffness uses one seed per size. The order of the octupolar transition at $zG = 0.5$ remains open.

X. CONCLUSION

Bodies made of antiferromagnetically coupled spins behave as polyhedra only as long as the inter-body coupling respects their multipole content. Decomposing all couplings into Legendre channels gives a uniform-state theorem, exact phase boundaries, and a parity rule: odd channels magnetize and even channels align. At




FIG. 11. Six-spin bodies. (a) Octupole A_3 and hexadecapole A_4 along the two-design family from the prism to the octahedron. (b) Within this family the inter-body energy selects P or O; on the line $zH_4 = \frac{3}{4} zG$ the whole family is degenerate.

finite temperature, the competition between a concave collinear energy and a convex label entropy, together with spin-wave entropy, produces an entropic ladder. It accounts quantitatively for the boundary slope, a reentrant nematic phase, and an equilibrium collinear ferrimagnet, all confirmed by pre-registered simulations in two and three dimensions. Transitions with a cubic Landau invariant are first order, while the tetrahedral octupolar transition is consistent with continuous behavior, unlike rigid polyhedral models. For six-spin bodies the framework predicts a four-step ladder, of which slab coexistence confirms the first collinear step (the M2 step is predicted, also after the anharmonic correction, but not resolved by our method), and a degenerate prism–octahedron family, and it provides a route to engineering non-Abelian orientational order from spins.

ACKNOWLEDGMENTS

Simulation code, decision rules, the full record of verdicts, and the derived data tables behind every figure are available at https://github.com/cywon-commits/Polyhedron_Spin_Group (directory `paper/`).

Appendix A: Harmonic free energy and label entropy

For a uniform collinear state with k reversed spins, the quadratic Hamiltonian in the $2nN$ transverse coordinates is block diagonal in $\mathbf{q}$. Its $2n \times 2n$ blocks $\mathcal{H}_k(\mathbf{q})$ are built from $\Phi'(\pm 1)$ for intra-body pairs and $g'(\pm 1)$ for inter-body pairs. The classical harmonic free energy is $F_k^{\mathrm{harm}} = \frac{T}{2} \sum_{\mathbf{q}} \ln \det{}' \mathcal{H}_k(\mathbf{q})$, where the prime removes the global zero modes (two for collinear states). Each body independently chooses which k of its n spins are reversed, because uniformity is enforced only for M_a. This gives the exact configurational entropy $\ln \binom{n}{k}$ per body. Low-temperature energies satisfy $e(T) = e_0 + \frac{n_{\mathrm{dof}}}{2} T(1 - n_0/(2N)) + O(T^2)$, with n_0 the number of global zero modes, which we verified to 1%.

[1] L. G. Fel, Phys. Rev. E **52**, 702 (1995).
[2] T. C. Lubensky and L. Radzihovsky, Phys. Rev. E **66**, 031704 (2002).
[3] G. W. Semenoff and F. Zhou, Phys. Rev. Lett. **98**, 100401 (2007).
[4] M. Kobayashi, Y. Kawaguchi, M. Nitta, and M. Ueda, Phys. Rev. Lett. **103**, 115301 (2009).
[5] Y. Kawaguchi and M. Ueda, Phys. Rep. **520**, 253 (2012).
[6] K. Liu, J. Nissinen, R.-J. Slager, K. Wu, and J. Zaanen, Phys. Rev. X **6**, 041025 (2016).
[7] J. Nissinen, K. Liu, R.-J. Slager, K. Wu, and J. Zaanen, Phys. Rev. E **94**, 022701 (2016).
[8] K. Liu, J. Greitemann, and L. Pollet, Phys. Rev. E **97**, 012706 (2018).
[9] M.-T. Suzuki, T. Koretsune, M. Ochi, and R. Arita, Phys. Rev. B **95**, 094406 (2017).
[10] Y. Kuramoto, H. Kusunose, and A. Kiss, J. Phys. Soc. Jpn. **78**, 072001 (2009).
[11] Y. Okamoto, G. J. Nilsen, J. P. Attfield, and Z. Hiroi, Phys. Rev. Lett. **110**, 097203 (2013).

[12] J. G. Rau, L. S. Wu, A. F. May, L. Poudel, B. Winn, V. O. Garlea, A. Huq, P. Whitfield, A. E. Taylor, M. D. Lumsden, M. J. P. Gingras, and A. D. Christianson, Phys. Rev. Lett. **116**, 257204 (2016).
[13] H. M. James and T. A. Keenan, J. Chem. Phys. **31**, 12 (1959).
[14] P. Delsarte, J. M. Goethals, and J. J. Seidel, Geom. Dedicata **6**, 363 (1977).
[15] C. N. Likos, A. Lang, M. Watzlawek, and H. Löwen, Phys. Rev. E **63**, 031206 (2001).
[16] V. Poénaru and G. Toulouse, J. Phys. (Paris) **38**, 887 (1977).
[17] N. D. Mermin, Rev. Mod. Phys. **51**, 591 (1979).
[18] K. Hukushima and K. Nemoto, J. Phys. Soc. Jpn. **65**, 1604 (1996).
[19] J. R. Espinosa, E. Sanz, C. Valeriani, and C. Vega, J. Chem. Phys. **139**, 144502 (2013).
[20] M. S. S. Challa, D. P. Landau, and K. Binder, Phys. Rev. B **34**, 1841 (1986).
[21] W. Kabsch, Acta Crystallogr. A **32**, 922 (1976).
[22] A. M. Polyakov, Phys. Lett. B **59**, 79 (1975).
[23] J. M. Kosterlitz and D. J. Thouless, J. Phys. C **6**, 1181 (1973).
[24] D. R. Nelson and J. M. Kosterlitz, Phys. Rev. Lett. **39**, 1201 (1977).
[25] N. D. Mermin and H. Wagner, Phys. Rev. Lett. **17**, 1133 (1966).
[26] H. Kawamura and S. Miyashita, J. Phys. Soc. Jpn. **53**, 4138 (1984).
[27] J. Villain, R. Bidaux, J.-P. Carton, and R. Conte, J. Phys. (Paris) **41**, 1263 (1980).
[28] C. L. Henley, Phys. Rev. Lett. **62**, 2056 (1989).
[29] L. J. Heyderman and R. L. Stamps, J. Phys.: Condens. Matter **25**, 363201 (2013).
[30] S. H. Skjærvø, C. H. Marrows, R. L. Stamps, and L. J. Heyderman, Nat. Rev. Phys. **2**, 13 (2020).