

STAT 215A Fall 2023

Week 9

Chengzhong Ye

Announcements

- Lab 3 and homework 3 due next **Monday 10/23 at 11:59pm**
 - Write-up: just answer each question
- Midterm: 10/26
- Practice midterm solutions will be released today
 - We'll go over it on the Tuesday 10/24 lecture
- Happy World Statistics Day!



Lab 3: Stability of K-means + Computability

- Ben-Hur, et al. notes that similarity can be computed in $O(k_1 k_2 n)$
 - This should be your goal (though not required)
- You can do better than the Figure 3 in Ben-Hur
- `foreach`
 - If you're having issues with `foreach`, try `future` or `parallel`
- Remember, no need to push a blinded version
- Make sure your `lab3` folder is well-organized and **only** contains the files I would need to reproduce your results (will be part of your grade)!

Outline for today

- Regularization Pt. 1: Ridge

Regularization Part I: Ridge

Thanks to Tiffany Tang for sharing her slides

Expected prediction error

Goal: Evaluate the *error* in estimating an unknown function f by the estimator \hat{f}

Expected prediction error

Goal: Evaluate the *error* in estimating an unknown function f by the estimator \hat{f}

- x_0 : observed “predictors” for a sample (assume fixed, non-random)
-

Expected prediction error

Goal: Evaluate the *error* in estimating an unknown function f by the estimator \hat{f}

- x_0 : observed “predictors” for a sample (assume fixed, non-random)
- $y_0 = f(x_0) + \varepsilon$: the “response” for x_0
 - $f(x_0)$: the “true” value of f at x_0
 - $\varepsilon \sim (0, \sigma^2)$: additive random error
-

Expected prediction error

Goal: Evaluate the *error* in estimating an unknown function f by the estimator \hat{f}

- x_0 : observed “predictors” for a sample (assume fixed, non-random)
- $y_0 = f(x_0) + \varepsilon$: the “response” for x_0
 - $f(x_0)$: the “true” value of f at x_0
 - $\varepsilon \sim (0, \sigma^2)$: additive random error
- $\hat{y}_0 = \hat{f}(x_0)$: the predicted response for x_0

Expected prediction error

Goal: Evaluate the *error* in estimating an unknown function f by the estimator \hat{f}

- x_0 : observed “predictors” for a sample (assume fixed, non-random)
- $y_0 = f(x_0) + \varepsilon$: the “response” for x_0
 - $f(x_0)$: the “true” value of f at x_0
 - $\varepsilon \sim (0, \sigma^2)$: additive random error
- $\hat{y}_0 = \hat{f}(x_0)$: the predicted response for x_0

Expected prediction error (EPE):

$$\mathbb{E}[y_0 - \hat{f}(x_0)]^2 = \underbrace{\text{Var}(\hat{f}(x_0)) + \text{Bias}^2(\hat{f}(x_0))}_{\text{Mean squared error (MSE)}}$$

Expected prediction error

Goal: Evaluate the *error* in estimating an unknown function f by the estimator \hat{f}

- x_0 : observed “predictors” for a sample (assume fixed, non-random)
- $y_0 = f(x_0) + \varepsilon$: the “response” for x_0
 - $f(x_0)$: the “true” value of f at x_0
 - $\varepsilon \sim (0, \sigma^2)$: additive random error
- $\hat{y}_0 = \hat{f}(x_0)$: the predicted response for x_0

Expected prediction error (EPE):

$$\mathbb{E}[y_0 - \hat{f}(x_0)]^2 = \sigma^2 + \underbrace{\text{Var}(\hat{f}(x_0)) + \text{Bias}^2(\hat{f}(x_0))}_{\text{Mean squared error (MSE)}}$$

“irreducible” error

Mean squared error (MSE)

The Bias-Variance Tradeoff


$$\text{MSE} = \text{Var}(\hat{f}(x_0)) + \text{Bias}^2(\hat{f}(x_0))$$

- **Bias:** On average, how wrong is your prediction?

$$\text{Bias}(\hat{f}(x)) = \mathbb{E}(\hat{f}(x)) - f(x)$$

- **Variance:** If you obtain a new, but similar dataset, how much does this change your predictions?

$$\text{Var}(\hat{f}(x)) = \mathbb{E}(\hat{f}(x)^2) - \mathbb{E}(\hat{f}(x))^2$$



The Bias-Variance Tradeoff


$$\text{MSE} = \text{Var}(\hat{f}(x_0)) + \text{Bias}^2(\hat{f}(x_0))$$

- **Bias:** On average, how wrong is your prediction?

$$\text{Bias}(\hat{f}(x)) = \mathbb{E}(\hat{f}(x)) - f(x)$$


- **Variance:** If you obtain a new, but similar dataset, how much does this change your predictions?

$$\text{Var}(\hat{f}(x)) = \mathbb{E}(\hat{f}(x)^2) - \mathbb{E}(\hat{f}(x))^2$$



The Bias-Variance Tradeoff



$$\text{MSE} = \text{Var}(\hat{f}(x_0)) + \text{Bias}^2(\hat{f}(x_0))$$



Recall Ordinary Least Squares

$$\hat{\boldsymbol{\beta}}_{OLS} = \operatorname{argmin}_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X} \boldsymbol{\beta}\|_2^2$$

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$$



OLS

Advantages

- Simple
- Closed-form solution
- Interpretable (?)
- Under some modeling assumptions, OLS has some desirable properties

OLS + Model

Assume $y = X\beta + \varepsilon$ with $\mathbb{E}(\varepsilon) = 0$.

- Then $\hat{\beta}_{\text{OLS}}$ is an unbiased estimator for β

OLS + Model

Assume $y = X\beta + \varepsilon$ with $\mathbb{E}(\varepsilon) = 0$.

- Then $\hat{\beta}_{\text{OLS}}$ is an unbiased estimator for β
- If we also assume that $\text{Var}(\varepsilon) = \sigma^2 \mathbf{I}$ then OLS is BLUE
 - **Best Linear Unbiased Estimator**, i.e. out of all linear unbiased estimators $\hat{\beta}_{\text{OLS}}$ has minimum variance (Gauss-Markov)

OLS + Model

Assume $y = X\beta + \varepsilon$ with $\mathbb{E}(\varepsilon) = 0$.

- Then $\hat{\beta}_{\text{OLS}}$ is an unbiased estimator for β
- If we also assume that $\text{Var}(\varepsilon) = \sigma^2 \mathbf{I}$ then OLS is BLUE
 - **Best Linear Unbiased Estimator**, i.e. out of all linear unbiased estimators $\hat{\beta}_{\text{OLS}}$ has minimum variance (Gauss-Markov)
- Say we have a new point x_0 and prediction $\hat{y}_0 = x_0^\top \hat{\beta}$.

OLS + Model

Assume $y = X\beta + \varepsilon$ with $\mathbb{E}(\varepsilon) = 0$.

- Then $\hat{\beta}_{\text{OLS}}$ is an unbiased estimator for β
- If we also assume that $\text{Var}(\varepsilon) = \sigma^2 \mathbf{I}$ then OLS is BLUE
 - **Best Linear Unbiased Estimator**, i.e. out of all linear unbiased estimators $\hat{\beta}_{\text{OLS}}$ has minimum variance (Gauss-Markov)
- Say we have a new point x_0 and prediction $\hat{y}_0 = x_0^\top \hat{\beta}$.
 - $\text{Bias}(\hat{y}_0) = 0$

OLS + Model

Assume $y = X\beta + \varepsilon$ with $\mathbb{E}(\varepsilon) = 0$.

- Then $\hat{\beta}_{\text{OLS}}$ is an unbiased estimator for β
- If we also assume that $\text{Var}(\varepsilon) = \sigma^2 \mathbf{I}$ then OLS is BLUE
 - **Best Linear Unbiased Estimator**, i.e. out of all linear unbiased estimators $\hat{\beta}_{\text{OLS}}$ has minimum variance (Gauss-Markov)
- Say we have a new point x_0 and prediction $\hat{y}_0 = x_0^\top \hat{\beta}$.
 - $\text{Bias}(\hat{y}_0) = 0$
 - For large n , and assuming $\mathbb{E}(X) = 0$:

$$\mathbf{EPE} = \mathbb{E}_{x_0} \mathbb{E}(y_0 - \hat{y}_0)^2 \approx \sigma^2 + \sigma^2(p/n)$$

See *The Elements of Statistical Learning*, Section 2.5 for details

OLS

Advantages

- Simple
- Closed-form solution
- Interpretable (?)
- Under some modeling assumptions, OLS has some desirable properties

Disadvantages

- No bias (assuming correctly specified), all variance... can lead to overfitting
- When $p > n$:
 - $X^T X$ singular


A possible solution

$$\text{MSE} = \text{Var}(\hat{f}(x_0)) + \text{Bias}^2(\hat{f}(x_0))$$

- We could try to sacrifice some of the bias to reduce the variance
- One way to reduce the variance of your predictions is to restrict the parameter space in the optimization $\underset{\beta}{\operatorname{argmin}} \|\mathbf{y} - \mathbf{X}\beta\|_2^2$
- In the linear setting, this motivates regularized linear regression methods such as ridge, Lasso, and elastic net

The Bias-Variance Tradeoff

$$\text{MSE} = \text{Var}(\hat{f}(x_0)) + \text{Bias}^2(\hat{f}(x_0))$$



Ridge regression

$$\hat{\boldsymbol{\beta}}^r = \underset{\boldsymbol{\beta}}{\operatorname{argmin}} \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2}_{\text{Loss function}} \quad \text{subject to} \quad \underbrace{\|\boldsymbol{\beta}\|_2^2 \leq \tau}_{\text{constraint}}$$

or equivalently

$$\hat{\boldsymbol{\beta}}^r = \underset{\boldsymbol{\beta}}{\operatorname{argmin}} \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2}_{\text{Loss function}} + \underbrace{\lambda \|\boldsymbol{\beta}\|_2^2}_{\text{penalty}}$$

Ridge regression

$$\hat{\beta}^r = \underset{\beta}{\operatorname{argmin}} \underbrace{\|\mathbf{y} - \mathbf{X}\beta\|_2^2}_{\text{Loss function}} \quad \text{subject to} \quad \underbrace{\|\beta\|_2^2 \leq \tau}_{\text{constraint}}$$

or equivalently

$$\hat{\beta}^r = \underset{\beta}{\operatorname{argmin}} \underbrace{\|\mathbf{y} - \mathbf{X}\beta\|_2^2}_{\text{Loss function}} + \underbrace{\lambda \|\beta\|_2^2}_{\text{penalty}}$$

$\lambda \geq 0$ is a tuning or penalty parameter and regulates how much *shrinkage* we introduce:

- $\lambda \rightarrow 0 \implies \hat{\beta}^r \rightarrow \text{OLS}$

Ridge regression

$$\hat{\boldsymbol{\beta}}^r = \underset{\boldsymbol{\beta}}{\operatorname{argmin}} \underbrace{||\mathbf{y} - \mathbf{X}\boldsymbol{\beta}||_2^2}_{\text{Loss function}} \quad \text{subject to} \quad \underbrace{||\boldsymbol{\beta}||_2^2 \leq \tau}_{\text{constraint}}$$

or equivalently

$$\hat{\boldsymbol{\beta}}^r = \underset{\boldsymbol{\beta}}{\operatorname{argmin}} \underbrace{||\mathbf{y} - \mathbf{X}\boldsymbol{\beta}||_2^2}_{\text{Loss function}} + \underbrace{\lambda ||\boldsymbol{\beta}||_2^2}_{\text{penalty}}$$

$\lambda \geq 0$ is a tuning or penalty parameter and regulates how much *shrinkage* we introduce:

- $\lambda \rightarrow 0 \implies \hat{\boldsymbol{\beta}}^r \rightarrow \text{OLS}$
- $\lambda \rightarrow \infty \implies \hat{\boldsymbol{\beta}}^r \rightarrow 0$

Ridge regression

$$\hat{\beta}^r = \underset{\beta}{\operatorname{argmin}} \underbrace{\|\mathbf{y} - \mathbf{X}\beta\|_2^2}_{\text{Loss function}} \quad \text{subject to} \quad \underbrace{\|\beta\|_2^2 \leq \tau}_{\text{constraint}}$$

or equivalently

$$\hat{\beta}^r = \underset{\beta}{\operatorname{argmin}} \underbrace{\|\mathbf{y} - \mathbf{X}\beta\|_2^2}_{\text{Loss function}} + \underbrace{\lambda \|\beta\|_2^2}_{\text{penalty}}$$

$\lambda \geq 0$ is a tuning or penalty parameter and regulates how much *shrinkage* we introduce:

- $\lambda \rightarrow 0 \implies \hat{\beta}^r \rightarrow \text{OLS}$
- $\lambda \rightarrow \infty \implies \hat{\beta}^r \rightarrow 0$
- Usually try to find an intermediate value that provides some shrinkage
- Can choose λ via CV

Why ridge?

$$\hat{\boldsymbol{\beta}}^r = \operatorname{argmin}_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_2^2$$

1. The solution exists and is unique even when $p > n$ (unlike OLS)

$$\hat{\boldsymbol{\beta}}^r = (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$$

- Exercise: show this

Why ridge?

$$\hat{\beta}^r = \underset{\beta}{\operatorname{argmin}} \quad ||\mathbf{y} - \mathbf{X}\beta||_2^2 + \lambda ||\beta||_2^2$$

1. The solution exists and is unique even when $p > n$ (unlike OLS)

$$\hat{\beta}^r = (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$$

- Exercise: show this

2. Assume the setup from before: $\mathbf{y} = \mathbf{X}\beta + \varepsilon$, $\mathbb{E}(\varepsilon) = 0$, $\operatorname{Var}(\varepsilon) = \sigma^2 \mathbf{I}$

There is **always** a value of λ where the ridge MSE is less than the OLS MSE:

$$\operatorname{MSE}(\mathbf{X}\hat{\beta}^r(\lambda)) < \operatorname{MSE}(\mathbf{X}\hat{\beta}^{OLS})$$

Farebrother, R. W. (1976) <https://www.jstor.org/stable/2984971>

Theobald, C. M. (1974) <https://www.jstor.org/stable/2984775>

Why ridge?

$$\hat{\beta}^r = \underset{\beta}{\operatorname{argmin}} \quad ||\mathbf{y} - \mathbf{X}\beta||_2^2 + \lambda ||\beta||_2^2$$

1. The solution exists and is unique even when $p > n$ (unlike OLS)

$$\hat{\beta}^r = (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$$

- Exercise: show this

2. Assume the setup from before: $\mathbf{y} = \mathbf{X}\beta + \varepsilon$, $\mathbb{E}(\varepsilon) = 0$, $\operatorname{Var}(\varepsilon) = \sigma^2 \mathbf{I}$

There is **always** a value of λ where the ridge MSE is less than the OLS MSE:

$$\operatorname{MSE}(\mathbf{X}\hat{\beta}^r(\lambda)) < \operatorname{MSE}(\mathbf{X}\hat{\beta}^{OLS})$$

3. Handles correlated features well -- features that are highly correlated tend to get shrunk together, i.e. they are given equal contribution to the linear model

Ridge in practice

- Don't want to penalize the intercept
 - Before applying regularization, center columns of X and y
- Most of the time, should also scale columns of X so that we don't penalize some coefficients more than others simply because of different scales
- These practical guidelines apply to all regularization methods

Regularization Part 2: Lasso

Next time...