


Supplemental Figure 14



Supplemental Figure 14. Orthotopic PDX mouse model of human MED-411 FH mCherry/Luc cell line to evaluate anti-tumor activity of CAR.GD2 T-cells. **(A)** Illustration of the experimental design in which Med-411 FH mCherry/Luc cells were stereotactically implanted into the cerebellum of eight NSG mice (Figure created using Biorender - <https://biorender.com>). At the time of tumor engraftment (day 14), the effector cells were administered through *i.v.* injection. **(B)** Time course of *in vivo* bioluminescence imaging of each treated NSG mice from day 14 (day of effector cells infusion) to day 58. **(C)** The graph shows the average bioluminescence of four mice treated with NT T-cells (black line) and four mice treated with CAR-GD2 T-cells (dotted black line). Data are shown as average $\pm$ SD. Two-way Anova was used as multi comparison test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. **(D)** The graph shows bioluminescence analysis of each tumor bearing mice treated with NT-T (black lines with circle) or CAR.GD2-T cells (dotted black lines with triangle). **(E)** Sixty-day Overall Survival (OS) of tumor bearing mice treated with NT T-cells (black line) and CAR.GD2 T-cells (dotted black line) cells. Two-way Anova was used as multi comparison test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. **(F-G)** Average of human circulating T-cells evaluated as % of CD45+CD3+ **(F)** and CD3+CAR+ cells **(G)** in NSG mice treated with NT T-cells (red line) or CAR.GD2 T-cells (dotted black line) at day 14, 28 and 42 after effector T-cell infusion. Data are shown as average $\pm$ SD. Two-way Anova was used as multi comparison test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.