

Supplemental Figure 2. GD2 evaluation in MB cell lines. (A) Cytofluorimetric analysis of GD2 expression on MB cell lines at 100% of confluency: D283 Med, DAOY, MED-411 FHTC and MED-411 FH mCherry/Luciferase. Fluorescence Minus One (FMO) for BV421 is shown in blue while isotype control is reported in green. (B) The percentage of GD2+ cells on MB cell lines (D283 Med, DAOY, MED 411 FHTC and MED-411 FH mCherry/Luciferase) were assessed in triplicate by flow-cytometry analyses. Data are expressed as column bar graph showing mean $\pm$ SD. (C) MFI for GD2 on MB cell lines (D283 Med, DAOY, MED 411 FHTC and MED-411 FH mCherry/Luciferase). MFI of MB cell lines was normalized on IgG2a isotype MFI. Data are expressed as column bar graph showing mean $\pm$ SD. * $p < 0.05$, ** $p < 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$. (D-E). Cell confluency influences GD2 expression in DAOY-GFP cell line in 2D and 3D in vitro models. (D) The DAOY-GFP cell line was seeded in 24-well plate, at densities of 25000 cells/well (confluency 30%), 50000 cells/well (confluency 50%), 100000 cells/well (confluency 75%) and 200000 (confluency 100%) cells/well respectively. After 72h tumor cells were collected and analysed by flow-cytometry for GD2 expression. (E) The DAOY-GFP cell line was seeded in 96-ULA-well plate at densities of 2000 cells/well, 4000 cells/well, 8000 cells/well and 16000 cells/well until spheroid was formed (about 3 days). The spheroid generation was monitored with the Incucyte Live cell analysis system (right images) during time and after 72h tumor cells collected and analysed by flow-cytometry for GD2 expression. (F) The MED-411 FHTC cell line was seeded in 24-well plate, at densities of $1.25 \times 10^4/\text{cm}^2$, $2.5 \times 10^4/\text{cm}^2$, $5.0 \times 10^4/\text{cm}^2$ and $1.0 \times 10^5/\text{cm}^2$ cells/well respectively and after 72h tumor cells collected and analysed by flow-cytometry for GD2 expression.