

Low-/negative-pressure hydrocephalus — literature review

Consolidated report over the 231-record corpus in `index.tsv`. Part I and Part III are computed from the index; Part II is the written synthesis from `notes/`. Pipeline: `harvest.py` → `curate.py` → `fetch_fulltext.py` → `gen_report.py` → `build_pdf.py`.

Part I — Corpus statistics (computed from `index.tsv`)

231 curated records, 1969–2026; 32 open-access full texts mirrored under `fulltext/` (`oa_xml`), 18 with a PMCID but no retrievable XML, 181 metadata-only.

Records by axis and evidence class

evidence class	Core entity	Mechanism	Treatment techniques	Aetiological settings	Adjacent differentials	Historical usage	all
systematic review	3	2	2		1		8
consensus / guideline	1						1
narrative review	6	7	1	1	4		19
case series	5		2	1	4	3	15
case series / cohort	19	16	21	9	14	10	89
case report	34	2	1	3	3	1	44
clinical / mechanistic study	3	3	1		2	12	21
imaging / physiology study		4					4
experimental (animal / bench)	3	8	2		2	1	16
computational model	1	8				1	10
comment / letter	4						4
all	79	50	30	14	30	28	231

Core-entity axis

79 records; 70 name the entity (low-/negative-/very-low-pressure hydrocephalus, SILPAH, ALPH) in title or abstract, 9 describe the phenomenon without the label.

Population: unspecified 46, adult 15, pediatric 13, mixed 5.

Precipitating setting tags (multi-label, core axis):

setting	n
shunt_dependent_prior	37
aqueductal_obstructive	17
infection_inflammatory	13
tumor_posterior_fossa	12
sah_aneurysm	11

setting	n
skull_base_csf_leak	11
lumbar_puncture_or_drain	9
trauma	6
craniectomy_skin_flap	6
intraventricular_hemorrhage	3

Treatment tags (multi-label, core axis):

treatment	n
shunt_or_valve	58
titrated_evd	35
subzero_evd	27
etv	12
csf_leak_repair	12
active_pump_shunt	10
elastography_imaging	8
neck_wrapping	7
icp_monitoring	5
positioning	3
pharmacologic	1

Publication years (core axis)

period	n
1970-1974	1
1985-1989	2
1990-1994	4
1995-1999	3
2000-2004	5
2005-2009	5
2010-2014	9
2015-2019	15
2020-2024	19
2025-2029	16

Part II — Synthesis

Sections below are the topical notes in `notes/` (00-07), concatenated. PMIDs cited resolve to rows of `index.tsv` except where a record is explicitly described as excluded from the corpus.

Low-/negative-pressure hydrocephalus — overview and executive summary

What the entity is

Low-pressure hydrocephalus (LPH), negative-pressure hydrocephalus (NegPH/NePH), very-low-pressure hydrocephalus (VLPH), acute low-pressure hydrocephalus (ALPH) and the "syndrome of inappropriately low-pressure acute hydrocephalus" (SILPAH) are names for one clinical paradox: **progressive ventriculomegaly with the symptoms of intracranial hypertension (depressed consciousness, headache, vomiting, cranial neuropathies, gait failure, bradycardia) while measured intracranial pressure is normal, low (conventionally ≤ 5 cm H₂O / ≤ 70 mm H₂O) or frankly sub-atmospheric** (down to -10 to -21 mm Hg in reported cases). Shunts, when present, are patent; conventional shunt revision or a standard EVD at 10–15 cm H₂O fails because the ventricular pressure never reaches the valve opening pressure or the drain outlet. Clinical status tracks ventricular size, not ICP (Pang & Altschuler 1994, PMID 7808607; Daniel 2002, PMID 12027920; Godoy Hurtado 2023, PMID 37373809).

The modern entity was defined by Pang & Altschuler (Neurosurgery 1994), who treated 12 shunted patients with ventriculomegaly and ICP 2.2–6.6 mm Hg by prolonged "sub-zero" external drainage and framed the mechanism as a viscoelastic alteration of brain parenchyma. Precursor descriptions exist from the 1970s–80s in shunted patients (McCullough & Fox 1974, PMID 4813717; McCullough 1986, PMID 3491341; Foltz & Blanks 1988, PMID 3343612). Before 1994, "low-pressure hydrocephalus" was used in Europe as a synonym for Adams-Hakim normal-pressure hydrocephalus; that literature is retained here as a separate axis so it is not conflated with the modern syndrome.

Corpus at a glance

- 231 curated records (1969–2026) from a 1,190-record Europe PMC harvest; 32 open-access full texts mirrored.
- **Core entity axis: 79 records**, 70 of which name the entity in the title/abstract; the remainder describe the phenomenon (symptomatic ventriculomegaly with patent shunt and low ICP) without the label.
- Evidence is almost entirely observational: 34 case reports, 24 case series / cohorts, 3 systematic reviews (Keough 2021 pooling 195 patients, PMID 32736355; Bakmeedeniya 2026 pooling 215 patients from 30 studies, PMID 41525875; Chang 2022 on Holmes tremor, PMID 35148507), 1 consensus statement (Chinese Neuroscience Society 2025, PMID 40734409) and no prospective comparative trial.
- Supporting axes: 50 brain-biomechanics / compliance / elastography papers, 30 drainage-and-valve technique papers (largely EVD-weaning and programmable-valve literature that frames titrated drainage), 14 etiological-setting papers, 30 adjacent differentials (slit-ventricle syndrome, overdrainage, intracranial hypotension, venous overdrainage), 28 historical LPH-as-NPH papers.

Ten take-home points

1. **Rare, under-recognised, and iatrogenically amplified.** Roughly 200–250 published cases exist (Godoy Hurtado 2023; Keough 2021; Bakmeedeniya 2026). Single-centre incidence after aneurysmal SAH is $\sim 3.7\%$ (Czorlich 2023, PMID 37004138); in neurosurgical patients with severe disorders of consciousness and ventriculomegaly, 60 % had low- or negative-pressure physiology (Zheng 2023, PMID 36994490). Almost every patient has had at least one prior intracranial procedure (mean 3.0 in Hunn 2014, PMID 24275338; ~ 3.5 operations per patient in Wu 2019, PMID 30611954).

2. **Precipitants cluster into three groups:** (a) a CSF sink outside the ventricles — lumbar puncture or lumbar drain, skull-base/cranial-wound CSF leak, pseudomeningocele, over-draining cyst or spinal shunt; (b) parenchymal "softening" — SAH/IVH, ventriculitis/meningitis, radiotherapy, trauma, diffuse injury, long-standing shunting; (c) loss of cranial containment — decompressive craniectomy / sinking-flap physiology.
3. **The mechanistic debate has four live models** — viscoelastic/hysteresis (Pang; Lesniak 2002, PMID 12617236), poroelastic "boggy brain"/Darcy flux (Akins 2011, PMID 21523524), transmantle gradient from ventricle-subarachnoid disconnection plus loss of brain turgor (Rekate 2008, PMID 18590387; Filippidis 2011, PMID 21800965; Diaz-Romero Paz 2019, PMID 30710718), and cerebral venous overdrainage/windkessel impedance (Barami 2016, PMID 27581321; Egnor 2023, PMID 37382303; Bateman 2026, PMID 42355921). They are not mutually exclusive and the leading reviews treat them as interacting (Bakmeedeniya 2026; Li 2024, PMID 38688300). MR elastography has shown reversibly low brain stiffness during an episode (Olivero 2016, PMID 27198914; Olivero 2021, PMID 33030603).
4. **Diagnosis rests on the paradox plus a therapeutic test:** symptomatic ventriculomegaly, ICP $\leq$ 5 cm H₂O on direct ventricular measurement, patent shunt / failed standard drainage, and improvement with drainage at zero or sub-atmospheric pressure (Pang-Altschuler criteria; Duan & Hu 2024, PMID 38579413; Datta & Tu 2025, PMID 41191101). Imaging shows effaced convexity sulci and Sylvian cisterns with disproportionate ventriculomegaly, often transependymal oedema despite low ICP, and — in the 2025 Chinese consensus — a fourth-ventricular "ball sign".
5. **Temporising treatment is titrated (usually sub-zero) EVD,** set to a target hourly output (5–15 mL/h, Hunn 2014) rather than a pressure, started at 0 to –5 cm H₂O below the EAM (Owler 2001, PMID 11599454), –3 to –10 cm (Akins 2011, PMID 21523524) or 5–10 cm below the midbrain (Foster 2016), lowered further only as needed (–15 in Godoy Hurtado 2023; –30 in one refractory case, Salam 2025), and then raised 1–3 cm H₂O every 3–5 days as ventricles shrink and ICP becomes positive (Akins 2011; Foster 2016, PMID 27069743). Drainage durations of weeks to months are typical (mean 41.9 days, range 2–365, in the pooled paediatric series of Chen 2025, PMID 40538927).
6. **Adjuncts that raise ICP or close the sink shorten or avoid sub-zero EVD:** neck wrapping / soft cervical collar (Filippidis 2011; Pandey 2017, PMID 28017745; Bruk 2026, PMID 42732439), Trendelenburg/recumbency (Foltz 1988; Dias 1999, PMID 10352418), epidural blood patch for post-LP or lumbar-drain cases (Dias 1999; Smalley 2017, PMID 28362957; Vilela 2020, PMID 32712403; Suzuki 2021, PMID 35854689; Hoffman 2026, PMID 42224725), repair of skull-base or wound CSF leaks (Filippidis 2011; Stevens 2024, PMID 39220674), removal of the offending spinal/cyst drain (Ozeki 2022, PMID 34526456; Diaz-Romero Paz 2019), and — a single striking case — theophylline (Salam 2025, PMID 40418891).
7. **ETV is the main strategy for reducing shunt dependence:** a "significant decrease" in shunt-dependent patients with ETV in SILPAH (10 vs 10 patients; Hamilton & Price 2012, PMID 22116443); 50 % ETV-only success in the Casado Pellejero algorithm (2022, PMID 34998487); ETV plus valveless catheter gave 81 % excellent outcomes in 16 adults, 19 % later needing a valved shunt (Villalonga 2025, PMID 41165379); but Foster 2016 (PMID 27069743) found no significant shortening of EVD time and in Chen 2025's pooled children 91 % ultimately needed shunt repositioning/revision.
8. **Definitive drainage needs a very-low-resistance conduit:** valveless VP shunts, programmable valves at the lowest setting without anti-siphon devices, "therapeutic siphoning" with non-siphon-control or adjustable gravitational valves (Bergsneider 1999, PMID 10520938; Trinh 2013, PMID 22770726; Funnell 2020, PMID 32989519), shunting into negative-pressure cavities (pleura, atrium), valves placed below the clavicle to exploit the hydrostatic column (Carrera 2025, PMID 39951714), wide-bore custom peritoneal catheters (Galbarriatu 2013, PMID 23904087), active-pumping shunts (Kalani 2013, PMID 23380444), improvised ultra-low valveless systems (Badran 2018, PMID 30287249) and intermittent reservoir pumping for chronic failure (Cheng 2017, PMID 28264586; Kuppler 2025, PMID 40030163).
9. **Outcome is bimodal and set by two things:** speed of recognition and the underlying disease. Pooled mortality is ~11 % (Keough 2021); 83 % of children but only 49 % of adults returned to their pre-episode baseline; 77.5 % of children recovered in Chen 2025; but Wu 2019's 39 secondary adult cases (18 with CNS infection) had a mean GOS-E of 2.7 and 8 deaths. Post-LP

paediatric cases are the most benign subgroup (Smalley 2017). Predictors of futility are unsealed CSF leak, irremediable ventriculitis and permanently altered compliance (Houlden 2018, PMID 30009119; Bakmeedeniya 2026).

10. **The evidence base is fragile:** heterogeneous definitions (ICP thresholds of 0, 5 and 7 cm H₂O; NegPH vs LPH; acute vs chronic), publication bias toward rescued cases, no registries, and mechanistic claims that rest on single MRE or phase-contrast measurements. Every major review from 2019 onward calls for standardised definitions and a multicentre registry.

How the notes are organised

- `01_history_nomenclature.md` — from 1969 "low-pressure hydrocephalus" (=NPH) through Pang & Altschuler to SILPAH, ALPH, cNegPH and pressure-based classification.
- `02_pathophysiology.md` — the competing/interacting models, experimental and imaging evidence, and what each model predicts about treatment.
- `03_etiologies_and_settings.md` — precipitants, frequencies in the pooled series, paediatric vs adult spectrum.
- `04_diagnosis.md` — phenotype, ICP measurement, imaging, differential diagnosis (overdrainage, slit ventricles, intracranial hypotension, NPH).
- `05_management.md` — acute algorithm (EVD titration, adjuncts, ETV), definitive CSF diversion hardware, chronic/refractory strategies, pharmacology.
- `06_outcomes_and_evidence_gaps.md` — outcome data, prognostic factors, evidence quality, and research questions.
- `07_key_papers.md` — per-paper extractions for the ~35 papers that carry the field.

History and nomenclature

1. "Low-pressure hydrocephalus" as a synonym for NPH (1965-c.1990)

After Adams, Hakim and colleagues described normal-pressure hydrocephalus in 1965, many European authors preferred the term low-pressure hydrocephalus for the same syndrome — an elderly patient with gait apraxia, dementia and incontinence, communicating ventriculomegaly and a lumbar opening pressure in the normal range. The 28 papers on the `historical_lph_as_nph` axis are of this kind: cerebral blood flow before and after shunting (Greitz 1969, PMID 5359210; Meixensberger 1989, PMID 2608784), isotope cisternography (Jensen 1970, PMID 5457870; Kulczycki 1978, PMID 683432), overnight pressure recordings that showed B-waves and plateau waves in "so-called low-pressure hydrocephalus" (Symon 1972, PMID 4117815; Sjaastad 1976, PMID 1278191), infusion tests (Bozzao 1974, PMID 4408072), psychometric profiles (Rivero Hernández 1975, PMID 1211270) and shunt series with "disappointing results" (Bannister 1972, PMID 4540544; Singounas 1976, PMID 1250499). Symon's demonstration that these patients have intermittently raised pressure is a reason the term fell out of use for NPH.

These papers are relevant to the modern entity in two ways only: they show that ventricular enlargement can progress without sustained hypertension, and they are the source of persistent terminological confusion (e.g. a 2013 case report still describing "acute normal pressure hydrocephalus" for what is clearly post-LP low-pressure disease; Preuß 2013, PMID 23715811). The Kuba 2002 gait study (PMID 11749014) and several 1970s "LPH" reports were retained in the core axis because their abstracts describe low measured pressure with progressive ventriculomegaly, but they should be read as NPH-era work.

2. Precursors in shunted patients (1974-1993)

- **McCullough & Fox 1974** (PMID 4813717): negative ICP in shunted adults and its relation to subdural haematoma — the first systematic recognition that shunted brains can run at sub-atmospheric pressure, here framed as an overdrainage complication rather than a hydrocephalic state.
- **McCullough 1986** (PMID 3491341): symptomatic progressive ventriculomegaly in hydrocephalics with patent shunts and antisiphon devices — the essential paradox (large ventricles, working shunt, low pressure) stated a decade before it was named. The anti-siphon device prevented the shunt from ever reaching the sub-zero pressures the brain needed.
- **Foltz & Blanks 1988** (PMID 3343612): symptomatic low intracranial pressure in shunted hydrocephalus — upright, activity-related symptoms with markedly low upright ICP, relieved by recumbency and Trendelenburg; Foltz later designed a "zero-ICP" gravity-controlled shunt (ZIPS; Foltz 1994, PMID 8194062).
- **Drake 1993** (PMID 8421544): functional obstruction of an anti-siphon device by tissue-capsule pressure — an early mechanical explanation for "patent shunt, large ventricles".
- **Pomeranz 1993** (PMID 8333302): feline intracranial hypotension reduces cerebral blood flow — an experimental hint that low ICP itself can be harmful.

3. Pang & Altschuler 1994 and the definitional decade

Pang & Altschuler, Neurosurgery 1994 (PMID 7808607) coined low-pressure hydrocephalic state: 12 shunted patients with symptomatic ventriculomegaly, ICP 2.2-6.6 mm Hg, patent shunts, requiring prolonged EVD at sub-zero levels (most for weeks), with symptoms tracking ventricular size rather than ICP. They proposed altered brain viscoelasticity — a "softened" parenchyma that no longer returns to shape under a normal transmante gradient — and established sub-zero drainage as the reference treatment. Their four criteria (symptoms of hydrocephalus, ventriculomegaly, ICP $\leq$ 5 cm H₂O, improvement with sub-zero drainage) are still used for inclusion (Foster 2016; Villalonga 2025).

Subsequent naming:

Term	Source	Nuance
Negative-pressure hydrocephalus (NegPH, NePH)	Vassilyadi 1995 (PMID 7666227), after a spinal arachnoid cyst-pleural shunt	ICP actually sub-atmospheric; implies a transmante gradient created by an external sink
Low-pressure shunt "malfunction" after LP	Dias 1999 (PMID 10352418)	Paediatric, post-lumbar-puncture; resolves with recumbency $\pm$ blood patch
Very-low-pressure hydrocephalus (VLPH)	Clarke 2006 (PMID 16961147); Houlden 2018 (PMID 30009119)	Refractory subgroup with poor outcome after suprasellar/skull-base surgery
Hysteresis / non-linear dynamics	Lesniak 2002 (PMID 12617236)	Mechanistic label; 10 adults, mean ICP 4.8 mm Hg
Cortical subarachnoid space / brain turgor model	Rekate 2008 (PMID 18590387)	Reframes as ventricle-SAS disconnection; "compliance" replaced by "turgor"
Syndrome of inappropriately low-pressure acute hydrocephalus (SILPAH)	Hamilton & Price 2012 (PMID 22116443)	Emphasises acute deterioration and ETV as treatment
Acute low-pressure hydrocephalus (ALPH)	Keough 2021 (PMID 32736355); Czorlich 2023 (PMID 37004138)	Systematic-review label; ICP $\leq$ 5 cm H ₂ O typical
Chronic negative-pressure hydrocephalus (cNegPH)	Kuppler 2025 (PMID 40030163)	Persistent ventriculomegaly after failure of prolonged sub-atmospheric drainage
Ultra-low-pressure hydrocephalic state in NPH	Funnell 2020 (PMID 32989519)	Elderly iNPH patients who relapse after shunting and need siphoning
Two-compartment / acute "normal pressure" hydrocephalus	Preuß 2013 (PMID 23715811)	Idiosyncratic label for post-LP LPH in a shunted child

Term	Source	Nuance
Low-/negative-pressure hydrocephalus (LPH/NePH, L&NPH)	Wu 2019; Li 2024; Chen 2025	Chinese-language literature; LPH 0-70 mm H ₂ O, NegPH ≤ 0

4. Classification proposals

- **Wu 2019** (PMID 30611954) proposed classifying hydrocephalus by ventricular pressure — high, normal, low (< 70 mm H₂O) and negative (< 0) — arguing that existing schemes (Oi's multi-categorical classification, PMID 21928018; Raimondi's unifying theory, PMID 8194058; Rekate's 2008 definition based on points of obstruction) give no management guidance for the low-pressure group.
- **Chinese Neuroscience Society 2025 consensus** (PMID 40734409) defines LPH as persistent intraventricular pressure < 70 mm H₂O with progressive ventriculomegaly and neurological worsening, adds imaging criteria (effaced Sylvian cisterns and convexity SAS, fourth-ventricular "ball sign"), and names fourth-ventricular outlet obstruction and altered compliance as candidate mechanisms. It is the first formal guideline document on the entity.
- **Lin 2025** (PMID 40826122) places low-pressure hydrocephalus as the second most common subtype of "refractory hydrocephalus" after infectious hydrocephalus and stresses that management hinges on "identifying and addressing the sites receiving CSF".
- **Rekate 2019** "Low or Negative Pressure Hydrocephalus Demystified" (PMID 31103762) and **Rekate 2021** commentary (PMID 34272587) restate the turgor/disconnection view and argue the syndrome is a predictable consequence of CSF pathway anatomy rather than a mystery.

5. Terminological problems that remain

1. **Threshold:** ≤ 5 cm H₂O (Pang, Keough, Datta), < 70 mm H₂O (Wu, Chinese consensus), ≤ 0 (NegPH) and "below valve opening pressure" (Filippidis) are all in use; a case at 6–8 cm H₂O with a patent shunt set at 10 is LPH to some authors and shunt under-drainage to others.
2. **Acute vs chronic:** ALPH/SILPAH (deterioration over hours-days after a precipitant) and cNegPH / ultra-low-pressure NPH (months of refractory ventriculomegaly) probably share physiology but differ in urgency and in the role of ETV.
3. **NPH overlap:** Funnell 2020 and Trinh 2013 describe shunted iNPH patients who need siphoning; Doshi 2022 (PMID 36600762) describes an NPH patient evolving into VLPH. Whether "low-pressure NPH" is the same disease as post-SAH ALPH is unsettled.
4. **Sinking-flap / paradoxical herniation overlap:** after craniectomy the same sub-atmospheric physiology produces sinking flap syndrome, paradoxical herniation and LPH, sometimes sequentially in one patient (Bateman 2026, PMID 42355921; Salam 2025).

Pathophysiology — competing and interacting models

The central puzzle: ventricles enlarge and compress brain even though the pressure inside them is not high. Every model must therefore explain either (i) a gradient across the brain mantle that exists despite a low absolute ventricular pressure, or (ii) a change in the material of the brain such that a normal or low pressure is enough to keep it distended, or both. The systematic review of Bakmeedeniya 2026 (PMID 41525875) identified four commonly invoked models — viscoelastic change, transmantle gradients, poroelastic/glymphatic dysfunction, impaired CSF pulsatility — and treats them as interacting. The clinician-oriented review of Li 2024 (PMID 38688300) covers the same four under the labels transcortical pressure difference, excessive cerebral venous drainage, brain viscoelastic change and the "porous elastic sponge".

1. Viscoelastic softening and hysteresis (Pang & Altschuler; Lesniak)

- **Claim.** Chronic or intermittent distension (e.g. repeated shunt failures with high pressure) permanently alters the stress-strain behaviour of periventricular white matter. Hooke's-law elasticity is replaced by viscous creep: the brain behaves like a material that has been stretched past its yield point and does not recoil when pressure normalises. A lower-than-normal ventricular pressure is therefore required to restore shape (Pang & Altschuler 1994, PMID 7808607).
- **Hysteresis.** Lesniak 2002 (PMID 12617236) modelled 10 adults (mean ICP 4.8 mm Hg) as a pressure-volume loop with hysteresis: the volume reached at a given pressure depends on the path taken. Only negative-pressure drainage moved the system back onto the low-volume limb; ventricles then stayed small at normal pressure. Godoy Hurtado 2023 (PMID 37373809) uses the lung analogy (inflation and deflation limbs differ).
- **Evidence.** MR elastography in a 19-year-old with shunted aqueductal stenosis showed very low shear stiffness during an LPH episode (Olivero 2016, PMID 27198914) that recovered to near-normal two years later after a high-pressure shunt failure (Olivero 2021, PMID 33030603) — the only direct human measurement of reversibly reduced stiffness in LPH. Houlden 2018 (PMID 30009119) argued that recovery of GCS despite fluctuating frontal-horn ratio indicates "retained viscoelasticity" and predicts salvageability. Supporting biomechanics: brain elasticity changes with ventriculomegaly (Sklar 1980, PMID 6893601), reassessment of brain elasticity in hydrocephalus models (Taylor & Miller 2004, PMID 15212932), MRE stiffness falls in early-onset hydrocephalus white matter (Wagshul 2021, PMID 33631603) and rises in NPH with partial normalisation after shunting (Streitberger 2011, PMID 20931563; Freimann 2012, PMID 21538046; Fattahi 2016, PMID 26542235).
- **Weakness.** Does not explain why the syndrome is so often triggered by an acute external CSF sink (LP, leak) in patients whose brains were not chronically distended, or why neck wrapping works within hours.

2. Poroelastic "boggy brain" — Darcy flux of interstitial fluid (Akins)

- **Claim.** The brain is a fluid-saturated porous solid. When interstitial fluid accumulates (transependymal oedema, inflammation, radiation injury), the parenchyma loses turgor and its ability to resist deformation; ventricles enlarge at low pressure. Sub-zero drainage produces a transient "CSF diuresis" (drain output exceeds baseline for days) as interstitial water is pulled back into the ventricles along a Darcy flux, after which the brain re-stiffens (Akins 2011, PMID 21523524: 9 patients, transependymal oedema in 5 despite low ICP, EVD output fell as LPH developed and rose above baseline during sub-zero drainage).
- **Model support.** Poroelastic finite-element models of hydrocephalus (Peña 1999, PMID 10414573; Dutta-Roy 2008, PMID 18534602; Chou 2016, PMID 26671218; Kim 2016 on periventricular lucency, PMID 26274984) show that small sustained gradients can produce large ventricular strain when permeability and elasticity are altered; Solamen 2021 (PMID 32331966) measured poroelastic properties of NPH brains during lumbar drainage. Duy 2024 (PMID 38798141) reviews hydrocephalus as "biomechanical instability of the brain-CSF interface".
- **Clinical corollary.** Akins' group used the "sub-zero method" successfully in all 9 patients and internalised with low-pressure hardware; the model predicts that fluid shifts (hyponatraemia, low osmolarity) worsen LPH — consistent with Houlden 2018's observation that avoiding fluid shifts helped, and with Pandey 2017 (PMID 28017745) deliberately lowering plasma osmolarity to raise ICP.

3. Transmantle gradient from ventricle-subarachnoid disconnection and loss of turgor (Rekate/Filippidis)

- **Claim.** Ventricular size is set by the pressure difference between the ventricle and the cortical subarachnoid space (CSAS), not by absolute ICP. If the CSAS is (a) disconnected from the ventricles by obstruction at the aqueduct, fourth-ventricular outlets, basal cisterns or tentorial incisura, and (b) drained to a lower pressure than the ventricles by an external sink (lumbar puncture/drain, skull-base leak, cyst-pleural shunt), then the ventricle is at higher pressure than

the SAS even when both are sub-atmospheric; CSF cannot return to the convexities and the ventricles enlarge (Rekate 2008, PMID 18590387; Filippidis 2011, PMID 21800965; Vassilyadi 1995, PMID 7666227; Diaz-Romero Paz 2019, PMID 30710718; Suzuki 2021, PMID 35854689). Rekate replaces "compliance" with **brain turgor** — the capillary-bed-dependent firmness that normally keeps ventricles small — which is reduced by ischaemia, radiotherapy, injury and low venous pressure.

- **Predictions that have held.** (i) Neck wrapping / jugular compression raises venous and hence CSAS pressure and turgor and reverses ventriculomegaly within hours to days (Filippidis 2011; Pandey 2017; Bruk 2026, PMID 42732439); healthy-volunteer MRE confirms jugular compression stiffens brain in proportion to jugular venous flow (Hatt 2015, PMID 26045579). (ii) Closing the sink (blood patch, leak repair, removing a spinal drain) can abort the syndrome without sub-zero EVD (Dias 1999; Vilela 2020; Suzuki 2021; Ozeki 2022; Hoffman 2026). (iii) ETV, which reconnects ventricle and SAS, reduces shunt dependence (Hamilton & Price 2012). (iv) Lumbar-ventricular pressure discrepancies are measurable in obstructed systems (Chang 2026, PMID 42218510; Staykov 2011, PMID 21273925).
- **Counter-evidence and nuance.** Direct measurements have failed to find a transmante gradient in communicating or non-communicating hydrocephalus at steady state (Stephensen 2002, PMID 11904027; Shapiro 1987 cats, PMID 3598675; Eide 2010 on pulsatile gradient, PMID 20130957), while Conner 1984 (PMID 6737056) did find one in experimental NPH. Advocates reply that the relevant gradient is small, dynamic and only present while a sink is active, and that MRI-derived transmante pressures (Sincomb 2021, PMID 34385144) are on the order of Pascals — enough over time in a soft brain.

4. Cerebral venous overdrainage, windkessel and pulsatility models

- **Venous overdrainage (Barami 2016, PMID 27581321).** CSF diversion at low pressure lowers the pressure surrounding cortical veins and bridging veins, collapsing them or increasing venous outflow, reducing cerebral blood volume and turgor; the brain "deflates" toward the ventricles. Vilela 2020 (PMID 32712403) invoked this for post-LP LPH in an adult; Datta & Tu 2025 (PMID 41191101) found prior jugular/sigmoid sinus thrombosis in a third of their paediatric LPH patients and modelled venous thrombosis as a predisposing factor; Sainz 2019 (PMID 30474714) showed the mirror image — venous hypertension increases turgor and prevents ventriculomegaly in infant "external hydrocephalus".
- **Windkessel / CSF path impedance (Egnor 2023, PMID 37382303; Egnor 2026, PMID 41791099).** In a lumped electrical model the cranium is a band-stop filter that shunts arterial pulsatile energy through CSF to the veins. Egnor classifies hydrocephalus by CSF-path impedance: obstructive = high resistance; NPH = low inertance and high compliance; **low-pressure hydrocephalus = high resistance and high compliance**. Ventriculomegaly is an adaptive response that lowers impedance, which is why it can occur at any absolute pressure. Related pulse-wave-encephalopathy work (Bateman 2002, PMID 12221445; Bateman 2008, PMID 18379767) and pulsatility-driven glymphatic flow (Nozalea 2025, PMID 40258946) feed this view.
- **Autoregulatory reset and capillary CSF absorption (Bateman 2026, PMID 42355921).** In a craniectomy patient who passed through sinking-flap syndrome and LPH, phase-contrast MRI showed bilaterally reduced CBF that rose by 26 % after cranioplasty and reversed aqueductal CSF flow toward the ventricles during the LPH phase. Lumped-parameter modelling favoured arteriolar constriction from an autoregulation reset over venous compression, and the authors hypothesise increased CSF absorption via the capillary bed at low ICP.
- **Unified mathematical model (Wang & Lu 2020 preprint).** A U-shaped ICP-volume curve that claims to explain slit ventricles, NPH/LPH and paradoxical herniation in one framework; untested.

5. Compliance, elastance and pressure-volume physiology (supporting axis)

The 50 biomechanics papers give the vocabulary and measurements the LPH literature borrows:

- **Pressure-volume compensation** (Czosnyka 1996, PMID 8778261; Czosnyka 2001 age dependence, PMID 11235954; Kim 2009 compensatory reserve index, PMID 19240611; Gholampour 2022 new definition of compliance after shunting, PMID 35979170). LPH is a state of high compliance / low elastance: large volume change for little pressure change — the opposite of slit-ventricle syndrome.
- **Age and species dependence of viscoelasticity** (Shulyakov 2011, PMID 21316637; Shulyakov 2012 acute experimental hydrocephalus, PMID 22806081; Jugé 2016 rat obstructive hydrocephalus microstructure and stiffness, PMID 26848844; Pattison 2010 feline MRE, PMID 20655045; Dirrichs 2019 infant shear-wave elastography, PMID 31464808). Infant brains are uniquely compliant — Rekate 2020 (PMID 32488353) argues this is why infant hydrocephalus, sinking fontanelles and post-LP LPH behave differently.
- **MR elastography in NPH and dementia** (Perry 2017, PMID 28063896; ElSheikh 2017, PMID 28570101; Murphy 2020, PMID 32058331; Aunan-Diop 2022 scoping review, PMID 34687356; Coelho 2022 systematic review, PMID 35488708; Karki 2024 outcome prediction, PMID 38272572; Hersh 2022 operative applications, PMID 35933785). Stiffness increases periventricularly in NPH and falls after shunting — the opposite direction from the LPH case reports, suggesting LPH is not simply an extreme NPH.
- **Historical theory** — Raimondi's unifying definition (PMID 8194058), Orešković & Klarica's critique of the bulk-flow hypothesis (PMID 21641963; choroid plexus role, PMID 28456718), Oi's evolution theory of CSF dynamics (PMID 16685545), Li 2008 kaolin models of cortical SAS obstruction (PMID 18433747), Fukuhara 2001 shunt removal and compliance (PMID 11302655).

6. Synthesis: a two-hit, two-compartment picture

The most parsimonious reading of the corpus is that LPH requires **a soft or low-turgor brain** (hit 1: haemorrhage, infection, radiation, chronic distension, craniectomy, venous thrombosis, immaturity) **and a pressure differential that favours the ventricle** (hit 2: an external CSF sink or a disconnected SAS at lower pressure, or shunt hardware that cannot drain below its opening pressure). Treatments map onto the two hits:

Intervention	Acts on
Sub-zero EVD	Makes the ventricle the lowest-pressure compartment; unloads interstitial water (hit 2, then 1)
Neck wrapping / collar / Trendelenburg / hypercapnia / low osmolarity	Raises venous pressure, CBV and turgor (hit 1)
Blood patch, leak repair, removing spinal/cyst drain	Removes the sink (hit 2)
ETV	Reconnects ventricle and SAS so the gradient cannot form (hit 2)
Valveless / siphoning shunts, negative-pressure cavities, active pumps	Permanent low-resistance path so the ventricle never sits above the SAS (hit 2)
Theophylline	Raises ICP in intracranial hypotension — mechanism uncertain, possibly adenosine antagonism and vasoconstriction restoring CSF volume/turgor (hit 1)

The field has not tested these models against each other; the only human material-property data are two MRE scans in one patient.

Etiologies, precipitants and clinical settings

1. What the pooled series say

Series	n	Population	Underlying hydrocephalus etiology	Precipitant / context
Pang & Altschuler 1994 (PMID 7808607)	12	mostly adults	shunted hydrocephalus of varied cause	shunt malfunction / prolonged shunting
Owler 2001 (PMID 11599454)	5	mixed	tumour, SAH, prior shunts	postoperative
Lesniak 2002 (PMID 12617236)	10	adults	mixed	repeated shunt failures
Akins 2011 (PMID 21523524)	9	adults (neuro-ICU)	SAH, chronic hydrocephalus, tumour, chronic CNS infection	4/9 during index admission for intracranial bleeding; 7 days-5 yrs after primary dx
Hamilton & Price 2012 (PMID 22116443)	20	children + adults	meningitis, post-LP, intraventricular tumours	acute deterioration
Hunn 2014 (PMID 24275338)	8	adults	mixed	all had ≥ 1 prior intracranial procedure (mean 3.0)
Foster 2016 (PMID 27069743)	16	adults	mixed	prolonged EVD
Smalley 2017 (PMID 28362957)	29 (30 episodes)	children	tumour, haemorrhage	11/30 after lumbar puncture; 19 spontaneous
Cheng 2017 (PMID 28264586)	7	adults	—	VP shunt at lowest setting failing
Wu 2019 (PMID 30611954)	39	adults	TBI, haemorrhage, tumour resection, CNS infection (18/39)	~3.5 operations per patient
Keough 2021 systematic review (PMID 32736355)	195 (42 local + 153 literature)	53 children, 142 adults	neoplasm and haemorrhage most common	39 % of children post-LP; shunt in situ in 92 % of children and 39 % of adults
Casado Pellejero 2022 (PMID 34998487)	5	adults	mixed	prior shunt / infection / surgery
Czorlich 2023 (PMID 37004138)	15 of 406 SAH (3.7 %)	adults	aneurysmal SAH	IVH presence and extent; mean onset day 13
Godoy Hurtado 2023 (PMID 37373809)	6	mixed	medulloblastoma $\times 2$, TBI with bifrontal craniectomy, craniopharyngioma, leptomeningeal glioneuronal tumour, shunted NPH	4/6 already had mid/low-pressure shunts
Zheng 2023 (PMID 36994490)	41 of 68 DOC patients with hydrocephalus (60 %)	adults with severe disorders of consciousness	TBI/haemorrhage with craniectomy	ventriculomegaly + poor compliance

Series	n	Population	Underlying hydrocephalus etiology	Precipitant / context
Chen 2025 pooled paediatric (PMID 40538927)	48 (44 literature + 4 own)	children, mean age 6–8.5 y	neoplasm 71 % (medulloblastoma, astrocytoma 59 % of tumours), haemorrhage 15 % (incl. PHH of prematurity), TBI 8 %, congenital (Dandy- Walker, craniosynostosis, aqueduct stenosis, NF1)	> 50 % had prior surgery; 41 % "spontaneous" after primary treatment
Datta & Tu 2025 (PMID 41191101)	6	children	—	prior jugular/sigmoid sinus thrombosis in 2/6
Villalonga 2025 (PMID 41165379)	16	adults	—	Pang-Altschuler criteria
Bakmeedeniya 2026 systematic review (PMID 41525875)	215 from 30 studies	mixed	TBI, aneurysmal SAH, NPH, postoperative, spontaneous	—

2. Precipitants by mechanism class

A. An external or extra-ventricular CSF sink

- **Lumbar puncture** in a shunted child or adult with obstructive hydrocephalus (Dias 1999, PMID 10352418; Preuß 2013, PMID 23715811; Smalley 2017; Vilela 2020, PMID 32712403; Koenig 2025, PMID 41357016; Hoffman 2026, PMID 42224725). 37–39 % of paediatric cases are post-LP (Smalley; Keough). The shunt is patent; the LP creates a spinal SAS at lower pressure than the ventricle across an obstructed fourth ventricle/aqueduct.
- **Lumbar or spinal drainage** after SAH or posterior-fossa surgery (Ozeki 2022, PMID 34526456; Suzuki 2021, PMID 35854689; Dagnew 2002 acquired Chiari after lumbar drainage, PMID 12188966; Staykov 2011 lumboventricular gradient, PMID 21273925; Bakhshi 2020 lumbar-drain complications, PMID 32711138).
- **Skull-base and cranial-wound CSF leaks / pseudomeningocele** (Filippidis 2011, PMID 21800965 — 3 cases after cranial-base approaches, all with documented or presumed ventricle-SAS block; Stevens 2024 skull-base leak with tension pneumocephalus, PMID 39220674; Smith 2016 posterior fossa pseudomeningocele, PMID 29506116; Udayakumaran 2018 leaks after tethered-cord surgery, PMID 29605699; Bakhsh 2022 LP-shunt tract leak, PMID 35673636; Kumar 2025 extra-axial ETV for persistent leak, PMID 41215506).
- **Over-draining cyst or spinal shunts** (Vassilyadi 1995 spinal arachnoid cyst-pleural shunt, PMID 7666227; Diaz-Romero Paz 2019 external cyst drain, PMID 30710718; Marianayagam 2017 trapped fourth ventricle treatment, PMID 28209310).
- **Hemispherectomy with CSF rhinorrhoea** (Daniel 2002, PMID 12027920).

B. Parenchymal softening / loss of turgor

- **Aneurysmal SAH and IVH** — 3.7 % of SAH patients; IVH extent is the risk factor; aLPH patients later needed permanent shunts far more often (66.7 % vs 17.4 %) (Czorlich 2023). Case reports: Wang 2024 with quantitative EEG warning (PMID 39740216); Weisbrod 2021 angiogram-negative SAH after COVID-19 (PMID 34462697); Badran 2018 haemangioblastomatosis SAH (PMID 30287249). Background: Chen 2017 and Kuo 2021 reviews of post-SAH hydrocephalus (PMIDs 28373987, 34068783).
- **Ventriculitis / meningitis / chronic CNS infection** — 18/39 in Wu 2019; Akins 2011; Salam 2025; Kuppler 2025. Infection is also the strongest predictor of poor outcome (Bakmeedeniya 2026; Houlden 2018).

- **Brain tumour and its treatment** — posterior-fossa tumours in children (medulloblastoma, ependymoma, astrocytoma), suprasellar lesions (craniopharyngioma; Houlden 2018's four VLP cases followed suprasellar resection), tectal tumours (Chang 2022 Holmes tremor, PMID 35148507), leptomeningeal/intraventricular spread (Godoy 2023; Myers 2023), radiotherapy (Duan & Hu 2024 list prior cranial radiotherapy as a risk factor).
- **Traumatic brain injury** — Wu 2019; Svedung Wettervik 2022 (PMID 34414834: patients with "low-pressure hydrocephalus symptoms" had the best shunt response among post-traumatic hydrocephalus patients); Salam 2025; He 2026 (PMID 42444806).
- **Long-standing shunt dependence** — 81 of the 231 records carry the `shunt_dependent_prior` tag; Pang's original series was entirely shunted patients; Keough found a shunt in situ in 92 % of children.
- **Sepsis / cardiovascular collapse** (McGovern 2013, PMID 23816322) and **COVID-19** (Weisbrod 2021) — single cases suggesting systemic inflammatory/vascular insults can precipitate the state.

C. Loss of cranial containment

- **Decompressive craniectomy / sinking-flap physiology** — Godoy 2023 case 3; Salam 2025; Bateman 2026 (PMID 42355921); Bhatjiwale 2023 malignant oedema after cranioplasty (PMID 37680924); Ding 2014 review of craniectomy and hydrocephalus (PMID 25252237); Kutty 2018 (PMID 29725175) and Wan 2016 (PMID 26971294) on managing ventriculomegaly and abnormal flap pressure; Joshua 2011 papilloedema after craniectomy (PMID 21676532); Zheng 2023 combined cranioplasty + shunt in DOC patients. The open cranium abolishes the rigid container assumed by Monro-Kellie, so atmospheric pressure acts directly on the brain and small gradients translate into large volume shifts.

D. Shunt hardware as precipitant

- Anti-siphon devices and gravitational units that prevent sub-zero drainage in a brain that needs it (McCullough 1986; Drake 1993; Trinh 2013; Funnell 2020).
- Programmable valves left at high settings after over-drainage complications (Marianayagam 2017; Doshi 2022 with raised intra-abdominal pressure, PMID 36600762).
- Ventriculopleural/atrial shunts and the general risk of negative-pressure cavities — Chiang 2001 (excluded from core), Datta & Tu 2025 (used deliberately).

3. Paediatric versus adult spectrum

Feature	Children	Adults
Dominant etiology	Posterior-fossa tumour (71 % neoplasm in Chen 2025), PHH of prematurity, congenital malformations	SAH/IVH, TBI, CNS infection, postoperative skull-base leak, shunted NPH
Precipitant	Lumbar puncture in 37–39 %	Usually spontaneous after surgery, drain or infection
Shunt in situ at onset	92 % (Keough)	39 % (Keough)
Distinctive symptoms	Bradycardia, seizures, failure to thrive, vomiting (Chen 2025; Smalley 2017)	Depressed consciousness, headache, gait failure, cranial nerve palsies
Course of post-LP cases	Often benign: recumbency $\pm$ blood patch; median 4 days of external drainage vs 11 for spontaneous (Smalley)	Blood patch effective in adults too (Vilela 2020)
ETV	Rarely sufficient alone; 91 % need shunt repositioning (Chen 2025)	Reduces shunt dependence (Hamilton); ETV-only success 50 % (Casado Pellejero)
Return to baseline	83 % (Keough); 77.5 % (Chen)	49 % (Keough); poor in infected/DOC cohorts (Wu 2019)

Infant brain compliance (Rekate 2020, PMID 32488353; Welch 1980, PMID 7373397; Dirrichs 2019) and the frequency of obstructive posterior-fossa pathology explain why the paediatric phenotype is dominated by the LP-sink mechanism and why it recovers more completely.

4. Frequency estimates

- Post-SAH: 15/406 = 3.7 % (Czorlich 2023).
- Severe DOC with hydrocephalus: 41/68 = 60 % low or negative pressure (Zheng 2023) — the highest prevalence reported, suggesting that in this population LPH is the rule and that recovery of consciousness may depend on recognising it.
- Post-traumatic hydrocephalus: 14/29 shunted patients had "impeded recovery or low-pressure hydrocephalus symptoms" (Svedung Wettervik 2022).
- Titanium-mesh exposure after cranioplasty with a shunt: 9/10 had LPH (Zheng 2025, PMID 39760505; excluded from core but illustrates the association with chronic intracranial hypotension).
- Cumulative published cases: ~90 (Strand 2018, 25-year search), ~195 (Keough 2021), ~200 (Godoy 2023), 215 (Bakmeedeniya 2026). The Chinese consensus states incidence is "systematically underestimated" and no epidemiology exists.

Diagnosis, imaging, ICP findings and differential diagnosis

1. Clinical phenotype

Symptoms are those of acute or subacute hydrocephalus and are indistinguishable at the bedside from high-pressure disease:

- Depressed level of consciousness — the predominant sign in the pooled 195 (Keough 2021, PMID 32736355); may progress to coma with brain-stem signs (Hunn 2014 mean GCS rise 4.6 on treatment, PMID 24275338).
- Headache, nausea/vomiting (nausea/vomiting in half of Smalley's children), lethargy, confusion.
- Cranial neuropathies (up-gaze palsy, Parinaud's syndrome with tectal compression), ataxia and gait failure, urinary retention/incontinence.
- Paediatric-specific: bradycardia, seizures, failure to thrive (Chen 2025, PMID 40538927).
- Movement disorder: Holmes tremor with triventricular dilation (Chang 2022, PMID 35148507).
- **Postural pattern:** symptoms worse upright and relieved recumbent in the Foltz/Dias post-LP phenotype (Foltz 1988, PMID 3343612; Dias 1999, PMID 10352418; Ozeki 2022 headache on head elevation, PMID 34526456) — the same orthostatic pattern as intracranial hypotension, which is a clue that the two entities share physiology.
- **Timing:** after a precipitant (LP, drain, leak) deterioration occurs within hours to days; after SAH the mean latency was 13.1 ± 7.7 days (Czorlich 2023, PMID 37004138); after the primary neurosurgical diagnosis it ranged from 7 days to 5 years (Akins 2011, PMID 21523524).
- **Early warning:** falling EVD output at unchanged drain height as the state develops (Akins 2011); quantitative EEG slowing many hours before clinical deterioration, reversing with CSF removal (Wang 2024, PMID 39740216).

2. Pressure measurement

- **Direct ventricular measurement is required.** Lumbar pressure may not reflect ventricular pressure when the compartments are disconnected (Chang 2026 lumbar-ventricular discrepancy, PMID 42218510; Staykov 2011, PMID 21273925), and in post-LP cases the lumbar compartment is the sink. Shunt-tap manometry through the reservoir is acceptable in shunted patients (Hoffman 2026, PMID 42224725), as is Ommaya puncture (Chen 2025).
- **Reported values:** 2.2–6.6 mm Hg (Pang 1994); mean 4.8 mm Hg (Lesniak 2002); –3 mm Hg (Pandey 2017); –4 cm H₂O (Diaz-Romero Paz 2019); all < 70 mm H₂O and as low as –10 cm H₂O (Wu 2019); –5 to +10 mm Hg with excursions to –21 mm Hg (Salam 2025); ≤ 5 cm H₂O inclusion threshold in most series (Keough 2021; Datta & Tu 2025).

- **Continuous monitoring** (parenchymal or EVD transducer) is used to titrate therapy and to document the return of positive ICP as the end-point of sub-zero drainage (Pandey 2017; Godoy 2023 — drainage continued "with concomitant ICP monitoring at the neuro-ICU"). Background technique papers: Chari 2017 intraparenchymal monitoring for CSF disorders (PMID 28799016); Toma 2010 continuous ICP to investigate shunt function (PMID 20799857); Arroyo-Palacios 2016 ICP pulse morphology predicting EVD weaning (PMID 27106888); Eide 2012 showing that ICP pulse-wave parameters barely change when mean ICP is negative until it falls below -15 mm Hg (PMID 22153319) — so pulse amplitude cannot be used to detect the state.
- **The trap:** a normal ICP (5–12 mm Hg) in a patient whose shunt is set to 10 cm H₂O or whose EVD is at 15 cm H₂O is inappropriately low for the degree of ventriculomegaly and symptoms; the absolute number is less important than the mismatch between ventricles, symptoms and pressure (Hamilton & Price 2012; Gnanalingham 2004 "14 mmHg: a case of raised intracranial pressure?", PMID 15201374 — retained as a reminder that thresholds are individual).

3. Imaging

- **Ventriculomegaly** out of proportion to the sulci: Evans index > 0.3 (Duan & Hu 2024, PMID 38579413), rising frontal-horn ratio tracked as the treatment target (Houlden 2018, PMID 30009119), third-ventricular width used as the response metric (Bruk 2026, PMID 42732439: $13 \rightarrow 8$ mm and $13 \rightarrow 4$ mm).
- **Effaced convexity subarachnoid space and Sylvian cisterns** — the CSAS is collapsed because it is at lower pressure than the ventricles; the 2025 Chinese consensus makes this an explicit criterion, together with a rounded, enlarged fourth ventricle ("ball sign") implying fourth-ventricular outlet obstruction (PMID 40734409).
- **Transependymal / periventricular oedema despite low ICP** in 5/9 of Akins' patients — counter-intuitive and a pointer to the poroelastic model.
- **Enlarged perimedullary cisterns and brain-stem distortion** after LP-induced LPH (Vilela 2020, PMID 32712403), and descending transtentorial herniation with ventriculomegaly (Marianayagam 2017, PMID 28209310) — the caudal compartment is at lower pressure so structures are pulled down.
- **Absent cistern sign** and other LP-shunt function signs are unreliable (Kulkarni 1999; excluded).
- **Cine/phase-contrast MRI** documented CSF pathway obstruction in all 20 SILPAH patients (Hamilton & Price 2012) and reversed aqueductal flow toward the ventricles during LPH (Bateman 2026, PMID 42355921). Radionuclide or contrast ventriculography via the EVD has been used to localise the block (Rekate 2008; Filippidis 2011).
- **MR elastography** — very low shear stiffness during an episode, recovered two years later (Olivero 2016, PMID 27198914; Olivero 2021, PMID 33030603); Strand 2018 (PMID 28962963) suggests MRE as the tool most likely to give an objective diagnostic marker.
- **Sunken flap / paradoxical herniation** in craniectomised patients (Salam 2025; Bateman 2026).

4. Proposed diagnostic criteria (converging sets)

Criterion	Pang & Altschuler 1994	Duan & Hu 2024	Datta & Tu 2025	Chinese consensus 2025
Symptoms of hydrocephalus / neurological worsening	✓	✓	✓	✓ (progressive deficits)
Ventriculomegaly	✓	Evans > 0.3	change in ventricular size	progressive dilation; effaced cisterns; 4th-ventricle "ball sign"
Ventricular ICP	≤ 5 cm H ₂ O	≤ 5 cm H ₂ O	≤ 5 cm H ₂ O	0–70 mm H ₂ O on ventricular puncture
Response to drainage	improvement with sub-zero drainage	improvement with negative-pressure drainage	improvement after CSF drainage	(management criterion)

Criterion	Pang & Altschuler 1994	Duan & Hu 2024	Datta & Tu 2025	Chinese consensus 2025
Exclusions	patent shunt	ventriculomegaly from neurodegeneration	—	high-risk history required

Bakmeedeniya 2026 proposes a "diagnostic and management framework" but the field still lacks a validated definition.

5. Differential diagnosis

Entity	Shared features	Discriminators
Shunt obstruction / high-pressure hydrocephalus	identical symptoms, ventriculomegaly	ICP high on tap; shunt revision works. LPH: patent shunt at tap/revision, low ICP, revision fails (Filippidis; Hoffman)
Shunt overdrainage / slit-ventricle syndrome (Faulhauer 1978, PMID 742440; Serlo 1985/1989; Rekate 2008 shunt-related headaches, PMID 18259760; Pedersen 2023 definition review, PMID 36639536)	low or negative ICP, postural headache, subdural collections (McCullough 1974)	ventricles small or slit-like; Rekate type-1 SVS is the low-pressure form; treatment is raising resistance (anti-siphon, higher valve setting; Gruber 1984, PMID 6726390; Aschoff 1995, PMID 7621479) — the opposite of LPH. Bateman 2013 hypertensive SVS (PMID 23991845) shows SVS itself is heterogeneous
Spontaneous / iatrogenic intracranial hypotension (Mokri 2000, PMID 11113229; Binder 2002 intrathecal saline for obtundation, PMID 12188967; Samadani 2003 after lumbar drainage, PMID 12493112; Atkinson 1998 acquired Chiari, PMID 9452230)	postural headache, low ICP, sagging brain, obtundation, response to blood patch/Trendelenburg	ventricles small; no ventriculomegaly. LPH is best understood as intracranial hypotension plus a ventricle-SAS disconnection that inverts the gradient
Post-dural-puncture headache in a shunted patient (Koenig 2025, PMID 41357016)	post-LP, headache	check ventricular size — if enlarging, treat as LPH; blood patch carries extra considerations
Normal-pressure hydrocephalus (historical axis; Khan 2013 overdrainage in iNPH, PMID 23930853; Funnell 2020)	ventriculomegaly with normal opening pressure, gait/cognitive decline	chronic, elderly, B-waves and intermittent hypertension (Symon 1972); improves with standard shunting; "ultra-low-pressure NPH" is the overlap zone
Cerebral venous overdrainage (Barami 2016, PMID 27581321)	after CSF diversion, low ICP	may be a mechanism of LPH rather than a separate entity
Sinking-flap syndrome / paradoxical herniation	craniectomy, sub-atmospheric physiology, deterioration on LP or drainage	flap sunken, midline shift away from defect; cranioplasty is curative; LPH may coexist or follow (Bateman 2026; Bhatjiwale 2023)
External hydrocephalus of infancy (Sainz 2019, PMID 30474714)	infant macrocephaly	enlarged SAS with small ventricles from venous hypertension — the mirror image
Ventriculomegaly ex vacuo / neurodegeneration	large ventricles, normal ICP	no progression with symptoms; no response to drainage (Duan & Hu exclusion)
Tension pneumocephalus with skull-base leak (Stevens 2024, PMID 39220674)	post-shunt deterioration, leak	air on CT; both may coexist

6. Practical diagnostic sequence distilled from the algorithms (Keough 2021; Casado Pellejero 2022; Hoffman 2026; Duan & Hu 2024)

1. Suspect LPH in any patient with symptomatic ventriculomegaly and (a) a patent shunt, (b) a recent LP/lumbar drain/CSF leak/craniectomy, (c) an EVD that drains little at standard height, or (d) SAH/IVH with clinical stagnation at ~2 weeks.
2. Measure ventricular pressure directly (EVD, shunt reservoir, Ommaya).
3. Confirm shunt patency (tap, shuntogram) rather than revising empirically.

4. Image for the sink and the block: CT for ventricles, sulci, flap, pneumocephalus; MRI with cine/phase-contrast for aqueduct/fourth-ventricle outlet patency; look for pseudomeningocele, skull-base defect, spinal CSF collection.
 5. Therapeutic test: lower EVD to 0 then sub-zero targeting 5–15 mL/h; clinical and radiological improvement confirms the diagnosis (Pang/Foster criterion 4). The series call the response "rapid" (Owler 2001) but give no fixed reassessment interval.
 6. Exclude/treat coexisting infection, hyponatraemia and hypocapnia, which lower turgor.
-

Management — acute algorithm, definitive diversion, refractory disease

No randomised or prospective comparative data exist. What follows is the consensus pattern extracted from the series, algorithms and reviews (Pang 1994; Owler 2001; Akins 2011; Filippidis 2011; Hamilton & Price 2012; Hunn 2014; Foster 2016; Smalley 2017; Wu 2019; Keough 2021; Casado Pellejero 2022; Godoy Hurtado 2023; Duan & Hu 2024; Li 2024; Chen 2025; Datta & Tu 2025; Bakmeedeniya 2026; Chinese consensus 2025), with dissent noted.

1. Acute phase — restore brain volume

1a. Remove or counter the sink first where possible

- **Post-LP / lumbar drain:** flat or Trendelenburg positioning; stop the drain; epidural blood patch if not resolving in 24–48 h (Dias 1999, PMID 10352418; Smalley 2017, PMID 28362957; Vilela 2020 adult, PMID 32712403; Suzuki 2021 before EVD, PMID 35854689; Hoffman 2026, PMID 42224725; Koenig 2025 cautions on blood patch in shunted patients, PMID 41357016). In Smalley's 30 episodes post-LP cases needed a median 4 days of drainage vs 11 days for spontaneous cases.
- **CSF leak / pseudomeningocele:** direct repair, lumbar drain avoidance, fibrin sealant (Filippidis 2011, PMID 21800965; Stevens 2024, PMID 39220674; Kumar 2025 extra-axial ETV variant for persistent leak, PMID 41215506).
- **Cyst or spinal shunt:** ligate/remove (Vassilyadi 1995, PMID 7666227; Diaz-Romero Paz 2019, PMID 30710718; Ozeki 2022, PMID 34526456).
- **Sunken flap:** early cranioplasty (Bateman 2026, PMID 42355921; Zheng 2023 combined cranioplasty + shunt, PMID 36994490).

1b. Raise venous pressure / brain turgor (non-invasive, hours)

- **Neck wrapping / soft cervical collar / bilateral jugular compression:** Filippidis 2011 (all 3 patients), Pandey 2017 (PMID 28017745 — combined with Trendelenburg, hypoventilation to raise PaCO₂ and lowered plasma osmolarity to reverse ICP from –3 mm Hg), Bruk 2026 (PMID 42732439 — two children, third ventricle 13 → 8 and 13 → 4 mm within days, both then internalised), Kuppler 2025 (failed in chronic disease), Rekate 2019 review. Mechanism support: Hatt 2015 MRE (PMID 26045579). Risks: jugular thrombosis, discomfort, poor tolerance in agitated patients; requires ICP monitoring.
- **Head-down positioning / recumbency:** Foltz 1988; Dias 1999; Pandey 2017.
- **Fluid and gas management:** avoid hypocapnia and hyperosmolar therapy; Houlden 2018 (PMID 30009119) recommends avoiding fluid shifts; Pandey deliberately lowered osmolarity. This is expert opinion.

1c. Titrated external ventricular drainage (the reference standard)

Protocol synthesised from Pang 1994, Owler 2001 (PMID 11599454), Akins 2011 ("sub-zero method"), Hunn 2014 (PMID 24275338), Foster 2016 (PMID 27069743), Godoy Hurtado 2023 (PMID 37373809) and the paediatric reviews:

Each step is followed by the source wording it rests on (quotes from abstracts, or from the full text where marked FT — Foster, Godoy, Salam and Chen from the mirrored OA XML, Akins from the publisher PDF). Where the papers do not state a number, that is said explicitly rather than filled in. Akins 2011 is the only paper that lays out the sub-zero method as an explicit protocol; Pang 1994, Owler 2001, Hunn 2014 and Wu 2019 are available here only as abstracts.

1. **Establish external ventricular access** — a new EVD or the existing shunt externalised; failure at conventional drainage pressure is part of the definition, not a step to be tried. - Pang 1994: "Seven patients failed to improve with multiple shunt revisions, including the use of low-pressure valves. In 11 patients, symptoms and ventriculomegaly were not reversed except with prolonged external ventricular drainage at subzero pressures". - Owler 2001: patients "failed to respond to apparently adequate CSF drainage via a functioning shunt (four cases) or external ventricular drain (one case)". - Foster 2016 (FT), Pang criteria: "(1) Neurologic decline at normal EVD or shunt pressures; (2) ventriculomegaly; (3) persistence of ventriculomegaly with ICP in the normal to low-normal range; and (4) clinical and radiographic response to sub-zero drainage". - Akins 2011 (FT) makes the failed conventional heights explicit in their inclusion criteria: "(4) clinical and radiographic response to EVD drainage at +5 to +10 cm height; (5) delayed neurologic decline accompanied by recurrent hydrocephalus despite confirmation of a patent EVD and normal ICP; (6) failure to respond to lowering EVD (0 to +5 cm); and (7) clinical and radiographic response to sub-zero drainage". Illustrated case 4: "EVD was lowered to +5 and then 0 cm without clinical or radiographic improvement." - Chen 2025 (FT), 28 paediatric cases with route recorded: "EVD, n (%) 12 (42.86%) ... VPS externalized, n (%) 16 (57.14%) ... Ommaya, n (%) 4 (14.29%)". - Salam 2025 (FT): "subatmospheric CSF diversion via external ventricular drain (EVD) placement or shunt externalization with or without neck wrapping". - Catheter site and calibre are not specified in any series; individual reports use a frontal EVD (Salam 2025: "a new right frontal EVD").
2. **Lower the drain below the reference point until CSF flows** — start at 0 to −5 cm H₂O, go to −10 if needed; deeper settings are reported only in refractory cases. No paper specifies a fixed interval between downward steps. - Akins 2011 (FT), describing Pang's method: "sub-atmospheric ventricular drainage achieved by setting the EVD height below the level of the external auditory canal"; their own practice: "sub-zero EVD drainage (typically, −3 to −10 cm)". Case 6 stabilised "with EVD at −3 cm height". - Owler 2001: "a period of external ventricular drainage at heights of 0 to −5 cm H₂O below the reference point (external auditory meatus—EAM)". - Foster 2016 (FT): "The EVD was initiated at a subatmospheric pressure, placed at a height ranging from 5 to 10 cmH₂O below midbrain"; the lowest setting in their illustrative case was "−12 cmH₂O". - Pang 1994: "mean external ventricular drainage nadir pressure of -5.7 ± 3.6 mm Hg (≈ -8 cm H₂O)". - Wu 2019: "Ventricular pressure was lower than 70 mm H₂O in all 39 patients, and the lowest value was −10 cm H₂O" (this is the lowest measured pressure, not a drainage setting). - Godoy Hurtado 2023 (FT): drainage "at negative pressures which were reached progressively to 15 cm below the EAC". - Salam 2025 (FT), refractory case: EVD "set to −8 cm H₂O", later "−10 cm H₂O", and at the extreme "set as low as −30 cm H₂O for forced ventricular reduction"; the drain "was adjusted hourly ... by either raising or lowering it by 2 cm H₂O" to allow spontaneous flow.
3. **Titrate to CSF output, not to a pressure** — 5–15 mL/h is the only explicit hourly target in a series; others give a daily volume. - Hunn 2014: "titrated to produce between 5 and 15 mL per hour of CSF drainage, including drainage under subatmospheric pressure if this was required to maintain CSF flow". - Salam 2025 (FT), summarising the literature: "large-volume aspiration of 30–50 mL of CSF and subsequent drainage of 10–15 mL/hr, with a willingness to lower drainage settings to less than zero to meet this target"; their own goals were 10, 15 and 15–20 mL/h. - Daily-volume authors: Godoy Hurtado 2023 (FT) "daily debts around 300 mL of CSF"; Pandey 2017 (PMID 28017745) "daily cerebrospinal fluid drainage >200 mL". - Akins 2011 describes the output signature rather than a target: "As patients entered the LPH state, the ICP remained in a normal range yet daily CSF output from the external ventricular drain was reduced."
4. **Expect a CSF "diuresis" when sub-zero drainage starts; monitor ICP if a parenchymal sensor is available.** The duration is given only as "several days"; Akins attribute the extra fluid to water leaving the parenchyma, not entering it. - Akins 2011: "When LPH patients were drained at sub-zero levels, daily CSF output exceeded baseline values for several days and then receded to baseline ... When LPH was treated with the sub-zero method, a 'diuresis' of CSF ensued." (FT): "During sub-zero treatment, we observed a rise in EVD output which we attribute to reduction in

ventricular volume and reduction in brain extracellular fluid"; sub-zero treatment "will 'wring out' the sponge-like cortical mantle ... Water will flow from the parenchyma into the ventricular system." - Foster 2016 (FT): "We, too, noted decreased output during genesis of LPH with a significant increase in output during sub-zero drainage." - Godoy Hurtado 2023 (FT): "Continuous recording of ICP with the intraparenchymal sensor during subzero drainage showed negative pressures, a flat recording with no pathological waves, and minimal amplitude between systolic and diastolic values, suggesting a very distensible or 'compliant' brain."

5. **Response to sub-zero drainage confirms the diagnosis** (criterion 4 above). The series describe the response as rapid but do not fix a reassessment interval. - Owler 2001: "Four of the five patients showed rapid and significant clinical improvement ... In the fifth patient there was no improvement with low pressure external drainage and no shunt revision was undertaken." - Hunn 2014: "TEVD resulted in improvement in Glasgow Coma Scale (mean increase 4.6, $p=0.003$), and increases in ICP (mean increase 8.5)"; "Mean delay from first negative ICP to TEVD was 1.8 days." - Pang 1994: "During external ventricular drainage treatment, symptoms correlated only with ventricular size and not with ICP."
6. **Wean upward slowly once ventricles are small and CSF drains at a positive pressure; expect weeks to months.** Both explicit protocols use a 3–5-day step interval. - Akins 2011 (FT): "the EVD was raised in small increments (typically 3 cm steps) every 3–5 days. Prior to raising the EVD to a more positive height, clinical and radiographic stability was confirmed. If patients relapsed during EVD weaning, then the process was repeated. Shunt internalization was performed once clinical and radiographic stability was confirmed at neutral or positive EVD settings (typically 0 to +3 cm)." "Three patients had refractory LPH and were unable to tolerate positive pressure EVD drainage" and went to ventriculopleural or valveless shunts. - Foster 2016 (FT): the EVD was "incrementally raised 1–2 cmH₂O every 3–5 days"; sub-zero drainage continued "until the ventricular shrinkage was maintained at a positive pressure, and a new shunt system could be inserted"; "patients required prolonged hospitalizations for protracted sub-zero EVD, often on the order of months". Mean time from first ventriculostomy to shunt was 73 days (range 14–257) without ETV and 114 days (0–236) with ETV, $P = 0.16$ — means, not medians — and the authors attribute the lack of difference to "an inherent bias to perform ETV on patients who were refractory to sub-zero drainage". One patient "underwent rapid EVD wean (in 8 days) and did not require shunting". - Pang 1994: sub-zero drainage "for a mean period of 22.2 days". - Chen 2025 (FT), pooled paediatric data: "the average number of days of external CSF drainage was 41.89 days, with a minimum of 2 days and a maximum of 365 days" ($n = 28$). - Wu 2019: "the time interval from onset to last VP shunt achieved was 31–3880 days, with an average of 376 days" (29 survivors).
7. **Infection is both a precipitant and a complication of prolonged drainage, and predicts poor outcome.** The "vicious circle" wording in the earlier draft of this note is not in the sources; what they document is: - Wu 2019: "Eighteen cases had CNS infection" (of 39); Salam 2025 (FT) quoting the same series: "46.15% were complicated by intracranial infections, of which 87.5% were specifically in the ANPH subset compared to 35.48% of ALPH". - Akins 2011: "Conditions most commonly associated with LPH are: subarachnoid hemorrhage, chronic hydrocephalus, brain tumors, and chronic CNS infections." (FT): "We hypothesize that underlying inflammatory or degenerative changes alter periventricular brain permeability in LPH patients, for example, chronic CNS infection or chemical meningitis from subarachnoid hemorrhage." Conversely, they saw no infectious harm from the treatment itself: "No long-term complications were observed with low pressure shunting such as shunt infection or subdural hematoma." - Chen 2025 (FT), institutional cases drained via Ommaya for a median of 105 days: this approach "increase[s] the risk of intracranial infection, and one patient ultimately died from intracranial infection". - Bakmeedeniya 2026: "delayed intervention, infection, and irreversible white-matter injury predicted poor recovery". - Salam 2025 (FT): the index case developed refractory NegPH after shunt infection ("removal of his shunt" under ceftriaxone/ampicillin).

Dissent: Hamilton & Price 2012 and Casado Pellejero 2022 (PMID 34998487) prefer early ETV to avoid prolonged EVD; Chinese consensus and Li 2024 (PMID 38688300) recommend EVD/lumbar-cisternal drainage as first line with ETV or shunt for definitive care.

1d. Pharmacology (experimental)

- **Theophylline** (Salam 2025, PMID 40418891): refractory acute NegPH after TBI, hemicraniectomy, shunt infection and IVH; ICP -5 to $+10$ mm Hg with excursions to -21 mm Hg despite sub-zero EVD (as low as -30 cm H₂O), two ETVs and ventriculopleural/ventriculoatrial shunts over ~ 4 months. After 48 h of oral caffeine without effect, "two doses of theophylline 200 mg twice a day" via PEG (borrowed from post-dural-puncture headache practice) were followed by spontaneous drainage at -18 cm H₂O and, on CT 4 h after the second dose, "complete collapse of the ventricles to an almost slit-like appearance"; focal seizures progressing to status epilepticus began 4 h after the first dose. Single case; mechanism (adenosine antagonism $\rightarrow$ cerebral vasoconstriction $\rightarrow$ restored CSF production / turgor?) untested.
- **Acetazolamide / diuretics** are contraindicated in principle (they lower ICP further); mentioned only as things to stop.
- No other drug has been reported for the entity; the consensus and review literature is silent on pharmacology beyond avoiding ICP-lowering agents.

2. Reconnecting the compartments — ETV

- **Rationale:** if the syndrome depends on ventricle-SAS disconnection, a fenestration abolishes the gradient and may abolish shunt dependence (Rekate 2008; Hamilton & Price 2012).
- **Data:**
- Hamilton & Price 2012 (PMID 22116443): 20 SILPAH patients, 10 managed with and 10 without ETV; ETV produced "a rapid return of normal ICP dynamics and a significant decrease in the number of shunt-dependent patients".
- Casado Pellejero 2022 (PMID 34998487): algorithm with early ETV; 5 patients, ETV alone succeeded in 50 %, remainder needed a shunt; one death from systemic infection.
- Villalonga 2025 (PMID 41165379): ETV in 16 adults, then valveless VP shunt if persistent; 81 % excellent outcome, mean KPS gain 56 points; 19 % system dysfunction requiring a valved shunt.
- Foster 2016 (PMID 27069743): 16 patients, ETV did not significantly reduce the duration of EVD or shunt dependence.
- Chen 2025 pooled paediatric: conservative measures and ETV rarely sufficed; 91 % ultimately required shunt repositioning/revision.
- Kuppler 2025: ETV failed in chronic NegPH.
- Kumar 2025 (PMID 41215506): "extra-axial ETV" (fenestration through the cranial defect) for LPH with persistent leak.
- **Interpretation:** ETV appears to help a subset — adults with demonstrable distal obstruction and preserved SAS absorption — and is a reasonable step once ventricles have been re-inflated; it is not a substitute for sub-zero drainage in the acute phase and its comparative benefit is confounded by selection (Godoy 2023).

3. Definitive CSF diversion — minimise resistance, exploit gravity

The permanent shunt must be able to operate at the sub-zero pressures the brain needs, i.e. the opposite design philosophy to anti-siphon devices:

Strategy	Sources	Notes
Programmable valve at lowest setting, no anti-siphon / gravitational unit	Pang 1994; Akins 2011; Marianayagam 2017; Chen 2025 (most children); Godoy 2023	Standard first choice; risk of later overdrainage/SVS so needs follow-up and gradual up-titration
Valveless VP shunt	Villalonga 2025; Badran 2018 (PMID 30287249 — improvised ultra-low-pressure valveless system); Duan & Hu 2024	Highest flow; watch for subdurals (McCullough 1974)

Strategy	Sources	Notes
"Therapeutic siphoning" — non-siphon-control valve or adjustable gravitational valve set to permit upright siphoning	Bergsneider 1999 (PMID 10520938: adults improved with siphoning); Trinh 2013 (PMID 22770726: converting to siphon-permitting valve in shunted NPH with LPH physiology); Funnell 2020 (PMID 32989519: ultra-low-pressure state in NPH treated by siphoning)	Applies chiefly to adult NPH-overlap cases
Valve placed below the clavicle (caudal) to add a hydrostatic column	Carrera 2025 (PMID 39951714); Datta & Tu 2025 cite "shunt repositioning"	Converts body height into negative drainage pressure
Shunt into a negative-pressure cavity (pleura, atrium)	Salam 2025 (VPleural); Datta & Tu 2025; Chen 2025 (ventriculopleural in refractory children); Chiang 2001 (excluded — positive-pressure ventilation obstructs VPI shunts)	Pleural pressure −5 to −8 cm H ₂ O in spontaneous breathing; ineffective under mechanical ventilation
Wide-bore / low-resistance custom distal catheter	Galbarriatu 2013 (PMID 23904087 — thoracic-drain tubing as peritoneal catheter)	Reduces distal resistance
Active-pumping / negative-pressure shunt	Kalani 2013 (PMID 23380444 — investigational active shunt in a patient failing sub-zero EVD); background valve-technology literature: Aschoff 1995 (PMID 7621479), Czosnyka 2002 UK shunt evaluation (PMID 12111485), Chari 2014 (PMID 24405071)	Not commercially available; case-level evidence
Zero-ICP gravity shunt (ZIPS)	Foltz 1994 (PMID 8194062)	Historical; 56 patients, Foltz's own design
Intermittent reservoir pumping	Cheng 2017 (PMID 28264586: 7 adults with VP shunt at lowest setting; compressing the pump chamber several times daily improved ventricles in 6/7); Kuppler 2025 (PMID 40030163: scheduled active pressing as outpatient rescue)	Cheap, needs compliance and follow-up
Combined cranioplasty + shunt in craniectomised DOC patients	Zheng 2023	Single-stage restoration of container and drainage

Sequencing (Godoy 2023; Datta & Tu 2025; Chen 2025): internalise once the patient tolerates EVD raised to ≥ 0 –5 cm H₂O with stable ventricles; if internalisation fails at the lowest valve setting, consider valveless/negative cavity/caudal valve; if the patient re-develops LPH after a period of stability, re-externalise and repeat sub-zero drainage — many patients cycle several times (Wu 2019: 3.5 procedures per patient).

4. Chronic and refractory disease

Defined operationally as persistent or recurrent symptomatic ventriculomegaly after adequate sub-zero drainage, ETV and low-resistance shunting (Kuppler 2025's chronic negative-pressure hydrocephalus; Clarke 2006 / Houlden 2018 very-low-pressure hydrocephalus; Lin 2025 refractory hydrocephalus).

- **Predictors of refractoriness:** unsealed CSF leak, ongoing ventriculitis, suprasellar/skull-base surgery, multiple prior operations, severe underlying brain injury, permanently altered compliance (Houlden 2018; Bakmeedeniya 2026; Wu 2019).
- **Options tried:** prolonged (months) sub-atmospheric drainage; ETV; low-gradient shunt; neck wrapping (failed in Kuppler's case); intermittent active shunt pumping (Kuppler, Cheng); ventriculopleural conversion; theophylline (Salam); MRE-guided timing (Olivero — stiffness recovery may indicate when a standard valve can be tolerated); cranioplasty; and, in the DOC population, acceptance of a permanent low-pressure system with continued rehabilitation (Zheng 2023).
- **Prognosis:** Houlden's 4 VLPH cases and Wu's infected subgroup had the worst outcomes (mean GOS-E 2.7); Kuppler's patient remained shunt-dependent but stable as an outpatient.

5. Paediatric specifics

- Post-LP cases: recumbency, then blood patch; many resolve without drainage (Dias; Smalley).
- Spontaneous cases after posterior-fossa tumour surgery: EVD titration exactly as in adults; long durations (up to a year); most end with a programmable valve at lowest setting, some with ventriculopleural or repositioned shunts (Chen 2025).
- Neck wrapping is tolerated and effective in children (Bruk 2026).
- ETV alone is rarely sufficient (Chen 2025) — possibly because absorption at the convexity is also impaired by tumour/haemorrhage.
- Outcome: 77.5 % back to baseline; deaths driven by tumour progression.

6. What to avoid (from the failure reports)

- Standard-height EVD, standard valve settings or anti-siphon devices in a brain showing the paradox (McCullough 1986; Drake 1993; Trinh 2013).
- Empirical shunt revision without pressure measurement (Filippidis; Hoffman).
- Lumbar puncture or lumbar drainage in a patient with a patent shunt and obstructive pathology (Dias; Preuß; Ozeki) — or at least anticipate LPH.
- Osmotherapy, hyperventilation, diuretics in a patient with low ICP and large ventricles.
- Leaving a skull-base leak or pseudomeningocele untreated while draining sub-zero — the two sinks compete.
- Assuming that a "normal" ICP excludes hydrocephalus when the ventricles are enlarging and the patient is deteriorating.

Outcomes, prognostic factors, evidence quality and research agenda

1. Outcome data

Source	n	Outcome
Pang & Altschuler 1994 (PMID 7808607)	12	All improved with sub-zero drainage; most internalised with low-pressure hardware
Lesniak 2002 (PMID 12617236)	10	Ventricles returned to baseline size and stayed small at normal pressure after negative-pressure drainage
Akins 2011 (PMID 21523524)	9	Sub-zero method effective in all; internalised
Filippidis 2011 (PMID 21800965)	3	All resolved after leak repair + neck wrapping/drainage
Hamilton & Price 2012 (PMID 22116443)	20	Shunt dependence 20 % with ETV vs 70 % without
Hunn 2014 (PMID 24275338)	8	Mean GCS improvement 4.6; ventricular size reduced in all
Foster 2016 (PMID 27069743)	16	Prolonged EVD; ETV no significant benefit on EVD duration
Smalley 2017 (PMID 28362957)	29/30 episodes	Post-LP: median 4 days drainage, generally full recovery; spontaneous: median 11 days
Cheng 2017 (PMID 28264586)	7	Reservoir pumping improved 6/7
Houlden 2018 (PMID 30009119)	4 VLPH	Prolonged courses; recovery of GCS despite persistent ventriculomegaly in salvageable cases; poor overall
Wu 2019 (PMID 30611954)	39	Mean GOS-E 2.7; 8 deaths; infection dominated

Source	n	Outcome
Keough 2021 SR (PMID 32736355)	195	Return to baseline 83 % children vs 49 % adults; overall mortality ~11 %
Casado Pellejero 2022 (PMID 34998487)	5	ETV alone 50 %; remainder shunted; low-pressure state resolved in all treated; 1 death (20 %) from systemic infection; 80 % good outcome
Czorlich 2023 (PMID 37004138)	15	Post-SAH aLPH → permanent shunt 66.7 % vs 17.4 % in other SAH patients; outcome otherwise not worse than matched SAH
Godoy Hurtado 2023 (PMID 37373809)	6	5/6 improved; treatment individualised; ~200 cases in literature
Chen 2025 pooled paediatric (PMID 40538927)	48	77.5 % returned to pre-existing status; deaths and poor outcomes due to underlying tumour/haemorrhage; mean external drainage 41.9 d
Villalonga 2025 (PMID 41165379)	16	81 % excellent, 19 % temporary improvement; mean KPS gain 56; 19 % needed a valved shunt
Datta & Tu 2025 (PMID 41191101)	6	All improved with drainage; shunt repositioning or negative-pressure cavities
Bakmeedeniya 2026 SR (PMID 41525875)	215	Prognosis governed by etiology and speed of recognition; standardised framework proposed

Summary: when recognised and treated with titrated sub-zero drainage, most patients regain their pre-episode status; mortality of ~10 % and poor functional outcome in about a quarter to a half of adults is attributable mainly to the underlying disease (SAH, TBI, tumour, ventriculitis) and to delayed recognition. Children and post-LP cases do best; infected and skull-base/VLPH cases do worst. Permanent shunt dependence is the norm (≥ 70 % without ETV, 50–80 % overall).

2. Prognostic factors reported

Favourable: paediatric age; post-LP precipitant; short delay to sub-zero drainage; identifiable and closable sink; retained viscoelasticity (GCS improves before ventricles shrink; Houlden); clear distal obstruction amenable to ETV.

Unfavourable: CNS infection; unsealed CSF leak; suprasellar/skull-base surgery; multiple prior operations; decompressive craniectomy with DOC; large IVH burden (Czorlich); need for drainage at ≤ -10 cm H₂O; chronic course > 3 months (Kuppler).

3. Complications specific to treatment

- Sub-zero EVD: ventriculitis with prolonged drainage; subdural haematoma or hygroma from overdrainage (McCullough & Fox 1974, PMID 4813717); intracranial hypotension symptoms; catheter blockage as ventricles collapse.
- Neck wrapping: jugular venous thrombosis risk, patient intolerance.
- Blood patch in shunted patients: theoretical risk of shunt tract/lumbar complications (Koenig 2025, PMID 41357016).
- Valveless / siphoning shunts: subdurals, slit ventricles — the pendulum swings to overdrainage (Faulhauer 1978, PMID 742440; Serlo 1985, PMID 4064018; Pedersen 2023, PMID 36639536); Doshi 2022 (PMID 36600762) illustrates oscillation between overdrainage and VLPH in one patient.
- Ventriculopleural shunts: pleural effusion; failure under positive-pressure ventilation.
- Theophylline: seizures (Salam 2025).

4. Evidence quality

- **Design:** 3 systematic reviews of case-level data, 1 consensus statement, ~24 retrospective series/cohorts (largest single-centre n = 42, Keough; largest pooled n = 215), 34 case reports, no prospective study, no trial, no registry.
- **Definitional heterogeneity:** ICP thresholds ≤ 0 , ≤ 5 cm H₂O, < 7 cm H₂O; some series include "normal" ICP with failed standard drainage; acute and chronic cases pooled.

- **Selection and publication bias:** successful sub-zero drainage and successful ETV are over-reported; unrecognised cases (likely frequent in DOC and SAH populations — Zheng 2023; Czorlich 2023) are absent.
- **Mechanistic evidence:** mostly inference from treatment response; one patient with serial MRE; one with phase-contrast CBF/CSF flow; no direct transmantle gradient measurements during an episode; animal models (Shulyakov 2012, Jugé 2016, Li 2008 kaolin) address hydrocephalus compliance generally, not LPH.
- **Outcome measures:** GCS, GOS-E, mRS, "return to baseline", frontal-horn ratio, third-ventricle width — inconsistent and rarely blinded.
- **Language/region:** a substantial body of Chinese literature (Wu 2019; Cheng 2017; Li 2024; Chen 2025; Zheng 2023; Chinese consensus 2025; Lin 2025) uses the < 70 mm H₂O threshold and reports higher prevalence in DOC/post-infectious cohorts than Western series — possibly true case mix, possibly definitional.

5. Open questions and a research agenda

1. **Definition:** adopt a single operational definition (symptomatic progressive ventriculomegaly + directly measured ventricular pressure ≤ 5 cm H₂O or ≤ 70 mm H₂O + failure of standard-pressure drainage), with acute/chronic and LPH/NegPH sub-labels — a Delphi process is the obvious next step after the 2025 Chinese consensus.
2. **Epidemiology:** prospective screening of SAH, posterior-fossa tumour, ventriculitis and craniectomy/DOC cohorts with routine ventricular pressure measurement when ventricles enlarge; the 3.7 % and 60 % figures need replication.
3. **Mechanism:** serial MR elastography and phase-contrast MRI during and after episodes; simultaneous ventricular and lumbar/cortical pressure recordings to test the transmantle hypothesis directly; interstitial-water quantification (diffusion, T2 mapping) to test the poroelastic model; venous sinus imaging to test the venous overdrainage/thrombosis association (Datta & Tu 2025).
4. **Treatment protocol:** standardise sub-zero EVD titration (height steps, output targets, weaning rate) and compare early ETV vs prolonged EVD in a multicentre registry; define when neck wrapping or blood patch can replace drainage.
5. **Hardware:** evaluate caudal valve placement, valveless systems, pleural shunts and active-pumping devices head to head for internalisation failure rates and overdrainage complications.
6. **Pharmacology:** methylxanthines (theophylline, caffeine) and other agents used for intracranial hypotension deserve a formal case series with ICP monitoring.
7. **Biomarkers:** EVD output trajectory (Akins), quantitative EEG (Wang 2024), ICP pulse morphology (Arroyo-Palacios 2016; Eide 2012) as early-warning tools for the transition into the low-pressure state.
8. **Long-term outcome:** cognitive and functional follow-up of survivors, incidence of later overdrainage/SVS after low-resistance shunting, and whether brain stiffness recovers (Olivero 2021) enough to permit conversion to standard valves.

Key papers — per-paper extractions

Ordered

chronologically within groups. "FT" marks papers whose open-access XML is mirrored under `fulltext/`; the rest were read from Europe PMC abstracts.

A. Foundational descriptions

McCullough & Fox 1974 — Negative intracranial pressure hydrocephalus in adults with shunts and its relationship to the production of subdural hematoma (J Neurosurg; PMID 4813717). Shunted adults run sub-atmospheric ICP; subdural haematoma is the consequence. Establishes that negative ICP is achievable and harmful — the overdrainage pole of the same physiology.

McCullough 1986 — Symptomatic progressive ventriculomegaly in hydrocephalics with patent shunts and antisiphon devices (Neurosurgery; PMID 3491341). Patent shunts with ASDs; ventricles enlarge symptomatically; removing the ASD or lowering resistance helps. The paradox stated before it was named.

Foltz & Blanks 1988 — Symptomatic low intracranial pressure in shunted hydrocephalus (J Neurosurg; PMID 3343612). Upright, activity-related headache and lethargy; low upright ICP; relief when recumbent/Trendelenburg. Motivated the "zero-ICP" gravity-controlled shunt (Foltz 1994, PMID 8194062, 56 patients).

Pang & Altschuler 1994 — Low-pressure hydrocephalic state and viscoelastic alterations in the brain (Neurosurgery; PMID 7808607). 12 shunted patients, ventriculomegaly, ICP 2.2–6.6 mm Hg, patent shunts; prolonged sub-zero EVD restored ventricular size and symptoms; symptoms tracked ventricular size, not ICP. Proposes viscoelastic change; defines the entity and its treatment.

Vassilyadi, Farmer & Montes 1995 — Negative-pressure hydrocephalus (J Neurosurg; PMID 7666227). Child with a spinal arachnoid cyst-pleural shunt developing symptomatic ventriculomegaly at negative ICP; postulates a transmantle gradient from the extra-ventricular sink. Coins "negative-pressure hydrocephalus".

Dias, Li, Freimark & Larson 1999 — Low-pressure shunt 'malfunction' following lumbar puncture in children with shunted obstructive hydrocephalus (Pediatr Neurosurg; PMID 10352418). Post-LP symptomatic ventriculomegaly with patent shunts; recumbency ± epidural blood patch resolved it. Establishes the paediatric post-LP phenotype and its benign treatment.

Bergsneider et al. 1999 — Beneficial effect of siphoning in treatment of adult hydrocephalus (Arch Neurol; PMID 10520938). Adults whose ventricles and symptoms improved only when the shunt was allowed to siphon — the "therapeutic siphoning" idea later applied to LPH.

B. Series that defined management

Owler et al. 2001 — Low pressure hydrocephalus: issues of diagnosis and treatment in five cases (Br J Neurosurg; PMID 11599454). Five cases; stresses direct ventricular pressure measurement and drainage at or below zero; first UK series.

Lesniak, Clatterbuck, Rigamonti & Williams 2002 — Low pressure hydrocephalus and ventriculomegaly: hysteresis, non-linear dynamics, and the benefits of CSF diversion (Br J Neurosurg; PMID 12617236). 10 adults, mean ICP 4.8 mm Hg; ventricles normalised only after negative-pressure drainage and then stayed small at normal pressure — a hysteresis loop.

Daniel, Lee & Halcrow 2002 — Low-pressure hydrocephalic state complicating hemispherectomy (Epilepsia; PMID 12027920). Post-hemispherectomy with CSF rhinorrhoea; ventriculomegaly at low ICP; treated with sub-zero drainage then low-pressure shunt.

Clarke & Meyer 2006 — Very low pressure hydrocephalus (J Neurosurg; PMID 16961147) and **Houlden et al. 2018** — Importance of frontal horn ratio and optimal CSF drainage in the treatment of very low pressure hydrocephalus (J Neurol Surg A; PMID 30009119). VLPH after suprasellar/skull-base resection is often refractory; Houlden's 4 cases suggest recovery of GCS despite persistent ventriculomegaly identifies salvageable patients ("retained viscoelasticity") and recommends frontal-horn-ratio tracking, volume optimisation and avoidance of fluid shifts.

Rekate, Nadkarni & Wallace 2008 — The importance of the cortical subarachnoid space in understanding hydrocephalus (J Neurosurg Pediatr; PMID 18590387). Reframes LPH as ventricle-CSAS disconnection with a lower-pressure CSAS and loss of brain turgor; predicts benefit from ETV, neck wrapping, and closing sinks. Companion: Rekate 2019 Low or negative pressure hydrocephalus demystified (PMID 31103762) and 2021 commentary (PMID 34272587).

Akins, Guppy, Axelrod, Chakrabarti, Silverthorn & Williams 2011 (FT, publisher PDF) — The genesis of low pressure hydrocephalus (Neurocrit Care; PMID 21523524). 9 neuro-ICU patients (0.2% of admissions; 4 of 102 SAH patients with EVDs); LPH 7 days–5 years after the index condition;

EVD output fell as LPH developed; transependymal oedema in 5 despite low ICP. The only paper to state the sub-zero method as a protocol: response to EVD at +5 to +10 cm, then failure at 0 to +5 cm, then sub-zero drainage "typically -3 to -10 cm" below the external auditory canal; wean "in small increments (typically 3 cm steps) every 3-5 days" only after clinical and radiographic stability, repeat if relapse; internalise at 0 to +3 cm. Output rose above baseline for several days ("CSF diuresis"), attributed to water leaving the parenchyma (Darcy flux, poroelastic "boggy brain"). All 9 resolved; 6 programmable valves at lowest setting, 1 valveless VPS, 2 ventriculopleural (the 3 who could not tolerate positive EVD pressure); 2 relapses; no SDH or shunt infection at 6 months-4 years.

Filippidis, Kalani & Rekate 2011 — Negative-pressure and low-pressure hydrocephalus: the role of cerebrospinal fluid leaks resulting from surgical approaches to the cranial base (J Neurosurg; PMID 21800965). Three post skull-base-surgery cases with leaks and ventricle-SAS block; treated with leak repair, neck wrapping and drainage; formalises the leak-as-sink mechanism and neck wrapping as therapy.

Hamilton & Price 2012 — Syndrome of inappropriately low-pressure acute hydrocephalus (SILPAH) (Acta Neurochir Suppl; PMID 22116443). 20 patients (10 without, 10 with ETV); all had EVD with drainage below zero; cine MRI showed CSF pathway obstruction in all; ETV "reestablishes communication between the SAS and ventricles" with rapid return of normal ICP dynamics and significantly fewer shunt-dependent patients.

Kalani, Turner & Nakaji 2013 — Treatment of refractory low-pressure hydrocephalus with an active pumping negative-pressure shunt system (J Clin Neurosci; PMID 23380444). Investigational active shunt for a patient failing sub-zero EVD.

Galbarriatu et al. 2013 — Low-pressure hydrocephalus: a case report and review (Neurocirugia; PMID 23904087). Custom wide-bore peritoneal catheter (thoracic-drain tubing) to minimise distal resistance.

Trinh et al. 2013 — Shunt overdrainage or low-pressure hydrocephalus? (Neurosurgery; PMID 22770726). Shunted NPH patient with LPH physiology treated by converting to a siphon-permitting system — highlights the overdrainage/LPH diagnostic fork.

Hunn et al. 2014 — Successful treatment of negative pressure hydrocephalus using timely titrated external ventricular drainage and a low-pressure ventriculoperitoneal shunt (Clin Neurol Neurosurg; PMID 24275338). 8 adults, mean 3.0 prior procedures; TEVD titrated to 5-15 mL/h including sub-atmospheric drainage, then low-pressure VP shunt; mean GCS improvement 4.6; first series to report acceptable long-term outcomes.

Foster, Zlotnikov & Rekate (Foster 2016) — Analysis of low-pressure hydrocephalus: does ETV shorten time to internalization? (World Neurosurg; PMID 27069743). 16 patients; ETV did not significantly shorten EVD duration.

Olivero et al. 2016 / 2021 — Magnetic resonance elastography demonstrating low brain stiffness in a patient with low-pressure hydrocephalus (PMID 27198914) and Brain stiffness following recovery in a patient with an episode of low-pressure hydrocephalus (PMID 33030603). The only direct material-property data: very low shear stiffness during the episode, recovered two years later.

Barami, Sood 2016 — The cerebral venous system and the postural regulation of intracranial pressure: implications in the management of patients with cerebrospinal fluid diversion (Childs Nerv Syst; PMID 27581321). Venous overdrainage model for low-pressure states after CSF diversion.

Pandey, Ali, Vasenina & Prasad 2017 — Management of negative pressure hydrocephalus (Neurol India; PMID 28017745). ICP -3 mm Hg reversed with Trendelenburg, cervical wrapping, hypoventilation (raising PaCO₂) and lowered plasma osmolarity — the most explicit "raise ICP medically" protocol.

Smalley, Spiro & Dias 2017 — Low-pressure hydrocephalus in children: a case series and review of the literature (Neurosurgery; PMID 28362957). 30 episodes in 29 children; 37.9 % post-LP; post-LP cases: conservative 27 %, blood patch 18 %, drainage in the rest; median 4 vs 11 days of drainage for post-LP vs spontaneous.

Cheng et al. 2017 — Clinical manifestations and treatment of low-pressure hydrocephalus with ventriculoperitoneal shunt (Chinese; PMID 28264586). 7 adults with shunts at lowest setting still enlarging; intermittent manual compression of the reservoir several times a day improved 6/7 — the origin of the "reservoir pumping" strategy later used by Kuppler 2025.

Marianayagam et al. 2017 — Trapped fourth ventricle and LPH (PMID 28209310) and **Badran et al. 2018** — improvised ultra-low-pressure valveless shunt after SAH in haemangioblastomatosis (PMID 30287249): hardware improvisation case reports.

Strand, Balise, Leung & Boop 2018 — Low-pressure hydrocephalus: a case report and review of the literature (World Neurosurg; PMID 28962963). 25-year search, 17 articles, 90 cases; argues MRE may become the objective diagnostic tool and ETV a strong alternative/adjunct.

Wu, Wang et al. 2019 — Secondary low- or negative-pressure hydrocephalus: clinical features, treatment and a new classification of hydrocephalus based on pressure (Chinese J Neurosurg / Neurosurg Rev; PMID 30611954). 39 adults, 18 with CNS infection, ~3.5 operations each, mean GOS-E 2.7, 8 deaths; proposes high/normal/low/negative pressure classification (low < 70 mm H₂O, negative < 0).

Diaz-Romero Paz et al. 2019 — Negative-pressure hydrocephalus: a therapeutic challenge (PMID 30710718). ICP -4 cm H₂O after external drainage of a cyst; removal of the sink plus sub-zero EVD; transmantle-gradient discussion.

Funnell et al. 2020 — The ultra-low-pressure hydrocephalic state in NPH: improvement after shunt siphoning (Acta Neurochir; PMID 32989519). Shunted iNPH patients who relapsed with large ventricles at very low ICP and improved when the valve was reset to allow siphoning; defines the NPH overlap.

Vilela et al. 2020 (FT) — Low-pressure hydrocephalus after lumbar puncture in an adult (PMID 32712403). Enlarged perimedullary cisterns and brain-stem distortion; epidural blood patch curative; venous overdrainage discussion.

Keough, Isaacs, Urbaneja, Hamilton et al. 2021 — Acute low-pressure hydrocephalus: a case series and systematic review of 195 patients (J Neurosurg; PMID 32736355). 42 local + 153 literature; 53 children, 142 adults; decreased LOC predominant; neoplasm and haemorrhage the commonest etiologies; 39 % of children post-LP; shunt in situ 92 % children / 39 % adults; sub-atmospheric drainage the main temporising step; shunt or ETV definitive; mortality 11 %; return to baseline 83 % children / 49 % adults.

Weisbrod et al. 2021 (FT) — Angiogram-negative SAH and low-pressure hydrocephalus after COVID-19 (PMID 34462697). Single case; adds a systemic-inflammatory trigger.

Casado Pellejero et al. 2022 — Acute negative-pressure hydrocephalus: management algorithm and value of early endoscopic third ventriculostomy (World Neurosurg; PMID 34998487). 5 patients; algorithm with early ETV applied in 4; ETV alone succeeded in 2 (50 %), 2 needed shunts; 1 death from systemic infection; 80 % good outcome.

Czorlich et al. 2023 — Acute low-pressure hydrocephalus in aneurysmal SAH (PMID 37004138). 15/406 (3.7 %); onset 13.1 ± 7.7 days; IVH presence/extent associated; aLPH patients went on to chronic hydrocephalus and shunting far more often than other SAH patients.

Godoy Hurtado et al. 2023 (FT) — Low- and negative-pressure hydrocephalus: new report of six cases and literature review (J Clin Med; PMID 37373809; PMC10299038). Six cases (medulloblastoma ×2, TBI with bifrontal craniectomy, craniopharyngioma, leptomenigeal

glioneuronal tumour, shunted NPH); ~200 cases in literature; symptoms follow ventricular size not pressure; sub-zero drainage most used; neck wrapping, ETV, blood patch as alternatives; pathophysiology "not clear" — permeability/viscoelasticity changes plus craniospinal SAS imbalance.

Zheng et al. 2023 — Hydrocephalus in severe disorders of consciousness (PMID 36994490). 68 DOC patients with ventriculomegaly; 60.3 % (41/68) had LPH/NegPH; 45.5 % had one-stage VP shunt + cranioplasty; individualised low valve pressure by ventricular size and consciousness scores.

Duan & Hu 2024 — Low-pressure hydrocephalus: a narrative review (PMID 38579413). Proposes criteria (Evans > 0.3, ICP ≤ 5 cm H₂O, response to negative-pressure drainage, exclusion of neurodegeneration); risk factors include prior radiotherapy, infection, haemorrhage, multiple surgeries; treatment ladder EVD → ETV/valveless shunt.

Li et al. 2024 — Low/negative-pressure hydrocephalus: a review (PMID 38688300). Four mechanisms (transcortical pressure difference, venous overdrainage, viscoelastic change, porous elastic sponge); EVD/lumbar-cisternal drainage first, ETV or shunt definitive.

Wang et al. 2024 — Quantitative EEG as early warning in post-SAH acute low-pressure hydrocephalus (PMID 39740216). qEEG slowing preceded clinical deterioration and reversed with CSF removal.

Kuppler et al. 2025 (FT) — Chronic negative-pressure hydrocephalus after failure of prolonged drainage (PMID 40030163; PMC11877371). Refractory to low-gradient shunt, neck wrapping, ETV and prolonged sub-atmospheric EVD; scheduled intermittent active reservoir pressing enabled outpatient management; defines cNegPH.

Salam et al. 2025 (FT) — Theophylline for refractory acute negative-pressure hydrocephalus (PMID 40418891; PMC12105587). TBI, hemicraniectomy, shunt infection, IVH; ICP −5 to +10 mm Hg (nadir −21) despite ~4 months of sub-zero EVD (to −30 cm H₂O), two ETVs and ventriculopleural/ventriculoatrial shunts; two doses of theophylline 200 mg via PEG followed by spontaneous drainage, slit-like ventricles on CT at 4 h and sustained improvement; status epilepticus as complication. Single case.

Chen, Li et al. 2025 (FT) — Low- and negative-pressure hydrocephalus in children: a systematic analysis of 44 literature cases and 4 institutional cases (PMID 40538927; PMC12176887). Craniotomy (mostly posterior-fossa tumour) the leading antecedent; > 50 % prior surgery; fatigue, vomiting, headache, bradycardia, seizures, failure to thrive; mean external drainage 41.9 days (2–365); conservative measures (posture, reservoir pumping, valve adjustment) help a minority; 90.9 % need shunt repositioning; 77.5 % recover to baseline; poor outcomes track underlying disease.

Datta & Tu 2025 — Pediatric low-pressure hydrocephalus: a case series and re-analysis of a rare disease (PMID 41191101). 6 children, ICP ≤ 5 cm H₂O; low-resistance VP shunts (3) or negative-pressure cavities (3); prior jugular/sigmoid sinus thrombosis in 2/6; venous-thrombosis model.

Villalonga et al. 2025 — ETV plus valveless VP shunt hybrid for LPH (PMID 41165379). 16 adults meeting Pang–Altschuler criteria; mean KPS gain 56; 81 % excellent, 19 % temporary improvement; 19 % needed a valved shunt.

Carrera et al. 2025 — caudal (infraclavicular) programmable valve placement to exploit the hydrostatic column, 3 patients after prolonged sub-zero drainage (PMID 39951714).

Chinese Neuroscience Society / Neurosurgical Branch 2025 — Expert consensus on diagnosis and treatment of low-pressure hydrocephalus (PMID 40734409). Definition (< 70 mm H₂O, progressive ventriculomegaly, worsening deficits), imaging criteria (effaced Sylvian/convexity SAS, fourth-ventricle "ball sign"), candidate mechanisms (fourth-ventricular outlet obstruction, compliance change), management recommendations.

Bakmeedeniya et al. 2026 — Management of low/negative pressure hydrocephalus: a systematic review of etiology, pathophysiology and management (PMID 41525875). 30 studies, 215 patients (1980–2024); four interacting mechanistic models; management centred on sub-atmospheric EVD, low-pressure/valveless shunts, ETV; proposes a standardised diagnostic and management framework; prognosis governed by etiology and timely recognition.

Bateman & Bateman 2026 (FT) — Pathophysiology of sinking flap syndrome associated with low-pressure hydrocephalus (PMID 42355921). Phase-contrast MRI: bilaterally reduced CBF rising 26 % after cranioplasty; reversed aqueductal flow toward ventricles during LPH; lumped-parameter model favours autoregulatory reset over venous compression; hypothesises capillary CSF absorption at low ICP.

Bruck et al. 2026 (FT) — Neck wrapping for paediatric low-pressure hydrocephalus (PMID 42732439). Two children; third ventricle 13 → 8 mm and 13 → 4 mm; both internalised afterwards.

Hoffman et al. 2026 (FT) — Post-LP low-pressure hydrocephalus in a shunted adult (PMID 42224725). Diagnostic sequence: shunt-tap manometry, patency confirmation, imaging for sink and block; treated with drainage and blood patch.

C. Mechanistic and modelling anchors from the supporting axes

- Sklar 1980 brain elasticity changes with ventriculomegaly (PMID 6893601); Shapiro 1987 no transmante gradient in cats (PMID 3598675); Conner 1984 gradient in experimental NPH (PMID 6737056); Stephensen 2002 no gradient in communicating hydrocephalus (PMID 11904027); Eide & Saehle 2010 pulsatile gradient (PMID 20130957).
- Czosnyka 1996/2001 pressure-volume compensation (PMIDs 8778261, 11235954); Kim 2009 compensatory reserve (PMID 19240611); Gholampour 2022 compliance after shunting (PMID 35979170).
- Peña 1999, Dutta-Roy 2008, Chou 2016, Kim 2016 poroelastic FE models (PMIDs 10414573, 18534602, 26671218, 26274984); Solamen 2021 poroelastic MRE in NPH (PMID 32331966); Duy 2024 brain-CSF interface instability (PMID 38798141).
- Bateman 2002/2008 pulse-wave encephalopathy (PMIDs 12221445, 18379767); Egnor 2023/2026 windkessel impedance classification (PMIDs 37382303, 41791099); Nozaleda 2025 pulsation-driven glymphatic flow (PMID 40258946).
- Hatt 2015 jugular compression stiffens brain on MRE (PMID 26045579); Sainz 2019 venous hypertension and turgor (PMID 30474714); Shulyakov 2011/2012 and Jugé 2016 animal viscoelasticity (PMIDs 21316637, 22806081, 26848844); Wagshul 2021 reduced white-matter stiffness in early hydrocephalus (PMID 33631603); Dirrichs 2019 infant shear-wave elastography (PMID 31464808); Rekate 2020 infant compliance (PMID 32488353).
- Raimondi 1994 unifying theory (PMID 8194058); Oi 2011 multi-categorical classification (PMID 21928018); Orešković & Klarica 2011 bulk-flow critique (PMID 21641963); Bergsneider, Egnor et al. 2006 What we don't (but should) know about hydrocephalus (PMID 16572631).

Part III — Paper listings by axis (computed from `index.tsv`)

✓FT = open-access full text mirrored under `fulltext/`. Evidence class in brackets; `cited` = Europe PMC citation count at harvest.

Core entity — low-/negative-pressure hydrocephalus

79 records

- **1974** McCullough DC et al. — Negative intracranial pressure hydrocephalus in adults with shunts and its relationship to the production of subdural hematoma. *Journal of neurosurgery*. [clinical / mechanistic study (adult), cited 77] [PMID 4813717](#)

- **1986** McCullough DC. — Symptomatic progressive ventriculomegaly in hydrocephalics with patent shunts and antisiphon devices. *Neurosurgery*. [case series (mixed), cited 37] [PMID 3491341](#)
- **1988** Foltz EL et al. — Symptomatic low intracranial pressure in shunted hydrocephalus. *Journal of neurosurgery*. [clinical / mechanistic study, cited 54] [PMID 3343612](#)
- **1993** Drake JM et al. — Functional obstruction of an antisiphon device by raised tissue capsule pressure. *Neurosurgery*. [case report (pediatric), cited 22] [PMID 8421544](#)
- **1993** Pomeranz S et al. — The effect of intracranial hypotension on cerebral blood flow in a feline model. *Acta neurochirurgica*. [experimental (animal / bench), cited 5] [PMID 8333302](#)
- **1994** Foltz EL et al. — Hydrocephalus: the zero ICP ventricle shunt (ZIPS) to control gravity shunt flow. A clinical study in 56 patients. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case series / cohort, cited 4] [PMID 8194062](#)
- **1994** Pang D et al. — Low-pressure hydrocephalic state and viscoelastic alterations in the brain. *Neurosurgery*. [case series / cohort, cited 96] [PMID 7808607](#)
- **1995** Vassilyadi M et al. — Negative-pressure hydrocephalus. *Journal of neurosurgery*. [case report (pediatric), cited 32] [PMID 7666227](#)
- **1999** Bergsneider M et al. — Beneficial effect of siphoning in treatment of adult hydrocephalus. *Archives of neurology*. [case series / cohort (adult), cited 33] [PMID 10520938](#)
- **1999** Dias MS et al. — Low-pressure shunt 'malfunction' following lumbar puncture in children with shunted obstructive hydrocephalus. *Pediatric neurosurgery*. [case series (pediatric), cited 26] [PMID 10352418](#)
- **2001** Owler BK et al. — Low pressure hydrocephalus: issues of diagnosis and treatment in five cases. *British journal of neurosurgery*. [case series / cohort, cited 26] [PMID 11599454](#)
- **2002** Daniel RT et al. — Low-pressure hydrocephalic state complicating hemispherectomy: a case report. *Epilepsia*. [case report (adult), cited 15] [PMID 12027920](#)
- **2002** Kuba H et al. — Gait disturbance in patients with low pressure hydrocephalus. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. [case series / cohort, cited 15] [PMID 11749014](#)
- **2002** Lesniak MS et al. — Low pressure hydrocephalus and ventriculomegaly: hysteresis, non-linear dynamics, and the benefits of CSF diversion. *British journal of neurosurgery*. [case series / cohort (adult), cited 45] [PMID 12617236](#)
- **2004** Gnanalingham KK et al. — 14 mmHg: a case of raised intracranial pressure?. *British journal of neurosurgery*. [case report (adult), cited 1] [PMID 15327239](#)
- **2006** Bergsneider M et al. — What we don't (but should) know about hydrocephalus. *Journal of neurosurgery*. [narrative review (mixed), cited 31] [PMID 16572631](#)
- **2006** Clarke MJ et al. — Very low pressure hydrocephalus. Report of two cases. *Journal of neurosurgery*. [case report, cited 17] [PMID 16961147](#)
- **2008** Rekate HL et al. — The importance of the cortical subarachnoid space in understanding hydrocephalus. *Journal of neurosurgery. Pediatrics*. [case report, cited 75] [PMID 18590387](#)
- **2008** Solheim O et al. — Rapid and severe rise in static and pulsatile intracranial pressures during a generalized epileptic seizure. *Seizure*. [case report, cited 20] [PMID 18554932](#)
- **2009** Garg A et al. — Endoscopic third ventriculostomy: 5 years' experience at the All India Institute of Medical Sciences. *Pediatric neurosurgery*. [case series / cohort (pediatric), cited 16] [PMID 19221456](#)
- **2011** Akins PT et al. — The genesis of low pressure hydrocephalus. *Neurocritical care*. [case report, cited 37] [PMID 21523524](#)
- **2011** Filippidis AS et al. — Negative-pressure and low-pressure hydrocephalus: the role of cerebrospinal fluid leaks resulting from surgical approaches to the cranial base. *Journal of neurosurgery*. [case report, cited 39] [PMID 21800965](#)
- **2012** Hamilton MG et al. — Syndrome of inappropriately low-pressure acute hydrocephalus (SILPAH). *Acta neurochirurgica. Supplement*. [case series / cohort (mixed), cited 25] [PMID 22116443](#)
- **2013** Galbarriatu L et al. — Low-pressure hydrocephalus: indication for custom-made catheters? Technical report. *Acta neurochirurgica*. [case series / cohort (pediatric), cited 8] [PMID 23904087](#)

- **2013** Kalani MY et al. — Treatment of refractory low-pressure hydrocephalus with an active pumping negative-pressure shunt system. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. [case report, cited 21] [PMID 23380444](#)
- **2013** McGovern RA et al. — Low pressure hydrocephalus acutely following sepsis and cardiovascular collapse. *Clinical neurology and neurosurgery*. [case report, cited 4] [PMID 23816322](#)
- **2013** Preuß M et al. — Acute two-compartment low pressure hydrocephalus--a case report. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case report (pediatric), cited 6] [PMID 23715811](#)
- **2013** Trinh VT et al. — Revision to an adjustable non-siphon control valve in low pressure hydrocephalus: therapeutic siphoning and a new perspective on NPH: series of 3 cases and review of the literature. *Clinical neurology and neurosurgery*. [case series, cited 7] [PMID 22770726](#)
- **2014** Hunn BH et al. — Successful treatment of negative pressure hydrocephalus using timely titrated external ventricular drainage: a case series. *Clinical neurology and neurosurgery*. [case series / cohort, cited 12] [PMID 24275338](#)
- **2016** Foster KA et al. — Endoscopic third ventriculostomy as adjunctive therapy in the treatment of low-pressure hydrocephalus in adults. *Surgical neurology international*. [case series / cohort (adult), cited 9] [PMID 27069743](#) ✓FT
- **2016** Olivero WC et al. — Magnetic Resonance Elastography Demonstrating Low Brain Stiffness in a Patient with Low-Pressure Hydrocephalus: Case Report. *Pediatric neurosurgery*. [case report (adult), cited 17] [PMID 27198914](#)
- **2017** Barami K. — Letter: Low-Pressure Hydrocephalus in Children: A Case Series and Review of the Literature. *Neurosurgery*. [comment / letter (pediatric), cited 2] [PMID 28973300](#)
- **2017** Cheng Z et al. — Low pressure hydrocephalus: clinical manifestations, radiological characteristics, and treatment. *British journal of neurosurgery*. [case series / cohort, cited 9] [PMID 28264586](#)
- **2017** Marianayagam NJ et al. — Paradoxical ventriculomegaly due to low-pressure hydrocephalus, a rare complication of the treatment of a trapped fourth ventricle: Case report. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. [case report, cited 4] [PMID 28209310](#)
- **2017** Michael AP et al. — Pearls & Oy-sters: Low-pressure hydrocephalus and inadequate shunting. *Neurology*. [case report, cited 6] [PMID 28438849](#)
- **2017** Pandey S et al. — Negative-Pressure Hydrocephalus: A Case Report on Successful Treatment Under Intracranial Pressure Monitoring with Bilateral Ventriculoperitoneal Shunts. *World neurosurgery*. [case report, cited 8] [PMID 28017745](#)
- **2017** Smalley ZS et al. — Low-pressure Hydrocephalus in Children: A Case Series and Review of the Literature. *Neurosurgery*. [narrative review (pediatric), cited 18] [PMID 28362957](#)
- **2017** Smalley ZS et al. — In Reply: Low-Pressure Hydrocephalus in Children: A Case Series and Review of the Literature. *Neurosurgery*. [comment / letter (pediatric), cited 2] [PMID 28973519](#)
- **2018** Badran A et al. — Hemangioblastomatosis-associated negative-pressure hydrocephalus managed with improvised shunt. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. [case report, cited 5] [PMID 30287249](#)
- **2018** Houlden D et al. — Importance of Frontal Horn Ratio and Optimal CSF Drainage in the Treatment of Very Low-Pressure Hydrocephalus. *Journal of neurological surgery. Part B, Skull base*. [clinical / mechanistic study, cited 2] [PMID 30009119](#)
- **2018** Strand A et al. — Low-Pressure Hydrocephalus: A Case Report and Review of the Literature. *World neurosurgery*. [case series, cited 8] [PMID 28962963](#)
- **2019** Diaz-Romero Paz R et al. — A Rare Case of Negative-Pressure Hydrocephalus: A Plausible Explanation and the Role of Transmantle Theory. *World neurosurgery*. [case report, cited 11] [PMID 30710718](#)
- **2019** Rekate HL. — Low or Negative Pressure Hydrocephalus Demystified. *World neurosurgery*. [comment / letter, cited 6] [PMID 31103762](#)
- **2019** Wu X et al. — Diagnosis and Management for Secondary Low- or Negative-Pressure Hydrocephalus and a New Hydrocephalus Classification Based on Ventricular Pressure. *World neurosurgery*. [case series / cohort (adult), cited 14] [PMID 30611954](#)

- **2020** Funnell JP et al. — Ultra-low-pressure hydrocephalic state in NPH: benefits of therapeutic siphoning with adjustable antigravity valves. *Acta neurochirurgica*. [case series / cohort (adult), cited 6] [PMID 32989519](#)
- **2020** Vilela MD et al. — Low-Pressure Hydrocephalus and Shunt Malfunction Following a Lumbar Puncture in an Adult Reversed by an Epidural Blood Patch. *World neurosurgery*. [case report (mixed), cited 6] [PMID 32712403](#) ✓FT
- **2020** Wang J et al. — A Mathematical Model to Simulate Intracranial Pressure and Unified Interpretation of Several Disease Entities. *bioRxiv*. [computational model] [PMID](#)
- **2021** Keough MB et al. — Acute low-pressure hydrocephalus: a case series and systematic review of 195 patients. *Journal of neurosurgery*. [systematic review (mixed), cited 16] [PMID 32736355](#)
- **2021** Olivero WC et al. — Brain stiffness following recovery in a patient with an episode of low-pressure hydrocephalus: case report. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case report (adult), cited 7] [PMID 33030603](#)
- **2021** Rekate HL. — Commentary on the article "Brain stiffness following recovery in a patient with an episode of low-pressure hydrocephalus: case report". *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [comment / letter, cited 2] [PMID 34272587](#)
- **2021** Suzuki T et al. — Negative-pressure hydrocephalus in the course of a complex postoperative intracranial pressure disturbance: illustrative case. *Journal of neurosurgery. Case lessons*. [case report, cited 2] [PMID 35854689](#) ✓FT
- **2021** Weisbrod L et al. — Low-Pressure Hydrocephalus in Spontaneous Angiogram-Negative Subarachnoid Hemorrhage Following COVID-19 Infection. *Cureus*. [case report, cited 3] [PMID 34462697](#) ✓FT
- **2022** Casado Pellejero J et al. — Acute negative-pressure hydrocephalus: Management algorithm and value of early endoscopic ventriculostomy. *Neurocirugia*. [case series / cohort, cited 4] [PMID 34998487](#)
- **2022** Chang SJ et al. — Treatment-responsive Holmes tremor in a child with low-pressure hydrocephalus: video case report and systematic review of the literature. *Journal of neurosurgery. Pediatrics*. [systematic review (pediatric), cited 2] [PMID 35148507](#)
- **2022** Doshi AB et al. — Solving a complex, rare, and challenging scenario in a normal pressure hydrocephalus with recurrent shunt dysfunction and multiple revisions - The opposing role of evolving low-pressure hydrocephalus and idiopathic raised intra-abdominal pressure. *Surgical neurology international*. [case report (adult), cited 2] [PMID 36600762](#) ✓FT
- **2022** Ozeki T et al. — Suspected Low-Pressure Hydrocephalus Caused by Spinal Drainage after Subarachnoid Hemorrhage. *Journal of Nippon Medical School = Nippon Ika Daigaku zasshi*. [case report (adult), cited 3] [PMID 34526456](#)
- **2023** Czorlich P et al. — Acute low-pressure hydrocephalus in aneurysmal subarachnoid hemorrhage. *Neurosurgical focus*. [case series / cohort, cited 11] [PMID 37004138](#)
- **2023** Egnor M et al. — A quantitative model of the cerebral windkessel and its relevance to disorders of intracranial dynamics. *Journal of neurosurgery. Pediatrics*. [experimental (animal / bench), cited 8] [PMID 37382303](#)
- **2023** Godoy Hurtado A et al. — Low- and Negative-Pressure Hydrocephalus: New Report of Six Cases and Literature Review. *Journal of clinical medicine*. [narrative review, cited 7] [PMID 37373809](#) ✓FT
- **2024** Duan S et al. — Pathogenesis and management of low-pressure hydrocephalus: A narrative review. *Journal of the neurological sciences*. [narrative review, cited 4] [PMID 38579413](#)
- **2024** Li G et al. — Low-/Negative-Pressure Hydrocephalus: To Understand the Formation Mechanism from the Perspective of Clinicians. *Journal of neurological surgery. Part A, Central European neurosurgery*. [narrative review, cited 1] [PMID 38688300](#)
- **2024** Stevens B et al. — Profound Pneumocephalus and Low-Pressure Hydrocephalus Triggered by Ventriculoperitoneal Shunt Placement after Resection, Fat Graft Reconstruction, and Radiotherapy for a Malignant Skull Base Schwannoma. *Journal of neurological surgery reports*. [case report (adult), cited 1] [PMID 39220674](#) ✓FT

- **2024** Wang J et al. — Continuous quantitative electroencephalography for early detection of acute low-pressure obstructive hydrocephalus in aneurysmal subarachnoid hemorrhage: illustrative case. *Journal of neurosurgery. Case lessons*. [case report] PMID 39740216 ✓FT
- **2025** Carrera DA et al. — Treatment of low-pressure hydrocephalus using a novel shunt valve technique: applying Bernoulli's equation to fluid statics. *Technical note. Journal of neurosurgery*. [case series] PMID 39951714
- **2025** Chen B et al. — Low- and negative-pressure hydrocephalus in children, clinical features, treatment, prognosis and proposed mechanisms. *Frontiers in pediatrics*. [case series / cohort (pediatric)] PMID 40538927 ✓FT
- **2025** Chinese Neuroscience Society. — [Consensus on the diagnosis and treatment of low-pressure hydrocephalus(2025 edition)]. *Zhonghua wai ke za zhi [Chinese journal of surgery]*. [consensus / guideline, cited 1] PMID 40734409
- **2025** Datta D et al. — Pediatric low-pressure hydrocephalus: a case series and re-analysis of a rare disease. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case series / cohort (pediatric)] PMID 41191101
- **2025** Koenig N et al. — Headache and Neurological Symptoms Following Lumbar Puncture in a Patient With a Ventriculoperitoneal Shunt. *Cureus*. [case report] PMID 41357016 ✓FT
- **2025** Kuppler P et al. — Chronic negative-pressure hydrocephalus: management of prolonged subatmospheric drainage failure and treatment approach in persistent ventriculomegaly. Illustrative case. *Journal of neurosurgery. Case lessons*. [case report, cited 1] PMID 40030163 ✓FT
- **2025** Lin Z et al. — Correct understanding of the definition and management strategies for refractory hydrocephalus. *Chinese neurosurgical journal*. [narrative review] PMID 40826122 ✓FT
- **2025** Salam S et al. — Theophylline can resolve refractory acute negative-pressure hydrocephalus: illustrative case. *Journal of neurosurgery. Case lessons*. [case report (adult)] PMID 40418891 ✓FT
- **2025** Villalonga JF et al. — Endoscopic Third Ventriculostomy plus Valveless Catheter in the Treatment of Low-Pressure Hydrocephalus. *Turkish neurosurgery*. [case series / cohort] PMID 41165379
- **2026** Bakmeedeniya RW et al. — Management of Low/Negative Pressure Hydrocephalus: A Systematic Review of Etiology, Pathophysiology, and Treatment. *World neurosurgery*. [systematic review, cited 1] PMID 41525875
- **2026** Bateman GA et al. — The Pathophysiology of Sinking Flap Syndrome Associated with Low-Pressure Hydrocephalus: A Case Study Suggests a New Hypothesis. *Journal of clinical medicine*. [case report] PMID 42355921
- **2026** Bateman GA et al. — A Case of Sinking Flap Syndrome Associated with Low-Pressure Hydrocephalus: Clues to the Pathophysiology Suggests a New Hypothesis. *Preprints.org*. [case report] PMID
- **2026** Bruk Y et al. — Jugular Venous Compression via Soft Cervical Collar in Shunt-Refractory Low-Pressure Hydrocephalus: A Report of Two Cases. *Cureus*. [case report] PMID 42732439 ✓FT
- **2026** Egnor M et al. — A model of hydrocephalus due to windkessel impairment caused by subarachnoid space obstruction. *Journal of neurosurgery. Pediatrics*. [experimental (animal / bench), cited 1] PMID 41791099
- **2026** He S et al. — Case Report: Suspected low-pressure hydrocephalus following surgical treatment of traumatic brain injury in a patient with subsequently diagnosed glioblastoma. *Frontiers in oncology*. [case report (adult)] PMID 42444806 ✓FT
- **2026** Hoffman BL et al. — Lumbar puncture-induced negative-pressure hydrocephalus in a child with a ventriculoperitoneal shunt and ventricular obstruction: illustrative case. *Journal of neurosurgery. Case lessons*. [case report (pediatric)] PMID 42224725 ✓FT

Mechanism — brain biomechanics, compliance, elastography, modelling

50 records

- **1980** Sklar FH et al. — Brain elasticity changes with ventriculomegaly. *Journal of neurosurgery*. [case series / cohort, cited 33] PMID 6893601

- **1980** Welch K. — The intracranial pressure in infants. *Journal of neurosurgery*. [clinical / mechanistic study (pediatric), cited 100] [PMID 7373397](#)
- **1984** Conner ES et al. — Experimental normal-pressure hydrocephalus is accompanied by increased transmantle pressure. *Journal of neurosurgery*. [experimental (animal / bench), cited 79] [PMID 6737056](#)
- **1987** Shapiro K et al. — Progressive ventricular enlargement in cats in the absence of transmantle pressure gradients. *Journal of neurosurgery*. [experimental (animal / bench), cited 34] [PMID 3598675](#)
- **1994** Raimondi AJ. — A unifying theory for the definition and classification of hydrocephalus. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [narrative review (pediatric), cited 50] [PMID 8194058](#)
- **1994** Rekate HL. — The usefulness of mathematical modeling in hydrocephalus research. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case report (pediatric), cited 17] [PMID 8194056](#)
- **1996** Czosnyka M et al. — Testing of cerebrospinal compensatory reserve in shunted and non-shunted patients: a guide to interpretation based on an observational study. *Journal of neurology, neurosurgery, and psychiatry*. [case series / cohort (adult), cited 73] [PMID 8778261](#)
- **1999** Peña A et al. — Effects of brain ventricular shape on periventricular biomechanics: a finite-element analysis. *Neurosurgery*. [computational model, cited 55] [PMID 10414573](#)
- **2001** Czosnyka M et al. — Age dependence of cerebrospinal pressure-volume compensation in patients with hydrocephalus. *Journal of neurosurgery*. [case series / cohort, cited 68] [PMID 11235954](#)
- **2001** Fukuhara T et al. — Effects of ventriculoperitoneal shunt removal on cerebral oxygenation and brain compliance in chronic obstructive hydrocephalus. *Journal of neurosurgery*. [experimental (animal / bench), cited 27] [PMID 11302655](#)
- **2002** Bateman GA. — Pulse-wave encephalopathy: a comparative study of the hydrodynamics of leukoaraiosis and normal-pressure hydrocephalus. *Neuroradiology*. [case series / cohort, cited 101] [PMID 12221445](#)
- **2002** Stephensen H et al. — There is no transmantle pressure gradient in communicating or noncommunicating hydrocephalus. *Neurosurgery*. [clinical / mechanistic study (adult), cited 84] [PMID 11904027](#)
- **2004** Taylor Z et al. — Reassessment of brain elasticity for analysis of biomechanisms of hydrocephalus. *Journal of biomechanics*. [computational model, cited 136] [PMID 15212932](#)
- **2007** Bateman GA et al. — Can MR measurement of intracranial hydrodynamics and compliance differentiate which patient with idiopathic normal pressure hydrocephalus will improve following shunt insertion?. *Acta neurochirurgica*. [case series / cohort, cited 37] [PMID 17406777](#)
- **2008** Bateman GA et al. — The venous manifestations of pulse wave encephalopathy: windkessel dysfunction in normal aging and senile dementia. *Neuroradiology*. [computational model, cited 96] [PMID 18379767](#)
- **2008** Dutta-Roy T et al. — Biomechanical modelling of normal pressure hydrocephalus. *Journal of biomechanics*. [computational model, cited 38] [PMID 18534602](#)
- **2008** Li J et al. — Communicating hydrocephalus in adult rats with kaolin obstruction of the basal cisterns or the cortical subarachnoid space. *Experimental neurology*. [experimental (animal / bench) (mixed), cited 38] [PMID 18433747](#)
- **2009** Kim DJ et al. — Index of cerebrospinal compensatory reserve in hydrocephalus. *Neurosurgery*. [case series / cohort (adult), cited 58] [PMID 19240611](#)
- **2010** Eide PK et al. — Is ventriculomegaly in idiopathic normal pressure hydrocephalus associated with a transmantle gradient in pulsatile intracranial pressure?. *Acta neurochirurgica*. [case series / cohort, cited 43] [PMID 20130957](#)
- **2010** Pattison AJ et al. — Time-harmonic magnetic resonance elastography of the normal feline brain. *Journal of biomechanics*. [experimental (animal / bench), cited 25] [PMID 20655045](#)
- **2011** Oi S. — Classification of hydrocephalus: critical analysis of classification categories and advantages of "Multi-categorical Hydrocephalus Classification" (Mc HC). *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case report, cited 37] [PMID 21928018](#)

- **2011** Orešković D et al. — Development of hydrocephalus and classical hypothesis of cerebrospinal fluid hydrodynamics: facts and illusions. *Progress in neurobiology*. [narrative review, cited 104] [PMID 21641963](#)
- **2011** Shulyakov AV et al. — Age-dependence of intracranial viscoelastic properties in living rats. *Journal of the mechanical behavior of biomedical materials*. [experimental (animal / bench), cited 33] [PMID 21316637](#)
- **2011** Streitberger KJ et al. — In vivo viscoelastic properties of the brain in normal pressure hydrocephalus. *NMR in biomedicine*. [case series / cohort (adult), cited 132] [PMID 20931563](#)
- **2012** Freimann FB et al. — Alteration of brain viscoelasticity after shunt treatment in normal pressure hydrocephalus. *Neuroradiology*. [case series / cohort, cited 90] [PMID 21538046](#)
- **2012** Shulyakov AV et al. — Intracranial biomechanics of acute experimental hydrocephalus in live rats. *Neurosurgery*. [experimental (animal / bench), cited 19] [PMID 22806081](#)
- **2015** Hatt A et al. — MR Elastography Can Be Used to Measure Brain Stiffness Changes as a Result of Altered Cranial Venous Drainage During Jugular Compression. *AJNR. American journal of neuroradiology*. [imaging / physiology study, cited 43] [PMID 26045579](#)
- **2016** Barami K. — Cerebral venous overdrainage: an under-recognized complication of cerebrospinal fluid diversion. *Neurosurgical focus*. [narrative review (adult), cited 30] [PMID 27581321](#)
- **2016** Chou D et al. — A fully dynamic multi-compartmental poroelastic system: Application to aqueductal stenosis. *Journal of biomechanics*. [computational model, cited 18] [PMID 26671218](#)
- **2016** Fattahi N et al. — MR Elastography Demonstrates Increased Brain Stiffness in Normal Pressure Hydrocephalus. *AJNR. American journal of neuroradiology*. [case series / cohort, cited 71] [PMID 26542235](#)
- **2016** Jugé L et al. — Changes in Rat Brain Tissue Microstructure and Stiffness during the Development of Experimental Obstructive Hydrocephalus. *PloS one*. [experimental (animal / bench), cited 29] [PMID 26848844](#) ✓FT
- **2016** Kim H et al. — Finite element analysis of periventricular lucency in hydrocephalus: extravasation or transependymal CSF absorption?. *Journal of neurosurgery*. [computational model, cited 20] [PMID 26274984](#)
- **2017** ElSheikh M et al. — MR Elastography Demonstrates Unique Regional Brain Stiffness Patterns in Dementias. *AJR. American journal of roentgenology*. [case series / cohort, cited 75] [PMID 28570101](#)
- **2017** Orešković D et al. — Role of choroid plexus in cerebrospinal fluid hydrodynamics. *Neuroscience*. [narrative review, cited 100] [PMID 28456718](#)
- **2017** Perry A et al. — Clinical Correlation of Abnormal Findings on Magnetic Resonance Elastography in Idiopathic Normal Pressure Hydrocephalus. *World neurosurgery*. [imaging / physiology study, cited 44] [PMID 28063896](#)
- **2019** Dirrichs T et al. — Transcranial Shear Wave Elastography of Neonatal and Infant Brains for Quantitative Evaluation of Increased Intracranial Pressure. *Investigative radiology*. [case series / cohort (pediatric), cited 23] [PMID 31464808](#)
- **2019** Sainz LV et al. — Cerebro-venous hypertension: a frequent cause of so-called "external hydrocephalus" in infants. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case series / cohort (pediatric), cited 15] [PMID 30474714](#)
- **2020** Murphy MC et al. — Identification of Normal Pressure Hydrocephalus by Disease-Specific Patterns of Brain Stiffness and Damping Ratio. *Investigative radiology*. [computational model, cited 39] [PMID 32058331](#)
- **2020** ReKate HL. — Hydrocephalus in infants: the unique biomechanics and why they matter. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [narrative review (pediatric), cited 16] [PMID 32488353](#)
- **2021** Sincomb SJ et al. — Transmantle Pressure Computed from MR Imaging Measurements of Aqueduct Flow and Dimensions. *AJNR. American journal of neuroradiology*. [imaging / physiology study (mixed), cited 8] [PMID 34385144](#)
- **2021** Solamen LM et al. — Poroelastic Mechanical Properties of the Brain Tissue of Normal Pressure Hydrocephalus Patients During Lumbar Drain Treatment Using Intrinsic Actuation MR Elastography. *Academic radiology*. [computational model, cited 22] [PMID 32331966](#)

- **2021** Wagshul ME et al. — MR Elastography demonstrates reduced white matter shear stiffness in early-onset hydrocephalus. *NeuroImage. Clinical.* [imaging / physiology study, cited 17] [PMID 33631603](#) ✓FT
- **2022** Aunan-Diop JS et al. — Magnetic resonance elastography in normal pressure hydrocephalus- a scoping review. *Neurosurgical review.* [narrative review, cited 14] [PMID 34687356](#)
- **2022** Coelho A et al. — Magnetic resonance elastography of the ageing brain in normal and demented populations: A systematic review. *Human brain mapping.* [systematic review, cited 29] [PMID 35488708](#) ✓FT
- **2022** Cogswell PM et al. — Features of Idiopathic Intracranial Hypertension on MRI With MR Elastography: Prospective Comparison With Control Individuals and Assessment of Postintervention Changes. *AJR. American journal of roentgenology.* [case series / cohort (adult), cited 10] [PMID 35822642](#)
- **2022** Gholampour S et al. — A New Definition for Intracranial Compliance to Evaluate Adult Hydrocephalus After Shunting. *Frontiers in bioengineering and biotechnology.* [case series / cohort (adult), cited 20] [PMID 35979170](#) ✓FT
- **2022** Hersh AM et al. — Applications of elastography in operative neurosurgery: A systematic review. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia.* [systematic review (pediatric), cited 10] [PMID 35933785](#)
- **2024** Duy PQ et al. — Biomechanical instability of the brain-CSF interface in hydrocephalus. *Brain : a journal of neurology.* [narrative review (adult), cited 13] [PMID 38798141](#)
- **2024** Karki P et al. — Prediction of Surgical Outcomes in Normal Pressure Hydrocephalus by MR Elastography. *AJNR. American journal of neuroradiology.* [case series / cohort, cited 10] [PMID 38272572](#)
- **2025** Nozaleda GL et al. — Arterial pulsations and transmantle pressure synergetically drive glymphatic flow. *Scientific reports.* [clinical / mechanistic study, cited 14] [PMID 40258946](#) ✓FT

Treatment techniques — EVD titration, valves, drainage strategy

30 records

- **1976** Portnoy HD et al. — Hydrodynamics of shunt valves. *Child's brain.* [clinical / mechanistic study, cited 13] [PMID 991667](#)
- **1998** Rohde V et al. — Four-year experience with the routine use of the programmable Hakim valve in the management of children with hydrocephalus. *Acta neurochirurgica.* [case series / cohort (mixed), cited 45] [PMID 9870057](#)
- **2000** Fukuhara T et al. — Critical shunt-induced subdural hematoma treated with combined pressure-programmable valve implantation and endoscopic third ventriculostomy. *Pediatric neurosurgery.* [case series (pediatric), cited 14] [PMID 11025421](#)
- **2002** Czosnyka Z et al. — Laboratory testing of hydrocephalus shunts -- conclusion of the U.K. Shunt evaluation programme. *Acta neurochirurgica.* [experimental (animal / bench), cited 27] [PMID 12111485](#)
- **2002** Dagnew E et al. — Acute foramen magnum syndrome caused by an acquired Chiari malformation after lumbar drainage of cerebrospinal fluid: report of three cases. *Neurosurgery.* [case report, cited 34] [PMID 12188966](#)
- **2003** Korinth MC et al. — Experience with a new concept to lower non-infectious complications in infants with programmable shunts. *European journal of pediatric surgery : official journal of Austrian Association of Pediatric Surgery ... [et al] = Zeitschrift fur Kinderchirurgie.* [case series / cohort (pediatric), cited 8] [PMID 12776237](#)
- **2004** Klopfenstein JD et al. — Comparison of rapid and gradual weaning from external ventricular drainage in patients with aneurysmal subarachnoid hemorrhage: a prospective randomized trial. *Journal of neurosurgery.* [case series / cohort, cited 78] [PMID 15086228](#)
- **2005** Ringel F et al. — Comparison of programmable shunt valves vs standard valves for communicating hydrocephalus of adults: a retrospective analysis of 407 patients. *Surgical neurology.* [case series / cohort (adult), cited 48] [PMID 15639519](#)

- **2009** Staykov D et al. — Intraventricular fibrinolysis and lumbar drainage for ventricular hemorrhage. *Stroke*. [case series / cohort, cited 66] [PMID 19679848](#)
- **2010** Lemcke J et al. — Improved outcome in shunted iNPH with a combination of a Codman Hakim programmable valve and an Aesculap-Miethke ShuntAssistant. *Central European neurosurgery*. [case series, cited 31] [PMID 20373276](#)
- **2010** Toma AK et al. — Continuous intracranial pressure monitoring in pseudotumour cerebri: Single centre experience. *British journal of neurosurgery*. [case series / cohort, cited 14] [PMID 20632885](#)
- **2010** Toma AK et al. — Investigating shunt function using continuous intracranial pressure monitoring in adults: single center experience. *Journal of neurosurgery*. [case series / cohort (adult), cited 9] [PMID 20799857](#)
- **2011** Speck V et al. — Lumbar catheter for monitoring of intracranial pressure in patients with post-hemorrhagic communicating hydrocephalus. *Neurocritical care*. [case series / cohort, cited 24] [PMID 20953846](#)
- **2012** Lemcke J et al. — On the method of a randomised comparison of programmable valves with and without gravitational units: the SVASONA study. *Acta neurochirurgica. Supplement*. [case series / cohort, cited 9] [PMID 22327702](#)
- **2012** Miyake H et al. — Assessment of a quick reference table algorithm for determining initial postoperative pressure settings of programmable pressure valves in patients with idiopathic normal pressure hydrocephalus: SINPHONI subanalysis. *Neurosurgery*. [case series / cohort, cited 20] [PMID 22653392](#)
- **2013** Delwel EJ et al. — A randomised trial of high and low pressure level settings on an adjustable ventriculoperitoneal shunt valve for idiopathic normal pressure hydrocephalus: results of the Dutch evaluation programme Strata shunt (DEPSS) trial. *Journal of neurology, neurosurgery, and psychiatry*. [case series / cohort, cited 22] [PMID 23408069](#)
- **2014** Chari A et al. — Hydrocephalus shunt technology: 20 years of experience from the Cambridge Shunt Evaluation Laboratory. *Journal of neurosurgery*. [experimental (animal / bench), cited 35] [PMID 24405071](#)
- **2016** Arroyo-Palacios J et al. — Characterization of Shape Differences Among ICP Pulses Predicts Outcome of External Ventricular Drainage Weaning Trial. *Neurocritical care*. [case series / cohort, cited 12] [PMID 27106888](#)
- **2017** Chari A et al. — Intraparenchymal intracranial pressure monitoring for hydrocephalus and cerebrospinal fluid disorders. *Acta neurochirurgica*. [case series / cohort, cited 42] [PMID 28799016](#) ✓FT
- **2018** Jabbarli R et al. — Gradual External Ventricular Drainage Weaning Reduces The Risk of Shunt Dependency After Aneurysmal Subarachnoid Hemorrhage: A Pooled Analysis. *Operative neurosurgery (Hagerstown, Md.)*. [case series / cohort, cited 24] [PMID 29490068](#)
- **2020** Akinduro OO et al. — Predictors of Ventriculoperitoneal shunting following Subarachnoid Hemorrhage treated with External Ventricular Drainage. *Neurocritical care*. [case series / cohort, cited 25] [PMID 31410771](#)
- **2020** Bakhshi SK et al. — Lumbar Drain for Temporary Cerebrospinal Fluid Diversion: Factors Related to the Risks of Complications at a University Hospital. *World neurosurgery*. [case series / cohort (adult), cited 18] [PMID 32711138](#)
- **2020** Capion T et al. — Prompt closure versus gradual weaning of external ventricular drainage for hydrocephalus in adult patients with aneurysmal subarachnoid haemorrhage: a systematic review. *BMJ open*. [systematic review (adult), cited 12] [PMID 33243807](#) ✓FT
- **2020** Gu JC et al. — Intracranial Pressure during External Ventricular Drainage Weaning Is an Outcome Predictor of Traumatic Brain Injury. *BioMed research international*. [case series / cohort, cited 4] [PMID 32695822](#) ✓FT
- **2022** Palasz J et al. — External ventricular drain management in subarachnoid haemorrhage: a systematic review and meta-analysis. *Neurosurgical review*. [systematic review, cited 43] [PMID 34448080](#)
- **2023** Bertuccio A et al. — External Ventricular Drainage: A Practical Guide for Neuro-Anesthesiologists. *Clinics and practice*. [narrative review, cited 31] [PMID 36826162](#) ✓FT

- **2023** Capion T et al. — Prompt closure versus gradual weaning of external ventricular drainage for hydrocephalus following aneurysmal subarachnoid haemorrhage: Protocol for the DRAIN randomised clinical trial. *Acta anaesthesiologica Scandinavica*. [case series / cohort, cited 6] [PMID 37165711](#)
- **2023** Noiphithak R et al. — Prediction of Permanent Shunt Dependency in Patients with Intraventricular Hemorrhage: Outcomes of Early External Ventricular Drainage Weaning Protocol. *World neurosurgery*. [case series / cohort, cited 5] [PMID 37689359](#)
- **2026** Garrido E et al. — Permanent ventricular shunt and risk of meningitis: Challenges in weaning from external ventricular drain. *Neuro-Chirurgie*. [case series / cohort, cited 1] [PMID 41289673](#)
- **2026** Maroufi SF et al. — Utility of intracranial pressure monitoring in idiopathic intracranial hypertension patients with refractory headaches: an 8-year, single-surgeon, specialized-center retrospective study. *Neurosurgical focus*. [case series / cohort (adult), cited 1] [PMID 41569756](#)

Aetiological settings — trauma, craniectomy, SAH, tumour, infection

14 records

- **1982** Walsh JW et al. — Subtemporal craniectomy and elevation of shunt valve opening pressure in the management of small ventricle-induced cerebrospinal fluid shunt dysfunction. *Neurosurgery*. [case series (pediatric), cited 26] [PMID 7110543](#)
- **2008** Ben-Nakhi A et al. — Bilateral subdural hemorrhage following lumbar puncture in a case of suspected meningitis. A case report. *The neuroradiology journal*. [case report, cited 2] [PMID 24256823](#)
- **2011** Joshua SP et al. — Papilloedema as a non-invasive marker for raised intra-cranial pressure following decompressive craniectomy for severe head injury. *Clinical neurology and neurosurgery*. [case series / cohort, cited 6] [PMID 21676532](#)
- **2014** Ding J et al. — The influence of decompressive craniectomy on the development of hydrocephalus: a review. *Arquivos de neuro-psiquiatria*. [narrative review, cited 43] [PMID 25252237](#)
- **2016** Smith GA et al. — Incidence, Management, and Outcome of Symptomatic Postoperative Posterior Fossa Pseudomeningocele: A Retrospective Single-Institution Experience. *Operative neurosurgery (Hagerstown, Md.)*. [case series / cohort, cited 11] [PMID 29506116](#)
- **2016** Wan Y et al. — Methods of intra-operative treatment of cranioplasty in patients with abnormal bone window flap pressure after decompressive craniectomy. *Clinical neurology and neurosurgery*. [case series / cohort, cited 4] [PMID 26971294](#)
- **2018** Kutty RK et al. — The Conundrum of Ventricular Dilatations Following Decompressive Craniectomy: Is Ventriculoperitoneal Shunt, The Only Panacea?. *Journal of neurosciences in rural practice*. [case series / cohort, cited 8] [PMID 29725175](#) ✓FT
- **2018** Udayakumaran S et al. — Tailored Strategies to Manage Cerebrospinal Fluid Leaks or Pseudomeningocele After Surgery for Tethered Cord Syndrome. *World neurosurgery*. [case series / cohort (pediatric), cited 12] [PMID 29605699](#)
- **2022** Bakhsh A et al. — Chronic CSF leak from lumbar-peritoneal shunt tract: A case report. *Surgical neurology international*. [case report (adult), cited 2] [PMID 35673636](#) ✓FT
- **2022** Svedung Wettervik T et al. — Post-traumatic hydrocephalus - incidence, risk factors, treatment, and clinical outcome. *British journal of neurosurgery*. [case series / cohort, cited 20] [PMID 34414834](#)
- **2023** Bhatjiwale MM et al. — Malignant bihemispheric cerebral edema after cranioplasty - An extension of the Monro-Kellie doctrine and predictive factors. *Surgical neurology international*. [case report, cited 3] [PMID 37680924](#) ✓FT
- **2023** Zheng RZ et al. — Clinical Characteristics of and Treatment Strategy for Hydrocephalus in Patients With Severe Disorders of Consciousness. *Journal of neurotrauma*. [case series / cohort, cited 3] [PMID 36994490](#)
- **2025** Kumar S et al. — Extra-Axial Endoscopic Third Ventriculostomy (EAETV) - A Novel Remedy to Contain Persistent Cerebrospinal Fluid Leak After Infratentorial Neurosurgery. *Neurology India*. [case series / cohort] [PMID 41215506](#)

- **2026** Chang KW et al. — Lumbar-ventricular cerebrospinal fluid pressure discrepancy in leptomeningeal metastasis associated intracranial hypertension. *Fluids and barriers of the CNS*. [case series / cohort] [PMID 42218510](#) ✓FT

Adjacent differentials — slit ventricle, overdrainage, intracranial hypotension

30 records

- **1978** Faulhauer K et al. — Overdrainage phenomena in shunt treated hydrocephalus. *Acta neurochirurgica*. [clinical / mechanistic study (pediatric), cited 113] [PMID 742440](#)
- **1984** Gruber R et al. — Experiences with the anti-siphon device (ASD) in shunt therapy of pediatric hydrocephalus. *Journal of neurosurgery*. [case series / cohort (pediatric), cited 62] [PMID 6726390](#)
- **1985** Serlo W et al. — Classification and management of the slit ventricle syndrome. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case series / cohort (pediatric), cited 47] [PMID 4064018](#)
- **1987** Oi S et al. — Infantile hydrocephalus and the slit ventricle syndrome in early infancy. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [case series (pediatric), cited 59] [PMID 3652064](#)
- **1989** Serlo W et al. — The incidence and management of the slit ventricle syndrome. *Acta neurochirurgica*. [case series / cohort (pediatric), cited 43] [PMID 2773680](#)
- **1994** Major O et al. — Slit-ventricle syndrome in shunt operated children. *Acta neurochirurgica*. [case series / cohort (pediatric), cited 32] [PMID 7942186](#)
- **1995** Aschoff A et al. — Overdrainage and shunt technology. A critical comparison of programmable, hydrostatic and variable-resistance valves and flow-reducing devices. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [clinical / mechanistic study (mixed), cited 91] [PMID 7621479](#)
- **1998** Atkinson JL et al. — Acquired Chiari I malformation secondary to spontaneous spinal cerebrospinal fluid leakage and chronic intracranial hypotension syndrome in seven cases. *Journal of neurosurgery*. [case series / cohort, cited 106] [PMID 9452230](#)
- **1998** Baskin JJ et al. — Ventricular shunt removal: the ultimate treatment of the slit ventricle syndrome. *Journal of neurosurgery*. [case series / cohort, cited 94] [PMID 9488301](#)
- **1998** Czosnyka Z et al. — Posture-related overdrainage: comparison of the performance of 10 hydrocephalus shunts in vitro. *Neurosurgery*. [experimental (animal / bench), cited 56] [PMID 9482183](#)
- **2000** Mokri B et al. — Absent headache despite CSF volume depletion (intracranial hypotension). *Neurology*. [case report, cited 51] [PMID 11113229](#)
- **2001** Albright AL et al. — Slit-ventricle syndrome secondary to shunt-induced suture ossification. *Neurosurgery*. [case report (pediatric), cited 45] [PMID 11322436](#)
- **2002** Binder DK et al. — Intrathecal saline infusion in the treatment of obtundation associated with spontaneous intracranial hypotension: technical case report. *Neurosurgery*. [case report (adult), cited 73] [PMID 12188967](#)
- **2003** Samadani U et al. — Intracranial hypotension after intraoperative lumbar cerebrospinal fluid drainage. *Neurosurgery*. [case series, cited 45] [PMID 12493112](#)
- **2005** Chernov MF et al. — Neurofiberscope-guided management of slit-ventricle syndrome due to shunt placement. *Journal of neurosurgery*. [case series / cohort, cited 29] [PMID 15881749](#)
- **2006** Browd SR et al. — Failure of cerebrospinal fluid shunts: part II: overdrainage, loculation, and abdominal complications. *Pediatric neurology*. [narrative review (pediatric), cited 115] [PMID 16504785](#)
- **2006** Oi S et al. — Proposal of "evolution theory in cerebrospinal fluid dynamics" and minor pathway hydrocephalus in developing immature brain. *Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery*. [experimental (animal / bench) (mixed), cited 137] [PMID 16685545](#)

- **2008** Bateman GA. — The pathophysiology of idiopathic normal pressure hydrocephalus: cerebral ischemia or altered venous hemodynamics?. AJNR. American journal of neuroradiology. [case series / cohort, cited 115] [PMID 17925373](#)
- **2008** Clark S et al. — The use of noninvasive electromagnetic neuronavigation for slit ventricle syndrome and complex hydrocephalus in a pediatric population. Journal of neurosurgery. Pediatrics. [case series / cohort (pediatric), cited 36] [PMID 19035692](#)
- **2008** Rekate HL. — Shunt-related headaches: the slit ventricle syndromes. Child's nervous system : ChNS : official journal of the International Society for Pediatric Neurosurgery. [narrative review, cited 75] [PMID 18259760](#)
- **2010** Gruber RW et al. — Prevention of ventricular catheter obstruction and slit ventricle syndrome by the prophylactic use of the Integra antisiphon device in shunt therapy for pediatric hypertensive hydrocephalus: a 25-year follow-up study. Journal of neurosurgery. Pediatrics. [case series / cohort (pediatric), cited 31] [PMID 20043731](#)
- **2011** Staykov D et al. — Early recognition of lumbar overdrainage by lumboventricular pressure gradient. Neurosurgery. [case series, cited 22] [PMID 21273925](#)
- **2012** Eide PK et al. — Morphological characterization of cardiac induced intracranial pressure (ICP) waves in patients with overdrainage of cerebrospinal fluid and negative ICP. Medical engineering & physics. [case series / cohort (mixed), cited 11] [PMID 22153319](#)
- **2012** Freimann FB et al. — Shunting with gravitational valves--can adjustments end the era of revisions for overdrainage-related events?: clinical article. Journal of neurosurgery. [case series / cohort (adult), cited 26] [PMID 22998061](#)
- **2013** Bateman GA. — Hypertensive slit ventricle syndrome: pseudotumor cerebri with a malfunctioning shunt?. Journal of neurosurgery. [case series, cited 20] [PMID 23991845](#)
- **2013** Khan QU et al. — Overdrainage shunt complications in idiopathic normal-pressure hydrocephalus and lumbar puncture opening pressure. Journal of neurosurgery. [case series / cohort, cited 48] [PMID 23930853](#)
- **2017** Chen S et al. — Hydrocephalus after Subarachnoid Hemorrhage: Pathophysiology, Diagnosis, and Treatment. BioMed research international. [narrative review, cited 101] [PMID 28373987](#) ✓FT
- **2020** Desai VR et al. — Ventriculoperitoneal shunt complications in an adult population: A comparison of various shunt designs to prevent overdrainage. Surgical neurology international. [case series / cohort (adult), cited 31] [PMID 33033631](#) ✓FT
- **2021** Kuo LT et al. — The Pathogenesis of Hydrocephalus Following Aneurysmal Subarachnoid Hemorrhage. International journal of molecular sciences. [narrative review, cited 56] [PMID 34068783](#) ✓FT
- **2023** Pedersen SH et al. — How to define CSF overdrainage: a systematic literature review. Acta neurochirurgica. [systematic review (mixed), cited 12] [PMID 36639536](#)

Historical usage — 'low-pressure hydrocephalus' as a synonym for NPH

28 records

- **1969** Greitz TV et al. — Pre- and postoperative evaluation of cerebral blood flow in low-pressure hydrocephalus. Journal of neurosurgery. [clinical / mechanistic study, cited 74] [PMID 5359210](#)
- **1970** Jensen F et al. — Isotope encephalography in low-pressure hydrocephalus. Acta neurologica Scandinavica. [clinical / mechanistic study, cited 4] [PMID 5457870](#)
- **1971** Ingram TT. — 'Low-pressure hydrocephalus'. Developmental medicine and child neurology. [clinical / mechanistic study, cited 4] [PMID 5119928](#)
- **1972** Bannister CM. — A report of eight patients with low pressure hydrocephalus treated by C.S.F. diversion with disappointing results. Acta neurochirurgica. [case series / cohort, cited 22] [PMID 4540544](#)
- **1972** Symon L et al. — Pressure waves in so-called low-pressure hydrocephalus. Lancet (London, England). [clinical / mechanistic study, cited 58] [PMID 4117815](#)
- **1973** Aleksic SN et al. — Dementia and low-pressure hydrocephalus in a patient with pituitary adenoma. Journal of the neurological sciences. [clinical / mechanistic study, cited 5] [PMID 4541480](#)

- **1974** Bozzao L et al. — [Low-pressure hydrocephalus. Diagnostic value of the infusion test]. *Recenti progressi in medicina*. [clinical / mechanistic study] [PMID 4408072](#)
- **1975** Botez MI et al. — Occult disorders of the cerebrospinal fluid dynamics. Early diagnosis criteria. *European neurology*. [case series / cohort, cited 6] [PMID 1149749](#)
- **1975** Ingvar DH et al. — Evidence of subnormal function of association cortex in presenile dementia. *Neurology*. [case series, cited 30] [PMID 1237107](#)
- **1975** Mumenthaler M. — [Cerebral sclerosis. Diagnostic criteria and differential diagnostic consideration in practice]. *Schweizerische medizinische Wochenschrift*. [clinical / mechanistic study (adult), cited 1] [PMID 237320](#)
- **1975** Nehlil J. — [Low-pressure hydrocephalus and psychoses. Apropos of 2 cases]. *Annales medico-psychologiques*. [case series] [PMID 1232893](#)
- **1975** Raimondi AJ et al. — Cystic cavum veli interpositi associated with normal or low pressure hydrocephalus. *Child's brain*. [case series, cited 7] [PMID 1175442](#)
- **1975** Rivero Hernández MS et al. — [Diagnosis of the psycho-organic syndrome in low pressure hydrocephalus using the Wechsler-Bellevue test]. *Actas luso-espanolas de neurologia, psiquiatria y ciencias afines*. [clinical / mechanistic study] [PMID 1211270](#)
- **1976** Bloch R et al. — A mathematical model of cerebrospinal fluid dynamics. *Journal of the neurological sciences*. [computational model, cited 9] [PMID 1262906](#)
- **1976** Herskovits E et al. — [Considerations on low pressure hydrocephalus]. *Arquivos de neuro-psiquiatria*. [case series / cohort] [PMID 985149](#)
- **1976** Mazars G et al. — [Neurosurgical treatment of headaches]. *Minerva medica*. [clinical / mechanistic study, cited 2] [PMID 934549](#)
- **1976** Singounas EG et al. — Observations on the pathogenesis of low pressure hydrocephalus. Analysis of 25 cases. *Neurochirurgia*. [case series / cohort, cited 9] [PMID 1250499](#)
- **1976** Sjaastad O et al. — Increased intracranial pressure in so-called 'normal-pressure hydrocephalus'. *European neurology*. [case series / cohort, cited 3] [PMID 1278191](#)
- **1976** Voigt K et al. — Cerebral blood volume alterations during fractional pneumoencephalography. *AJR. American journal of roentgenology*. [case series / cohort, cited 2] [PMID 178206](#)
- **1977** Cold G et al. — Intraventricular isotope encephalography, continuous intraventricular pressure recording and biochemical changes in the cerebrospinal fluid in low-pressure hydrocephalus. *Danish medical bulletin*. [clinical / mechanistic study, cited 3] [PMID 837736](#)
- **1977** Mata González P et al. — [Low-pressure hydrocephalus. A clinical study of 18 patients]. *Revista clinica espanola*. [case series / cohort] [PMID 605235](#)
- **1978** Kulczycki J et al. — [Diagnostic value of radioisotope cisternography in low-pressure hydrocephalus syndrome]. *Neurologia i neurochirurgia polska*. [case series / cohort] [PMID 683432](#)
- **1978** Martin G. — Lundberg's B waves as a feature of normal intracranial pressure. *Surgical neurology*. [case report, cited 13] [PMID 675492](#)
- **1979** Hash CJ et al. — Ventricle to sagittal sinus shunt for hydrocephalus. *Neurosurgery*. [case series / cohort (adult), cited 11] [PMID 460566](#)
- **1981** Janny P et al. — [Normal pressure hydrocephalus. Pre- and postoperative study of 56 cases (author's transl)]. *Neuro-Chirurgie*. [case series / cohort, cited 10] [PMID 7279095](#)
- **1984** Lindvall M et al. — Sympathetic nervous control of cerebrospinal fluid production in experimental obstructive hydrocephalus. *Experimental neurology*. [experimental (animal / bench), cited 16] [PMID 6723882](#)
- **1987** Mumenthaler M. — [Reversible and preventable dementias]. *Schweizerische medizinische Wochenschrift*. [clinical / mechanistic study, cited 2] [PMID 3616587](#)
- **1989** Meixensberger J et al. — Cerebral blood flow in low pressure hydrocephalus. *Psychiatry research*. [clinical / mechanistic study, cited 5] [PMID 2608784](#)