

Supplementary materials

Prognostic and therapeutic insights from lactate metabolism and tumor immune microenvironment in head and neck squamous cell carcinoma

Authors: Huanyu Jiang^{a,b}, Lijuan Zhou^b, Haidong Zhang^{a,b}, Shanchun Gong^b, Zhenkun Yu^{a,b*}

^aSchool of Medicine, Southeast University, Nanjing 210009, Jiangsu, China


^bDepartment of Otolaryngology Head and Neck Surgery, The Affiliated BenQ Hospital of Nanjing Medical University, Nanjing 210019, Jiangsu, China

Corresponding author: Zhenkun Yu


Email: yuzhenkun65@hotmail.com

Supplementary Table S1. Baseline characteristics of patients with HNSCC in TCGA


Characteristics	Count	Percentage
Age		
≥60	278	56.28%
<60	216	43.72%
Sex		
male	362	73.28%
female	132	26.72%
Clinical_T		
T1	33	6.68%
T2	141	28.54%
T3	129	26.11%
T4	176	35.63%
Not Reported	15	3.04%
Clinical_N		
N0	237	47.98%
N1	80	16.19%
N2	150	30.36%
N3	5	1.01%
Not Reported	22	4.45%
Clinical_M		
M0	464	93.93%
M1	5	1.01%
Not Reported	25	5.06%
Clinical_stage		
Stage I	19	3.85%
Stage II	93	18.83%
Stage III	102	20.65%
Stage IV	266	53.85%
Not Reported	14	2.83%
Grade		
G1	61	12.35%
G2	294	59.51%
G3	118	23.89%
G4	2	0.40%
Not Reported	19	3.85%




Supplementary Fig. S1. The LRS formula was constructed using a Lasso regression.




Supplementary Fig. S2. Positive and negative controls for monocarboxylate transporter 4 (MCT4) immunohistochemistry. (A) Positive control using human prostate tissue. (B) Negative control without the primary antibody.



Supplementary Fig. S3. Positive and negative controls for glycogen phosphorylase L (PYGL) immunohistochemistry. (A) Positive control using human kidney tissue. (B) Negative control without the primary antibody.



Supplementary Fig. S4. Positive and negative controls for CXCL9 immunostaining. (A) Positive control using human prostate cancer tissue. (B) Negative control processed without the primary antibody.



Supplementary Fig. S5. Positive and negative controls for CXCL10 immunostaining. (A) Positive control using human nasopharyngeal carcinoma tissue. (B) Negative control processed without the primary antibody.