

INTRODUCTION TO THE PRIME NUMBER THEORY

FROM SPECIAL FUNCTIONS TO A SIMPLIFIED PROOF

Anonymous

January 13, 2021

Institute = None

TABLE OF CONTENTS

1. Introduction to PNT
2. Propositions of $\Gamma(s)$ and $\zeta(s)$
3. Prime-related Propositions of Special Functions
4. Simplified Proof of PNT based on Newman's Work
5. References

INTRO TO PNT



HOW PRIME NUMBERS ASYMPTOTICALLY DISTRIBUTE
AMONG LARGE POSITIVE INTEGERS?

The prime-counting function $\pi(x)$ is introduced to count the number of primes less than x , i.e. $\pi(x) = \sum_{p \leq x} 1$.

WHAT PNT IS

The prime-counting function $\pi(x)$ is introduced to count the number of primes less than x , i.e. $\pi(x) = \sum_{p \leq x} 1$.




Figure 1: The values of $\pi(x)$ for the first 60 positive integers.

WHAT PNT IS

The prime-counting function $\pi(x)$ is introduced to count the number of primes less than x , i.e. $\pi(x) = \sum_{p \leq x} 1$.




Figure 2: The values of $\pi(x)$ for the first 60 positive integers.

It was conjectured in the end of the 18th century by Gauss that $\lim_{x \rightarrow \infty} \frac{\pi(x)}{x/\log(x)} = 1$, known as the prime number theorem (PNT).

- The PNT was first proved in 1896 by Jacques Hadamard and by Charles de la Vallée Poussin independently, using **bulk of properties of the Riemann zeta function**.

- The PNT was first proved in 1896 by Jacques Hadamard and by Charles de la Vallée Poussin independently, using **bulk of properties of the Riemann zeta function**.
- The "elementary" proof were found around 1948 by Atle Selberg and by Paul Erdős. (**Does not use any complex analysis.**)

- The PNT was first proved in 1896 by Jacques Hadamard and by Charles de la Vallée Poussin independently, using **bulk of properties of the Riemann zeta function**.
- The "elementary" proof were found around 1948 by Atle Selberg and by Paul Erdős. (**Does not use any complex analysis.**)
- D. J. Newman found a very simple version of the Tauberian argument needed for an analytic proof of the PNT.

We will describe the resulting proof later, which has a beautifully simple structure and uses **hardly anything beyond Cauchy's theorem**.

**IN THIS PRESENTATION, WE WILL INTRODUCE A
SIMPLIFIED PROOF BASED ON NEWMAN'S WORK.**

MAIN STRUCTURE OF A SIMPLIFIED PROOF

$$\left. \begin{array}{l} \Gamma(s) \text{ on } \mathbb{C} \\ \text{extension of } \zeta(s) \text{ on } \Re(s) > 0 \\ \text{the functional equation} \end{array} \right\} \Rightarrow \zeta(s) \text{ on } \mathbb{C}$$

$$\left. \begin{array}{l} \zeta(s) \text{ on } \mathbb{C} \\ \text{zeros of } \zeta(s) \text{ when } \Re(s) = 1 \end{array} \right\} \Rightarrow \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1$$

$$\left. \begin{array}{l} \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1 \\ \text{Analytic Theorem} \end{array} \right\} \Rightarrow \frac{\vartheta(x)}{x} \sim 1 \Leftrightarrow \frac{\pi(x)}{x/\log x} \sim 1 \text{ (PNT)}$$

Figure 3: Main Structure of the Simplified Proof

PROPOSITIONS OF $\Gamma(s)$ AND $\zeta(s)$

HOW $\Gamma(s)$ IS DEFINED ON \mathbb{C}

Gamma function is one possible extension of factorial, i.e.

$\Gamma(n) = (n - 1)!$ when $n \in \mathbb{N}$. Furthermore, via following propositions, $\Gamma(s)$ can be extended to a meromorphic function on \mathbb{C} .

HOW $\Gamma(s)$ IS DEFINED ON \mathbb{C}

Gamma function is one possible extension of factorial, i.e.

$\Gamma(n) = (n-1)!$ when $n \in \mathbb{N}$. Furthermore, via following propositions, $\Gamma(s)$ can be extended to a meromorphic function on \mathbb{C} .

- (Definition of $\Gamma(s)$ when $\Re(s) > 1$) $\Gamma(s) := \int_0^\infty t^{s-1} e^{-t} dt;$

HOW $\Gamma(s)$ IS DEFINED ON \mathbb{C}

Gamma function is one possible extension of factorial, i.e.

$\Gamma(n) = (n-1)!$ when $n \in \mathbb{N}$. Furthermore, via following propositions, $\Gamma(s)$ can be extended to a meromorphic function on \mathbb{C} .

- (Definition of $\Gamma(s)$ when $\Re(s) > 1$) $\Gamma(s) := \int_0^\infty t^{s-1} e^{-t} dt$;
- Since $\Gamma(s)$ is the extension of factorial, $\Gamma(s+1) = s\Gamma(s)$ for $\Re(1+s) > 1$.

HOW $\Gamma(s)$ IS DEFINED ON \mathbb{C}

Gamma function is one possible extension of factorial, i.e.

$\Gamma(n) = (n-1)!$ when $n \in \mathbb{N}$. Furthermore, via following propositions, $\Gamma(s)$ can be extended to a meromorphic function on \mathbb{C} .

- (Definition of $\Gamma(s)$ when $\Re(s) > 1$) $\Gamma(s) := \int_0^\infty t^{s-1} e^{-t} dt$;
- Since $\Gamma(s)$ is the extension of factorial, $\Gamma(s+1) = s\Gamma(s)$ for $\Re(1+s) > 1$.

Therefore, $\Gamma(s)$ is analytic on $\mathbb{C} \setminus \{0, -1, -2, \dots\}$.

SOME PROPOSITIONS OF $\Gamma(s)$

From the previous analysis, $\Gamma(s)$ is analytic on $\mathbb{C} \setminus \{0, -1, -2, \dots\}$.

Furthermore, $\Gamma(s)$ satisfies the following propositions

SOME PROPOSITIONS OF $\Gamma(s)$

From the previous analysis, $\Gamma(s)$ is analytic on $\mathbb{C} \setminus \{0, -1, -2, \dots\}$.

Furthermore, $\Gamma(s)$ satisfies the following propositions

- The product form of $\Gamma(s)$ when $s \notin \{0, -1, -2, \dots\}$ lies that:

$$\Gamma(s) = \frac{1}{s} \prod_{k=1}^{\infty} \left[\left(1 + \frac{1}{k}\right)^s \left(1 + \frac{s}{k}\right)^{-1} \right] = \lim_{n \rightarrow \infty} \frac{n! n^s}{s(s+1) \cdots (s+n)}$$

SOME PROPOSITIONS OF $\Gamma(s)$

From the previous analysis, $\Gamma(s)$ is analytic on $\mathbb{C} \setminus \{0, -1, -2, \dots\}$.

Furthermore, $\Gamma(s)$ satisfies the following propositions

- The product form of $\Gamma(s)$ when $s \notin \{0, -1, -2, \dots\}$ lies that:

$$\Gamma(s) = \frac{1}{s} \prod_{k=1}^{\infty} \left[\left(1 + \frac{1}{k}\right)^s \left(1 + \frac{s}{k}\right)^{-1} \right] = \lim_{n \rightarrow \infty} \frac{n! n^s}{s(s+1) \cdots (s+n)}$$

- $\Gamma(s)$ never vanishes, thus $\frac{1}{\Gamma(s)}$ is analytic on \mathbb{C} .

SOME PROPOSITIONS OF $\Gamma(s)$

From the previous analysis, $\Gamma(s)$ is analytic on $\mathbb{C} \setminus \{0, -1, -2, \dots\}$.

Furthermore, $\Gamma(s)$ satisfies the following propositions

- The product form of $\Gamma(s)$ when $s \notin \{0, -1, -2, \dots\}$ lies that:

$$\Gamma(s) = \frac{1}{s} \prod_{k=1}^{\infty} \left[\left(1 + \frac{1}{k}\right)^s \left(1 + \frac{s}{k}\right)^{-1} \right] = \lim_{n \rightarrow \infty} \frac{n! n^s}{s(s+1) \cdots (s+n)}$$

- $\Gamma(s)$ never vanishes, thus $\frac{1}{\Gamma(s)}$ is analytic on \mathbb{C} .
- (Euler's reflection formula) $\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin(\pi s)}$.

SOME PROPOSITIONS OF $\Gamma(s)$

From the previous analysis, $\Gamma(s)$ is analytic on $\mathbb{C} \setminus \{0, -1, -2, \dots\}$.

Furthermore, $\Gamma(s)$ satisfies the following propositions

- The product form of $\Gamma(s)$ when $s \notin \{0, -1, -2, \dots\}$ lies that:

$$\Gamma(s) = \frac{1}{s} \prod_{k=1}^{\infty} \left[\left(1 + \frac{1}{k}\right)^s \left(1 + \frac{s}{k}\right)^{-1} \right] = \lim_{n \rightarrow \infty} \frac{n! n^s}{s(s+1) \cdots (s+n)}$$

- $\Gamma(s)$ never vanishes, thus $\frac{1}{\Gamma(s)}$ is analytic on \mathbb{C} .
- (Euler's reflection formula) $\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin(\pi s)}$.

The following propositions has been proven in the LECTURE.

HOW ζ IS DEFINED ON \mathbb{C}

One method to the analytic extension of $\zeta(s)$ is presented as follows

HOW ζ IS DEFINED ON \mathbb{C}

One method to the analytic extension of $\zeta(s)$ is presented as follows

1. (Definition of $\zeta(s)$ when $\Re(s) > 1$) $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s};$

HOW ζ IS DEFINED ON \mathbb{C}

One method to the analytic extension of $\zeta(s)$ is presented as follows

1. (Definition of $\zeta(s)$ when $\Re(s) > 1$) $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$;
2. An equivalent definition for $\Re(s) > 1$ lies that

$$\zeta(s) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \quad \Re(s) > 1$$

HOW ζ IS DEFINED ON \mathbb{C}

One method to the analytic extension of $\zeta(s)$ is presented as follows

1. (Definition of $\zeta(s)$ when $\Re(s) > 1$) $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$;
2. An equivalent definition for $\Re(s) > 1$ lies that

$$\zeta(s) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \quad \Re(s) > 1$$

3. The extension of $\zeta(s)$ on \mathbb{C}_+ can be written as

$$\zeta(s) = \frac{1}{\Gamma(s)} \left(\int_0^1 \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) t^{s-1} dt + (s-1)^{-1} + \int_1^{\infty} \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \right)$$

HOW ζ IS DEFINED ON \mathbb{C}

One method to the analytic extension of $\zeta(s)$ is presented as follows

1. (Definition of $\zeta(s)$ when $\Re(s) > 1$) $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$;
2. An equivalent definition for $\Re(s) > 1$ lies that

$$\zeta(s) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \quad \Re(s) > 1$$

3. The extension of $\zeta(s)$ on \mathbb{C}_+ can be written as

$$\zeta(s) = \frac{1}{\Gamma(s)} \left(\int_0^1 \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) t^{s-1} dt + (s-1)^{-1} + \int_1^{\infty} \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \right)$$

4. Via the functional equation $\zeta(1-s) = 2^{1-s} \pi^{-s} \zeta(s) \Gamma(s) \cos(s\pi/2)$, $\zeta(s)$ can be meromorphically extended on \mathbb{C} .

HOW ζ IS DEFINED ON \mathbb{C}

Formulas/Equations

$$\zeta(1-s) = 2^{1-s} \pi^{-s} \zeta(s) \Gamma(s) \cos(s\pi/2)$$

$$\zeta(s) = \frac{1}{\Gamma(s)} \left(\int_0^1 \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) t^{s-1} dt + (s-1)^{-1} + \int_1^\infty \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \right)$$

HOW ζ IS DEFINED ON \mathbb{C}

Formulas/Equations

$$\zeta(1-s) = 2^{1-s} \pi^{-s} \zeta(s) \Gamma(s) \cos(s\pi/2)$$

$$\zeta(s) = \frac{1}{\Gamma(s)} \left(\int_0^1 \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) t^{s-1} dt + (s-1)^{-1} + \int_1^\infty \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \right)$$

Furthermore, $\zeta(s) - \frac{1}{s-1}$ is analytic on \mathbb{C}_+ .

HOW ζ IS DEFINED ON \mathbb{C}

Formulas/Equations

$$\zeta(1-s) = 2^{1-s} \pi^{-s} \zeta(s) \Gamma(s) \cos(s\pi/2)$$

$$\zeta(s) = \frac{1}{\Gamma(s)} \left(\int_0^1 \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) t^{s-1} dt + (s-1)^{-1} + \int_1^\infty \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \right)$$

Furthermore, $\zeta(s) - \frac{1}{s-1}$ is analytic on \mathbb{C}_+ .

The functional equation shows that the only singularity of $\zeta(s)$ is at $s = 1$, which is a single pole with residue 1.

HOW ζ IS DEFINED ON \mathbb{C}

Formulas/Equations

$$\zeta(1-s) = 2^{1-s} \pi^{-s} \zeta(s) \Gamma(s) \cos(s\pi/2)$$

$$\zeta(s) = \frac{1}{\Gamma(s)} \left(\int_0^1 \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) t^{s-1} dt + (s-1)^{-1} + \int_1^\infty \frac{t^{s-1} e^{-t}}{1 - e^{-t}} dt \right)$$

Furthermore, $\zeta(s) - \frac{1}{s-1}$ is analytic on \mathbb{C}_+ .

The functional equation shows that the only singularity of $\zeta(s)$ is at $s = 1$, which is a single pole with residue 1.

Significant conclusions

- $\zeta(s) - \frac{1}{s-1}$ is analytic on \mathbb{C} ;
- $\zeta(s)$ never vanishes when $\Re(s) > 1$.

HOW ζ IS DEFINED ON \mathbb{C}

Significant conclusions

- $\zeta(s) - \frac{1}{s-1}$ is analytic on \mathbb{C} ;
- $\zeta(s)$ never vanishes when $\Re(s) > 1$ ($\Re(s) \geq 1$ indeed).




Figure 4: Critical strip of Riemann zeta function

$$\left. \begin{array}{l} \Gamma(s) \text{ on } \mathbb{C} \\ \text{extension of } \zeta(s) \text{ on } \Re(s) > 0 \\ \text{the functional equation} \end{array} \right\} \Rightarrow \zeta(s) \text{ on } \mathbb{C}$$

$$\left. \begin{array}{l} \zeta(s) \text{ on } \mathbb{C} \\ \text{zeros of } \zeta(s) \text{ when } \Re(s) = 1 \end{array} \right\} \Rightarrow \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1$$

$$\left. \begin{array}{l} \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1 \\ \text{Analytic Theorem} \end{array} \right\} \Rightarrow \frac{\vartheta(x)}{x} \sim 1 \Leftrightarrow \frac{\pi(x)}{x/\log x} \sim 1 \text{ (PNT)}$$

Figure 5: Proof Progress

PRIME-RELATED PROPOSITIONS OF SPECIAL FUNCTIONS

Definitions of some prime-related functions

- $\pi(x)$, prime-counting function, is defined as $\pi(x) = \sum_{p \leq x} 1$;
- $\Phi(s)$ denotes $\sum_p \frac{\log p}{p^s} \quad \Re(s) > 1$;
- Λ function is defined as $\Lambda : \mathbb{N} \rightarrow \mathbb{R}, n \mapsto \begin{cases} \log p & n = p^k, k \in \mathbb{N} \\ 0 & \text{otherwise.} \end{cases}$;
- $\vartheta(x)$ is defined as $\vartheta(x) = \sum_{p \leq x} \log p$;
- (Well-known proposition) $\zeta(s) = \prod_p (1 - p^{-s})^{-1} \quad \Re(s) > 1$.

REMARK: $x \in \mathbb{R}$ and $s \in \mathbb{C}$.

SPECIAL FUNCTIONS & PRIMES

Proposition: $-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n \geq 2} \Lambda(n) e^{-s \log n} \quad \Re(s) > 1$

Proof.

Recall that $\zeta(s) = \prod_p (1 - p^{-1})^{-1}$, then

$$-\frac{\zeta'(s)}{\zeta(s)} = -(\log \zeta(s))' = \sum_p \frac{e^{-s \log p} \log p}{1 - e^{-s \log p}} = \sum_p \left(\log p \sum_{k=1}^{\infty} (e^{-s \log p})^k \right)$$

Since the infinite summation uniformly converges for any p ,

$$\sum_p \left(\log p \sum_{k=1}^{\infty} (e^{-s \log p})^k \right) = \sum_{k=1}^{\infty} \sum_p (\log p \cdot e^{-s \log p^k}) = \sum_{n=2}^{\infty} \Lambda(n) e^{-s \log n}$$

□

NOTE: $\Lambda : \mathbb{N} \rightarrow \mathbb{R}, n \mapsto \begin{cases} \log p & n = p^k, k \in \mathbb{N} \\ 0 & \text{otherwise.} \end{cases}$

Proposition: $\zeta(s)$ has no zeros on $\Re(s) = 1$

PROOF SKETCH Suppose that $\exists t_0$ such that $\zeta(1 + it_0) = 0$.

Set $F(s) = \zeta^4(s)\zeta^5(s + it_0)\zeta^2(s + 2it_0)$. Since $s = 1$ is the zero of $F(s)$, $F(s)$ is analytic in the neighbourhood of $s = 1$. For any $x \in (1, 1 + \delta)$ where δ is small enough, we define $f(x) := F(s)|_{\mathbb{R}}$. By definition,

Proposition: $\zeta(s)$ has no zeros on $\Re(s) = 1$

PROOF SKETCH Suppose that $\exists t_0$ such that $\zeta(1 + it_0) = 0$.

Set $F(s) = \zeta^4(s)\zeta^5(s + it_0)\zeta^2(s + 2it_0)$. Since $s = 1$ is the zero of $F(s)$, $F(s)$ is analytic in the neighbourhood of $s = 1$. For any $x \in (1, 1 + \delta)$ where δ is small enough, we define $f(x) := F(s)|_{\mathbb{R}}$. By definition,

$$\begin{aligned}\Re\left(\frac{f'(x)}{f(x)}\right) &= \Re\left(4\frac{\zeta'(x)}{\zeta(x)} + 5\frac{\zeta'(x + it_0)}{\zeta(x + it_0)} + 2\frac{\zeta'(x + 2it_0)}{\zeta(x + 2it_0)}\right) \\ &= -\Re\sum_{n \geq 2} \Lambda(n)(4e^{x \log n} 5e^{(x+it_0) \log n} + 2e^{(x+2it_0) \log n}) \\ &= -\sum_{n \geq 2} \Lambda(n)e^{-x \log n}(4 + 5\cos(t_0 \log n) + 2\cos(2t_0 \log n)) \\ &= -\sum_{n \geq 2} \Lambda(n)e^{-x \log n}((2\cos(t_0 \log n) + 5/4)^2 + 7/16) < 0\end{aligned}$$

NOTE: $-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n \geq 2} \Lambda(n)e^{-s \log n}.$

SPECIAL FUNCTIONS & PRIMES

Proposition: $\zeta(s)$ has no zeros on $\Re(s) = 1$

PROOF SKETCH Suppose that $\exists t_0$ such that $\zeta(1 + it_0) = 0$.

Set $F(s) = \zeta^4(s)\zeta^5(s + it_0)\zeta^2(s + 2it_0)$. Since $s = 1$ is the zero of $F(s)$, $F(s)$ is analytic in the neighbourhood of $s = 1$. For any $x \in (1, 1 + \delta)$ where δ is small enough, we define $f(x) := F(s)|_{\mathbb{R}}$. By definition,

$$\begin{aligned}\Re\left(\frac{f'(x)}{f(x)}\right) &= \Re\left(4\frac{\zeta'(x)}{\zeta(x)} + 5\frac{\zeta'(x + it_0)}{\zeta(x + it_0)} + 2\frac{\zeta'(x + 2it_0)}{\zeta(x + 2it_0)}\right) \\ &= -\Re\sum_{n \geq 2} \Lambda(n)(4e^{x \log n} 5e^{(x+it_0) \log n} + 2e^{(x+2it_0) \log n}) \\ &= -\sum_{n \geq 2} \Lambda(n)e^{-x \log n}(4 + 5\cos(t_0 \log n) + 2\cos(2t_0 \log n)) \\ &= -\sum_{n \geq 2} \Lambda(n)e^{-x \log n}((2\cos(t_0 \log n) + 5/4)^2 + 7/16) < 0\end{aligned}$$

Whereas, $\lim_{x \rightarrow 1^+} (x-1)\frac{f'(x)}{f(x)} = 1$ implies that $\left(\frac{f'(x)}{f(x)}\right) \geq 0!$

NOTE: $-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n \geq 2} \Lambda(n)e^{-s \log n}.$

$$\left. \begin{array}{l} \Gamma(s) \text{ on } \mathbb{C} \\ \text{extension of } \zeta(s) \text{ on } \Re(s) > 0 \\ \text{the functional equation} \end{array} \right\} \Rightarrow \zeta(s) \text{ on } \mathbb{C}$$

$$\left. \begin{array}{l} \zeta(s) \text{ on } \mathbb{C} \\ \text{zeros of } \zeta(s) \text{ when } \Re(s) = 1 \end{array} \right\} \Rightarrow \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1$$

$$\left. \begin{array}{l} \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1 \\ \text{Analytic Theorem} \end{array} \right\} \Rightarrow \frac{\vartheta(x)}{x} \sim 1 \Leftrightarrow \frac{\pi(x)}{x/\log x} \sim 1 \text{ (PNT)}$$

Figure 6: Proof Progress

Proposition: $\Phi(s) - \frac{1}{s-1}$ is meromorphic on \mathbb{C}_+ , analytic when $\Re(s) \geq 1$.

Proposition: $\Phi(s) - \frac{1}{s-1}$ is meromorphic on \mathbb{C}_+ , analytic when $\Re(s) \geq 1$.

Proof.

$$\text{When } \Re(s) > 1, -\frac{\zeta'(s)}{\zeta(s)} = \sum_{k=1}^{\infty} \sum p(\log p \cdot e^{-s \log p^k}) = \sum_{k=1}^{\infty} \Phi(ks).$$

□

Proposition: $\Phi(s) - \frac{1}{s-1}$ is meromorphic on \mathbb{C}_+ , analytic when $\Re(s) \geq 1$.

Proof.

$$\text{When } \Re(s) > 1, -\frac{\zeta'(s)}{\zeta(s)} = \sum_{k=1}^{\infty} \sum p(\log p \cdot e^{-s \log p^k}) = \sum_{k=1}^{\infty} \Phi(ks).$$

Since $\Phi(s)$ is analytic when $\Re(s) > 1$, $\Phi(s/2)$ is meromorphic when $\Re(s) > 1$, so is $\Phi(s/4)$, $\Phi(s/8)$, etc. Therefore $\Phi(s)$ is meromorphic when $\Re(s) > 0$, whose poles are all simple at the zeros of $\zeta(s)$. \square

Proposition: $\Phi(s) - \frac{1}{s-1}$ is meromorphic on \mathbb{C}_+ , analytic when $\Re(s) \geq 1$.

Proof.

$$\text{When } \Re(s) > 1, -\frac{\zeta'(s)}{\zeta(s)} = \sum_{k=1}^{\infty} \sum p(\log p \cdot e^{-s \log p^k}) = \sum_{k=1}^{\infty} \Phi(ks).$$

Since $\Phi(s)$ is analytic when $\Re(s) > 1$, $\Phi(s/2)$ is meromorphic when $\Re(s) > 1$, so is $\Phi(s/4)$, $\Phi(s/8)$, etc. Therefore $\Phi(s)$ is meromorphic when $\Re(s) > 0$, whose poles are all simple at the zeros of $\zeta(s)$.

We claim that the only pole of $-\frac{\zeta'(s)}{\zeta(s)}$ when $\Re(s) = 1$ is at $s = 1$, since $\zeta(s)$ has neither zeros nor singularities when $\Re(s) = 1$ and $\Im(s) \neq 0$.

Hence $\Phi(s) - \frac{1}{s-1}$ is analytic on $\{s : \Re(s) \geq 1\}$ due to the residue of $\zeta(s)$ at $s = 1$ is 1. □

SHORT PROOF OF THE PNT

DECLARATION

THIS PROOF IS SIMPLIFIED FROM NEWMAN'S ORIGINAL WORK, ALSO REFERS
TO A PROOF SKETCH BY TERENCE TAO. (SEE REFERENCES PART)

Theorem

Let $f(t)(t \geq 0)$ be a bounded and locally integrable function and suppose that the function

$$g(s) = \int_0^{\infty} f(t)e^{-st}dt \quad \Re(s) > 0$$

extends analytically to $\Re(s) \geq 0$. Then $\int_0^{\infty} f(t)dt$ exists (and valued $g(0)$).

The full proof has been presented in the LECTURE.

ANALYTIC THEOREM

Theorem

Let $f(t)$ ($t \geq 0$) be a bounded and locally integrable function and suppose that the function

$$g(s) = \int_0^{\infty} f(t)e^{-st}dt \quad \Re(s) > 0$$

extends analytically to $\Re(s) \geq 0$. Then $\int_0^{\infty} f(t)dt$ exists (and valued $g(0)$).

PROOF SKETCH

PROOF PROGRESS

$$\left. \begin{array}{l} \Gamma(s) \text{ on } \mathbb{C} \\ \text{extension of } \zeta(s) \text{ on } \Re(s) > 0 \\ \text{the functional equation} \end{array} \right\} \Rightarrow \zeta(s) \text{ on } \mathbb{C}$$

$$\left. \begin{array}{l} \zeta(s) \text{ on } \mathbb{C} \\ \text{zeros of } \zeta(s) \text{ when } \Re(s) = 1 \end{array} \right\} \Rightarrow \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1$$

$$\left. \begin{array}{l} \Phi(s) - \frac{1}{s-1} \text{ is analytic when } \Re(s) \geq 1 \\ \text{Analytic Theorem} \end{array} \right\} \Rightarrow \frac{\vartheta(x)}{x} \sim 1 \Leftrightarrow \frac{\pi(x)}{x/\log x} \sim 1 \text{ (PNT)}$$

Figure 7: Proof Progress

Theorem

$$\lim_{x \rightarrow \infty} \frac{\vartheta(x)}{x} = 1$$

Theorem

$$\lim_{x \rightarrow \infty} \frac{\vartheta(x)}{x} = 1$$

NOTE: $\vartheta(x) = \sum_{p \leq x} \log p$.

Theorem

$$\lim_{x \rightarrow \infty} \frac{\vartheta(x)}{x} = 1$$

Proof.

For $\Re(s) > 1$, we have

$$\Phi(s) = \sum_p \frac{\log p}{p^s} = s \sum_p \int_p^\infty \frac{\log p}{x^{s+1}} dx = s \int_1^\infty \frac{\vartheta(x)}{x^{s+1}} dx = s \int_0^\infty e^{-st} \vartheta(e^t) dt$$

$$\text{Set } g(s) := \frac{\Phi(s+1)}{s+1} - \frac{1}{s} = \int_0^\infty e^{-st} \left(\frac{\vartheta(e^t)}{e^t} - 1 \right) dt, f(x) = \frac{\vartheta(e^t)}{e^t} - 1.$$

Since

$$\begin{aligned} \vartheta(x) &= \sum_{n=0}^{\infty} \left(\vartheta\left(\frac{x}{2^n}\right) - \vartheta\left(\frac{x}{2^{n+1}}\right) \right) \leq \sum_{n=0}^{\infty} \left(\log\left(\frac{[x/2^n]}{[x/2^{n+1}]}\right) \right) \\ &\leq \sum_{n=1}^{\infty} \log 2^{[x/2^n]} \leq \sum_{n=1}^{\infty} \frac{x \log 2}{2^n} \leq (2 \log 2)x \end{aligned}$$

$f(x)$ is bounded when $x \in [0, \infty)$. Hence $\int_0^\infty \frac{\vartheta(e^t)}{e^t} - 1 dt = \int_0^\infty \frac{\vartheta(x) - x}{x^2} dx$ exists.

ANALYTIC THEOREM

Theorem

$$\lim_{x \rightarrow \infty} \frac{\vartheta(x)}{x} = 1$$

Proof.

We claim that $\vartheta(x) \sim x$ when x is large enough. Otherwise, we have $\limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x} > 1$ or $\liminf_{x \rightarrow \infty} \frac{\vartheta(x)}{x} < 1$.

Without the loss of generality, suppose that $\limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x} > 1$. Equivalently speaking, there exists $\lambda > 1$ and an increasing sequence $\{x_n\} \xrightarrow{n \rightarrow \infty} \infty$ such that $\frac{\vartheta(x)}{x} > \lambda$. For any $x \in \{x_n\}$, we have

$$\int_x^{\lambda x} \frac{\vartheta(t) - t}{t^2} dt \geq \int_x^{\lambda x} \frac{\lambda x - t}{t^2} dt = \int_1^\lambda \frac{\lambda - t}{t^2} dt > 0$$

As a result, it contradicts to the existence of $\int_1^\infty \frac{\vartheta(t) - t}{t^2} dt$. □

The Final Theorem (PNT)

$$\lim_{n \rightarrow \infty} \frac{\pi(x)}{x} = 1$$

ANALYTIC THEOREM

The Final Theorem (PNT)

$$\lim_{n \rightarrow \infty} \frac{\pi(x)}{x} = 1$$

Proof.

$\forall \varepsilon \in (0, 1)$, we have

$$\vartheta(x) \leq \sum_{p \leq x} \log p = \pi(x) \log x$$

and

$$\vartheta(x) \geq \sum_{x^{1-\varepsilon} < p \leq x} \log p \geq (1 - \varepsilon)(\pi(x) - \pi(x^{1-\varepsilon})) \log x$$

$$\text{Therefore } (1 - \varepsilon)[\pi(x) - \pi(x^{1-\varepsilon})] \leq \frac{\vartheta(x)}{\log x} \leq \pi(x).$$

The PNT follows easily since $\lim_{x \rightarrow \infty} \frac{\vartheta(x)}{x}$.

□

THANKS FOR LISTENING!

REFERENCES



R. E. Greene and S. G. Krantz.

Function Theory of One Complex Variable (third edition).

American Mathematical Society Providence, Rhode Island, 2006.



A. E. Ingham.

The Distribution of Prime Numbers.

Cambridge University Press, 1990.



D. J. Newman.

Simple analytic proof of the prime number theorem.

Amer. Math. Monthly, 87:693–696, 1980.



T. Tao.

254a, notes 2: Complex-analytic multiplicative number theory.

Terence Tao's blog, 12 2014.