

# CS161 - Quiz 6

Started: Mar 10 at 6:06pm

## Quiz Instructions

---

### Question 1

1 pts

$Pr(\alpha) = Pr(\alpha \wedge \beta) Pr(\alpha \wedge \neg\beta)$  for any events  $\alpha$  and  $\beta$

- ☐ True
- ☒ False

### Question 2

1 pts

If X and Y are probabilistically independent, then they must continue to be independent given any variable Z.

- ☒ True
- ☐ False

### Question 3

1 pts

The local semantics of Bayesian network says:

- ☐ Every node is independent of its parents given its non-descendants
- ☐ Every node is independent of its descendants given its parents
- ☐ None of the others
- ☐ Every node is independent of its parents given its descendants

- ☒ Every node is independent of its non-descendants given its parents

**Question 4****1 pts**

Consider a Bayesian network  $X_1 \rightarrow X_2 \rightarrow \dots \rightarrow X_n$  (a chain with  $n$  nodes). Assume that each variable  $X_i$  has only two values  $x_i$  and  $\overline{x_i}$ . Then  $Pr(x_3 \mid \overline{x_1})$  is equal to:

- ☒  $Pr(x_3 \mid x_2) Pr(x_2 \mid \overline{x_1}) + Pr(x_3 \mid \overline{x_2}) Pr(\overline{x_2} \mid \overline{x_1})$
- ☐  $Pr(x_3, \overline{x_1}) Pr(\overline{x_1})$
- ☐ None of the others
- ☐  $Pr(x_3)$
- ☐  $Pr(x_3 \mid x_2, \overline{x_1}) + Pr(x_3 \mid \overline{x_2}, \overline{x_1})$

Quiz saved at 6:08pm

[Submit Quiz](#)