

## Question 1

Not yet answered

Marked out of 2.00

Consider the following distributions for a random variable  $X$ . In each case, match the distribution with the expectation of

$$Y = 10 - 2X^2$$

Geometric( $p=0.6$ )

Choose...

Binomial( $n=10, p=0.3$ )

Choose...

Poisson distribution with

$$\lambda = 5$$

Choose...

$$f(x) = \frac{1}{2}e^{-x/2}dx, \quad 0 \leq x \leq \infty$$

Choose...

## Question 2

Answer saved

Marked out of 1.00

An  $n$ -tuple is a sequence of  $n$  numbers ordered in a particular way. For example,  $(2,2,3,4)$  is a 3-tuple. And  $(3,4,2,2)$  is a different 3-tuple. The  $n$ -tuple concept is different from the concept of set.

So thinking about  $n$ -tuples, the difference between the sample space of a binomial experiment and the sample space of a geometric experiment is, as we saw in lectures,

- ☐ a. There is no difference between the outcomes of the sample space in a binomial experiment and a geometric experiment, because they are all Bernoulli trials.
- ☐ b. All outcomes in the sample space of the binomial experiment are  $n$ -tuples of fixed length  $n$ , but in a geometric experiment all outcomes  $n$ -tuples differ in length  $n$ , some are length  $n-1$ , others length different from 1, etc.
- ☐ c. The random variable in a Binomial experiment has values  $0,1,2,\dots$  and so does the random variable in a Geometric experiment, so since the random variable is the outcome in the sample space both the binomial and the geometric have identical  $n$ -tuples.
- ☐ d. Outcomes in the sample space of the binomial experiment differ in length, but in a geometric experiment all outcomes  $n$ -tuples have the same length  $n$ .

**Question 3**

Answer saved

Marked out of 1.00

Two different manufacturers supply a component with an exponentially distributed lifetime, that is, the length of service the component gives until it fails is an exponentially distributed random variable. Manufacturer A's device has expected lifetime 4 months and manufacturer B's has 10 months. A particular user has a batch of devices of which 40% came from manufacturer A and 60% from manufacturer B. If a randomly selected device from this batch is used, what is the probability that the lifetime of this device is more than 2 months?

- ☐ a. 0.4
- ☐ b. 0.6
- ☐ c. 0.18127
- ☐ d. 0.73385

## Question 4

Answer saved

Marked out of 1.00

A very complicated task, when given by a psychologist to randomly chosen children, takes an average 5 seconds to complete. But there is variability, and we know that there is a standard deviation of 1 second. The probability that a randomly chosen child takes between 4 and 7 seconds to complete the task is

Select one:

- ☐ a. 0.6148
- ☐ b. 0.00001
- ☐ c. 0.8185
- ☐ d. 0.8185

## Question 5

Answer saved

Marked out of 3.00

This is a question that requires showing work. That means, you must define your notation, must tell us what you are calculating, must show detailed work and provide final answers and explain.

No attachments are allowed for this question. You must answer in the space provided.

Not all taxpayers want to finance new infrastructure projects. For that reason, public opinion is constantly measured by local governments in order to determine the chance of new projects implementation. Public opinion in a city regarding the opening of a car pool lane in its most congested highway is reflected in the following table.

	Yes	No
Center of the city	0.150	0.250
Suburbs	0.250	0.150
Rural areas	0.050	0.150

The table reflects the opinion of adults eligible to vote and is saying, for example, that 15% of the town adults eligible to vote live in the center of the city and are in favor of the car pool lane.

With this information, answer the following questions:

- (i) What is the probability that a randomly chosen eligible voter disapproves of the car pool lane?
- (ii) What is the probability that a randomly chosen eligible voter does not live in the center of the city and disapproves of the car pool lane?
- (iii) What is the probability that a voter from the suburbs disapproves of the car pool lane?



Maximum size for new files: 64MB

[Files](#)

You can drag and drop files here to add them.

## Question 6

Answer saved

Marked out of 3.00

Consider a random variable  $X$  with density function

$$f(x) = \frac{1}{25}(10 - 2x), \quad 0 \leq x \leq 5$$

Match the expressions.

$$\frac{1}{25}(10x - x^2), \quad 0 \leq x \leq 5$$

0.669873

1.66

## Question 7

Answer saved

Marked out of 3.00

This is a question that requires showing work. That means, you must define your notation, must tell us what you are calculating, must show detailed work and provide final answers and explain.

-----

A complex system requires that a certain function be performed with reliability 0.9999. Devices that perform the function have reliability only 0.8, and so it is necessary to build redundancy. Several of the devices are to be used, and if at least one works then the function will be performed. Furthermore, we are willing to assume that the redundant devices function independently of each other. How many of the devices must be installed in order that the system will have the desired reliability?

*No attachments are allowed for this question. The question must be answered in the space provided.*

↵

A ▾

B

I

≡

≡

≡

≡

🔗

🔄

🖼️

## Question 8

Answer saved

Marked out of 1.00

(This exercise is based on Horgan 2009, Section 6.3.) A microprocessor chip is a very important component of every computer. Once in a while tech news magazines report a defect in a chip and producers have to respond by giving some indication of the damage that we should expect due to the defect. In 1994, a flaw was discovered in the Intel Pentium chip. The chip would give an incorrect result when dividing two numbers. But Intel initially announced that such an error would occur in 1 in 9 billion divides. Consequently, it did not immediately offer to replace the chip. Horgan (2010) demonstrates what a bad decision that was. She shows, using the product rule for independent events, that the probability of an error can be as large as 0.28 in just 3 billion divides, which is not uncommon in many computer operations.

Calculate the probability of at least one error in 5 billion divides.

Show work. No attachments are allowed. Answer in the space provided.





## Question 9

Answer saved

Marked out of 7.00

Suppose you were going to sample the opinion of 200 registered voters in the United States as a gauge of public opinion on some issue. Would it make a significant difference whether you chose sampling with replacement or sampling without replacement? Why or why not? Explain. File attachments not allowed in this question.

↵

A ▾

B

I

≡

≡

≡

≡

↺

↻

🖼️

## Question 10

Answer saved

Marked out of 3.00

Suppose that  $X$  is a random variable with Expected value  $\mu_X$  and variance  $\sigma_X^2$ . Use rules of expectations to show what the following is equal to

$$E$$

$$(3X - \mu_{3X})$$

$$V$$

↵
A ▾
B
I
≡
≡
≡
≡
🔗
🔄
🖼️

## Question 11

Not yet answered

Marked out of 1.00

In the factories and defective part example of one of lecture 5, there are several probabilities. When calculating

$$P(B \cap D)$$

we are calculating the  probability that a part randomly chosen among the  parts has the two characteristics of being from factory  and being . On the other hand, when we are calculating

$$P(D | B)$$

we are calculating the probability that a part randomly chosen among  parts is defective. That last probability is . The

$$P(B \cap D)$$

equals  but the

$$P(D | B)$$

equals . If we were just calculating the probability that a randomly chosen part is defective, that probability is , and that is a .

To make a decision as to which factory a defective part comes from we need to compare the conditional probabilities , , . The defective part, we decide, comes from the factory that has the highest probability among those. This does not mean we know exactly where the defective part came from, but we are less likely to be wrong if we make our prediction using this method.

The example just illustrated here and in the lectures resembles de example of

defective	100	$P(A D)$	$P(B D)$	joint
total probability	$P(C D)$	Ryan Voda	Mehul Jain	0.09
a conditional probability	Noah Gardner	B	30	0.19
0.3				

## Question 12

Answer saved

Marked out of 1.00

Let

$$f(x) = 3x^2, \quad 0 \leq x \leq 1,$$

and  $f(x) = 0$  for any other value of  $x$  in the real line.The cumulative distribution function of  $X$  (cdf) is

Select one:

☐ a.

$$F(x) = x^3, \quad 0 \leq x \leq 1$$

☐ b.

$$F(x) = 2x + 1, \quad 0 \leq x \leq 1$$

☐ c.

$$F(x) = 3, \quad 0 \leq x \leq 1$$

☐ d.

$$F(x) = 6x, \quad 0 \leq x \leq 1$$

## Question 13

Not yet answered

Marked out of 4.00

According to <https://data.census.gov/cedsci/profile?q&g=0100000US>, 32.9% of the population 25 years and older in the United States have a Bachelor's Degree or Higher Education. We are going to sample from that population at random, without replacement, but being such a large population, we can assume independence. Answer the following questions:

The probability that in a sample of 300 hundred people there are at least 105 with a Bachelor's degree or higher education

The probability that we need to sample less than 15 people to find the first 3 adult with a Bachelor's degree or higher education

In calculating the probability that we need to sample less than 5 people to find the first person with a Bachelor's degree or higher education we are using a

The probability that we need to sample less than 5 people to find the first person with a Bachelor's degree or higher education

## Question 14

Answer saved

Marked out of 1.00

According to the LOTUS results seen in lecture, the expected value of  $g(X) = 100 + 2X$  equals (select what it equals) for  $X$  in the following families

$X$  lognormal, with parameters  $\mu = 10$  and  $\sigma = 2$

$X$  exponential with parameter  $\lambda$  equal 5

$X$  Poisson with parameter  $\lambda$  equal 5

## Question 15

Answer saved

Marked out of 1.00

An analog signal received at a detector (measured in microvolts) is normally distributed with a mean of 100 and a variance of 256. What is the probability that the signal will be less than 120 microvolts given that it is larger than 110 microvolts.

Select one:

- ☐ a. 0.60538
- ☐ b. 2.58
- ☐ c. 0.7324
- ☐ d. 0.2211

## Question 16

Incomplete answer

Marked out of 3.00

This is a question that requires showing work. That means, you must define your notation, must tell us what you are calculating, must show detailed work and provide final answers and explain.

*For submission of your work, you may do the work on paper and upload a pdf file. Alternatively, if you know how to use the equation editor, you may use it and write your answers in the space provided below. If you do that, the work and equations must show clearly.*

-----

X is a continuous random variable with density

$$f(x) = \frac{1}{26}(4x + 1), \quad 2 \leq x \leq 4,$$

- (a) Find the 90th percentile of X
- (b) Calculate the expected value of X
- (c) Calculate the probability that X is smaller than 3.2

↵
A ▾
B
I
☰
☷
☰
☷
🔗
🔄
🖼️

Maximum file size: 64MB, maximum number of files: 1

📄
📁
📋
📂

[Files](#)

You can drag and drop files here to add them.

Accepted file types

Document files .doc .docx .epub .gdoc .odt .oth .ott .pdf .rtf

PDF document .pdf



## Question 17

Answer saved

Marked out of 1.00

Los Angeles County had 10.4 million people in 2019, and possibly much more in 2022. That is a very large population. And yet, some individuals are called for Jury Duty pretty often, almost every 2 or 3 years. More specifically, they are called to be in a panel from which jurors are chosen. So sometimes do not get selected to be in the jury. But they have to go and be in the panel for a day nonetheless. The article in our course web site titled: "The Binomial and Hypergeometric Probability Distributions in Jury Selection" by Jude T. Sommerfeld (a copy of which is available for view for your convenience and only for this exam) talks about panels and jury selection. The article considers that it is appropriate to use the hypergeometric distribution in which of the following scenarios?

- ☐ a. Both of the cases presented in the other two choices given in this question.
- ☐ b. Calculating the probability of having 8 black persons in a jury pool of 100 people drawn from a total population of 16000 men.
- ☐ c. Calculating the probability of having 9 women in the choice of 100 potential jurors out of a jury panel of 350 people consisting of 102 women from a district's population which was 53% female.

## Question 18

Answer saved

Marked out of 1.00

The Pell grant example in Lecture 5 talks about 2 types of probability. Later on in the lecture and the material studied this week, we have also talked about different probabilities. In the Pell grant example, the 0.32 is a  and the 0.34 is a . If we had been interested in calculating the probability that a Pell grant receiver was from UCLA, we could not calculate it because we do not have the information about , which is needed to apply  theorem.

**Question 19**

Not yet answered

Marked out of 1.00

The Maxwell-Boltzman probability density function for the speed of air particles in a room does the following when the room temperature increases. Choose all that applies

- ☐ a. it shifts to the left of the speed horizontal axis
- ☐ b. Is such that the speed has a higher expected value
- ☐ c. It shifts to the right of the speed horizontal axis
- ☐ d. Is such that the speed has a lower expected value

## Question 20

Not yet answered

Marked out of 3.00

This is a question that requires showing work. That means, you must define your notation, must tell us what you are calculating, must show detailed work and provide final answers and explain.

*For submission of your work, you may do the work on paper and upload a pdf file. Alternatively, if you know how to use the equation editor, you may use it and write your answers in the space provided below. If you do that, the work and equations must show clearly.*

-----

A box contains 6 bags of Crunchy Cheetos and 3 bags of Doritos. At the currently happening DataFest at UCLA, 3 bags are going to be randomly chosen without replacement and given to the student that guesses the answer to the midnight quiz question. We are interested in the number of Cheetos bags in the sample.

(a) Write a probability distribution table that contains in one column the possible values of the random variable of interest, in another column the probability of those values, and in another the work showing the mathematical calculations that are used to calculate the probabilities in each case. Work must be shown for full credit.

(b) After you obtain the table, show how to calculate the expected value of the random variable and its standard deviation using the values in your table. Use the appropriate Greek letter symbols to refer to the expected value and the standard deviation. Say in plain words and referring to the context (Cheetos and Doritos ) of this problem to describe what the values you obtain are conveying.

(c) If you had drawn with replacement, what would the expected value have been. Use appropriate formulas for the context of this problem. You do not need to construct a table for this part.

*For submission of your work, you may do the work on paper and upload a pdf file. Alternatively, if you know how to use the equation editor, you may use it and write your answers in the space provided below. If you do that, the work and equations must show clearly.*



Maximum file size: 64MB, maximum number of files: 1

[Files](#)

You can drag and drop files here to add them.

**Question 21**

Not yet answered

Marked out of 1.00

The weight of anodized reciprocating pistons produced by Brown Company follows a normal distribution with mean 10 pounds and standard deviation 0.2 pounds.

The heaviest 2.5% of the pistons produced are rejected as overweight.

What is the probability that of 10 randomly chosen pistons at least 2 have weight larger than 10.392?

Select one:

- ☐ a. 0.0246115
- ☐ b. 0.981
- ☐ c. 0.3389
- ☐ d. 0.577

## Question 22

Not yet answered

Marked out of 3.00

This is a question that requires showing work. That means, you must define your notation, must tell us what you are calculating, must show detailed work and provide final answers and explain.

For submission of your work, you may do the work on paper and upload a pdf file. Alternatively, if you know how to use the equation editor, you may use it and write your answers in the space provided below. If you do that, the work and equations must show clearly.

-----

-----

The National Health and Nutrition Examination Survey (NHANES) examines about 5000 persons per year <https://www.cdc.gov/nchs/nhanes/index.htm> and constantly screens diabetic people to detect people with type I diabetes. Let  $X$  be a random variable measuring the number of diabetic people it takes to find the first Type I diabetic person. Assume that the probability of a diabetic person having type I diabetes is 0.05.

- (i) List five or six outcomes of the sample space. Use as notation: 1 for type 1 and 0 for not type 1 diabetic. Also provide the probability of each of these outcomes and write a small table containing the first values of  $X$  and  $P(X)$ . Indicate also what probability rule are you using to calculate the probabilities of these outcomes and what assumptions you are making to be allowed to apply the rule.
- (ii) Granted that populations are finite. But would a finite number of outcomes in this sample space suffice to guarantee that all the axioms of probability hold? Explain why or why not.
- (iii) Write in a table the cumulative probability mass function for the number of people it takes to find the first diabetic persons. Your table must have values of  $X$  in one column and the cumulative probability in the other column, with the column well labelled as cumulative probability. You may write the table for  $X=1$  to  $X=5$  and add another row to complete the table by writing the probability of  $X$  greater than 5, because you will not write a table that goes from  $X=1$  to infinity, of course. Can you find a formula for that last probability,  $P(X>5)$  in terms of the parameter of the distribution? Hint: there is a handout posted with our lecture notes that will allow you to deduce the formula from one of the formulas given there and your knowledge of probability rules.

For submission of your work, you may do the work on paper and upload a pdf file. Alternatively, if you know how to use the equation editor, you may use it and write your answers in the space provided below. If you do that, the work and equations must show clearly.



Maximum file size: 64MB, maximum number of files: 1



Files

You can drag and drop files here to add them.

## Question 23

Not yet answered

Marked out of 3.00

This is a question that requires showing work. That means, you must define your notation, must tell us what you are calculating, must show detailed work and provide final answers and explain.

No attachments are allowed in this question. You must enter your answer in the space provided.

-----

We are now in a scenario where there is a disease floating around and a diagnostic test for the disease. Something like COVID and testing for COVID. In this scenario, 1.7% of the population are True Positive, 10% of the population test positive, and 80% of the population are True Negatives. What is the specificity of this test? Use a table to demonstrate your work.



Question **24**

Not yet answered

Marked out of 1.00

The following expression

$$\int_x 2\mu_x^2 x f(x) dx$$

, where  $f(x)$  is a density function and the integration is over all the domain of the random variable  $X$ , equals

Select one:

☐ a.

$$2\mu_x^2$$

☐ b.

$$\sigma_x^2$$

☐ c.

$$2\mu_x^3$$

☐ d. 1



## Question 25

Not yet answered

Marked out of 3.00

In this question you must show work and explain your work, like in all the work questions. Attachments are not allowed.

We observed several days at the door of the UCLA bookstore how many people entering the store it took to observe a famous person that we recognized entering the textbook store. This is the data we collected.

34, 21, 10, 108, 130, 66, 21, 56, 104, 147

Use this data to estimate the probability that it will take 100 people to observe a famous one that we recognize entering the store.

Describe the steps of your calculations and justify them based on methods used in lecture. No attachments are allowed in this question. Use the space provided below.



Question **26**

Not yet answered

Marked out of 1.00

A self driving car makes a forbidden U-turn and a regular car crashes against it. Three witnesses, whom we call Bison, Rabindranath, and Tom saw what happened. Suppose the reliability of the witnesses is estimated by having the witnesses observe a number of similar scenes. It is found that Bison has probability 0.9 of stating that the self-driving car did a forbidden U-turn, Rabindranath has probability 0.8 of stating that, and Tom has probability 0.7 of stating that. Let A, B, and C denote, respectively, the events that persons Bison, Rabindranath and Tom will state that the self driving car did the forbidden U-turn. Assuming that these events are independent events, calculate the probability that all three witnesses will testify that the self-driving car did the forbidden U-turn.

- ☐ a. 0.504
- ☐ b. 0.902
- ☐ c. 0.398
- ☐ d. 0.5

**Question 27**

Not yet answered

Marked out of 1.00

According to the lognormal model fitted to radon data from Minnesota, the probability that a randomly chosen household has radon level larger than 4 is estimated as

- ☐ a. 0.574
- ☐ b. approximately 0
- ☐ c. 0.426
- ☐ d. 0.41006

## Question 28

Not yet answered

Marked out of 1.00

The Old Faithful is a famous geyser in Yellowstone (<https://www.yellowstonepark.com/things-to-do/geysers-hot-springs/about-old-faithful/>)

The geyser varies in the time it makes visitors wait for its next eruption. Using past data we calculated that it takes an average of 35 minute waiting time from the moment the last eruption occurred to a new eruption, when the last eruption was short. You arrive at the site where you can watch Old Faithful, but you arrive right at the end of a short eruption. What is the probability that you will have to wait less than 30 minutes for its next eruption?

- ☐ a. 0.57563
- ☐ b.  $1 - e^{-35}$
- ☐ c. 1
- ☐ d. 0.42437

## Question 29

Not yet answered

Marked out of 1.00

If  $X$  is uniformly distributed over  $(0,1)$ , find the density function of

$$Y = e^x$$

.

Select one:

☐ a.

$$1/y, \quad y \in [1, e]$$

☐ b. exponential with  $\lambda=1$ .

☐ c.

$$N(\mu = 1/2, \sigma^2 = 1/12)$$

Question **30**

Not yet answered

Marked out of 3.00

A box contains 6 bags of Crunchy Cheetos and 3 bags of Doritos. At the currently happening DataFest at UCLA, 3 bags are going to be randomly chosen without replacement and given to the student that guesses the answer to the midnight quiz question. Match the following probabilities

Probability that two bags are Crunchy Cheetos bags.

The expected number of Doritos bags

The probability that one bag is Crunchy Cheetos and the other are Doritos.