

## RWorksheet\_Jacildo#4a

Czharina Mae Jacildo

2024-10-18

1. The table below shows the data about shoe size and height. Create a data frame.

```
shoe_size <- c(6.5, 9.0, 8.5, 8.5, 10.5, 7.0, 9.5, 9.0, 13.0, 7.5, 10.5, 8.5, 12.0, 10.5, 13.0, 11.5, 8.5, 10.5, 13.0, 11.5, 8.5)
height <- c(66.0, 68.0, 64.5, 65.0, 70.0, 64.0, 70.0, 71.0, 72.0, 64.0, 74.5, 67.0, 71.0, 71.0, 77.0, 71.0, 77.0, 71.0, 77.0, 71.0, 77.0)
gender <- c("F", "F", "F", "F", "M", "F", "F", "F", "M", "F", "M", "F", "M", "M", "M", "M", "F", "F", "F", "F", "F")
data <- data.frame(Shoe_size = shoe_size, Height = height, Gender = gender)
print(data)
```

##	Shoe_size	Height	Gender
## 1	6.5	66.0	F
## 2	9.0	68.0	F
## 3	8.5	64.5	F
## 4	8.5	65.0	F
## 5	10.5	70.0	M
## 6	7.0	64.0	F
## 7	9.5	70.0	F
## 8	9.0	71.0	F
## 9	13.0	72.0	M
## 10	7.5	64.0	F
## 11	10.5	74.5	M
## 12	8.5	67.0	F
## 13	12.0	71.0	M
## 14	10.5	71.0	M
## 15	13.0	77.0	M
## 16	11.5	72.0	M
## 17	8.5	59.0	F
## 18	5.0	62.0	F
## 19	10.0	72.0	M
## 20	6.5	66.0	F
## 21	7.5	64.0	F
## 22	8.5	67.0	M
## 23	10.5	73.0	M
## 24	8.5	69.0	F
## 25	10.5	72.0	M
## 26	11.0	70.0	M
## 27	9.0	69.0	M
## 28	13.0	70.0	M

a. Describe the data.

- It shows the data about shoe size, height, and gender.

- b. Create a subset by males and females with their corresponding shoe size and height. What its result?  
Show the R scripts.

```
male <- subset(data, gender == "M")
female <- subset(data, gender == "F")

print(male)
```

```
##      Shoe_size Height Gender
## 5          10.5   70.0      M
## 9          13.0   72.0      M
## 11         10.5   74.5      M
## 13         12.0   71.0      M
## 14         10.5   71.0      M
## 15         13.0   77.0      M
## 16         11.5   72.0      M
## 19         10.0   72.0      M
## 22          8.5   67.0      M
## 23         10.5   73.0      M
## 25         10.5   72.0      M
## 26         11.0   70.0      M
## 27          9.0   69.0      M
## 28         13.0   70.0      M
```

```
print(female)
```

```
##      Shoe_size Height Gender
## 1           6.5   66.0      F
## 2           9.0   68.0      F
## 3           8.5   64.5      F
## 4           8.5   65.0      F
## 6           7.0   64.0      F
## 7           9.5   70.0      F
## 8           9.0   71.0      F
## 10          7.5   64.0      F
## 12          8.5   67.0      F
## 17          8.5   59.0      F
## 18          5.0   62.0      F
## 20          6.5   66.0      F
## 21          7.5   64.0      F
## 24          8.5   69.0      F
```

- c. Find the mean of shoe size and height of the respondents. Write the R scripts and its result.

```
shoe_size_mean <- mean(shoe_size)
height_mean <- mean(height)

print(shoe_size_mean)
```

```
## [1] 9.410714
```

```
print(height_mean)
```

```
## [1] 68.57143
```

- d. Is there a relationship between shoe size and height? Why?

- Yes, usually taller people have longer feet.

2. Construct character vector months to a factor with factor() and assign the result to factor\_months\_vector. Print out factor\_months\_vector and assert that R prints out the factor levels below the actual values.

```
months_vector <- c("March", "April", "January", "November", "January",
"September", "October", "September", "November", "August",
"January", "November", "November", "February", "May", "August", "July", "December", "August", "August", "September")

factor_months_vector <- factor(months_vector)
print(factor_months_vector)
```

```
## [1] March      April      January   November  January   September October
## [8] September November August     January   November  November  February
## [15] May        August    July      December  August    August    September
## [22] November  February  April
## 11 Levels: April August December February January July March May ... September
```

3. Then check the summary() of the months\_vector and factor\_months\_vector. | Interpret the results of both vectors. Are they both equally useful in this case?

```
summary_months_vector <- summary(months_vector)
summary_factor_months <- summary(factor_months_vector)

print(summary_months_vector)
```

```
##      Length      Class      Mode
##          24 character character
```

```
print(summary_factor_months)
```

```
##      April      August  December  February   January      July      March      May
##          2          4          1          2          3          1          1          1
## November  October September
##          5          1          3
```

4. Create a vector and factor for the table below.

```
direction <- c("East", "West", "North")
frequency <- c(1, 4, 3)

new_order_data <- factor(direction, levels = c("East", "West", "North"))
print(new_order_data)
```

```
## [1] East West North
## Levels: East West North
```

5. Enter the data below in Excel with file name = import\_march.csv

- a. Import the excel file into the Environment Pane using read.table() function. Write the code.

```
library(readxl)
excelData <- read.table("/cloud/project/worksheet#4/import_march.csv", header = TRUE, sep = ",")
```

- b. View the dataset. Write the R scripts and its result.

```
print(excelData)
```

```
##      Students Strategy.1 Strategy.2 Strategy.3
## 1      Male          8          10          8
## 2          4          8          6
```

## 3		0	6	4
## 4	Female	14	4	15
## 5		10	2	12
## 6		6	0	9