

# Gauss Factorials: Properties and Applications

Karl Dilcher

Dalhousie University, Halifax, Nova Scotia

Joint work with



John B. Cosgrave

Dublin, Ireland



# 1. Introduction

Recall Wilson's Theorem:

$p$  is a prime if and only if

$$(p - 1)! \equiv -1 \pmod{p}.$$

(Converse is due to Lagrange).

# 1. Introduction

Recall Wilson's Theorem:

$p$  is a prime if and only if

$$(p - 1)! \equiv -1 \pmod{p}.$$

(Converse is due to Lagrange). A proof depends on the fact that any integer  $a$  with  $1 < a < p - 1$  has its inverse  $a^{-1} \not\equiv a \pmod{p}$ .

If we write out the factorial  $(p-1)!$  and exploit symmetry modulo  $p$ , we obtain

$$1 \cdot 2 \cdot \dots \cdot \frac{p-1}{2} \frac{p+1}{2} \cdot \dots \cdot (p-1) \equiv \left(\frac{p-1}{2}\right)! (-1)^{\frac{p-1}{2}} \left(\frac{p-1}{2}\right)! \pmod{p},$$

If we write out the factorial  $(p-1)!$  and exploit symmetry modulo  $p$ , we obtain

$$1 \cdot 2 \cdot \dots \cdot \frac{p-1}{2} \cdot \frac{p+1}{2} \cdot \dots \cdot (p-1) \equiv \left(\frac{p-1}{2}\right)! (-1)^{\frac{p-1}{2}} \left(\frac{p-1}{2}\right)! \pmod{p},$$

and thus, with Wilson's Theorem,

$$\left(\frac{p-1}{2}\right)!^2 \equiv (-1)^{\frac{p+1}{2}} \pmod{p}.$$

If we write out the factorial  $(p-1)!$  and exploit symmetry modulo  $p$ , we obtain

$$1 \cdot 2 \cdot \dots \cdot \frac{p-1}{2} \frac{p+1}{2} \cdot \dots \cdot (p-1) \equiv \left(\frac{p-1}{2}\right)! (-1)^{\frac{p-1}{2}} \left(\frac{p-1}{2}\right)! \pmod{p},$$

and thus, with Wilson's Theorem,

$$\left(\frac{p-1}{2}\right)!^2 \equiv (-1)^{\frac{p+1}{2}} \pmod{p}.$$

This was apparently first observed by Lagrange (1773).



If we write out the factorial  $(p-1)!$  and exploit symmetry modulo  $p$ , we obtain

$$1 \cdot 2 \cdot \dots \cdot \frac{p-1}{2} \cdot \frac{p+1}{2} \cdot \dots \cdot (p-1) \equiv \left(\frac{p-1}{2}\right)! (-1)^{\frac{p-1}{2}} \left(\frac{p-1}{2}\right)! \pmod{p},$$

and thus, with Wilson's Theorem,

$$\left(\frac{p-1}{2}\right)!^2 \equiv (-1)^{\frac{p+1}{2}} \pmod{p}.$$

This was apparently first observed by Lagrange (1773).

For  $p \equiv 1 \pmod{4}$  the RHS is  $-1$ , so

$$\text{ord}_p \left( \left( \frac{p-1}{2} \right)! \right) = 4 \quad \text{for} \quad p \equiv 1 \pmod{4}.$$

In the case  $p \equiv 3 \pmod{4}$  we get

$$\left(\frac{p-1}{2}\right)! \equiv \pm 1 \pmod{p}.$$

In the case  $p \equiv 3 \pmod{4}$  we get

$$\left(\frac{p-1}{2}\right)! \equiv \pm 1 \pmod{p}.$$

What is the sign on the right?

In the case  $p \equiv 3 \pmod{4}$  we get

$$\left(\frac{p-1}{2}\right)! \equiv \pm 1 \pmod{p}.$$

What is the sign on the right?

### Theorem 1 (Mordell, 1961)

For a prime  $p \equiv 3 \pmod{4}$ ,

$$\left(\frac{p-1}{2}\right)! \equiv -1 \pmod{p} \iff h(-p) \equiv 1 \pmod{4},$$

where  $h(-p)$  is the class number of  $\mathbb{Q}(\sqrt{-p})$ .

In the case  $p \equiv 3 \pmod{4}$  we get

$$\left(\frac{p-1}{2}\right)! \equiv \pm 1 \pmod{p}.$$

What is the sign on the right?

### Theorem 1 (Mordell, 1961)

For a prime  $p \equiv 3 \pmod{4}$ ,

$$\left(\frac{p-1}{2}\right)! \equiv -1 \pmod{p} \iff h(-p) \equiv 1 \pmod{4},$$

where  $h(-p)$  is the class number of  $\mathbb{Q}(\sqrt{-p})$ .

Discovered independently by Chowla.

This completely determines the order  $\pmod{p}$  of  $\left(\frac{p-1}{2}\right)!$

Now consider the two halves of the product

$$1 \cdot 2 \cdots \frac{p-1}{2} \frac{p+1}{2} \cdots (p-1)$$

and denote them, respectively, by

$$\Pi_1^{(2)}, \quad \Pi_2^{(2)}.$$

Now consider the two halves of the product

$$1 \cdot 2 \cdots \frac{p-1}{2} \frac{p+1}{2} \cdots (p-1)$$

and denote them, respectively, by

$$\Pi_1^{(2)}, \quad \Pi_2^{(2)}.$$

By Wilson's theorem:

$$\Pi_1^{(2)} \Pi_2^{(2)} \equiv -1 \pmod{p},$$

Now consider the two halves of the product

$$1 \cdot 2 \cdots \frac{p-1}{2} \frac{p+1}{2} \cdots (p-1)$$

and denote them, respectively, by

$$\Pi_1^{(2)}, \quad \Pi_2^{(2)}.$$

By Wilson's theorem:

$$\Pi_1^{(2)} \Pi_2^{(2)} \equiv -1 \pmod{p},$$

and by symmetry:

$$\Pi_2^{(2)} \equiv (-1)^{\frac{p-1}{2}} \Pi_1^{(2)} \pmod{p}.$$



What can we say about the three partial products

$$\Pi_1^{(3)}, \quad \Pi_2^{(3)}, \quad \Pi_3^{(3)}$$

obtained by dividing the entire product  $(p - 1)!$  into *three* equal parts?

What can we say about the three partial products

$$\Pi_1^{(3)}, \quad \Pi_2^{(3)}, \quad \Pi_3^{(3)}$$

obtained by dividing the entire product  $(p - 1)!$  into *three* equal parts?

We require  $p \equiv 1 \pmod{3}$ ; in fact,  $p \equiv 1 \pmod{6}$ .

What can we say about the three partial products

$$\Pi_1^{(3)}, \quad \Pi_2^{(3)}, \quad \Pi_3^{(3)}$$

obtained by dividing the entire product  $(p-1)!$  into *three* equal parts?

We require  $p \equiv 1 \pmod{3}$ ; in fact,  $p \equiv 1 \pmod{6}$ .

Then

$$\Pi_1^{(3)} = 1 \cdot 2 \cdots \frac{p-1}{3}, \quad \Pi_2^{(3)} = \frac{p+2}{3} \cdots \frac{2p-2}{3}, \quad \Pi_3^{(3)} = \frac{2p+1}{3} \cdots (p-1).$$

What can we say about the three partial products

$$\Pi_1^{(3)}, \quad \Pi_2^{(3)}, \quad \Pi_3^{(3)}$$

obtained by dividing the entire product  $(p-1)!$  into *three* equal parts?

We require  $p \equiv 1 \pmod{3}$ ; in fact,  $p \equiv 1 \pmod{6}$ .

Then

$$\Pi_1^{(3)} = 1 \cdot 2 \cdots \frac{p-1}{3}, \quad \Pi_2^{(3)} = \frac{p+2}{3} \cdots \frac{2p-2}{3}, \quad \Pi_3^{(3)} = \frac{2p+1}{3} \cdots (p-1).$$

Once again there is an obvious symmetry:

$$\Pi_3^{(3)} \equiv \Pi_1^{(3)} \pmod{p},$$

(without a power of  $-1$  since  $\frac{p-1}{3}$  is always even.)

What can we say about the three partial products

$$\Pi_1^{(3)}, \quad \Pi_2^{(3)}, \quad \Pi_3^{(3)}$$

obtained by dividing the entire product  $(p-1)!$  into *three* equal parts?

We require  $p \equiv 1 \pmod{3}$ ; in fact,  $p \equiv 1 \pmod{6}$ .

Then

$$\Pi_1^{(3)} = 1 \cdot 2 \cdots \frac{p-1}{3}, \quad \Pi_2^{(3)} = \frac{p+2}{3} \cdots \frac{2p-2}{3}, \quad \Pi_3^{(3)} = \frac{2p+1}{3} \cdots (p-1).$$

Once again there is an obvious symmetry:

$$\Pi_3^{(3)} \equiv \Pi_1^{(3)} \pmod{p},$$

(without a power of  $-1$  since  $\frac{p-1}{3}$  is always even.)

No obvious relation between  $\Pi_1^{(3)}$  and the “middle third”  $\Pi_2^{(3)}$ .

Natural extension:

For any  $M \geq 2$  and for a prime  $p \equiv 1 \pmod{M}$ , divide  $(p-1)!$  into the products

Natural extension:

For any  $M \geq 2$  and for a prime  $p \equiv 1 \pmod{M}$ , divide  $(p-1)!$  into the products

$$\Pi_j^{(M)} = \prod_{i=1}^{\frac{p-1}{M}} \left( (j-1)\frac{p-1}{M} + i \right), \quad (j = 1, 2, \dots, M).$$

Natural extension:

For any  $M \geq 2$  and for a prime  $p \equiv 1 \pmod{M}$ , divide  $(p-1)!$  into the products

$$\Pi_j^{(M)} = \prod_{i=1}^{\frac{p-1}{M}} \left( (j-1)\frac{p-1}{M} + i \right), \quad (j = 1, 2, \dots, M).$$

Once again, clear that

$$\Pi_{M-j}^{(M)} \equiv \pm \Pi_j^{(M)} \pmod{p}, \quad j = 1, 2, \dots, \lfloor \frac{M-1}{2} \rfloor.$$



Natural extension:

For any  $M \geq 2$  and for a prime  $p \equiv 1 \pmod{M}$ , divide  $(p-1)!$  into the products

$$\Pi_j^{(M)} = \prod_{i=1}^{\frac{p-1}{M}} \left( (j-1)\frac{p-1}{M} + i \right), \quad (j = 1, 2, \dots, M).$$

Once again, clear that

$$\Pi_{M-j}^{(M)} \equiv \pm \Pi_j^{(M)} \pmod{p}, \quad j = 1, 2, \dots, \lfloor \frac{M-1}{2} \rfloor.$$

Example:

$p$	$\Pi_1^{(3)}$	$\Pi_2^{(3)}$	$\Pi_3^{(3)}$	$p$	$\Pi_1^{(4)}$	$\Pi_2^{(4)}$	$\Pi_3^{(4)}$	$\Pi_4^{(4)}$
7	<b>2</b>	<b>-2</b>	<b>2</b>	5	1	2	-2	-1
13	-2	3	-2	13	6	3	-3	-6
19	-2	-5	-2	17	7	-3	-3	7
31	2	-8	2	29	-6	-2	2	6
37	7	3	7	37	-16	5	-5	16
43	-3	19	-3	41	13	7	7	13
61	<b>-14</b>	<b>14</b>	<b>-14</b>	53	26	7	-7	-26
67	-20	-33	-20	61	19	7	-7	-19
73	33	-12	33	73	18	-35	-35	18
79	-37	3	-37	89	22	42	42	22
97	21	-11	21	97	20	-28	-28	20

$p$	$\Pi_1^{(3)}$	$\Pi_2^{(3)}$	$\Pi_3^{(3)}$	$p$	$\Pi_1^{(4)}$	$\Pi_2^{(4)}$	$\Pi_3^{(4)}$	$\Pi_4^{(4)}$
7	<b>2</b>	<b>-2</b>	<b>2</b>	5	1	2	-2	-1
13	-2	3	-2	13	6	3	-3	-6
19	-2	-5	-2	17	7	-3	-3	7
31	2	-8	2	29	-6	-2	2	6
37	7	3	7	37	-16	5	-5	16
43	-3	19	-3	41	13	7	7	13
61	<b>-14</b>	<b>14</b>	<b>-14</b>	53	26	7	-7	-26
67	-20	-33	-20	61	19	7	-7	-19
73	33	-12	33	73	18	-35	-35	18
79	-37	3	-37	89	22	42	42	22
97	21	-11	21	97	20	-28	-28	20

We observe:

- The obvious symmetries.

$p$	$\Pi_1^{(3)}$	$\Pi_2^{(3)}$	$\Pi_3^{(3)}$	$p$	$\Pi_1^{(4)}$	$\Pi_2^{(4)}$	$\Pi_3^{(4)}$	$\Pi_4^{(4)}$
7	<b>2</b>	<b>-2</b>	<b>2</b>	5	1	2	-2	-1
13	-2	3	-2	13	6	3	-3	-6
19	-2	-5	-2	17	7	-3	-3	7
31	2	-8	2	29	-6	-2	2	6
37	7	3	7	37	-16	5	-5	16
43	-3	19	-3	41	13	7	7	13
61	<b>-14</b>	<b>14</b>	<b>-14</b>	53	26	7	-7	-26
67	-20	-33	-20	61	19	7	-7	-19
73	33	-12	33	73	18	-35	-35	18
79	-37	3	-37	89	22	42	42	22
97	21	-11	21	97	20	-28	-28	20

We observe:

- The obvious symmetries.
- $\Pi_1^{(3)} \equiv -\Pi_2^{(3)} \pmod{p}$  for  $p = 7$  and  $p = 61$ .

Are these last cases just coincidences?

Are these last cases just coincidences?

It turns out:

$\Pi_1^{(3)} \equiv -\Pi_2^{(3)} \pmod{p}$  also for  $p = 331$ ,  $p = 547$ ,  $p = 1951$ ,  
and further relatively rare primes (explained later).

Are these last cases just coincidences?

It turns out:

$\Pi_1^{(3)} \equiv -\Pi_2^{(3)} \pmod{p}$  also for  $p = 331, p = 547, p = 1951$ ,  
and further relatively rare primes (explained later).

In contrast: No primes  $p$  for which

$$\Pi_1^{(3)} \equiv \Pi_2^{(3)} \pmod{p}, \quad p \equiv 1 \pmod{6}, \text{ or}$$

$$\Pi_1^{(4)} \equiv \pm \Pi_2^{(4)} \pmod{p}, \quad p \equiv 1 \pmod{4}.$$

Are these last cases just coincidences?

It turns out:

$\Pi_1^{(3)} \equiv -\Pi_2^{(3)} \pmod{p}$  also for  $p = 331, p = 547, p = 1951$ ,  
and further relatively rare primes (explained later).

In contrast: No primes  $p$  for which

$$\Pi_1^{(3)} \equiv \Pi_2^{(3)} \pmod{p}, \quad p \equiv 1 \pmod{6}, \text{ or}$$

$$\Pi_1^{(4)} \equiv \pm \Pi_2^{(4)} \pmod{p}, \quad p \equiv 1 \pmod{4}.$$

This will be explained later.



## 2. Composite Moduli

Define the *Gauss factorial* by

$$N_n! = \prod_{\substack{1 \leq j \leq N \\ \gcd(j,n)=1}} j.$$

## 2. Composite Moduli

Define the *Gauss factorial* by

$$N_n! = \prod_{\substack{1 \leq j \leq N \\ \gcd(j,n)=1}} j.$$

Analogue of Wilson's theorem for composite moduli:

### Theorem 2 (Gauss)

For any integer  $n \geq 2$  we have

$$(n-1)_n! \equiv \begin{cases} -1 \pmod{n} & \text{for } n = 2, 4, p^\alpha, \text{ or } 2p^\alpha, \\ 1 \pmod{n} & \text{otherwise,} \end{cases}$$

where  $p$  is an odd prime and  $\alpha$  is a positive integer.

## 2. Composite Moduli

Define the *Gauss factorial* by

$$N_n! = \prod_{\substack{1 \leq j \leq N \\ \gcd(j,n)=1}} j.$$

Analogue of Wilson's theorem for composite moduli:

### Theorem 2 (Gauss)

For any integer  $n \geq 2$  we have

$$(n-1)_n! \equiv \begin{cases} -1 \pmod{n} & \text{for } n = 2, 4, p^\alpha, \text{ or } 2p^\alpha, \\ 1 \pmod{n} & \text{otherwise,} \end{cases}$$

where  $p$  is an odd prime and  $\alpha$  is a positive integer.

The first case indicates exactly those  $n$  that have primitive roots.

Everything we did before can be extended to composite moduli.

Everything we did before can be extended to composite moduli.

E.g., the multiplicative orders of  $(\frac{n-1}{2})_n!$  modulo  $n$  were completely determined; only orders 1, 2, 4 occur. (JBC & KD, 2008).

Everything we did before can be extended to composite moduli.

E.g., the multiplicative orders of  $(\frac{n-1}{2})_n!$  modulo  $n$  were completely determined; only orders 1, 2, 4 occur. (JBC & KD, 2008).

Next, divide the product  $(n-1)_n!$  into  $M \geq 2$  partial products:

Everything we did before can be extended to composite moduli.

E.g., the multiplicative orders of  $(\frac{n-1}{2})_n!$  modulo  $n$  were completely determined; only orders 1, 2, 4 occur. (JBC & KD, 2008).

Next, divide the product  $(n-1)_n!$  into  $M \geq 2$  partial products:  
For  $n \equiv 1 \pmod{M}$ , set

$$\Pi_j^{(M)} := \prod_{i \in I_j^{(M)}} i, \quad (j = 1, 2, \dots, M),$$

where, for  $j = 1, 2, \dots, M$ ,

$$I_j^{(M)} := \left\{ i \mid (j-1)\frac{n-1}{M} + 1 \leq i \leq j\frac{n-1}{M}, \gcd(i, n) = 1 \right\}.$$

Everything we did before can be extended to composite moduli.

E.g., the multiplicative orders of  $(\frac{n-1}{2})_n!$  modulo  $n$  were completely determined; only orders 1, 2, 4 occur. (JBC & KD, 2008).

Next, divide the product  $(n-1)_n!$  into  $M \geq 2$  partial products:  
For  $n \equiv 1 \pmod{M}$ , set

$$\Pi_j^{(M)} := \prod_{i \in I_j^{(M)}} i, \quad (j = 1, 2, \dots, M),$$

where, for  $j = 1, 2, \dots, M$ ,

$$I_j^{(M)} := \left\{ i \mid (j-1)\frac{n-1}{M} + 1 \leq i \leq j\frac{n-1}{M}, \gcd(i, n) = 1 \right\}.$$

- Dependence on  $n$  is implied in the notation;



Everything we did before can be extended to composite moduli.

E.g., the multiplicative orders of  $(\frac{n-1}{2})_n!$  modulo  $n$  were completely determined; only orders 1, 2, 4 occur. (JBC & KD, 2008).

Next, divide the product  $(n-1)_n!$  into  $M \geq 2$  partial products:  
For  $n \equiv 1 \pmod{M}$ , set

$$\Pi_j^{(M)} := \prod_{i \in I_j^{(M)}} i, \quad (j = 1, 2, \dots, M),$$

where, for  $j = 1, 2, \dots, M$ ,

$$I_j^{(M)} := \left\{ i \mid (j-1)\frac{n-1}{M} + 1 \leq i \leq j\frac{n-1}{M}, \gcd(i, n) = 1 \right\}.$$

- Dependence on  $n$  is implied in the notation;
- When  $n = p$ : reduces to previous case.

Example:

$n$	$\Pi_1^{(3)}$	$\Pi_2^{(3)}$	$\Pi_3^{(3)}$	$n$	$\Pi_1^{(4)}$	$\Pi_2^{(4)}$	$\Pi_3^{(4)}$	$\Pi_4^{(4)}$
70	29	1	29	61	19	7	-7	-19
73	33	-12	33	65	<b>8</b>	<b>8</b>	<b>8</b>	<b>8</b>
76	-29	-15	-29	69	31	-26	-26	31
79	-37	3	-37	73	18	-35	-35	18
82	-33	-25	-33	77	16	31	31	16
85	-28	9	-28	81	2	40	-40	2
88	5	-7	5	85	<b>13</b>	<b>13</b>	<b>13</b>	<b>13</b>
91	<b>29</b>	<b>29</b>	<b>29</b>	89	22	42	42	22
94	-23	43	-23	93	34	-10	-10	34
97	21	-11	21	97	20	-28	-28	20

Example:

$n$	$\Pi_1^{(3)}$	$\Pi_2^{(3)}$	$\Pi_3^{(3)}$	$n$	$\Pi_1^{(4)}$	$\Pi_2^{(4)}$	$\Pi_3^{(4)}$	$\Pi_4^{(4)}$
70	29	1	29	61	19	7	-7	-19
73	33	-12	33	65	<b>8</b>	<b>8</b>	<b>8</b>	<b>8</b>
76	-29	-15	-29	69	31	-26	-26	31
79	-37	3	-37	73	18	-35	-35	18
82	-33	-25	-33	77	16	31	31	16
85	-28	9	-28	81	2	40	-40	2
88	5	-7	5	85	<b>13</b>	<b>13</b>	<b>13</b>	<b>13</b>
91	<b>29</b>	<b>29</b>	<b>29</b>	89	22	42	42	22
94	-23	43	-23	93	34	-10	-10	34
97	21	-11	21	97	20	-28	-28	20

We see: In contrast to prime case, it *can* happen that all partial products are congruent to each other.

### 3. The Distribution of Totatives

We pause to consider the *number* of elements in our subintervals:

$$\phi_{M,j}(n) := \#I_j^{(M)}.$$

(Called *totatives* by J. J. Sylvester and later D. H. Lehmer).

### 3. The Distribution of Totatives

We pause to consider the *number* of elements in our subintervals:

$$\phi_{M,j}(n) := \#I_j^{(M)}.$$

(Called *totatives* by J. J. Sylvester and later D. H. Lehmer).

- When  $n = p$  is a prime, then  $\phi_{M,j}(n) = \frac{p-1}{M}$  for all  $j$ .

### 3. The Distribution of Totatives

We pause to consider the *number* of elements in our subintervals:

$$\phi_{M,j}(n) := \#I_j^{(M)}.$$

(Called *totatives* by J. J. Sylvester and later D. H. Lehmer).

- When  $n = p$  is a prime, then  $\phi_{M,j}(n) = \frac{p-1}{M}$  for all  $j$ .
- When  $M = 1$ , then  $\phi_{1,1}(n) = \phi(n)$ .

### 3. The Distribution of Totatives

We pause to consider the *number* of elements in our subintervals:

$$\phi_{M,j}(n) := \#I_j^{(M)}.$$

(Called *totatives* by J. J. Sylvester and later D. H. Lehmer).

- When  $n = p$  is a prime, then  $\phi_{M,j}(n) = \frac{p-1}{M}$  for all  $j$ .
- When  $M = 1$ , then  $\phi_{1,1}(n) = \phi(n)$ .
- When  $M = 2$ , then  $\phi_{2,1}(n) = \phi_{2,2}(n) = \frac{1}{2}\phi(n)$ .

### 3. The Distribution of Totatives

We pause to consider the *number* of elements in our subintervals:

$$\phi_{M,j}(n) := \#I_j^{(M)}.$$

(Called *totatives* by J. J. Sylvester and later D. H. Lehmer).

- When  $n = p$  is a prime, then  $\phi_{M,j}(n) = \frac{p-1}{M}$  for all  $j$ .
- When  $M = 1$ , then  $\phi_{1,1}(n) = \phi(n)$ .
- When  $M = 2$ , then  $\phi_{2,1}(n) = \phi_{2,2}(n) = \frac{1}{2}\phi(n)$ .

In general, the situation is less straightforward.



### 3. The Distribution of Totatives

We pause to consider the *number* of elements in our subintervals:

$$\phi_{M,j}(n) := \#I_j^{(M)}.$$

(Called *totatives* by J. J. Sylvester and later D. H. Lehmer).

- When  $n = p$  is a prime, then  $\phi_{M,j}(n) = \frac{p-1}{M}$  for all  $j$ .
- When  $M = 1$ , then  $\phi_{1,1}(n) = \phi(n)$ .
- When  $M = 2$ , then  $\phi_{2,1}(n) = \phi_{2,2}(n) = \frac{1}{2}\phi(n)$ .

In general, the situation is less straightforward.

E.g., for  $n = 4$ :

$$\phi_{3,1}(n) = \phi_{3,3}(n) = 1, \text{ but } \phi_{3,2}(n) = 0.$$

When do we have equal distribution?

When do we have equal distribution?

**Theorem 3 (Lehmer, 1955)**

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least one prime factor  $p \equiv 1 \pmod{M}$ , then*

$$\phi_{M,j}(n) = \frac{1}{M} \phi(n), \quad (j = 1, 2, \dots, M).$$

When do we have equal distribution?

**Theorem 3 (Lehmer, 1955)**

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least one prime factor  $p \equiv 1 \pmod{M}$ , then*

$$\phi_{M,j}(n) = \frac{1}{M} \phi(n), \quad (j = 1, 2, \dots, M).$$

Note: Condition is sufficient, but not necessary.

When do we have equal distribution?

**Theorem 3 (Lehmer, 1955)**

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least one prime factor  $p \equiv 1 \pmod{M}$ , then*

$$\phi_{M,j}(n) = \frac{1}{M} \phi(n), \quad (j = 1, 2, \dots, M).$$

Note: Condition is sufficient, but not necessary.

E.g.,  $M = 8$  and  $n = 105 = 3 \cdot 5 \cdot 7$ .

None of the prime factors are  $\equiv 1 \pmod{8}$ , but

$$\phi_{M,j}(n) = \frac{1}{8} \phi(105) = 6 \text{ for } j = 1, \dots, 8.$$

## 4. When Are the Partial Products Congruent?

Return to our table:

$n$	$\pi_1^{(3)}$	$\pi_2^{(3)}$	$\pi_3^{(3)}$	$n$	$\pi_1^{(4)}$	$\pi_2^{(4)}$	$\pi_3^{(4)}$	$\pi_4^{(4)}$
70	29	1	29	61	19	7	-7	-19
73	33	-12	33	65	<b>8</b>	<b>8</b>	<b>8</b>	<b>8</b>
76	-29	-15	-29	69	31	-26	-26	31
79	-37	3	-37	73	18	-35	-35	18
82	-33	-25	-33	77	16	31	31	16
85	-28	9	-28	81	2	40	-40	2
88	5	-7	5	85	<b>13</b>	<b>13</b>	<b>13</b>	<b>13</b>
91	<b>29</b>	<b>29</b>	<b>29</b>	89	22	42	42	22
94	-23	43	-23	93	34	-10	-10	34
97	21	-11	21	97	20	-28	-28	20

Note:

$$91 = 7 \cdot 13, \quad 65 = 5 \cdot 13, \quad 85 = 5 \cdot 17.$$

This observation holds in general:

This observation holds in general:

#### Theorem 4

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least two distinct prime factors  $\equiv 1 \pmod{M}$ , then*

$$\Pi_j^{(M)} \equiv \left(\frac{n-1}{M}\right)_n! \pmod{n}, \quad j = 1, 2, \dots, M.$$



This observation holds in general:

#### Theorem 4

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least two distinct prime factors  $\equiv 1 \pmod{M}$ , then*

$$\Pi_j^{(M)} \equiv \left(\frac{n-1}{M}\right)_n! \pmod{n}, \quad j = 1, 2, \dots, M.$$

Result is best possible:

E.g.,  $M = 3$  and  $n = 70 = 2 \cdot 5 \cdot 7$ .

- Only one factor  $\equiv 1 \pmod{3}$ ,
- $\Pi_1^{(3)} \equiv 29 \pmod{70}$ ,  $\Pi_2^{(3)} \equiv 1 \pmod{70}$ .

This observation holds in general:

#### Theorem 4

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least two distinct prime factors  $\equiv 1 \pmod{M}$ , then*

$$\Pi_j^{(M)} \equiv \left(\frac{n-1}{M}\right)_n! \pmod{n}, \quad j = 1, 2, \dots, M.$$

Result is best possible:

E.g.,  $M = 3$  and  $n = 70 = 2 \cdot 5 \cdot 7$ .

- Only one factor  $\equiv 1 \pmod{3}$ ,
- $\Pi_1^{(3)} \equiv 29 \pmod{70}$ ,  $\Pi_2^{(3)} \equiv 1 \pmod{70}$ .

On the other hand, condition is sufficient but not necessary.

E.g.,  $M = 3$  and  $n = 2^2 \cdot 61$ ; statement still holds.

Proof is based on an observation:

$$\Pi_j^{(M)} = \frac{\left(j \frac{n-1}{M}\right)_n!}{\left((j-1) \frac{n-1}{M}\right)_n!}, \quad j = 1, 2, \dots, M,$$

Proof is based on an observation:

$$\Pi_j^{(M)} = \frac{(j^{\frac{n-1}{M}})_n!}{((j-1)^{\frac{n-1}{M}})_n!}, \quad j = 1, 2, \dots, M,$$

and a lemma:

### Lemma 5

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ ,  $n = p^\alpha q^\beta w$  for distinct prime  $p, q \equiv 1 \pmod{M}$ ,  $\alpha, \beta \geq 1$ , and  $\gcd(pq, w) = 1$ . Then for  $j = 1, 2, \dots, M$ ,*

$$(j^{\frac{n-1}{M}})_n! \equiv \frac{\varepsilon j^{\frac{p-1}{M}}}{p^{jA}} \pmod{q^\beta w}, \quad A = \frac{p^{\alpha-1}}{M} \phi(q^\beta w),$$

*where  $\varepsilon = -1$  if  $w = 1$ , and  $\varepsilon = 1$  if  $w > 1$ .*

Proof is based on an observation:

$$\prod_j^{(M)} = \frac{(j^{\frac{n-1}{M}})_n!}{((j-1)^{\frac{n-1}{M}})_n!}, \quad j = 1, 2, \dots, M,$$

and a lemma:

### Lemma 5

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ ,  $n = p^\alpha q^\beta w$  for distinct prime  $p, q \equiv 1 \pmod{M}$ ,  $\alpha, \beta \geq 1$ , and  $\gcd(pq, w) = 1$ . Then for  $j = 1, 2, \dots, M$ ,*

$$(j^{\frac{n-1}{M}})_n! \equiv \frac{\varepsilon j^{\frac{p-1}{M}}}{p^{jA}} \pmod{q^\beta w}, \quad A = \frac{p^{\alpha-1}}{M} \phi(q^\beta w),$$

*where  $\varepsilon = -1$  if  $w = 1$ , and  $\varepsilon = 1$  if  $w > 1$ .*

To prove the Theorem, use this and the Chinese Remainder Theorem; dependence on  $j$  disappears.

Idea of proof:

Idea of proof:

Break the range of the product in  $(j^{\frac{n-1}{M}})_n!$  into

- a number of products of approximately equal length,
- a shorter "tail."

Idea of proof:

Break the range of the product in  $(j^{\frac{n-1}{M}})_n!$  into

- a number of products of approximately equal length,
- a shorter "tail."

Then evaluate the products of the first type using the Gauss-Wilson theorem  $(\text{mod } q^\beta w)$ .



Idea of proof:

Break the range of the product in  $(j^{\frac{n-1}{M}})_n!$  into

- a number of products of approximately equal length,
- a shorter "tail."

Then evaluate the products of the first type using the Gauss-Wilson theorem  $(\text{mod } q^\beta w)$ .

Carefully count which elements to include/exclude.

## 5. Some Consequences

1. We saw:

If  $n$  has at least two distinct prime factors  $\equiv 1 \pmod{M}$ ,  
then the  $\Pi_j^{(M)}$  are congruent to each other.

## 5. Some Consequences

1. We saw:

If  $n$  has at least two distinct prime factors  $\equiv 1 \pmod{M}$ ,  
then the  $\Pi_j^{(M)}$  are congruent to each other.

Since their product is  $(n-1)_n!$ , we have by Gauss-Wilson,

$$\left(\frac{n-1}{M}\right)_n!^M \equiv 1 \pmod{n}.$$

## 5. Some Consequences

1. We saw:

If  $n$  has at least two distinct prime factors  $\equiv 1 \pmod{M}$ ,  
then the  $\Pi_j^{(M)}$  are congruent to each other.

Since their product is  $(n-1)_n!$ , we have by Gauss-Wilson,

$$\left(\frac{n-1}{M}\right)_n!^M \equiv 1 \pmod{n}.$$

This implies:

### Corollary 6

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least two distinct prime factors  $\equiv 1 \pmod{M}$ , then the multiplicative order of  $(\frac{n-1}{M})_n!$  modulo  $n$  is a divisor of  $M$ .*

2. In the case of at least *three* distinct prime factors  
 $\equiv 1 \pmod{M}$  we can say more:

2. In the case of at least *three* distinct prime factors  $\equiv 1 \pmod{M}$  we can say more:

### Theorem 7

Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .

If  $n$  has at least three distinct prime factors  $\equiv 1 \pmod{M}$ , then

$$\Pi_j^{(M)} \equiv 1 \pmod{n}, \quad j = 1, 2, \dots, M.$$

2. In the case of at least *three* distinct prime factors  $\equiv 1 \pmod{M}$  we can say more:

### Theorem 7

*Let  $M \geq 2$  and  $n \equiv 1 \pmod{M}$ .*

*If  $n$  has at least three distinct prime factors  $\equiv 1 \pmod{M}$ , then*

$$\prod_j^{(M)} \equiv 1 \pmod{n}, \quad j = 1, 2, \dots, M.$$

Method of proof is similar to that of the previous lemma.

Summary:

# of prime factors $\equiv 1 \pmod{M}$	All $\Pi_1^{(M)}, \dots, \Pi_M^{(M)}$ :
1	have the same number of factors
2	are congruent to each other $\pmod{M}$
3	are congruent to 1 $\pmod{M}$



## 6. The Gauss and Jacobi Theorems

Return to the question of how  $\Pi_1^{(4)}$  and  $\Pi_2^{(4)}$  are related, for prime moduli  $p \equiv 1 \pmod{4}$ .

## 6. The Gauss and Jacobi Theorems

Return to the question of how  $\Pi_1^{(4)}$  and  $\Pi_2^{(4)}$  are related, for prime moduli  $p \equiv 1 \pmod{4}$ .

Consider their quotient:

$$Q_4(p) := \frac{\Pi_2^{(4)}}{\Pi_1^{(4)}}$$

## 6. The Gauss and Jacobi Theorems

Return to the question of how  $\Pi_1^{(4)}$  and  $\Pi_2^{(4)}$  are related, for prime moduli  $p \equiv 1 \pmod{4}$ .

Consider their quotient:

$$Q_4(p) := \frac{\Pi_2^{(4)}}{\Pi_1^{(4)}} = \frac{\Pi_1^{(4)} \Pi_2^{(4)}}{\left(\Pi_1^{(4)}\right)^2}$$

## 6. The Gauss and Jacobi Theorems

Return to the question of how  $\Pi_1^{(4)}$  and  $\Pi_2^{(4)}$  are related, for prime moduli  $p \equiv 1 \pmod{4}$ .

Consider their quotient:

$$Q_4(p) := \frac{\Pi_2^{(4)}}{\Pi_1^{(4)}} = \frac{\Pi_1^{(4)} \Pi_2^{(4)}}{\left(\Pi_1^{(4)}\right)^2} = \frac{\frac{p-1}{2}!}{\left(\frac{p-1}{4}!\right)^2}$$

## 6. The Gauss and Jacobi Theorems

Return to the question of how  $\Pi_1^{(4)}$  and  $\Pi_2^{(4)}$  are related, for prime moduli  $p \equiv 1 \pmod{4}$ .

Consider their quotient:

$$Q_4(p) := \frac{\Pi_2^{(4)}}{\Pi_1^{(4)}} = \frac{\Pi_1^{(4)} \Pi_2^{(4)}}{\left(\Pi_1^{(4)}\right)^2} = \frac{\frac{p-1}{2}!}{\left(\frac{p-1}{4}!\right)^2} = \binom{\frac{p-1}{2}}{\frac{p-1}{4}}.$$

## 6. The Gauss and Jacobi Theorems

Return to the question of how  $\Pi_1^{(4)}$  and  $\Pi_2^{(4)}$  are related, for prime moduli  $p \equiv 1 \pmod{4}$ .

Consider their quotient:

$$Q_4(p) := \frac{\Pi_2^{(4)}}{\Pi_1^{(4)}} = \frac{\Pi_1^{(4)} \Pi_2^{(4)}}{(\Pi_1^{(4)})^2} = \frac{\frac{p-1}{2}!}{\left(\frac{p-1}{4}!\right)^2} = \binom{\frac{p-1}{2}}{\frac{p-1}{4}}.$$

There exists a celebrated congruence for this binomial coefficient:

Fix  $p$ ,  $a$ , and  $b$  such that

$$p \equiv 1 \pmod{4}, \quad p = a^2 + b^2, \quad a \equiv 1 \pmod{4}.$$

Fix  $p$ ,  $a$ , and  $b$  such that

$$p \equiv 1 \pmod{4}, \quad p = a^2 + b^2, \quad a \equiv 1 \pmod{4}.$$

### Theorem 8 (Gauss, 1828)

*Let  $p$  and  $a$  be as above. Then*

$$\left( \frac{\frac{p-1}{2}}{\frac{p-1}{4}} \right) \equiv 2a \pmod{p}.$$



Fix  $p$ ,  $a$ , and  $b$  such that

$$p \equiv 1 \pmod{4}, \quad p = a^2 + b^2, \quad a \equiv 1 \pmod{4}.$$

### Theorem 8 (Gauss, 1828)

*Let  $p$  and  $a$  be as above. Then*

$$\left( \frac{\frac{p-1}{2}}{\frac{p-1}{4}} \right) \equiv 2a \pmod{p}.$$

As an easy application we get

$$\pi_2^{(4)} \not\equiv \pm \pi_1^{(4)} \pmod{p} \quad \text{for all } p \equiv 1 \pmod{4}.$$

Fix  $p$ ,  $a$ , and  $b$  such that

$$p \equiv 1 \pmod{4}, \quad p = a^2 + b^2, \quad a \equiv 1 \pmod{4}.$$

### Theorem 8 (Gauss, 1828)

*Let  $p$  and  $a$  be as above. Then*

$$\left( \frac{\frac{p-1}{2}}{\frac{p-1}{4}} \right) \equiv 2a \pmod{p}.$$

As an easy application we get

$$\Pi_2^{(4)} \not\equiv \pm \Pi_1^{(4)} \pmod{p} \quad \text{for all } p \equiv 1 \pmod{4}.$$

A similar theorem, due to Jacobi (1837), implies that

$$\Pi_2^{(3)} \not\equiv \Pi_1^{(3)} \pmod{p} \quad \text{for all } p \equiv 1 \pmod{6}.$$

A number of further consequences can be derived from these classical theorems.

A number of further consequences can be derived from these classical theorems. Two samples:

### Corollary 9

*For a prime  $p \equiv 1 \pmod{6}$  we have*

$$\Pi_2^{(3)} \equiv -\Pi_1^{(3)} \pmod{p} \Leftrightarrow p = 27x^2 + 27x + 7, x \in \mathbb{Z}.$$

A number of further consequences can be derived from these classical theorems. Two samples:

### Corollary 9

*For a prime  $p \equiv 1 \pmod{6}$  we have*

$$\pi_2^{(3)} \equiv -\pi_1^{(3)} \pmod{p} \Leftrightarrow p = 27x^2 + 27x + 7, x \in \mathbb{Z}.$$

The first such primes are 7, 61 (seen earlier), 331, 547, 1951.

## Corollary 10

*Let  $p \equiv 1 \pmod{4}$ . Then*

(a)  $\frac{p-1}{4}! \equiv 1 \pmod{p}$  *only if*  $p = 5$ .

## Corollary 10

*Let  $p \equiv 1 \pmod{4}$ . Then*

- (a)  $\frac{p-1}{4}! \equiv 1 \pmod{p}$  *only if*  $p = 5$ .
- (b)  $\left(\frac{p-1}{4}!\right)^k \not\equiv -1 \pmod{p}$  *for*  $k = 1, 2, 4$ .

## Corollary 10

Let  $p \equiv 1 \pmod{4}$ . Then

- (a)  $\frac{p-1}{4}! \equiv 1 \pmod{p}$  only if  $p = 5$ .
- (b)  $\left(\frac{p-1}{4}!\right)^k \not\equiv -1 \pmod{p}$  for  $k = 1, 2, 4$ .
- (c)  $\left(\frac{p-1}{4}!\right)^8 \equiv -1 \pmod{p}$  holds for  
 $p = 17, 241, 3361, 46817, 652081, \dots$



### Corollary 10

Let  $p \equiv 1 \pmod{4}$ . Then

- (a)  $\frac{p-1}{4}! \equiv 1 \pmod{p}$  only if  $p = 5$ .
- (b)  $\left(\frac{p-1}{4}!\right)^k \not\equiv -1 \pmod{p}$  for  $k = 1, 2, 4$ .
- (c)  $\left(\frac{p-1}{4}!\right)^8 \equiv -1 \pmod{p}$  holds for  
 $p = 17, 241, 3361, 46817, 652081, \dots$

Part (c) is related to the solution of a certain Pell equation.

## 7. Extensions of Gauss' Theorem

Recall:

Theorem 11 (Gauss, 1828)

*If  $p$  and  $a$  are such that  $p = a^2 + b^2$ ,  $a \equiv 1 \pmod{4}$ , then*

$$\left( \begin{array}{c} \frac{p-1}{2} \\ \frac{p-1}{4} \end{array} \right) \equiv 2a \pmod{p}.$$

## 7. Extensions of Gauss' Theorem

Recall:

Theorem 11 (Gauss, 1828)

*If  $p$  and  $a$  are such that  $p = a^2 + b^2$ ,  $a \equiv 1 \pmod{4}$ , then*

$$\left(\frac{\frac{p-1}{2}}{\frac{p-1}{4}}\right) \equiv 2a \pmod{p}.$$

Theorem 12 (Chowla, Dwork, Evans, 1986)

*With  $p$  and  $a$  as above,*

$$\left(\frac{\frac{p-1}{2}}{\frac{p-1}{4}}\right) \equiv \left(1 + \frac{1}{2}pq_p(2)\right)\left(2a - \frac{p}{2a}\right) \pmod{p^2},$$

*where  $q_p(2) := (2^{p-1} - 1)/p$  is the Fermat quotient to base 2.*

The concept of Gauss factorial was essential in the proof of the following further extension:

### Theorem 13

*With  $p$  and  $a$  as above,*

$$\begin{aligned} \left(\frac{p-1}{2}\right) &\equiv \left(2a - \frac{p}{2a} - \frac{p^2}{8a^3}\right) \\ &\times \left(1 + \frac{1}{2}pq_p(2) + \frac{1}{8}p^2 \left(2E_{p-3} - q_p(2)^2\right)\right) \pmod{p^3}, \end{aligned}$$

*where  $E_n$  is the  $n$ th Euler number.*

The concept of Gauss factorial was essential in the proof of the following further extension:

### Theorem 13

*With  $p$  and  $a$  as above,*

$$\begin{aligned} \binom{\frac{p-1}{2}}{\frac{p-1}{4}} &\equiv \left( 2a - \frac{p}{2a} - \frac{p^2}{8a^3} \right) \\ &\times \left( 1 + \frac{1}{2}pq_p(2) + \frac{1}{8}p^2 \left( 2E_{p-3} - q_p(2)^2 \right) \right) \pmod{p^3}, \end{aligned}$$

*where  $E_n$  is the  $n$ th Euler number.*

Similar extensions were also obtained for Jacobi's theorem, concerning the binomial coefficient

$$\binom{\frac{2(p-1)}{3}}{\frac{p-1}{3}} \quad (p \equiv 1 \pmod{6}).$$

