

some tests



# Contents

1	Calc 5.1 – Area between two curves	5
---	------------------------------------	---



# Chapter 1

## Calc 5.1 – Area between two curves




Figure 1.1: area-between-curves-1

Area between  $f(x)$  &  $g(x)$ .

$A = \text{Area under } f(x) - \text{Area under } g(x).$

$$A = \int_a^b f(x)dx - \int_a^b g(x)dx.$$

$$A = \int_a^b [f(x) - g(x)] dx.$$

Notitz:

$$\begin{aligned} f(x) &\geq g(x) \\ \forall x &\in [a, b]. \\ (f(x) &\text{ is above } g(x)). \end{aligned}$$

$$\begin{aligned} \int_a^c [f(x) - 0] dx + \int_c^b [0 - f(x)] dx \\ \int_a^c f(x)dx - \int_c^b f(x)dx \end{aligned}$$




Figure 1.2: are-between-curves-2

Beispiel:

Find the area bounded above by  $y = 2x + 5$ , and bounded below by  $y = x^3$  on  $[0, 2]$

$$\begin{aligned}
 A &= \int_0^2 (2x + 5) - x^3 dx \\
 &= \int_0^2 2x + 5 - x^3 dx \\
 &= x^2 + 5x - \frac{x^4}{4} \Big|_0^2 \\
 A &= \left[ 2^2 + 5 \cdot 2 - \frac{2^4}{4} \right] - [0] \\
 A &= 10
 \end{aligned}$$

Beispiel:

Find area between  $y = x^2$  and  $y = x + 6$ .




Figure 1.3: area-between-curves-3

Notitz:

**Steps:**

1. Find x-cords of the Intersection of the curves. ( Set  $f(x) = g(x)$  )
2. Which function is on the top?  
(Pick one point for each interval)
3. set-up and solve

$$\begin{aligned}
 x + 6 &= x^2 \\
 x^2 - x - 6 &= 0 \\
 (x - 3)(x + 2) \\
 x - 3 &= 0 & x + 2 &= 0 \\
 x &= 3 & x &= -2
 \end{aligned}$$

Those are the only places  $f(x)$  and  $g(x)$  are intercepting, so those are the bounds of integration.

$$A = \int_{-2}^3.$$




Figure 1.4: figures/area-between-curves-4

$$\begin{aligned}
 A &= \int_{-2}^3 (x + 6) - (x^2) dx \Rightarrow \int_{-2}^3 x + 6 - x^2 dx \\
 &= \left[ \frac{x^2}{2} + 6x - \frac{x^3}{3} \right]_{-2}^3 = \left[ \frac{3^2}{2} + 6 \cdot 3 - \frac{3^3}{3} \right] - \left[ \frac{(-2)^2}{2} + 6(-2) - \frac{-2^3}{3} \right] \\
 &= \left[ \frac{9}{2} + 18 - 9 \right] - \left[ 2 - 12 + \frac{8}{3} \right] = \left[ \frac{9}{2} + 9 \right] - \left[ -10 + \frac{8}{3} \right] \\
 &= \frac{27}{2} - \frac{-22}{3} \rightarrow \frac{27}{2} + \frac{22}{3} = \frac{125}{6}
 \end{aligned}$$

Beispiel:

Find area bound by  $y = x^3$  and  $y = x$ .

$$\begin{aligned}
 x^3 &= x \\
 x^3 - x &= 0 \\
 x(x^2 - 1) &= 0 \\
 x(x + 1)(x - 1) &= 0 \\
 x &= 0, \quad x = 1, \quad x = -1
 \end{aligned}$$

regel

Here we test again which function is where above the other, to know, in which direction to integrate. (always the one on the top minus the one on the bottom, as shown at the start of this chapter.)




Figure 1.5: figures/area-between-curves-5

Here we add those integrals together, because we want both areas together.

$$\begin{aligned}
 A &= \int_{-1}^0 x^3 - x \, dx + \int_0^1 x - x^3 \, dx \\
 &= \left[ \frac{x^4}{4} - \frac{x^2}{2} \right]_{-1}^0 + \left[ \frac{x^2}{2} - \frac{x^4}{4} \right]_0^1 \\
 &= \left[ \left( \frac{0}{4} - \frac{0}{2} \right) - \left( \frac{(-1)^4}{4} - \frac{(-1)^2}{2} \right) \right] + \left[ \left( \frac{1^2}{2} - \frac{1^4}{4} \right) - \left( \frac{0}{2} - \frac{0}{4} \right) \right] \\
 &= \left[ 0 - \left( \frac{1}{4} - \frac{1}{2} \right) \right] + \left[ \left( \frac{1}{2} - \frac{1}{4} \right) - 0 \right] \\
 &= \frac{1}{4} + \frac{1}{4} = \frac{1}{2}
 \end{aligned}$$




Figure 1.6: figures/area-between-curves-6

In this case, the area represents the distance car 'me' is ahead of car 'you'.

This can then be solved by:

$$A = \int_0^b V_1(t) - V_2(t) \, dt.$$

Beispiel:

Find the area bound by  $x = y^2$  and  $y = x - 2$ .








Figure 1.9: figures/area-between-curves-9

$$\begin{aligned}
 A &= \int_0^1 \sqrt{x} - (-\sqrt{x}) \, dx + \int_1^4 \sqrt{x} - (x - 2) \, dx \\
 &= \int_0^1 2\sqrt{x} \, dx + \int_1^4 \sqrt{x} - x + 2 \, dx \\
 &= 2 \int_0^1 x^{\frac{1}{2}} \, dx + \int_1^4 x^{\frac{1}{2}} - x + 2 \, dx \\
 &= \left[ 2 \frac{x^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^1 + \left[ \frac{x^{\frac{3}{2}}}{\frac{3}{2}} - \frac{x^2}{2} + 2x \right]_1^4 \\
 &= \left[ \frac{4x^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^1 + \left[ \frac{2x^{\frac{3}{2}}}{3} - \frac{x^2}{2} + 2x \right]_1^4 \\
 &\quad \downarrow \text{evaluating} \quad \downarrow \\
 &= \left( \frac{4 \cdot 1 \cdot \frac{3}{2}}{3} - 0 \right) + \left[ \left( \frac{2(4)^{\frac{3}{2}}}{3} - \frac{4^2}{2} + 2 \cdot 4 \right) - \left( \frac{2}{3} - \frac{1}{2} + 2 \right) \right] \\
 &= \frac{4}{3} + \left[ \left( \frac{16}{3} - 8 + 8 \right) - \frac{13}{6} \right] \\
 &= \frac{4}{3} + \frac{16}{3} - \frac{13}{6} = \frac{27}{6} \\
 &= \frac{9}{2}
 \end{aligned}$$

Notitz:

am too tired, video stopped at 1:27:12, in the following it is also made in respect to y, now it was x, which you can see, at a headturn, 90 degrees to the right, consists of some less ints.