# The Abstraction Gap Between Mathematics and Physics: A Geometric Perspective on Dimensional Analysis

Fredrik Bierich
fredrik.bierich@gmail.com

November 10, 2026

**Abstract**

This mathematical corpus attempts to bridge the abstraction gap between mathematics and physics by questioning the concept of angles and redefining the circular functions as direct functional mappings from the physical parameters of a circular sector – namely the radius and the arc length – to its orthogonal projections. The utility of this framework is then demonstrated by modeling simple harmonic motion and uniform circular motion through these geometric functions, providing an alternative and exploratory foundation for evaluating formal logical reasoning and dimensional correctness in computational models.

## 0.1 Circular Sector

### 0.1.1 Arc Length

$$r \in \mathbb{R}^+ \tag{1}$$

$$x \in [-r, r] \tag{2}$$

$$r \mapsto \overline{OA} \tag{3}$$

$$x \mapsto \overline{OB_0} \tag{4}$$

$$r \mapsto \overline{OC_n}, n \geq 0 \tag{5}$$

$$a_n \mapsto \overline{OB_n}, n \geq 0 \tag{6}$$

$$s_n \mapsto \overline{AC_n}, n \geq 0 \tag{7}$$

$$s \mapsto \operatorname{arc} AC_0 \tag{8}$$

$$s = \lim_{n \to \infty} 2^n s_n \tag{9a}$$

$$s_n = \sqrt{r^2 + r^2 - 2ra_n} \tag{9b}$$

$$a_n = \frac{r\sqrt{r^2 - a_{n-1}^2}}{\sqrt{r^2 + r^2 - 2ra_{n-1}}}, n \geq 1 \tag{9c}$$

$$a_0 = x, x \neq r \tag{9d}$$

$$s = \lim_{n \to \infty} 2^n s_n \tag{10a}$$

$$s_n = \sqrt{2r^2 - 2ra_n} \tag{10b}$$

$$a_n = \frac{r\sqrt{r^2 - a_{n-1}^2}}{\sqrt{r^2 + r^2 - 2ra_{n-1}}}, n \geq 1 \tag{10c}$$

$$a_0 = x, x \neq r \tag{10d}$$

$$s = \lim_{n \to \infty} 2^n s_n \tag{11a}$$

$$s_n = \sqrt{2r^2 - 2ra_n} \tag{11b}$$

$$a_n = \frac{r\sqrt{r^2 - a_{n-1}^2}}{\sqrt{2r^2 - 2ra_{n-1}}}, n \geq 1 \tag{11c}$$

$$a_0 = x, x \neq r \tag{11d}$$

$$s = \lim_{n \to \infty} 2^n s_n \tag{12a}$$

$$s_n = \sqrt{2r^2 - 2ra_n} \tag{12b}$$

$$a_n = r \cdot \frac{\sqrt{r^2 - a_{n-1}^2}}{\sqrt{2r^2 - 2ra_{n-1}}}, n \geq 1 \tag{12c}$$

$$a_0 = x, x \neq r \tag{12d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{13a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{13b}$$
$$a_n = r\sqrt{\frac{r^2 - a_{n-1}^2}{2r^2 - 2ra_{n-1}}}, n \geq 1 \tag{13c}$$
$$a_0 = x, x \neq r \tag{13d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{14a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{14b}$$
$$a_n = r\sqrt{\frac{(r + a_{n-1})(r - a_{n-1})}{2r^2 - 2ra_{n-1}}}, n \geq 1 \tag{14c}$$
$$a_0 = x, x \neq r \tag{14d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{15a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{15b}$$
$$a_n = r\sqrt{\frac{(r + a_{n-1})(r - a_{n-1})}{2r(r - a_{n-1})}}, n \geq 1 \tag{15c}$$
$$a_0 = x, x \neq r \tag{15d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{16a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{16b}$$
$$a_n = r\sqrt{\frac{r + a_{n-1}}{2r}}, n \geq 1 \tag{16c}$$
$$a_0 = x \tag{16d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{17a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{17b}$$
$$a_n = r\sqrt{\frac{2r(r + a_{n-1})}{(2r)^2}}, n \geq 1 \tag{17c}$$
$$a_0 = x \tag{17d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{18a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{18b}$$
$$a_n = r\sqrt{\frac{2r^2 + 2ra_{n-1}}{(2r)^2}}, n \geq 1 \tag{18c}$$
$$a_0 = x \tag{18d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{19a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{19b}$$
$$a_n = r \cdot \frac{\sqrt{2r^2 + 2ra_{n-1}}}{\sqrt{(2r)^2}}, n \geq 1 \tag{19c}$$
$$a_0 = x \tag{19d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{20a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{20b}$$
$$a_n = r \cdot \frac{\sqrt{2r^2 + 2ra_{n-1}}}{|2r|}, n \geq 1 \tag{20c}$$
$$a_0 = x \tag{20d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{21a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{21b}$$
$$a_n = r \cdot \frac{\sqrt{2r^2 + 2ra_{n-1}}}{2r}, n \geq 1 \tag{21c}$$
$$a_0 = x \tag{21d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{22a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{22b}$$
$$a_n = \frac{r\sqrt{2r^2 + 2ra_{n-1}}}{2r}, n \geq 1 \tag{22c}$$
$$a_0 = x \tag{22d}$$

$$s = \lim_{n\to\infty} 2^n s_n \tag{23a}$$
$$s_n = \sqrt{2r^2 - 2ra_n} \tag{23b}$$
$$a_n = \frac{\sqrt{2r^2 + 2ra_{n-1}}}{2}, n \geq 1 \tag{23c}$$
$$a_0 = x \tag{23d}$$

$$s = \lim_{n\to\infty} 2^n \sqrt{2r^2 - 2ra_n} \tag{24a}$$
$$a_n = \frac{\sqrt{2r^2 + 2ra_{n-1}}}{2}, n \geq 1 \tag{24b}$$
$$a_0 = x \tag{24c}$$

#### 0.1.1.1 Quadrant

$$Q_r \stackrel{\text{def}}{=} \lim_{n\to\infty} 2^n \sqrt{2r^2 - 2ra_n} \tag{25a}$$

$$a_n = \frac{\sqrt{2r^2 + 2ra_{n-1}}}{2}, n \geq 1 \tag{25b}$$

$$a_0 = 0 \tag{25c}$$

### 0.1.2 Area

$$r \in \mathbb{R}^+ \tag{26}$$

$$s \in (0, 4Q_r] \tag{27}$$

$$r \mapsto \overline{OA} \tag{28}$$

$$s \mapsto \operatorname{arc} AC_0 \tag{29}$$

$$A \mapsto \operatorname{area} \triangledown OAC_0 \tag{30}$$

$$A = \frac{rs}{2} \tag{31}$$

### 0.1.3 Circular Functions

#### 0.1.3.1 Arccosine

$$\begin{aligned} \arccos\colon \{(r,x) \in \mathbb{R}^+ \times \mathbb{R} \mid -r \leq x \leq r\} &\to \mathbb{R} \\ (r,x) &\mapsto s \in [0, 2Q_r] \end{aligned} \tag{32}$$

$$\arccos(r,x) = \lim_{n\to\infty} 2^n \sqrt{2r^2 - 2ra_n} \tag{33a}$$

$$a_n = \frac{\sqrt{2r^2 + 2ra_{n-1}}}{2}, n \geq 1 \tag{33b}$$

$$a_0 = x \tag{33c}$$

##### 0.1.3.1.1 Scalability

$$\arccos(nr,nx) = \lim_{m\to\infty} 2^m \sqrt{2(nr)^2 - 2(nr)a_m} \tag{34a}$$

$$a_m = \frac{\sqrt{2(nr)^2 + 2(nr)a_{m-1}}}{2}, m \geq 1 \tag{34b}$$

$$a_0 = nx, n > 0 \tag{34c}$$

$$\arccos(nr,nx) = \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2(nr)a_m} \tag{35a}$$

$$a_m = \frac{\sqrt{2(nr)^2 + 2(nr)a_{m-1}}}{2}, m \geq 1 \tag{35b}$$

$$a_0 = nx, n > 0 \tag{35c}$$

$$\arccos(nr, nx) = \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2nra_m} \tag{36a}$$

$$a_m = \frac{\sqrt{2(nr)^2 + 2(nr)a_{m-1}}}{2}, m \geq 1 \tag{36b}$$

$$a_0 = nx, n > 0 \tag{36c}$$

$$\arccos(nr, nx) = \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2nra_m} \tag{37a}$$

$$a_m = \frac{\sqrt{2n^2r^2 + 2(nr)a_{m-1}}}{2}, m \geq 1 \tag{37b}$$

$$a_0 = nx, n > 0 \tag{37c}$$

$$\arccos(nr, nx) = \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2nra_m} \tag{38a}$$

$$a_m = \frac{\sqrt{2n^2r^2 + 2nra_{m-1}}}{2}, m \geq 1 \tag{38b}$$

$$a_0 = nx, n > 0 \tag{38c}$$

$$\arccos(nr, nx) = \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2nr(nb_m)} \tag{39a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2nr(nb_{m-1})}}{2}, m \geq 1 \tag{39b}$$

$$nb_0 = nx, n > 0 \tag{39c}$$

$$\arccos(nr, nx) = \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2n^2rb_m} \tag{40a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2nr(nb_{m-1})}}{2}, m \geq 1 \tag{40b}$$

$$nb_0 = nx, n > 0 \tag{40c}$$

$$\arccos(nr, nx) = \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2n^2rb_m} \tag{41a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{41b}$$

$$nb_0 = nx, n > 0 \tag{41c}$$

$$\arccos(nr, nx) = \frac{n \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2n^2rb_m}}{n} \tag{42a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{42b}$$

$$nb_0 = nx, n > 0 \tag{42c}$$

$$\arccos(nr, nx) = \frac{n \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2n^2rb_m}}{|n|} \tag{43a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{43b}$$

$$nb_0 = nx, n > 0 \tag{43c}$$

$$\arccos(nr, nx) = \frac{n \lim_{m\to\infty} 2^m \sqrt{2n^2r^2 - 2n^2rb_m}}{\sqrt{n^2}} \tag{44a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{44b}$$

$$nb_0 = nx, n > 0 \tag{44c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \cdot \frac{\sqrt{2n^2r^2 - 2n^2rb_m}}{\sqrt{n^2}} \tag{45a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{45b}$$

$$nb_0 = nx, n > 0 \tag{45c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{\frac{2n^2r^2 - 2n^2rb_m}{n^2}} \tag{46a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{46b}$$

$$nb_0 = nx, n > 0 \tag{46c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{\frac{2n^2r^2}{n^2} - \frac{2n^2rb_m}{n^2}} \tag{47a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{47b}$$

$$nb_0 = nx, n > 0 \tag{47c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - \frac{2n^2rb_m}{n^2}} \tag{48a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{48b}$$

$$nb_0 = nx, n > 0 \tag{48c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{49a}$$

$$nb_m = \frac{\sqrt{2n^2r^2 + 2n^2rb_{m-1}}}{2}, m \geq 1 \tag{49b}$$

$$nb_0 = nx, n > 0 \tag{49c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{50a}$$

$$\frac{nb_m}{n} = \frac{\left(\frac{\sqrt{2n^2r^2+2n^2rb_{m-1}}}{2}\right)}{n}, m \geq 1 \tag{50b}$$

$$nb_0 = nx, n > 0 \tag{50c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{51a}$$

$$b_m = \frac{\left(\frac{\sqrt{2n^2r^2+2n^2rb_{m-1}}}{2}\right)}{n}, m \geq 1 \tag{51b}$$

$$nb_0 = nx, n > 0 \tag{51c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{52a}$$

$$b_m = \frac{\left(\frac{\sqrt{2n^2r^2+2n^2rb_{m-1}}}{n}\right)}{2}, m \geq 1 \tag{52b}$$

$$nb_0 = nx, n > 0 \tag{52c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{53a}$$

$$b_m = \frac{\left(\frac{\sqrt{2n^2r^2+2n^2rb_{m-1}}}{|n|}\right)}{2}, m \geq 1 \tag{53b}$$

$$nb_0 = nx, n > 0 \tag{53c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{54a}$$

$$b_m = \frac{\left(\frac{\sqrt{2n^2r^2+2n^2rb_{m-1}}}{\sqrt{n^2}}\right)}{2}, m \geq 1 \tag{54b}$$

$$nb_0 = nx, n > 0 \tag{54c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{55a}$$

$$b_m = \frac{\sqrt{\frac{2n^2r^2+2n^2rb_{m-1}}{n^2}}}{2}, m \geq 1 \tag{55b}$$

$$nb_0 = nx, n > 0 \tag{55c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{56a}$$

$$b_m = \frac{\sqrt{\frac{2n^2r^2}{n^2} + \frac{2n^2rb_{m-1}}{n^2}}}{2}, m \geq 1 \tag{56b}$$

$$nb_0 = nx, n > 0 \tag{56c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{57a}$$

$$b_m = \frac{\sqrt{2r^2 + \frac{2n^2 r b_{m-1}}{n^2}}}{2}, m \geq 1 \tag{57b}$$

$$nb_0 = nx, n > 0 \tag{57c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{58a}$$

$$b_m = \frac{\sqrt{2r^2 + 2rb_{m-1}}}{2}, m \geq 1 \tag{58b}$$

$$nb_0 = nx, n > 0 \tag{58c}$$

$$\arccos(nr, nx) = n \lim_{m\to\infty} 2^m \sqrt{2r^2 - 2rb_m} \tag{59a}$$

$$b_m = \frac{\sqrt{2r^2 + 2rb_{m-1}}}{2}, m \geq 1 \tag{59b}$$

$$b_0 = x, n > 0 \tag{59c}$$

$$\arccos(nr, nx) = n \arccos(r, x), n > 0 \tag{60}$$

#### 0.1.3.1.2 Negation

$$\arccos(r, -x) + \arcsin(r, -x) = Q_r \tag{61}$$

$$\arccos(r, -x) + \arcsin(r, -x) - \arcsin(r, -x) = Q_r - \arcsin(r, -x) \tag{62}$$

$$\arccos(r, -x) = Q_r - \arcsin(r, -x) \tag{63}$$

$$\arccos(r, -x) = Q_r - (-\arcsin(r, x)) \tag{64}$$

$$\arccos(r, -x) = Q_r + \arcsin(r, x) \tag{65}$$

$$\arccos(r, -x) = Q_r + \arccos(r, x) + \arcsin(r, x) - \arccos(r, x) \tag{66}$$

$$\arccos(r, -x) = Q_r + Q_r - \arccos(r, x) \tag{67}$$

$$\arccos(r, -x) = 2Q_r - \arccos(r, x) \tag{68}$$

#### 0.1.3.1.3 Partial Derivative

$$s = \arccos(r, x) \tag{69}$$

$$\cos(r, s) = x \tag{70}$$

$$\frac{\partial \cos(r, s)}{\partial x} \cdot \frac{\partial s}{\partial x} = \frac{\partial x}{\partial x} \tag{71}$$

$$\frac{-\sin(r, s)}{r} \cdot \frac{\partial s}{\partial x} = \frac{\partial x}{\partial x} \tag{72}$$

$$\frac{-\sin(r, s)}{r} \cdot \frac{\partial s}{\partial x} = 1 \tag{73}$$

$$\frac{\left(\frac{-\sin(r,s)}{r} \cdot \frac{\partial s}{\partial x}\right)}{\left(\frac{-\sin(r,s)}{r}\right)} = \frac{1}{\left(\frac{-\sin(r,s)}{r}\right)} \tag{74}$$

$$\frac{\partial s}{\partial x} = \frac{1}{\left(\frac{-\sin(r,s)}{r}\right)} \tag{75}$$

$$\frac{\partial s}{\partial x} = \frac{r}{-\sin(r,s)} \tag{76}$$

$$\frac{\partial s}{\partial x} = \frac{-r}{-(-\sin(r,s))} \tag{77}$$

$$\frac{\partial s}{\partial x} = \frac{-r}{\sin(r,s)} \tag{78}$$

$$\frac{\partial s}{\partial x} = \frac{-r}{\sqrt{r^2 - \cos^2(r,s)}} \tag{79}$$

$$\frac{\partial \arccos(r,x)}{\partial x} = \frac{-r}{\sqrt{r^2 - x^2}}, -r < x < r \tag{80}$$

#### 0.1.3.1.4 Partial Indefinite Integral

$$\int \arccos(r,x)\ \partial x = x\arccos(r,x) - \int x \cdot \frac{\partial \arccos(r,x)}{\partial x}\ \partial x, -r < x < r \tag{81}$$

$$\int \arccos(r,x)\ \partial x = x\arccos(r,x) - \int x \cdot \frac{-r}{\sqrt{r^2 - x^2}}\ \partial x, -r < x < r \tag{82}$$

$$\int \arccos(r,x)\ \partial x = x\arccos(r,x) - \int \frac{-rx}{\sqrt{r^2 - x^2}}\ \partial x, -r < x < r \tag{83}$$

$$\int \arccos(r,x)\ \partial x = x\arccos(r,x) - r\sqrt{r^2 - x^2} + C(r), -r \leq x \leq r \tag{84}$$

### 0.1.3.2 Cosine

$$\begin{aligned} \cos\colon \mathbb{R}^+ \times \mathbb{R} &\to \mathbb{R} \\ (r,s) &\mapsto x \in [-r,r] \end{aligned} \tag{85}$$

$$s = \lim_{n\to\infty} 2^n \sqrt{2r^2 - 2ra_n} \tag{86a}$$

$$a_n = \frac{\sqrt{2r^2 + 2ra_{n-1}}}{2}, n \geq 1 \tag{86b}$$

$$a_0 = x \tag{86c}$$

$$m \in \mathbb{N}_0 \tag{87a}$$

$$s \approx 2^m \sqrt{2r^2 - 2rb_{m,m}} \tag{87b}$$

$$b_{m,n} = \frac{\sqrt{2r^2 + 2rb_{m,n-1}}}{2}, 1 \leq n \leq m \tag{87c}$$

$$b_{m,0} = x \tag{87d}$$

$$m \in \mathbb{N}_0 \tag{88a}$$
$$x \approx b_{m,m} \tag{88b}$$
$$b_{m,n-1} = \frac{\sqrt{2r^2 + 2rb_{m,n}}}{2}, 1 \leq n \leq m \tag{88c}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s \tag{88d}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{89a}$$
$$b_{m,n-1} = \frac{\sqrt{2r^2 + 2rb_{m,n}}}{2}, 1 \leq n \leq m \tag{89b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{89c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{90a}$$
$$2b_{m,n-1} = 2 \cdot \frac{\sqrt{2r^2 + 2rb_{m,n}}}{2}, 1 \leq n \leq m \tag{90b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{90c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{91a}$$
$$2b_{m,n-1} = \frac{2\sqrt{2r^2 + 2rb_{m,n}}}{2}, 1 \leq n \leq m \tag{91b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{91c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{92a}$$
$$2b_{m,n-1} = \sqrt{2r^2 + 2rb_{m,n}}, 1 \leq n \leq m \tag{92b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{92c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{93a}$$
$$(2b_{m,n-1})^2 = \left(\sqrt{2r^2 + 2rb_{m,n}}\right)^2, 1 \leq n \leq m \tag{93b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{93c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{94a}$$
$$2^2 b_{m,n-1}^2 = \left(\sqrt{2r^2 + 2rb_{m,n}}\right)^2, 1 \leq n \leq m \tag{94b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{94c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{95a}$$
$$4b_{m,n-1}^2 = \left(\sqrt{2r^2 + 2rb_{m,n}}\right)^2, 1 \leq n \leq m \tag{95b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{95c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{96a}$$
$$4b_{m,n-1}^2 = 2r^2 + 2rb_{m,n}, 1 \leq n \leq m \tag{96b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{96c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{97a}$$
$$4b_{m,n-1}^2 - 2r^2 = 2r^2 + 2rb_{m,n} - 2r^2, 1 \leq n \leq m \tag{97b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{97c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{98a}$$
$$4b_{m,n-1}^2 - 2r^2 = 2rb_{m,n}, 1 \leq n \leq m \tag{98b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{98c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{99a}$$
$$\frac{4b_{m,n-1}^2 - 2r^2}{2r} = \frac{2rb_{m,n}}{2r}, 1 \leq n \leq m \tag{99b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{99c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{100a}$$
$$\frac{4b_{m,n-1}^2}{2r} - \frac{2r^2}{2r} = \frac{2rb_{m,n}}{2r}, 1 \leq n \leq m \tag{100b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{100c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{101a}$$
$$\frac{2b_{m,n-1}^2}{r} - \frac{2r^2}{2r} = \frac{2rb_{m,n}}{2r}, 1 \leq n \leq m \tag{101b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{101c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{102a}$$
$$\frac{2b_{m,n-1}^2}{r} - r = \frac{2rb_{m,n}}{2r}, 1 \leq n \leq m \tag{102b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{102c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{103a}$$
$$\frac{2b_{m,n-1}^2}{r} - r = b_{m,n}, 1 \leq n \leq m \tag{103b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{103c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{104a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{104b}$$
$$2^m\sqrt{2r^2 - 2rb_{m,0}} = s, m \geq 0 \tag{104c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{105a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{105b}$$
$$\frac{2^m\sqrt{2r^2 - 2rb_{m,0}}}{2^m} = \frac{s}{2^m}, m \geq 0 \tag{105c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{106a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{106b}$$
$$\sqrt{2r^2 - 2rb_{m,0}} = \frac{s}{2^m}, m \geq 0 \tag{106c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{107a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{107b}$$
$$\left(\sqrt{2r^2 - 2rb_{m,0}}\right)^2 = \left(\frac{s}{2^m}\right)^2, m \geq 0 \tag{107c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{108a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{108b}$$
$$2r^2 - 2rb_{m,0} = \left(\frac{s}{2^m}\right)^2, m \geq 0 \tag{108c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{109a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{109b}$$
$$2r^2 - 2rb_{m,0} = \frac{s^2}{\left(2^m\right)^2}, m \geq 0 \tag{109c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{110a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{110b}$$
$$2r^2 - 2rb_{m,0} = \frac{s^2}{2^{2m}}, m \geq 0 \tag{110c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{111a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{111b}$$
$$2r^2 - 2rb_{m,0} - 2r^2 = \frac{s^2}{2^{2m}} - 2r^2, m \geq 0 \tag{111c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{112a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{112b}$$
$$-2rb_{m,0} = \frac{s^2}{2^{2m}} - 2r^2, m \geq 0 \tag{112c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{113a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{113b}$$
$$\frac{-2rb_{m,0}}{-2r} = \frac{\left(\frac{s^2}{2^{2m}}\right) - 2r^2}{-2r}, m \geq 0 \tag{113c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{114a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{114b}$$
$$b_{m,0} = \frac{\left(\frac{s^2}{2^{2m}}\right) - 2r^2}{-2r}, m \geq 0 \tag{114c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{115a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{115b}$$
$$b_{m,0} = \frac{\left(\frac{s^2}{2^{2m}}\right)}{-2r} - \frac{2r^2}{-2r}, m \geq 0 \tag{115c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{116a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{116b}$$
$$b_{m,0} = \frac{s^2}{2^{2m}(-2r)} - \frac{2r^2}{-2r}, m \geq 0 \tag{116c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{117a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \leq n \leq m \tag{117b}$$
$$b_{m,0} = \frac{s^2}{-2^{2m+1}r} - \frac{2r^2}{-2r}, m \geq 0 \tag{117c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{118a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \le n \le m \tag{118b}$$
$$b_{m,0} = \frac{s^2}{-2^{2m+1}r} + \frac{-2r^2}{-2r}, m \ge 0 \tag{118c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{119a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \le n \le m \tag{119b}$$
$$b_{m,0} = \frac{s^2}{-2^{2m+1}r} + r, m \ge 0 \tag{119c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{120a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \le n \le m \tag{120b}$$
$$b_{m,0} = r + \frac{s^2}{-2^{2m+1}r}, m \ge 0 \tag{120c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{121a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \le n \le m \tag{121b}$$
$$b_{m,0} = r - \left(-\frac{s^2}{-2^{2m+1}r}\right), m \ge 0 \tag{121c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{122a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \le n \le m \tag{122b}$$
$$b_{m,0} = r - \left(\frac{-s^2}{-2^{2m+1}r}\right), m \ge 0 \tag{122c}$$

$$x = \lim_{m\to\infty} b_{m,m} \tag{123a}$$
$$b_{m,n} = \frac{2b_{m,n-1}^2}{r} - r, 1 \le n \le m \tag{123b}$$
$$b_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{123c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{124a}$$
$$r\hat{b}_{m,n} = \frac{2(r\hat{b}_{m,n-1})^2}{r} - r, 1 \le n \le m \tag{124b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{124c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{125a}$$
$$r\hat{b}_{m,n} = \frac{2r^2\hat{b}^2_{m,n-1}}{r} - r, 1 \le n \le m \tag{125b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{125c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{126a}$$
$$r\hat{b}_{m,n} = 2r\hat{b}^2_{m,n-1} - r, 1 \le n \le m \tag{126b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{126c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{127a}$$
$$\frac{r\hat{b}_{m,n}}{r} = \frac{2r\hat{b}^2_{m,n-1} - r}{r}, 1 \le n \le m \tag{127b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{127c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{128a}$$
$$\hat{b}_{m,n} = \frac{2r\hat{b}^2_{m,n-1} - r}{r}, 1 \le n \le m \tag{128b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{128c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{129a}$$
$$\hat{b}_{m,n} = \frac{2r\hat{b}^2_{m,n-1}}{r} - \frac{r}{r}, 1 \le n \le m \tag{129b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{129c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{130a}$$
$$\hat{b}_{m,n} = 2\hat{b}^2_{m,n-1} - \frac{r}{r}, 1 \le n \le m \tag{130b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{130c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{131a}$$
$$\hat{b}_{m,n} = 2\hat{b}^2_{m,n-1} - 1, 1 \le n \le m \tag{131b}$$
$$r\hat{b}_{m,0} = r - \frac{s^2}{2^{2m+1}r}, m \ge 0 \tag{131c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{132a}$$
$$\hat{b}_{m,n} = 2\hat{b}_{m,n-1}^2 - 1, 1 \leq n \leq m \tag{132b}$$
$$\frac{r\hat{b}_{m,0}}{r} = \frac{r - \left(\frac{s^2}{2^{2m+1}r}\right)}{r}, m \geq 0 \tag{132c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{133a}$$
$$\hat{b}_{m,n} = 2\hat{b}_{m,n-1}^2 - 1, 1 \leq n \leq m \tag{133b}$$
$$\hat{b}_{m,0} = \frac{r - \left(\frac{s^2}{2^{2m+1}r}\right)}{r}, m \geq 0 \tag{133c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{134a}$$
$$\hat{b}_{m,n} = 2\hat{b}_{m,n-1}^2 - 1, 1 \leq n \leq m \tag{134b}$$
$$\hat{b}_{m,0} = \frac{r}{r} - \frac{\left(\frac{s^2}{2^{2m+1}r}\right)}{r}, m \geq 0 \tag{134c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{135a}$$
$$\hat{b}_{m,n} = 2\hat{b}_{m,n-1}^2 - 1, 1 \leq n \leq m \tag{135b}$$
$$\hat{b}_{m,0} = 1 - \frac{\left(\frac{s^2}{2^{2m+1}r}\right)}{r}, m \geq 0 \tag{135c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{136a}$$
$$\hat{b}_{m,n} = 2\hat{b}_{m,n-1}^2 - 1, 1 \leq n \leq m \tag{136b}$$
$$\hat{b}_{m,0} = 1 - \frac{s^2}{2^{2m+1}r^2}, m \geq 0 \tag{136c}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{137a}$$
$$\hat{b}_{m,n} = T_2^{\circ n}\left(1 - \frac{s^2}{2^{2m+1}r^2}\right), 0 \leq n \leq m \tag{137b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{138a}$$
$$\hat{b}_{m,n} = T_{2^n}\left(1 - \frac{s^2}{2^{2m+1}r^2}\right), 0 \leq n \leq m \tag{138b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{139a}$$
$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{T_{2^n}^{(k)}(x_0)}{k!}\left(\left(1 - \frac{s^2}{2^{2m+1}r^2}\right) - x_0\right)^k\Bigg|_{x_0=1}, 0 \leq n \leq m \tag{139b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{140a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{T_{2^n}^{(k)}(1)}{k!} \left( \left( 1 - \frac{s^2}{2^{2m+1}r^2} \right) - 1 \right)^k, 0 \le n \le m \tag{140b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{141a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{2^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!} \left( \left( 1 - \frac{s^2}{2^{2m+1}r^2} \right) - 1 \right)^k, 0 \le n \le m \tag{141b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{142a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{2^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!} \left( 1 - \frac{s^2}{2^{2m+1}r^2} - 1 \right)^k, 0 \le n \le m \tag{142b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{143a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{2^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!} \left( -\frac{s^2}{2^{2m+1}r^2} \right)^k, 0 \le n \le m \tag{143b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{144a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{2^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!} \left( \frac{-s^2}{2^{2m+1}r^2} \right)^k, 0 \le n \le m \tag{144b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{145a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{2^k \left( \frac{-s^2}{2^{2m+1}r^2} \right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{145b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{146a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left( 2 \cdot \frac{-s^2}{2^{2m+1}r^2} \right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{146b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{147a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left( \frac{-2s^2}{2^{2m+1}r^2} \right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{147b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{148a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left(\frac{-s^2}{2^{2m}r^2}\right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{148b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{149a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left(\frac{-\left(\frac{s^2}{r^2}\right)}{2^{2m}}\right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{149b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{150a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left(\frac{-\left(\frac{s}{r}\right)^2}{2^{2m}}\right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{150b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{151a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left(\frac{-\left(\frac{s}{r}\right)^2}{(2^2)^m}\right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{151b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{152a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left(\frac{-\left(\frac{s}{r}\right)^2}{4^m}\right)^k \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{152b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{153a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left(-\left(\frac{s}{r}\right)^2\right)^k}{(4^m)^k} \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{153b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{154a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\frac{(-1)^k\left(\left(\frac{s}{r}\right)^2\right)^k}{(4^m)^k} \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{154b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{155a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(4^m)^k} \prod_{j=0}^{k-1}((2^n)^2 - j^2)}{(2k)!}, 0 \le n \le m \tag{155b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{156a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{\left( \frac{(-1)^k \left(\frac{s}{r}\right)^{2k} \prod_{j=0}^{k-1} ((2^n)^2 - j^2)}{(4^m)^k} \right)}{(2k)!}, 0 \leq n \leq m \tag{156b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{157a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k} \prod_{j=0}^{k-1} ((2^n)^2 - j^2)}{(4^m)^k (2k)!}, 0 \leq n \leq m \tag{157b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{158a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k} \prod_{j=0}^{k-1} ((2^n)^2 - j^2)}{(2k)!(4^m)^k}, 0 \leq n \leq m \tag{158b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{159a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \frac{\prod_{j=0}^{k-1} ((2^n)^2 - j^2)}{(4^m)^k}, 0 \leq n \leq m \tag{159b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{160a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \frac{\prod_{j=0}^{k-1} ((2^n)^2 - j^2)}{\prod_{j=0}^{k-1} 4^m}, 0 \leq n \leq m \tag{160b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{161a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \frac{(2^n)^2 - j^2}{4^m}, 0 \leq n \leq m \tag{161b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{162a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \frac{2^{2n} - j^2}{4^m}, 0 \leq n \leq m \tag{162b}$$

$$x = \lim_{m\to\infty} r\hat{b}_{m,m} \tag{163a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \frac{(2^2)^n - j^2}{4^m}, 0 \leq n \leq m \tag{163b}$$

$$x = \lim_{m \to \infty} r \hat{b}_{m,m} \tag{164a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \frac{4^n - j^2}{4^m}, 0 \leq n \leq m \tag{164b}$$

$$x = \lim_{m \to \infty} r \hat{b}_{m,m} \tag{165a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \left( \frac{4^n}{4^m} - \frac{j^2}{4^m} \right), 0 \leq n \leq m \tag{165b}$$

$$x = \lim_{m \to \infty} r \hat{b}_{m,m} \tag{166a}$$

$$\hat{b}_{m,n} = \sum_{k=0}^{2^n} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \left( 4^{n-m} - \frac{j^2}{4^m} \right), 0 \leq n \leq m \tag{166b}$$

$$x = \lim_{m \to \infty} r \sum_{k=0}^{2^m} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \left( 4^{m-m} - \frac{j^2}{4^m} \right) \tag{167}$$

$$x = \lim_{m \to \infty} r \sum_{k=0}^{2^m} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \left( 4^0 - \frac{j^2}{4^m} \right) \tag{168}$$

$$x = \lim_{m \to \infty} r \sum_{k=0}^{2^m} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \prod_{j=0}^{k-1} \left( 1 - \frac{j^2}{4^m} \right) \tag{169}$$

$$0 \leq j \leq k - 1 < 2^m \tag{170}$$

$$0 \leq j < 2^m \tag{171}$$

$$0^2 \leq j^2 < (2^m)^2 \tag{172}$$

$$0 \leq j^2 < (2^m)^2 \tag{173}$$

$$0 \leq j^2 < 2^{2m} \tag{174}$$

$$0 \leq j^2 < (2^2)^m \tag{175}$$

$$0 \leq j^2 < 4^m \tag{176}$$

$$\frac{0}{4^m} \leq \frac{j^2}{4^m} < \frac{4^m}{4^m} \tag{177}$$

$$0 \leq \frac{j^2}{4^m} < \frac{4^m}{4^m} \tag{178}$$

$$0 \leq \frac{j^2}{4^m} < 1 \tag{179}$$

$$-0 \geq -\frac{j^2}{4^m} > -1 \tag{180}$$

$$0 \geq -\frac{j^2}{4^m} > -1 \tag{181}$$

$$0+1 \geq -\frac{j^2}{4^m}+1 > -1+1 \tag{182}$$

$$1 \geq -\frac{j^2}{4^m}+1 > -1+1 \tag{183}$$

$$1 \geq 1-\frac{j^2}{4^m} > -1+1 \tag{184}$$

$$1 \geq 1-\frac{j^2}{4^m} > 0 \tag{185}$$

$$0 < 1-\frac{j^2}{4^m} \leq 1 \tag{186}$$

$$0 < \prod_{j=0}^{k-1}\left(1-\frac{j^2}{4^m}\right) \leq 1 \tag{187}$$

$$\left|\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right| = \frac{\left|(-1)^k\left(\frac{s}{r}\right)^{2k}\right|}{|(2k)!|} \tag{188}$$

$$\left|\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right| = \frac{\left|(-1)^k\right|\left|\left(\frac{s}{r}\right)^{2k}\right|}{|(2k)!|} \tag{189}$$

$$\left|\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right| = \frac{\left|\left(\frac{s}{r}\right)^{2k}\right|}{|(2k)!|} \tag{190}$$

$$\left|\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right| = \frac{\left(\frac{s}{r}\right)^{2k}}{|(2k)!|} \tag{191}$$

$$\left|\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right| = \frac{\left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{192}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{\left(\frac{-\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| \tag{193}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{\left(\frac{-\left(\frac{s}{r}\right)^{2k+2}}{(2(k+1))!}\right)}{\left(\frac{\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| \tag{194}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^k\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{\left(\frac{-\left(\frac{s}{r}\right)^{2k+2}}{(2k+2)!}\right)}{\left(\frac{\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| \tag{195}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{\left(\frac{-\left(\frac{s}{r}\right)^{2k+2}}{(2k+2)(2k+1)(2k)!}\right)}{\left(\frac{\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| \tag{196}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{\left(\frac{-(2k)!\left(\frac{s}{r}\right)^{2k+2}}{(2k+2)(2k+1)(2k)!}\right)}{\left(\frac{s}{r}\right)^{2k}}\right| \tag{197}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{\left(\frac{-\left(\frac{s}{r}\right)^{2k+2}}{(2k+2)(2k+1)}\right)}{\left(\frac{s}{r}\right)^{2k}}\right| \tag{198}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{-\left(\frac{s}{r}\right)^{2k+2-2k}}{(2k+2)(2k+1)}\right| \tag{199}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \left|\frac{-\left(\frac{s}{r}\right)^{2}}{(2k+2)(2k+1)}\right| \tag{200}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \frac{\left|-\left(\frac{s}{r}\right)^{2}\right|}{|(2k+2)(2k+1)|} \tag{201}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \frac{\left(\frac{s}{r}\right)^{2}}{|(2k+2)(2k+1)|} \tag{202}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \frac{\left(\frac{s}{r}\right)^{2}}{|(2k+2)|\,|(2k+1)|} \tag{203}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \frac{\left(\frac{s}{r}\right)^{2}}{(2k+2)\,|(2k+1)|} \tag{204}$$

$$\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \frac{\left(\frac{s}{r}\right)^{2}}{(2k+2)(2k+1)} \tag{205}$$

$$\lim_{k\to\infty}\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = \lim_{k\to\infty}\frac{\left(\frac{s}{r}\right)^{2}}{(2k+2)(2k+1)} \tag{206}$$

$$\lim_{k\to\infty}\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| = 0 \tag{207}$$

$$\lim_{k\to\infty}\left|\frac{\left(\frac{(-1)^{k+1}\left(\frac{s}{r}\right)^{2(k+1)}}{(2(k+1))!}\right)}{\left(\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\right)}\right| < 1 \tag{208}$$

$$\sum_{k=0}^{\infty}\frac{\left(\frac{s}{r}\right)^{2k}}{(2k)!} < \infty \tag{209}$$

$$\left|\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\prod_{j=0}^{k-1}\left(1-\frac{j^{2}}{4^{m}}\right)\right| \leq \frac{\left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{210}$$

$$x = r\sum_{k=0}^{\infty}\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\lim_{m\to\infty}\prod_{j=0}^{k-1}\left(1-\frac{j^{2}}{4^{m}}\right) \tag{211}$$

$$x = r\sum_{k=0}^{\infty}\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\prod_{j=0}^{k-1}(1-0) \tag{212}$$

$$x = r\sum_{k=0}^{\infty}\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\prod_{j=0}^{k-1}1 \tag{213}$$

$$x = r\sum_{k=0}^{\infty}\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\cdot 1^{k} \tag{214}$$

$$x = r\sum_{k=0}^{\infty}\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!}\cdot 1 \tag{215}$$

$$x = r\sum_{k=0}^{\infty}\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{216}$$

$$\cos(r,s) = r\sum_{k=0}^{\infty}\frac{(-1)^{k}\left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{217}$$

#### 0.1.3.2.1 Scalability

$$\cos(nr, ns) = nr \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{ns}{nr}\right)^{2k}}{(2k)!}, n > 0 \tag{218}$$

$$\cos(nr, ns) = nr \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!}, n > 0 \tag{219}$$

$$\cos(nr, ns) = n \cos(r, s), n > 0 \tag{220}$$

#### 0.1.3.2.2 Negation

$$\cos(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{-s}{r}\right)^{2k}}{(2k)!} \tag{221}$$

$$\cos(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^k \left(-\frac{s}{r}\right)^{2k}}{(2k)!} \tag{222}$$

$$\cos(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^k (-1)^{2k} \left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{223}$$

$$\cos(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^{k+2k} \left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{224}$$

$$\cos(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^{3k} \left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{225}$$

$$\cos(r, -s) = r \sum_{k=0}^{\infty} \frac{((-1)^3)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{226}$$

$$\cos(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{227}$$

$$\cos(r, -s) = \cos(r, s) \tag{228}$$

#### 0.1.3.2.3 Subtraction

$$\sin^2(r,\alpha-\beta)+(r-\cos(r,\alpha-\beta))^2=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2 \tag{229}$$

$$\cos^2(r,\alpha-\beta)+\sin^2(r,\alpha-\beta)+(r-\cos(r,\alpha-\beta))^2-\cos^2(r,\alpha-\beta)=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2 \tag{230}$$

$$r^2+(r-\cos(r,\alpha-\beta))^2-\cos^2(r,\alpha-\beta)=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2 \tag{231}$$

$$r^2+r^2-2r\cos(r,\alpha-\beta)+\cos^2(r,\alpha-\beta)-\cos^2(r,\alpha-\beta)=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2 \tag{232}$$

$$2r^2-2r\cos(r,\alpha-\beta)+\cos^2(r,\alpha-\beta)-\cos^2(r,\alpha-\beta)=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2 \tag{233}$$

$$2r^2-2r\cos(r,\alpha-\beta)=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2 \tag{234}$$

$$2r^2-2r\cos(r,\alpha-\beta)-2r^2=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2-2r^2 \tag{235}$$

$$-2r\cos(r,\alpha-\beta)=(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2-2r^2 \tag{236}$$

$$\frac{-2r\cos(r,\alpha-\beta)}{-2r}=\frac{(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2-2r^2}{-2r} \tag{237}$$

$$\cos(r,\alpha-\beta)=\frac{(\cos(r,\alpha)-\cos(r,\beta))^2+(\sin(r,\alpha)-\sin(r,\beta))^2-2r^2}{-2r} \tag{238}$$

$$\cos(r,\alpha-\beta)=\frac{\cos^2(r,\alpha)-2\cos(r,\alpha)\cos(r,\beta)+\cos^2(r,\beta)+(\sin(r,\alpha)-\sin(r,\beta))^2-2r^2}{-2r} \tag{239}$$

$$\cos(r,\alpha-\beta)=\frac{\cos^2(r,\alpha)-2\cos(r,\alpha)\cos(r,\beta)+\cos^2(r,\beta)+\sin^2(r,\alpha)-2\sin(r,\alpha)\sin(r,\beta)+\sin^2(r,\beta)-2r^2}{-2r} \tag{240}$$

$$\cos(r,\alpha-\beta)=\frac{(\cos^2(r,\alpha)+\sin^2(r,\alpha))-2\cos(r,\alpha)\cos(r,\beta)+\cos^2(r,\beta)-2\sin(r,\alpha)\sin(r,\beta)+\sin^2(r,\beta)-2r^2}{-2r} \tag{241}$$

$$\cos(r,\alpha-\beta)=\frac{(\cos^2(r,\alpha)+\sin^2(r,\alpha))-2\cos(r,\alpha)\cos(r,\beta)+(\cos^2(r,\beta)+\sin^2(r,\beta))-2\sin(r,\alpha)\sin(r,\beta)-2r^2}{-2r} \tag{242}$$

$$\cos(r,\alpha-\beta)=\frac{r^2-2\cos(r,\alpha)\cos(r,\beta)+(\cos^2(r,\beta)+\sin^2(r,\beta))-2\sin(r,\alpha)\sin(r,\beta)-2r^2}{-2r} \tag{243}$$

$$\cos(r,\alpha-\beta)=\frac{r^2-2\cos(r,\alpha)\cos(r,\beta)+r^2-2\sin(r,\alpha)\sin(r,\beta)-2r^2}{-2r} \tag{244}$$

$$\cos(r, \alpha - \beta) = \frac{2r^2 - 2\cos(r, \alpha)\cos(r, \beta) - 2\sin(r, \alpha)\sin(r, \beta) - 2r^2}{-2r} \tag{245}$$

$$\cos(r, \alpha - \beta) = \frac{-2\cos(r, \alpha)\cos(r, \beta) - 2\sin(r, \alpha)\sin(r, \beta)}{-2r} \tag{246}$$

$$\cos(r, \alpha - \beta) = \frac{-2\left(\cos(r, \alpha)\cos(r, \beta) + \sin(r, \alpha)\sin(r, \beta)\right)}{-2r} \tag{247}$$

$$\cos(r, \alpha - \beta) = \frac{\cos(r, \alpha)\cos(r, \beta) + \sin(r, \alpha)\sin(r, \beta)}{r} \tag{248}$$

#### 0.1.3.2.4 Addition

$$\cos(r, \alpha + \beta) = \cos(r, \alpha - (-\beta)) \tag{249}$$

$$\cos(r, \alpha + \beta) = \frac{\cos(r, \alpha)\cos(r, -\beta) + \sin(r, \alpha)\sin(r, -\beta)}{r} \tag{250}$$

$$\cos(r, \alpha + \beta) = \frac{\cos(r, \alpha)\cos(r, \beta) + \sin(r, \alpha)\sin(r, -\beta)}{r} \tag{251}$$

$$\cos(r, \alpha + \beta) = \frac{\cos(r, \alpha)\cos(r, \beta) + \sin(r, \alpha)(-\sin(r, \beta))}{r} \tag{252}$$

$$\cos(r, \alpha + \beta) = \frac{\cos(r, \alpha)\cos(r, \beta) - \sin(r, \alpha)\sin(r, \beta)}{r} \tag{253}$$

#### 0.1.3.2.5 Periodicity

$$\cos(r, s + Q_r) = \frac{\cos(r, s)\cos(r, Q_r) - \sin(r, s)\sin(r, Q_r)}{r} \tag{254}$$

$$\cos(r, s + Q_r) = \frac{\cos(r, s) \cdot 0 - \sin(r, s)\sin(r, Q_r)}{r} \tag{255}$$

$$\cos(r, s + Q_r) = \frac{0 - \sin(r, s)\sin(r, Q_r)}{r} \tag{256}$$

$$\cos(r, s + Q_r) = \frac{-\sin(r, s)\sin(r, Q_r)}{r} \tag{257}$$

$$\cos(r, s + Q_r) = \frac{-\sin(r, s) \cdot r}{r} \tag{258}$$

$$\cos(r, s + Q_r) = -\sin(r, s) \tag{259}$$

$$\cos(r, s + Q_r + Q_r) = -\sin(r, s + Q_r) \tag{260}$$

$$\cos(r, s + 2Q_r) = -\sin(r, s + Q_r) \tag{261}$$

$$\cos(r, s + 2Q_r) = -\frac{\sin(r, s)\cos(r, Q_r) + \cos(r, s)\sin(r, Q_r)}{r} \tag{262}$$

$$\cos(r, s + 2Q_r) = -\frac{\sin(r, s) \cdot 0 + \cos(r, s)\sin(r, Q_r)}{r} \tag{263}$$

$$\cos(r, s + 2Q_r) = -\frac{0 + \cos(r, s)\sin(r, Q_r)}{r} \tag{264}$$

$$\cos(r, s + 2Q_r) = -\frac{\cos(r, s)\sin(r, Q_r)}{r} \tag{265}$$

$$\cos(r, s + 2Q_r) = -\frac{\cos(r, s) \cdot r}{r} \tag{266}$$

$$\cos(r, s + 2Q_r) = \frac{-\cos(r, s) \cdot r}{r} \tag{267}$$

$$\cos(r, s + 2Q_r) = -\cos(r, s) \tag{268}$$

$$\cos(r, s + 2Q_r + Q_r) = -\cos(r, s + Q_r) \tag{269}$$

$$\cos(r, s + 3Q_r) = -\cos(r, s + Q_r) \tag{270}$$

$$\cos(r, s + 3Q_r) = -(-\sin(r, s)) \tag{271}$$

$$\cos(r, s + 3Q_r) = \sin(r, s) \tag{272}$$

$$\cos(r, s + 3Q_r + Q_r) = \sin(r, s + Q_r) \tag{273}$$

$$\cos(r, s + 4Q_r) = \sin(r, s + Q_r) \tag{274}$$

$$\cos(r, s + 4Q_r) = \frac{\sin(r, s)\cos(r, Q_r) + \cos(r, s)\sin(r, Q_r)}{r} \tag{275}$$

$$\cos(r, s + 4Q_r) = \frac{\sin(r, s) \cdot 0 + \cos(r, s)\sin(r, Q_r)}{r} \tag{276}$$

$$\cos(r, s + 4Q_r) = \frac{0 + \cos(r, s)\sin(r, Q_r)}{r} \tag{277}$$

$$\cos(r, s + 4Q_r) = \frac{\cos(r, s)\sin(r, Q_r)}{r} \tag{278}$$

$$\cos(r, s + 4Q_r) = \frac{\cos(r, s) \cdot r}{r} \tag{279}$$

$$\cos(r, s + 4Q_r) = \cos(r, s) \tag{280}$$

#### 0.1.3.2.6 Partial Derivative

$$\frac{\partial \cos(r, s)}{\partial s} = \frac{\partial \left( r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \right)}{\partial s} \tag{281}$$

$$\frac{\partial \cos(r, s)}{\partial s} = \frac{\partial \left( r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \frac{s^{2k}}{r^{2k}}}{(2k)!} \right)}{\partial s} \tag{282}$$

$$\frac{\partial \cos(r, s)}{\partial s} = \frac{\partial \left( r \sum\limits_{k=0}^{\infty} \frac{(-1)^k s^{2k}}{r^{2k}(2k)!} \right)}{\partial s} \tag{283}$$

$$\frac{\partial \cos(r, s)}{\partial s} = \frac{\partial \left( \sum\limits_{k=0}^{\infty} \frac{(-1)^k r s^{2k}}{r^{2k}(2k)!} \right)}{\partial s} \tag{284}$$

$$\frac{\partial \cos(r, s)}{\partial s} = \frac{\partial \left( \sum\limits_{k=0}^{\infty} \frac{(-1)^k s^{2k}}{r^{2k-1}(2k)!} \right)}{\partial s} \tag{285}$$

$$\frac{\partial \cos(r, s)}{\partial s} = \sum_{k=0}^{\infty} \frac{(-1)^k \frac{\partial s^{2k}}{\partial s}}{r^{2k-1}(2k)!} \tag{286}$$

$$\frac{\partial \cos(r, s)}{\partial s} = \sum_{k=1}^{\infty} \frac{(-1)^k (2k) s^{2k-1}}{r^{2k-1}(2k)!} \tag{287}$$

$$\frac{\partial \cos(r, s)}{\partial s} = \sum_{k=1}^{\infty} \frac{(-1)^k (2k) \frac{s^{2k-1}}{r^{2k-1}}}{(2k)!} \tag{288}$$

$$\frac{\partial \cos(r,s)}{\partial s} = \sum_{k=1}^{\infty} \frac{(-1)^k (2k) \left(\frac{s}{r}\right)^{2k-1}}{(2k)!} \tag{289}$$

$$\frac{\partial \cos(r,s)}{\partial s} = \sum_{k=1}^{\infty} \frac{(-1)^k (2k) \left(\frac{s}{r}\right)^{2k-1}}{(2k)(2k-1)!} \tag{290}$$

$$\frac{\partial \cos(r,s)}{\partial s} = \sum_{k=1}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k-1}}{(2k-1)!} \tag{291}$$

$$\frac{\partial \cos(r,s)}{\partial s} = \sum_{k=1}^{\infty} \frac{-(-1)^{k-1} \left(\frac{s}{r}\right)^{2k-1}}{(2k-1)!} \tag{292}$$

$$\frac{\partial \cos(r,s)}{\partial s} = -\sum_{k=1}^{\infty} \frac{(-1)^{k-1} \left(\frac{s}{r}\right)^{2k-1}}{(2k-1)!} \tag{293}$$

$$\frac{\partial \cos(r,s)}{\partial s} = -\sum_{k=1}^{\infty} \frac{(-1)^{k-1} \left(\frac{s}{r}\right)^{2(k-1)+1}}{(2k-1)!} \tag{294}$$

$$\frac{\partial \cos(r,s)}{\partial s} = -\sum_{k=1}^{\infty} \frac{(-1)^{k-1} \left(\frac{s}{r}\right)^{2(k-1)+1}}{(2(k-1)+1)!} \tag{295}$$

$$\frac{\partial \cos(r,s)}{\partial s} = -\sum_{k=0}^{\infty} \frac{(-1)^{k} \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{296}$$

$$\frac{\partial \cos(r,s)}{\partial s} = \frac{-r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!}}{r} \tag{297}$$

$$\frac{\partial \cos(r,s)}{\partial s} = \frac{-\sin(r,s)}{r} \tag{298}$$

#### 0.1.3.2.7 Partial Indefinite Integral

$$\int \cos(r,s)\ \partial s = \int \frac{r \cos(r,s)}{r}\ \partial s \tag{299}$$

$$\int \cos(r,s)\ \partial s = \int r \cdot \frac{\cos(r,s)}{r}\ \partial s \tag{300}$$

$$\int \cos(r,s)\ \partial s = r \int \frac{\cos(r,s)}{r}\ \partial s \tag{301}$$

$$\int \cos(r,s)\ \partial s = r \int \frac{\partial \sin(r,s)}{\partial s}\ \partial s \tag{302}$$

$$\int \cos(r,s)\ \partial s = r \sin(r,s) + C(r) \tag{303}$$

#### 0.1.3.3 Arcsine

$$\begin{aligned}\arcsin\colon \{(r,y) \in \mathbb{R}^+ \times \mathbb{R} \mid -r \leq y \leq r\} &\to \mathbb{R} \\ (r,y) &\mapsto s \in [-Q_r, Q_r]\end{aligned} \tag{304}$$

$$\arcsin(r,y) = \frac{\arcsin(r,y) + \arcsin(r,y)}{2} \tag{305}$$

$$\arcsin(r,y) = \frac{\arcsin(r,y) + \arcsin(r,y) + \arccos(r,y) - \arccos(r,y)}{2} \tag{306}$$

$$\arcsin(r,y) = \frac{\arcsin(r,y) + \arcsin(r,-y) + \arccos(r,-y) - \arccos(r,y)}{2} \tag{307}$$

$$\arcsin(r,y) = \frac{\arcsin(r,y) - \arcsin(r,y) + \arccos(r,-y) - \arccos(r,y)}{2} \tag{308}$$

$$\arcsin(r,y) = \frac{\arccos(r,-y) - \arccos(r,y)}{2} \tag{309}$$

##### 0.1.3.3.1 Scalability

$$\arcsin(nr,ny) = \frac{\arccos(nr,-ny) - \arccos(nr,ny)}{2}, n > 0 \tag{310}$$

$$\arcsin(nr,ny) = \frac{\arccos(nr,n(-y)) - \arccos(nr,ny)}{2}, n > 0 \tag{311}$$

$$\arcsin(nr,ny) = \frac{n\arccos(r,-y) - \arccos(nr,ny)}{2}, n > 0 \tag{312}$$

$$\arcsin(nr,ny) = \frac{n\arccos(r,-y) - n\arccos(r,y)}{2}, n > 0 \tag{313}$$

$$\arcsin(nr,ny) = \frac{n(\arccos(r,-y) - \arccos(r,y))}{2}, n > 0 \tag{314}$$

$$\arcsin(nr,ny) = n \cdot \frac{\arccos(r,-y) - \arccos(r,y)}{2}, n > 0 \tag{315}$$

$$\arcsin(nr,ny) = n\arcsin(r,y), n > 0 \tag{316}$$

##### 0.1.3.3.2 Negation

$$s = \arcsin(r,-y) \tag{317}$$

$$\sin(r,s) = -y \tag{318}$$

$$-\sin(r,s) = -(-y) \tag{319}$$

$$\sin(r,-s) = -(-y) \tag{320}$$

$$\sin(r,-s) = y \tag{321}$$

$$-s = \arcsin(r,y) \tag{322}$$

$$-(-s) = -\arcsin(r,y) \tag{323}$$

$$s = -\arcsin(r,y) \tag{324}$$

$$\arcsin(r,-y) = -\arcsin(r,y) \tag{325}$$

#### 0.1.3.3.3 Partial Derivative

$$s = \arcsin(r, y) \tag{326}$$

$$\sin(r, s) = y \tag{327}$$

$$\frac{\partial \sin(r, s)}{\partial y} \cdot \frac{\partial s}{\partial y} = \frac{\partial y}{\partial y} \tag{328}$$

$$\frac{\cos(r, s)}{r} \cdot \frac{\partial s}{\partial y} = \frac{\partial y}{\partial y} \tag{329}$$

$$\frac{\cos(r, s)}{r} \cdot \frac{\partial s}{\partial y} = 1 \tag{330}$$

$$\frac{\left(\frac{\cos(r,s)}{r} \cdot \frac{\partial s}{\partial y}\right)}{\left(\frac{\cos(r,s)}{r}\right)} = \frac{1}{\left(\frac{\cos(r,s)}{r}\right)} \tag{331}$$

$$\frac{\partial s}{\partial y} = \frac{1}{\left(\frac{\cos(r,s)}{r}\right)} \tag{332}$$

$$\frac{\partial s}{\partial y} = \frac{r}{\cos(r, s)} \tag{333}$$

$$\frac{\partial s}{\partial y} = \frac{r}{\sqrt{r^2 - \sin^2(r, s)}} \tag{334}$$

$$\frac{\partial \arcsin(r, y)}{\partial y} = \frac{r}{\sqrt{r^2 - y^2}}, -r < y < r \tag{335}$$

#### 0.1.3.3.4 Partial Indefinite Integral

$$\int \arcsin(r, y) \, \partial y = y \arcsin(r, y) - \int y \cdot \frac{\partial \arcsin(r, y)}{\partial y} \, \partial y, -r < y < r \tag{336}$$

$$\int \arcsin(r, y) \, \partial y = y \arcsin(r, y) - \int y \cdot \frac{r}{\sqrt{r^2 - y^2}} \, \partial y, -r < y < r \tag{337}$$

$$\int \arcsin(r, y) \, \partial y = y \arcsin(r, y) - \int \frac{ry}{\sqrt{r^2 - y^2}} \, \partial y, -r < y < r \tag{338}$$

$$\int \arcsin(r, y) \, \partial y = y \arcsin(r, y) + \int -\frac{ry}{\sqrt{r^2 - y^2}} \, \partial y, -r < y < r \tag{339}$$

$$\int \arcsin(r, y) \, \partial y = y \arcsin(r, y) + \int \frac{-ry}{\sqrt{r^2 - y^2}} \, \partial y, -r < y < r \tag{340}$$

$$\int \arcsin(r, y) \, \partial y = y \arcsin(r, y) + r\sqrt{r^2 - y^2} + C(r), -r \leq y \leq r \tag{341}$$

#### 0.1.3.4 Sine

$$\begin{aligned} \sin\colon \mathbb{R}^+ \times \mathbb{R} &\to \mathbb{R} \\ (r, s) &\mapsto y \in [-r, r] \end{aligned} \tag{342}$$

$$(\alpha, \beta) \in \mathbb{R}^2 \tag{343}$$

$$\cos(r, \alpha - \beta) = r \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha-\beta}{r}\right)^{2k}}{(2k)!} \tag{344}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} \frac{(-1)^k \left(\frac{\alpha-\beta}{r}\right)^{2k}}{(2k)!} \tag{345}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} \frac{(-1)^k \left(\frac{\alpha}{r} - \frac{\beta}{r}\right)^{2k}}{(2k)!} \tag{346}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} \frac{(-1)^k \left(\frac{\alpha}{r} + \frac{-\beta}{r}\right)^{2k}}{(2k)!} \tag{347}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} \frac{(-1)^k \sum_{j=0}^{2k} \binom{2k}{j} \left(\frac{\alpha}{r}\right)^{2k-j} \left(\frac{-\beta}{r}\right)^{j}}{(2k)!} \tag{348}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} \frac{(-1)^k \sum_{j=0}^{2k} \frac{(2k)!}{j!(2k-j)!} \left(\frac{\alpha}{r}\right)^{2k-j} \left(\frac{-\beta}{r}\right)^{j}}{(2k)!} \tag{349}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} \frac{(-1)^k \sum_{j=0}^{2k} (2k)! \frac{\left(\frac{\alpha}{r}\right)^{2k-j} \left(\frac{-\beta}{r}\right)^{j}}{j!(2k-j)!}}{(2k)!} \tag{350}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} \frac{(-1)^k (2k)! \sum_{j=0}^{2k} \frac{\left(\frac{\alpha}{r}\right)^{2k-j} \left(\frac{-\beta}{r}\right)^{j}}{j!(2k-j)!}}{(2k)!} \tag{351}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} (-1)^k \sum_{j=0}^{2k} \frac{\left(\frac{\alpha}{r}\right)^{2k-j} \left(\frac{-\beta}{r}\right)^{j}}{j!(2k-j)!} \tag{352}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} (-1)^k \sum_{j=0}^{2k} \frac{\left(\frac{\alpha}{r}\right)^{2k-j} \left(\frac{-\beta}{r}\right)^{j}}{(2k-j)!j!} \tag{353}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \sum_{k=0}^{n} (-1)^k \sum_{j=0}^{2k} \frac{\left(\frac{\alpha}{r}\right)^{2k-j}}{(2k-j)!} \frac{\left(\frac{-\beta}{r}\right)^{j}}{j!} \tag{354}$$

$$\cos(r, \alpha - \beta) = r \lim_{n\to\infty} \left( \sum_{k=0}^{n} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k}}{(2k)!} \sum_{j=0}^{n-k} \frac{(-1)^j \left(\frac{\beta}{r}\right)^{2j}}{(2j)!} + \sum_{k=0}^{n-1} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!} \sum_{j=0}^{n-1-k} \frac{(-1)^j \left(\frac{\beta}{r}\right)^{2j+1}}{(2j+1)!} \right) \tag{355}$$

$$\cos(r, \alpha - \beta) = r \left( \lim_{n\to\infty} \left( \sum_{k=0}^{n} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k}}{(2k)!} \sum_{j=0}^{n-k} \frac{(-1)^j \left(\frac{\beta}{r}\right)^{2j}}{(2j)!} \right) + \lim_{n\to\infty} \left( \sum_{k=0}^{n-1} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!} \sum_{j=0}^{n-1-k} \frac{(-1)^j \left(\frac{\beta}{r}\right)^{2j+1}}{(2j+1)!} \right) \right) \tag{356}$$

$$\cos(r, \alpha - \beta) = r \left( \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k}}{(2k)!} \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k}}{(2k)!} + \lim_{n\to\infty} \left( \sum_{k=0}^{n-1} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!} \sum_{j=0}^{n-1-k} \frac{(-1)^j \left(\frac{\beta}{r}\right)^{2j+1}}{(2j+1)!} \right) \right) \tag{357}$$

$$\cos(r, \alpha - \beta) = r \left( \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k}}{(2k)!} \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k}}{(2k)!} + \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!} \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!} \right) \tag{358}$$

$$\cos(r, \alpha - \beta) = r \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k}}{(2k)!} \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k}}{(2k)!} + r \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!} \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!} \tag{359}$$

$$\cos(r, \alpha - \beta) = \frac{r \left( r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k}}{(2k)!} \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k}}{(2k)!} + r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!} \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!} \right)}{r} \tag{360}$$

$$\cos(r, \alpha - \beta) = \frac{r^2 \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k}}{(2k)!} \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k}}{(2k)!} + r^2 \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!} \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!}}{r} \tag{361}$$

$$\cos(r,\alpha-\beta)=\frac{\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k}}{(2k)!}\right)\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k}}{(2k)!}\right)+r^2\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!}\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!}}{r} \tag{362}$$

$$\cos(r,\alpha-\beta)=\frac{\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k}}{(2k)!}\right)\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k}}{(2k)!}\right)+\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!}\right)\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!}\right)}{r} \tag{363}$$

$$\cos(r,\alpha-\beta)=\frac{\cos(r,\alpha)\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k}}{(2k)!}\right)+\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!}\right)\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!}\right)}{r} \tag{364}$$

$$\cos(r,\alpha-\beta)=\frac{\cos(r,\alpha)\cos(r,\beta)+\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!}\right)\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!}\right)}{r} \tag{365}$$

$$\frac{\cos(r,\alpha)\cos(r,\beta)+\sin(r,\alpha)\sin(r,\beta)}{r}=\frac{\cos(r,\alpha)\cos(r,\beta)+\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!}\right)\left(r\sum\limits_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!}\right)}{r} \tag{366}$$

$$\sin(r,\alpha)\sin(r,\beta)=\left(r\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\alpha}{r}\right)^{2k+1}}{(2k+1)!}\right)\left(r\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{\beta}{r}\right)^{2k+1}}{(2k+1)!}\right) \tag{367}$$

$$\sin(r,s)\overset{\{\alpha,\beta\}\mapsto s}{=}r\sum_{k=0}^{\infty}\frac{(-1)^k\left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{368}$$

#### 0.1.3.4.1 Scalability

$$\sin(nr, ns) = nr \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{ns}{nr}\right)^{2k+1}}{(2k+1)!}, n > 0 \tag{369}$$

$$\sin(nr, ns) = nr \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!}, n > 0 \tag{370}$$

$$\sin(nr, ns) = n \sin(r, s), n > 0 \tag{371}$$

#### 0.1.3.4.2 Negation

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{-s}{r}\right)^{2k+1}}{(2k+1)!} \tag{372}$$

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^k \left(-\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{373}$$

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^k (-1)^{2k+1} \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{374}$$

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^{k+2k+1} \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{375}$$

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{(-1)^{3k+1} \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{376}$$

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{-(-1)^{3k} \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{377}$$

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{-((-1)^3)^k \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{378}$$

$$\sin(r, -s) = r \sum_{k=0}^{\infty} \frac{-(-1)^k \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{379}$$

$$\sin(r, -s) = -r \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \tag{380}$$

$$\sin(r, -s) = -\sin(r, s) \tag{381}$$

#### 0.1.3.4.3 Addition/Subtraction

$$\sin(r, \alpha \pm \beta) = \cos(r, Q_r - (\alpha \pm \beta)) \tag{382}$$

$$\sin(r, \alpha \pm \beta) = \cos(r, Q_r - \alpha \mp \beta) \tag{383}$$

$$\sin(r, \alpha \pm \beta) = \cos(r, (Q_r - \alpha) \mp \beta) \tag{384}$$

$$\sin(r, \alpha \pm \beta) = \frac{\cos(r, Q_r - \alpha)\cos(r, \beta) \pm \sin(r, Q_r - \alpha)\sin(r, \beta)}{r} \tag{385}$$

$$\sin(r, \alpha \pm \beta) = \frac{\sin(r, \alpha)\cos(r, \beta) \pm \sin(r, Q_r - \alpha)\sin(r, \beta)}{r} \tag{386}$$

$$\sin(r, \alpha \pm \beta) = \frac{\sin(r, \alpha)\cos(r, \beta) \pm \cos(r, \alpha)\sin(r, \beta)}{r} \tag{387}$$

#### 0.1.3.4.4 Periodicity

$$\sin(r, s + Q_r) = \frac{\sin(r, s)\cos(r, Q_r) + \cos(r, s)\sin(r, Q_r)}{r} \tag{388}$$

$$\sin(r, s + Q_r) = \frac{\sin(r, s) \cdot 0 + \cos(r, s)\sin(r, Q_r)}{r} \tag{389}$$

$$\sin(r, s + Q_r) = \frac{0 + \cos(r, s)\sin(r, Q_r)}{r} \tag{390}$$

$$\sin(r, s + Q_r) = \frac{\cos(r, s)\sin(r, Q_r)}{r} \tag{391}$$

$$\sin(r, s + Q_r) = \frac{\cos(r, s) \cdot r}{r} \tag{392}$$

$$\sin(r, s + Q_r) = \cos(r, s) \tag{393}$$

$$\sin(r, s + Q_r + Q_r) = \cos(r, s + Q_r) \tag{394}$$

$$\sin(r, s + 2Q_r) = \cos(r, s + Q_r) \tag{395}$$

$$\sin(r, s + 2Q_r) = \frac{\cos(r, s)\cos(r, Q_r) - \sin(r, s)\sin(r, Q_r)}{r} \tag{396}$$

$$\sin(r, s + 2Q_r) = \frac{\cos(r, s) \cdot 0 - \sin(r, s)\sin(r, Q_r)}{r} \tag{397}$$

$$\sin(r, s + 2Q_r) = \frac{0 - \sin(r, s)\sin(r, Q_r)}{r} \tag{398}$$

$$\sin(r, s + 2Q_r) = \frac{-\sin(r, s)\sin(r, Q_r)}{r} \tag{399}$$

$$\sin(r, s + 2Q_r) = \frac{-\sin(r, s) \cdot r}{r} \tag{400}$$

$$\sin(r, s + 2Q_r) = -\sin(r, s) \tag{401}$$

$$\sin(r, s + 2Q_r + Q_r) = -\sin(r, s + Q_r) \tag{402}$$

$$\sin(r, s + 3Q_r) = -\sin(r, s + Q_r) \tag{403}$$

$$\sin(r, s + 3Q_r) = -\cos(r, s) \tag{404}$$

$$\sin(r, s + 3Q_r + Q_r) = -\cos(r, s + Q_r) \tag{405}$$

$$\sin(r, s + 4Q_r) = -\cos(r, s + Q_r) \tag{406}$$

$$\sin(r, s + 4Q_r) = -\frac{\cos(r, s)\cos(r, Q_r) - \sin(r, s)\sin(r, Q_r)}{r} \tag{407}$$

$$\sin(r, s + 4Q_r) = -\frac{\cos(r, s) \cdot 0 - \sin(r, s)\sin(r, Q_r)}{r} \tag{408}$$

$$\sin(r, s + 4Q_r) = -\frac{0 - \sin(r, s)\sin(r, Q_r)}{r} \tag{409}$$

$$\sin(r, s + 4Q_r) = -\frac{-\sin(r, s)\sin(r, Q_r)}{r} \tag{410}$$

$$\sin(r, s + 4Q_r) = -\frac{-\sin(r, s) \cdot r}{r} \tag{411}$$

$$\sin(r, s + 4Q_r) = \frac{-(-\sin(r, s) \cdot r)}{r} \tag{412}$$

$$\sin(r, s + 4Q_r) = \frac{-(-\sin(r, s)) \cdot r}{r} \tag{413}$$

$$\sin(r, s + 4Q_r) = \frac{\sin(r, s) \cdot r}{r} \tag{414}$$

$$\sin(r, s + 4Q_r) = \sin(r, s) \tag{415}$$

#### 0.1.3.4.5 Partial Derivative

$$\frac{\partial \sin(r, s)}{\partial s} = \frac{\partial \left( r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k+1}}{(2k+1)!} \right)}{\partial s} \tag{416}$$

$$\frac{\partial \sin(r, s)}{\partial s} = \frac{\partial \left( r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \frac{s^{2k+1}}{r^{2k+1}}}{(2k+1)!} \right)}{\partial s} \tag{417}$$

$$\frac{\partial \sin(r, s)}{\partial s} = \frac{\partial \left( r \sum\limits_{k=0}^{\infty} \frac{(-1)^k s^{2k+1}}{r^{2k+1}(2k+1)!} \right)}{\partial s} \tag{418}$$

$$\frac{\partial \sin(r, s)}{\partial s} = \frac{\partial \left( \sum\limits_{k=0}^{\infty} \frac{(-1)^k r s^{2k+1}}{r^{2k+1}(2k+1)!} \right)}{\partial s} \tag{419}$$

$$\frac{\partial \sin(r, s)}{\partial s} = \frac{\partial \left( \sum\limits_{k=0}^{\infty} \frac{(-1)^k s^{2k+1}}{r^{2k}(2k+1)!} \right)}{\partial s} \tag{420}$$

$$\frac{\partial \sin(r, s)}{\partial s} = \sum_{k=0}^{\infty} \frac{(-1)^k \frac{\partial s^{2k+1}}{\partial s}}{r^{2k}(2k+1)!} \tag{421}$$

$$\frac{\partial \sin(r, s)}{\partial s} = \sum_{k=0}^{\infty} \frac{(-1)^k (2k+1) s^{2k}}{r^{2k}(2k+1)!} \tag{422}$$

$$\frac{\partial \sin(r,s)}{\partial s} = \sum_{k=0}^{\infty} \frac{(-1)^k (2k+1) \frac{s^{2k}}{r^{2k}}}{(2k+1)!} \tag{423}$$

$$\frac{\partial \sin(r,s)}{\partial s} = \sum_{k=0}^{\infty} \frac{(-1)^k (2k+1) \left(\frac{s}{r}\right)^{2k}}{(2k+1)!} \tag{424}$$

$$\frac{\partial \sin(r,s)}{\partial s} = \sum_{k=0}^{\infty} \frac{(-1)^k (2k+1) \left(\frac{s}{r}\right)^{2k}}{(2k+1)(2k)!} \tag{425}$$

$$\frac{\partial \sin(r,s)}{\partial s} = \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!} \tag{426}$$

$$\frac{\partial \sin(r,s)}{\partial s} = \frac{r \sum\limits_{k=0}^{\infty} \frac{(-1)^k \left(\frac{s}{r}\right)^{2k}}{(2k)!}}{r} \tag{427}$$

$$\frac{\partial \sin(r,s)}{\partial s} = \frac{\cos(r,s)}{r} \tag{428}$$

#### 0.1.3.4.6 Partial Indefinite Integral

$$\int \sin(r,s)\ \partial s = \int \frac{r \sin(r,s)}{r}\ \partial s \tag{429}$$

$$\int \sin(r,s)\ \partial s = \int \frac{-(-r \sin(r,s))}{r}\ \partial s \tag{430}$$

$$\int \sin(r,s)\ \partial s = \int \frac{-r(-\sin(r,s))}{r}\ \partial s \tag{431}$$

$$\int \sin(r,s)\ \partial s = \int -r \cdot \frac{-\sin(r,s)}{r}\ \partial s \tag{432}$$

$$\int \sin(r,s)\ \partial s = -r \int \frac{-\sin(r,s)}{r}\ \partial s \tag{433}$$

$$\int \sin(r,s)\ \partial s = -r \int \frac{\partial \cos(r,s)}{\partial s}\ \partial s \tag{434}$$

$$\int \sin(r,s)\ \partial s = -r \cos(r,s) + C(r) \tag{435}$$





Figure 1: Circular sector.

## 0.2 Circular Sector Sequence

### 0.2.1 Apothem

$$\text{area}\,\triangle OAC_{n-1} = \frac{\overline{AC_{n-1}OB_n}}{2}, n \geq 1 \tag{436}$$

$$\text{area}\,\triangle OAC_{n-1} = \frac{\overline{OC_{n-1}AB_{n-1}}}{2}, n \geq 1 \tag{437}$$

$$\frac{\overline{AC_{n-1}OB_n}}{2} = \frac{\overline{OC_{n-1}AB_{n-1}}}{2}, n \geq 1 \tag{438}$$

$$\overline{AC_{n-1}OB_n} = \overline{OC_{n-1}AB_{n-1}}, n \geq 1 \tag{439}$$

$$\frac{\overline{AC_{n-1}OB_n}}{\overline{AC_{n-1}}} = \frac{\overline{OC_{n-1}AB_{n-1}}}{\overline{AC_{n-1}}}, n \geq 1 \tag{440}$$

$$\overline{OB_n} = \frac{\overline{OC_{n-1}AB_{n-1}}}{\overline{AC_{n-1}}}, n \geq 1 \tag{441}$$

$$\overline{OB_n} = \frac{\overline{OC_{n-1}}\sqrt{\overline{OA}^2 - \overline{OB_{n-1}}^2}}{\overline{AC_{n-1}}}, n \geq 1 \tag{442}$$

$$\overline{OB_n} = \frac{\overline{OC_{n-1}}\sqrt{\overline{OA}^2 - \overline{OB_{n-1}}^2}}{\sqrt{\overline{OA}^2 + \overline{OC_{n-1}}^2 - 2\overline{OC_{n-1}OB_{n-1}}}}, n \geq 1 \tag{443}$$

### 0.2.2 Half-chord Length

$$\overline{OB_n}^2 + \overline{AB_n}^2 = \overline{OA}^2 \tag{444}$$

$$\overline{OB_n}^2 + \overline{AB_n}^2 - \overline{OB_n}^2 = \overline{OA}^2 - \overline{OB_n}^2 \tag{445}$$

$$\overline{AB_n}^2 = \overline{OA}^2 - \overline{OB_n}^2 \tag{446}$$

$$\overline{AB_n} = \sqrt{\overline{OA}^2 - \overline{OB_n}^2} \tag{447}$$

### 0.2.3 Chord Length

$$\overline{AC_n} = \sqrt{\overline{AB_n}^2 + \overline{B_nC_n}^2} \tag{448}$$

$$\overline{AC_n} = \sqrt{\overline{OA}^2 - \overline{OB_n}^2 + \overline{B_nC_n}^2} \tag{449}$$

$$\overline{AC_n} = \sqrt{\overline{OA}^2 - \overline{OB_n}^2 + \left(\overline{OC_n} - \overline{OB_n}\right)^2} \tag{450}$$

$$\overline{AC_n} = \sqrt{\overline{OA}^2 - \overline{OB_n}^2 + \overline{OC_n}^2 - 2\overline{OC_nOB_n} + \overline{OB_n}^2} \tag{451}$$

$$\overline{AC_n} = \sqrt{\overline{OA}^2 + \overline{OC_n}^2 - 2\overline{OC_nOB_n}} \tag{452}$$

### 0.2.4 Arc Length

$$\text{arc}\,AC_0 \stackrel{\text{def}}{=} \lim_{n\to\infty} 2^n\overline{AC_n} \tag{453}$$

### 0.2.5 Area

$$\operatorname{area} \triangledown OAC_0 = \lim_{n\to\infty} 2^{n-1} \operatorname{area} OAC_nC_{n-1} \tag{454}$$

$$\operatorname{area} \triangledown OAC_0 = \lim_{n\to\infty} 2^{n-1} \frac{\overline{AC_{n-1}OC_n}}{2} \tag{455}$$

$$\operatorname{area} \triangledown OAC_0 = \frac{\lim_{n\to\infty} 2^{n-1}\overline{AC_{n-1}OC_n}}{2} \tag{456}$$


$$\operatorname{area} \triangledown OAC_0 = \frac{\lim_{n\to\infty} \left(2^{n-1}\overline{AC_{n-1}}\right) \overline{OC_n}}{2} \tag{457}$$

$$\operatorname{area} \triangledown OAC_0 = \frac{\lim_{n\to\infty} \overline{OC_n} \left(2^{n-1}\overline{AC_{n-1}}\right)}{2} \tag{458}$$

$$\operatorname{area} \triangledown OAC_0 = \frac{\left(\lim_{n\to\infty} \overline{OC_n}\right) \left(\lim_{n\to\infty} 2^{n-1}\overline{AC_{n-1}}\right)}{2} \tag{459}$$


$$\operatorname{area} \triangledown OAC_0 = \frac{\overline{OA} \lim_{n\to\infty} 2^{n-1}\overline{AC_{n-1}}}{2} \tag{460}$$

$$\operatorname{area} \triangledown OAC_0 = \frac{\overline{OA} \operatorname{arc} AC_0}{2} \tag{461}$$




(a) Recursive step.

Figure 2: Circular sector sequence.




(b) Base case.

Figure 2: Circular sector sequence.

## 0.3 Simple Harmonic Motion

$$[m] = \text{kg} \tag{462}$$

$$[\omega] = \text{m/s} \tag{463}$$

$$[k] = \text{N/m} \tag{464}$$

$$[r] = \text{m} \tag{465}$$

$$s = \omega t \tag{466}$$

$$\omega = \lim_{|\Delta t| \to 0} \frac{|\Delta s|}{|\Delta t|} \tag{467}$$

$$(\Delta s)^2 \approx (\Delta x)^2 + (\Delta y)^2 \tag{468}$$

$$\frac{(\Delta s)^2}{(\Delta x)^2} \approx \frac{(\Delta x)^2}{(\Delta x)^2} + \frac{(\Delta y)^2}{(\Delta x)^2} \tag{469}$$

$$\frac{(\Delta s)^2}{(\Delta x)^2} \approx 1 + \frac{(\Delta y)^2}{(\Delta x)^2} \tag{470}$$

$$\frac{(\Delta s)^2}{(\Delta x)^2} \approx 1 + \left(\frac{\Delta y}{\Delta x}\right)^2 \tag{471}$$

$$\frac{(\Delta s)^2}{(\Delta x)^2} = 1 + 0 \iff x \to 0 \tag{472}$$

$$\frac{(\Delta s)^2}{(\Delta x)^2} = 1 \iff x \to 0 \tag{473}$$

$$(\Delta x)^2 \frac{(\Delta s)^2}{(\Delta x)^2} = (\Delta x)^2 \iff x \to 0 \tag{474}$$

$$\frac{(\Delta x)^2 (\Delta s)^2}{(\Delta x)^2} = (\Delta x)^2 \iff x \to 0 \tag{475}$$

$$(\Delta s)^2 = (\Delta x)^2 \iff x \to 0 \tag{476}$$

$$|\Delta s| = |\Delta x| \iff x \to 0 \tag{477}$$

$$\omega = \lim_{|\Delta t| \to 0} \frac{|\Delta x|}{|\Delta t|} \iff x \to 0 \tag{478}$$

$$F = ma \tag{479}$$

$$F = -kx \tag{480}$$

$$ma = -kx \tag{481}$$

$$x = \cos(r, s) \tag{482}$$

$$x(t) = \cos(r, \omega t) \tag{483}$$

$$v(t) = \frac{dx}{dt} \tag{484}$$

$$v(t) = \left.\frac{\partial \cos(r, s)}{\partial s}\right|_{s=\omega t} \cdot \frac{d(\omega t)}{dt} \tag{485}$$

$$v(t) = \frac{-\sin(r, \omega t)}{r} \cdot \frac{d(\omega t)}{dt} \tag{486}$$

$$v(t) = \frac{-\sin(r, \omega t)}{r} \cdot \omega \tag{487}$$

$$v(t) = \frac{-\omega \sin(r, \omega t)}{r} \tag{488}$$

$$a(t) = \frac{dv}{dt} \tag{489}$$

$$a(t) = \frac{d\left(\frac{\partial \cos(r,s)}{\partial s}\Big|_{s=\omega t} \frac{d(\omega t)}{dt}\right)}{dt} \tag{490}$$

$$a(t) = \left(\frac{\partial^2 \cos(r, s)}{\partial s^2} \cdot \frac{ds}{dt}\Big|_{s=\omega t}\right) \cdot \frac{d(\omega t)}{dt} + \frac{\partial \cos(r, s)}{\partial s}\Big|_{s=\omega t} \cdot \frac{d^2(\omega t)}{dt^2} \tag{491}$$

$$a(t) = \left(\frac{-\cos(r, \omega t)}{r^2} \cdot \frac{d(\omega t)}{dt}\right) \cdot \frac{d(\omega t)}{dt} + \frac{\partial \cos(r, s)}{\partial s}\Big|_{s=\omega t} \cdot \frac{d^2(\omega t)}{dt^2} \tag{492}$$

$$a(t) = \frac{-\cos(r, \omega t)}{r^2} \cdot \left(\frac{d(\omega t)}{dt}\right)^2 + \frac{\partial \cos(r, s)}{\partial s}\Big|_{s=\omega t} \cdot \frac{d^2(\omega t)}{dt^2} \tag{493}$$

$$a(t) = \frac{-\cos(r, \omega t)}{r^2} \cdot \omega^2 + \frac{\partial \cos(r, s)}{\partial s}\Big|_{s=\omega t} \cdot \frac{d^2(\omega t)}{dt^2} \tag{494}$$

$$a(t) = \frac{-\omega^2 \cos(r, \omega t)}{r^2} + \frac{\partial \cos(r, s)}{\partial s}\Big|_{s=\omega t} \cdot \frac{d^2(\omega t)}{dt^2} \tag{495}$$

$$a(t) = \frac{-\omega^2 \cos(r, \omega t)}{r^2} + \frac{-\sin(r, \omega t)}{r} \cdot \frac{d^2(\omega t)}{dt^2} \tag{496}$$

$$a(t) = \frac{-\omega^2 \cos(r, \omega t)}{r^2} + \frac{-\sin(r, \omega t)}{r} \cdot 0 \tag{497}$$

$$a(t) = \frac{-\omega^2 \cos(r, \omega t)}{r^2} + 0 \tag{498}$$

$$a(t) = \frac{-\omega^2 \cos(r, \omega t)}{r^2} \tag{499}$$

$$a(x) = \frac{-\omega^2 x}{r^2} \tag{500}$$

$$m\frac{-\omega^2 x}{r^2} = -kx \tag{501}$$

$$\frac{m(-\omega^2 x)}{r^2} = -kx \tag{502}$$

$$\frac{-m\omega^2 x}{r^2} = -kx \tag{503}$$

$$\frac{-m\omega^2 x}{r^2} = \frac{-kr^2 x}{r^2} \tag{504}$$

$$m\omega^2 = kr^2 \tag{505}$$

$$K(t) = \frac{mv^2(t)}{2} \tag{506}$$

$$K(t) = \frac{m\left(\frac{-\omega \sin(r,\omega t)}{r}\right)^2}{2} \tag{507}$$

$$K(t) = \frac{m\frac{(-\omega \sin(r,\omega t))^2}{r^2}}{2} \tag{508}$$

$$K(t) = \frac{m\frac{(-\omega)^2 \sin^2(r,\omega t)}{r^2}}{2} \tag{509}$$

$$K(t) = \frac{m\frac{\omega^2 \sin^2(r,\omega t)}{r^2}}{2} \tag{510}$$

$$K(t) = \frac{\left(\frac{m\omega^2 \sin^2(r,\omega t)}{r^2}\right)}{2} \tag{511}$$

$$K(t) = \frac{\left(\frac{kr^2 \sin^2(r,\omega t)}{r^2}\right)}{2} \tag{512}$$

$$K(t) = \frac{k \sin^2(r,\omega t)}{2} \tag{513}$$

$$U(t) = \frac{kx^2(t)}{2} \tag{514}$$

$$U(t) = \frac{k \cos^2(r,\omega t)}{2} \tag{515}$$

$$E = K + U \tag{516}$$

$$E = \frac{k \sin^2(r,\omega t)}{2} + \frac{k \cos^2(r,\omega t)}{2} \tag{517}$$

$$E = \frac{k \sin^2(r,\omega t) + k \cos^2(r,\omega t)}{2} \tag{518}$$

$$E = \frac{k(\sin^2(r,\omega t) + \cos^2(r,\omega t))}{2} \tag{519}$$

$$E = \frac{kr^2}{2} \tag{520}$$

$$E = \frac{m\omega^2}{2} \tag{521}$$

## 0.4 Uniform Circular Motion

$$[m] = \text{kg} \tag{522}$$

$$[\omega] = \text{m/s} \tag{523}$$

$$[r] = \text{m} \tag{524}$$

$$s = \omega t \tag{525}$$

$$\omega = \lim_{|\Delta t| \to 0} \frac{|\Delta s|}{|\Delta t|} \tag{526}$$

$$x = \cos(r, s) \tag{527}$$

$$x(t) = \cos(r, \omega t) \tag{528}$$

$$y = \sin(r, s) \tag{529}$$

$$y(t) = \sin(r, \omega t) \tag{530}$$

$$\vec{r}(t) = (x, y) \tag{531}$$

$$\vec{r}(t) = (\cos(r, \omega t), \sin(r, \omega t)) \tag{532}$$

$$\vec{v}(t) = \frac{d\vec{r}}{dt} \tag{533}$$

$$\vec{v}(t) = \frac{d(x, y)}{dt} \tag{534}$$

$$\vec{v}(t) = \left( \frac{dx}{dt}, \frac{dy}{dt} \right) \tag{535}$$

$$\frac{dx}{dt} = \left. \frac{\partial \cos(r, s)}{\partial s} \right|_{s=\omega t} \cdot \frac{d(\omega t)}{dt} \tag{536}$$

$$\frac{dx}{dt} = \frac{-\sin(r, \omega t)}{r} \cdot \frac{d(\omega t)}{dt} \tag{537}$$

$$\frac{dx}{dt} = \frac{-\sin(r, \omega t)}{r} \cdot \omega \tag{538}$$

$$\frac{dx}{dt} = \frac{-\omega \sin(r, \omega t)}{r} \tag{539}$$

$$\frac{dy}{dt} = \left. \frac{\partial \sin(r, s)}{\partial s} \right|_{s=\omega t} \cdot \frac{d(\omega t)}{dt} \tag{540}$$

$$\frac{dy}{dt} = \frac{\cos(r, \omega t)}{r} \cdot \frac{d(\omega t)}{dt} \tag{541}$$

$$\frac{dy}{dt} = \frac{\cos(r, \omega t)}{r} \cdot \omega \tag{542}$$

$$\frac{dy}{dt} = \frac{\omega \cos(r, \omega t)}{r} \tag{543}$$

$$\vec{v}(t) = \left( \frac{-\omega \sin(r, \omega t)}{r}, \frac{\omega \cos(r, \omega t)}{r} \right) \tag{544}$$

$$\vec{v}(t) = \frac{\omega \left( -\sin(r, \omega t), \cos(r, \omega t) \right)}{r} \tag{545}$$

$$v = |\vec{v}| \tag{546}$$

$$v = \left| \frac{\omega \left( -\sin(r, \omega t), \cos(r, \omega t) \right)}{r} \right| \tag{547}$$

$$v = \frac{|\omega \left( -\sin(r, \omega t), \cos(r, \omega t) \right)|}{|r|} \tag{548}$$

$$v = \frac{|\omega| \, |(-\sin(r, \omega t), \cos(r, \omega t))|}{|r|} \tag{549}$$

$$v = \frac{\omega \, |(-\sin(r, \omega t), \cos(r, \omega t))|}{|r|} \tag{550}$$

$$v = \frac{\omega \sqrt{(-\sin(r, \omega t))^2 + \cos^2(r, \omega t)}}{|r|} \tag{551}$$

$$v = \frac{\omega \sqrt{\sin^2(r, \omega t) + \cos^2(r, \omega t)}}{|r|} \tag{552}$$

$$v = \frac{\omega \sqrt{r^2}}{|r|} \tag{553}$$

$$v = \frac{\omega \, |r|}{|r|} \tag{554}$$

$$v = \omega \tag{555}$$

$$E = \frac{mv^2}{2} \tag{556}$$

$$E = \frac{m\omega^2}{2} \tag{557}$$

$$\vec{a}(t) = \frac{d^2 \vec{r}}{dt^2} \tag{558}$$

$$\vec{a}(t) = \frac{d^2 (x, y)}{dt^2} \tag{559}$$

$$\vec{a}(t) = \left( \frac{d^2 x}{dt^2}, \frac{d^2 y}{dt^2} \right) \tag{560}$$

$$\frac{d^2 x}{dt^2} = \frac{d \left( \frac{\partial \cos(r,s)}{\partial s} \Big|_{s=\omega t} \frac{d(\omega t)}{dt} \right)}{dt} \tag{561}$$

$$\frac{d^2 x}{dt^2} = \left( \frac{\partial^2 \cos(r, s)}{\partial s^2} \cdot \frac{ds}{dt} \Big|_{s=\omega t} \right) \cdot \frac{d(\omega t)}{dt} + \frac{\partial \cos(r, s)}{\partial s} \Big|_{s=\omega t} \cdot \frac{d^2 (\omega t)}{dt^2} \tag{562}$$

$$\frac{d^2 x}{dt^2} = \left( \frac{-\cos(r, \omega t)}{r^2} \cdot \frac{d(\omega t)}{dt} \right) \cdot \frac{d(\omega t)}{dt} + \frac{\partial \cos(r, s)}{\partial s} \Big|_{s=\omega t} \cdot \frac{d^2 (\omega t)}{dt^2} \tag{563}$$

$$\frac{d^2 x}{dt^2} = \frac{-\cos(r, \omega t)}{r^2} \cdot \left( \frac{d(\omega t)}{dt} \right)^2 + \frac{\partial \cos(r, s)}{\partial s} \Big|_{s=\omega t} \cdot \frac{d^2 (\omega t)}{dt^2} \tag{564}$$

$$\frac{d^2 x}{dt^2} = \frac{-\cos(r, \omega t)}{r^2} \cdot \omega^2 + \frac{\partial \cos(r, s)}{\partial s} \Big|_{s=\omega t} \cdot \frac{d^2 (\omega t)}{dt^2} \tag{565}$$

$$\frac{d^2x}{dt^2} = \frac{-\omega^2\cos(r,\omega t)}{r^2} + \left.\frac{\partial\cos(r,s)}{\partial s}\right|_{s=\omega t} \cdot \frac{d^2(\omega t)}{dt^2} \tag{566}$$

$$\frac{d^2x}{dt^2} = \frac{-\omega^2\cos(r,\omega t)}{r^2} + \frac{-\sin(r,\omega t)}{r} \cdot \frac{d^2(\omega t)}{dt^2} \tag{567}$$

$$\frac{d^2x}{dt^2} = \frac{-\omega^2\cos(r,\omega t)}{r^2} + \frac{-\sin(r,\omega t)}{r} \cdot 0 \tag{568}$$

$$\frac{d^2x}{dt^2} = \frac{-\omega^2\cos(r,\omega t)}{r^2} + 0 \tag{569}$$

$$\frac{d^2x}{dt^2} = \frac{-\omega^2\cos(r,\omega t)}{r^2} \tag{570}$$

$$\frac{d^2y}{dt^2} = \frac{d\left(\left.\frac{\partial\sin(r,s)}{\partial s}\right|_{s=\omega t}\frac{d(\omega t)}{dt}\right)}{dt} \tag{571}$$

$$\frac{d^2y}{dt^2} = \left(\left.\frac{\partial^2\sin(r,s)}{\partial s^2}\cdot\frac{ds}{dt}\right|_{s=\omega t}\right)\cdot\frac{d(\omega t)}{dt} + \left.\frac{\partial\sin(r,s)}{\partial s}\right|_{s=\omega t}\cdot\frac{d^2(\omega t)}{dt^2} \tag{572}$$

$$\frac{d^2y}{dt^2} = \left(\frac{-\sin(r,\omega t)}{r^2}\cdot\frac{d(\omega t)}{dt}\right)\cdot\frac{d(\omega t)}{dt} + \left.\frac{\partial\sin(r,s)}{\partial s}\right|_{s=\omega t}\cdot\frac{d^2(\omega t)}{dt^2} \tag{573}$$

$$\frac{d^2y}{dt^2} = \frac{-\sin(r,\omega t)}{r^2}\cdot\left(\frac{d(\omega t)}{dt}\right)^2 + \left.\frac{\partial\sin(r,s)}{\partial s}\right|_{s=\omega t}\cdot\frac{d^2(\omega t)}{dt^2} \tag{574}$$

$$\frac{d^2y}{dt^2} = \frac{-\sin(r,\omega t)}{r^2}\cdot\omega^2 + \left.\frac{\partial\sin(r,s)}{\partial s}\right|_{s=\omega t}\cdot\frac{d^2(\omega t)}{dt^2} \tag{575}$$

$$\frac{d^2y}{dt^2} = \frac{-\omega^2\sin(r,\omega t)}{r^2} + \left.\frac{\partial\sin(r,s)}{\partial s}\right|_{s=\omega t}\cdot\frac{d^2(\omega t)}{dt^2} \tag{576}$$

$$\frac{d^2y}{dt^2} = \frac{-\omega^2\sin(r,\omega t)}{r^2} + \frac{\cos(r,\omega t)}{r}\cdot\frac{d^2(\omega t)}{dt^2} \tag{577}$$

$$\frac{d^2y}{dt^2} = \frac{-\omega^2\sin(r,\omega t)}{r^2} + \frac{\cos(r,\omega t)}{r}\cdot 0 \tag{578}$$

$$\frac{d^2y}{dt^2} = \frac{-\omega^2\sin(r,\omega t)}{r^2} + 0 \tag{579}$$

$$\frac{d^2y}{dt^2} = \frac{-\omega^2\sin(r,\omega t)}{r^2} \tag{580}$$

$$\vec{a}(t) = \left(\frac{-\omega^2\cos(r,\omega t)}{r^2}, \frac{-\omega^2\sin(r,\omega t)}{r^2}\right) \tag{581}$$

$$\vec{a}(t) = \frac{-\omega^2\left(\cos(r,\omega t), \sin(r,\omega t)\right)}{r^2} \tag{582}$$

$$\vec{a} = \frac{-\omega^2\vec{r}}{r^2} \tag{583}$$

$$a = |\vec{a}| \tag{584}$$

$$a = \left|\frac{-\omega^2\vec{r}}{r^2}\right| \tag{585}$$

$$a = \frac{|-\omega^2 \vec{r}|}{|r^2|} \tag{586}$$

$$a = \frac{|-\omega^2| \, |\vec{r}|}{|r^2|} \tag{587}$$

$$a = \frac{\omega^2 \, |\vec{r}|}{|r^2|} \tag{588}$$

$$a = \frac{\omega^2 r}{|r^2|} \tag{589}$$

$$a = \frac{\omega^2 r}{r^2} \tag{590}$$

$$a = \frac{\omega^2}{r} \tag{591}$$

$$F = ma \tag{592}$$

$$F = m\frac{\omega^2}{r} \tag{593}$$

$$F = \frac{m\omega^2}{r} \tag{594}$$