

Math Cheatsheet

Math Cheatsheet

Trigonometry

Sine

Right Angles

Cosine

Right Angles

Tangent

Right Angles

Geometry

Similarity

Triangle

Equilateral

Right Angle

Circle

Pyramid

Tetrahedron

Kegel

Kugel

Vector Geometry

Definition

Getting a Vector from Two Points

Stretching

Linear Combination

Linear Dependency

Scalar Absolute Value (Length)

Unit vector

Dot product

Angles

Vector Shadow

Straights

Parameter Definition

Positional Relationships

Exponential functions

Logistical functions

Acknowledgements

Trigonometry

Remember: *SOH CAH TOA*

Sine

$$\frac{a}{\sin(\alpha)} = \frac{b}{\sin(\beta)} = \frac{c}{\sin(\gamma)}$$

Right Angles

$$\sin(\theta) = \frac{o}{h} = \frac{opposite}{hypotenuse}$$

Cosine

$$\begin{aligned} a^2 &= b^2 + c^2 - 2bc \cdot \cos(\alpha) \\ b^2 &= a^2 + c^2 - 2ac \cdot \cos(\beta) \\ c^2 &= a^2 + b^2 - 2ab \cdot \cos(\gamma) \end{aligned}$$

Right Angles

$$\cos(\theta) = \frac{a}{h} = \frac{adjacent}{hypotenuse}$$

Tangent

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$$

Right Angles

$$\tan(\theta) = \frac{o}{a} = \frac{opposite}{adjacent}$$

Geometry

Similarity

$$\begin{aligned} s &= k \cdot s_0 \\ A &= k^2 \cdot A_0 \\ V &= k^3 \cdot V_0 \end{aligned}$$

Triangle

Equilateral

$$\begin{aligned} h &= \frac{\sqrt{3}}{2} a \\ A &= \frac{\sqrt{3}}{4} a^2 \end{aligned}$$

Right Angle

$$A = \frac{a \cdot b}{2}$$


Circle

$$\begin{aligned} A &= \pi \cdot r^2 \\ U &= \pi \cdot 2r = \pi \cdot d \end{aligned}$$

Pyramid


$$V = \frac{G \cdot h}{3}$$


Tetrahedron



$$V = \frac{\sqrt{2}}{12} a^3$$

Kegel





$$V = \frac{G \cdot h}{3} = \frac{r^2 \pi \cdot h}{3}$$

$$b = 2\pi \cdot r$$

$$M = \frac{b \cdot m}{2} = r\pi \cdot m$$

$$m = G + M$$

$$m^2 = h^2 + r^2$$

$$\alpha = 360^\circ \cdot \frac{r}{m}$$

Kugel

$$V = \frac{4}{3}r^3\pi$$

$$A = 4\pi r^2$$

Vector Geometry

Definition

$$\vec{a} = \begin{pmatrix} a_x \\ a_y \end{pmatrix}$$

$$\vec{a} = \begin{pmatrix} a_x \\ a_y \\ a_z \end{pmatrix}$$

Getting a Vector from Two Points

$$B = (11;-2) \quad C = (8;6)$$

$$\vec{BC} = \begin{pmatrix} C_x \\ C_y \end{pmatrix} - \begin{pmatrix} B_x \\ B_y \end{pmatrix}$$

Stretching

$$k \cdot \vec{a}$$

$$k \cdot \begin{pmatrix} a_x \\ a_y \end{pmatrix}$$

$$\begin{pmatrix} k \cdot a_x \\ k \cdot a_y \end{pmatrix}$$

$$\begin{pmatrix} a'_x \\ a'_y \end{pmatrix}$$

Linear Combination

AKA aligning the tip of each vector to the tail of another vector, resulting in a chain, which then gives you the starting point of the combination of vectors.

$$\vec{x} = k_1 \vec{a} + k_2 \vec{b} + k_3 \vec{c}$$

Linear Dependency

2 Vectors are linearly dependent if you can achieve one of the vectors by multiplying the other vector with any value of k, where k can't be 0.

$$\vec{a} = k \cdot \vec{b}$$

If there is a solution where k_1, k_2, k_3, \dots are all $\neq 0$ then the three vectors are linearly dependent.

$$\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = k_1 \cdot \vec{a} + k_2 \cdot \vec{b} + k_3 \cdot \vec{c}$$

Scalar Absolute Value (Length)

$$\begin{aligned} |\vec{a}| &= \sqrt{a_x^2 + a_y^2} \\ |\vec{b}| &= \sqrt{b_x^2 + b_y^2 + b_z^2} \\ &\vdots \end{aligned}$$

Unit vector

A unit vector is a vector that equals to 1 in its absolute value. The following formula can be used to determine the unit vector pointing in the same direction as the vector a .

$$\vec{e}_a = \frac{1}{|\vec{a}|} * \vec{a} = \frac{\vec{a}}{|\vec{a}|}$$

You can use the following formula to figure out what the new vector should be if it's to point in the same direction as the given one a but have a specific length l .

$$\vec{a}_l = \frac{l}{|\vec{a}|} * \vec{a}$$

Dot product

$$\begin{aligned} \vec{a} \cdot \vec{b} &= a_x b_x + a_y b_y \\ \vec{a} \cdot \vec{b} &= a_x b_x + a_y b_y + a_z b_z \\ &\vdots \end{aligned}$$

OR

$$\vec{a} \cdot \vec{b} = \cos(\alpha) \cdot |\vec{a}| \cdot |\vec{b}|$$

The Scalar Absolute value can be calculated with the use of the `dotp` function using the TI-Nspire CX CAS as follows.

```
function dotp(veca, vecb):
    return veca.x * vecb.x + veca.y * vecb.y

function dotp(veca, vecb):
    return veca.x * vecb.x + veca.y * vecb.y + veca.z * vecb.z
```

$$|\vec{AB}| = \sqrt{\text{dotp}(\vec{AB}, \vec{AB})}$$


Angles

Vectors need to point **AWAY** from angle!

$$\begin{aligned}\vec{AC} \cdot \vec{AB} &= \cos(\alpha) \cdot |\vec{AC}| \cdot |\vec{AB}| \\ \cos(\alpha) &= \frac{\vec{AC} \cdot \vec{AB}}{|\vec{AC}| \cdot |\vec{AB}|} \\ \alpha &= \cos^{-1}\left(\frac{\vec{AC} \cdot \vec{AB}}{|\vec{AC}| \cdot |\vec{AB}|}\right)\end{aligned}$$

Vector Shadow

$$\begin{aligned}|\vec{a_b}| &= |\vec{a}| \cdot \cos(\alpha) \\ \vec{a_b} &= |\vec{a_b}| \cdot \vec{e_b} \\ \vec{a_b} &= |\vec{a}| \cdot \cos(\alpha) \cdot \vec{e_b}\end{aligned}$$



Straights

Parameter Definition

$$g: \vec{r} = \vec{a} + t \cdot \vec{u}$$

Positional Relationships

Two straights can be in one of four relations: *identical*, *intersecting in one point*, *parallel and non-intersecting*, *not parallel*. By defining both of the straights generally

$$\begin{aligned}g: \vec{r} &= \vec{a} + t \cdot \vec{u} \\ h: \vec{r} &= \vec{b} + s \cdot \vec{v}\end{aligned}$$

we can make some determinations about when which of these cases are true.

If the equation $\vec{a} + t \cdot \vec{u} = \vec{b} + s \cdot \vec{v}$ has

1. one solution then g and h intersect in one point.
2. no solutions
 1. and $\vec{u} = k\vec{v}$ is valid the straights g and h are parallel.
 2. and $\vec{u} = k\vec{v}$ is invalid the straights g and h are not parallel.
3. infinite solutions then g and h are the same.

For straights in a 2D space the case 2.2 is not possible.

Exponential functions

$$E = S \cdot q^t$$

You get the resulting value E by multiplying the starting value s with the result of the growth factor q to the power of time t that has passed. This can be more generally described as:

$$f(x) = b \cdot a^x$$

Logistical functions

$$f(t) = \frac{a \cdot S}{a + (S - a)e^{-kt}}$$

Here a is defined as the y-axis intercept of the function, or the starting value if you will. S is defined as the value of $f(t)$ when t is infinitely big or otherwise referred to as carrying capacity. This can be more generally described as (might be inaccurate as I can't find a concise formula from different sources):

$$f(x) = \frac{a \cdot S}{a + (S - a)e^{-kx}}$$

Acknowledgements

Author(s): d20cay, (SirW)Johnson)

Modified: 2020-11-30