# The Open Pediatric Cancer Project

This manuscript (<u>permalink</u>) was automatically generated from <u>d3b-center/OpenPedCan-methods@3c232b8</u> on 2024-05-17.

# **Authors**

#### • Eric Wafula

**(D)** 0000-0001-8073-3797

Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Sangeeta Shukla

© 0000-0002-3727-9602

Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### · Aditya Lahiri

© 0000-0001-9352-1312

Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Krutika S. Gaonkar

© 0000-0003-0838-2405

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Run Jin

© 0000-0002-8958-9266

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Komal S. Rathi

**D** 0000-0001-5534-6904

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Yuankun Zhu

© 0000-0002-2455-9525

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Bailey K. Farrow

**(D)** 0000-0001-6727-6333

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Daniel P. Miller

**(D)** 0000-0002-2032-4358

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Mariarita Santi

© 0000-0002-6728-3450

Department of Pathology and Laboratory Medicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Department of Pathology and Laboratory Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, 19104, USA

### · Adam A. Kraya

### © 0000-0002-8526-5694

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Xiaoyan Huang

### **(D)** 0000-0001-7267-4512

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Bo Zhang

### **D** 0000-0002-0743-5379

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Chuwei Zhong

## © 0000-0003-2406-2735

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Zhuangzhuang Geng

### **D** 0009-0007-6883-0691

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Brian M. Ennis

### © 0000-0002-2653-5009

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Ryan J. Corbett

### **(D)** <u>0000-0002-3478-0784</u>

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Sharon J. Diskin

#### **(D)** 0000-0002-7200-8939

Division of Oncology, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Department of Pediatrics, University of Pennsylvania, Philadelphia, PA, 19104, USA

### Nicholas Van Kuren

#### **(D)** 0000-0002-7414-9516

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Matthew R. Lueder

#### 🔟 0009-0002-7370-102X

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA;

Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Department of Pathology and Laboratory Medicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Noel Coleman

**(D)** 0000-0001-6454-1285

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### · Christopher Blackden

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### • Jennifer L. Mason

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Saksham Phul

**(D)** 0000-0002-2771-2572

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### • Miguel A. Brown

**1** 0000-0001-6782-1442

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Alex Sickler

**(D)** 0000-0001-7830-7537

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Kelsey Keith

**(iii)** 0000-0002-7451-5117

Department of Biomedical and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

#### Dave Hill

© 0000-0002-1337-1789

Department of Biomedical and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### • Asif T Chinwalla

© 0000-0001-7831-3996

Department of Biomedical and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### Yuanchao Zhang

Department of Biomedical and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

### • Adam C. Resnick

© 0000-0003-0436-4189

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA · Funded by Children's Brain Tumor Network; NIH 3P30 CA016520-44S5, U2C HL138346-03, U24 CA220457-03; NCI/NIH Contract No. 75N91019D00024, Task Order No. 75N91020F00003; Children's Hospital of Philadelphia Division of Neurosurgery

#### Alvin Farrel

### **(iii)** 0000-0003-1087-9840

Department of Biomedical and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Oncology, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Center for Childhood Cancer Research, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA · Funded by NCI/NIH Contract No. 75N91019D00024, Task Order No. 75N91020F00003

### Deanne Taylor

### © 0000-0002-3302-4610

Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Department of Pediatrics, University of Pennsylvania Perelman Medical School, Philadelphia, PA, 19104, USA · Funded by NCI/NIH Contract No. 75N91019D00024, Task Order No. 75N91020F00003

## • Jo Lynne Rokita^ <sup>™</sup>

### © 0000-0003-2171-3627

Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Division of Neurosurgery, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA; Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA · Funded by NCI/NIH Contract No. 75N91019D00024, Task Order No. 75N91020F00003

# **Contact information**

^Lead Contact: Jo Lynne Rokita <a href="mailto:rokita@chop.edu">rokita@chop.edu</a>

⊠Correspondence: Jo Lynne Rokita <u>rokita@chop.edu</u>

### In Brief

# **Highlights**

# **Summary**

# **Keywords**

# Introduction

The Open Pediatric Cancer (OpenPedCan) project at the Children's Hospital of Philadelphia is an open analysis effort in which we harmonize pediatric cancer data from multiple sources, perform downstream cancer analyses on these data, and provide them on PedcBioPortal and v2.1 of NCI's <a href="Pediatric Molecular Targets Platform (MTP)">Pediatric Molecular Targets Platform (MTP)</a>. The OpenPedCan analyses currently include the following datasets, described more fully below:

- TARGET
- Kids First Neuroblastoma
- OpenPBTA
- GTEx

- TCGA
- DGD (CHOP P30 Panel)

**Open Pediatric Brain Tumor Atlas (OpenPBTA)** In September of 2018, the <u>Children's Brain Tumor Network (CBTN)</u> released the <u>Pediatric Brain Tumor Atlas (PBTA)</u>, a genomic dataset (whole genome sequencing, whole exome sequencing, RNA sequencing, proteomic, and clinical data) for nearly 1,000 tumors, available from the <u>Gabriella Miller Kids First Portal</u>. In September of 2019, the Open Pediatric Brain Tumor Atlas (OpenPBTA) Project was launched. OpenPBTA was a global open science initiative to comprehensively define the molecular landscape of tumors of 943 patients from the CBTN and the PNOC003 DIPG clinical trial from the <u>Pediatric Pacific Neuro-oncology Consortium</u> through real-time, collaborative analyses and <u>collaborative manuscript writing</u> on GitHub, now published in <u>Cell Genomics</u>. Additional PBTA data has been, and will be continually added to, OpenPedCan.

Therapeutically Applicable Research to Generate Effective Treatments (TARGET) The Therapeutically Applicable Research to Generate Effective Treatments (TARGET) Initiative is an NCI-funded collection of disease-specific projects that seeks to identify the genomic changes of pediatric cancers. The overall goal is to collect genomic data to accelerate the development of more effective therapies. OpenPedCan analyses include the seven diseases present in the TARGET dataset: Acute Lymphoblastic Leukemia (ALL), Acute Myeloid Leukemia (AML), Clear cell sarcoma of the kidney, Neuroblastoma, Osteosarcoma, Rhabdoid tumor, and Wilm's Tumor.

**Gabriella Miller Kids First Neuroblastoma** (Kids First) The Gabriella Miller Kids First Pediatric Research Program (Kids First) is a large-scale effort to accelerate research and gene discovery in pediatric cancers and structural birth defects. The program includes whole genome sequencing (WGS) from patients with pediatric cancers and structural birth defects and their families. OpenPedCan analyses include Neuroblastoma data from the Kids First project.

**The Genotype-Tissue Expression (GTEX)** GTEx project is an ongoing effort to build a comprehensive public data resource and tissue bank to study tissue-specific gene expression, regulation and their relationship with genetic variants. Samples were collected from 54 non-diseased tissue sites across nearly 1000 individuals, primarily for molecular assays including WGS, WES, and RNA-Seq. OpenPedCan project includes 17,382 GTEx RNA-Seq samples from GTEx v8 release, which span across 31 GTEx groups in the v12 release.

**The Cancer Genome Atlas Program** (TCGA) TCGA is a landmark cancer genomics program that molecularly characterized over 20,000 primary cancer and matched normal samples spanning 33 cancer types. It is a joint effort between NCI and the National Human Genome Research Institute. OpenPedCan project includes 10,414 TCGA RNA-Seq samples (716 normal and 9698 tumor) from 33 cancer types in the v12 release.

**DGD** (CHOP P30 Panel) CHOP's <u>Division of Genome Diagnostics</u> has partnered with CCDI to add somatic panel sequencing data to OpenPedCan and the Molecular Targets Platform.

The OpenPedCan operates on a pull request model to accept contributions from community participants. The maintainers have set up continuous integration software via GitHub Actions to confirm the reproducibility of analyses within the project's Docker container.

The project maintainers include scientists from the <u>Center for Data-Driven Discovery in Biomedicine at the Children's Hospital of Philadelphia</u>. We invite researchers to join OpenPedCan to help rigorously characterize the genomic landscape of these diseases to enable more rapid discovery of additional mechanisms contributing to the pathogenesis of pediatric brain and spinal cord tumors and overall accelerate clinical translation on behalf of patients.

# Results

# **Discussion**

# Acknowledgments

# **Author Contributions**

Author	Contributions
Eric Wafula	Formal analysis, Software
Sangeeta Shukla	Data curation, Formal analysis, Investigation, Methodology, Software, Writing – Original draft, Writing - Review and editing
Aditya Lahiri	Data curation, Formal analysis, Investigation, Methodology, Software, Writing – Original draft, Writing - Review and editing
Krutika S. Gaonkar	Data curation, Formal analysis, Investigation, Methodology, Software, Writing – Original draft, Writing - Review and editing
Run Jin	Data curation, Formal analysis, Visualization, Writing – Original draft, Writing - Review and editing
Komal S. Rathi	Formal analysis, Investigation, Methodology, Writing – Original draft
Yuankun Zhu	Data curation, Formal analysis, Investigation, Methodology, Supervision
Bailey K. Farrow	Data curation, Software
Daniel P. Miller	Formal analysis
Mariarita Santi	Investigation, Validation, Writing - Review and editing
Adam A. Kraya	Methodology
Xiaoyan Huang	Formal analysis
Bo Zhang	Data curation, Formal analysis
Chuwei Zhong	Formal analysis
Zhuangzhuang Geng	Data curation, Formal analysis, Investigation, Methodology, Software, Writing – Original draft
Brian M. Ennis	Data curation, Formal analysis
Ryan J. Corbett	Formal analysis
Sharon J. Diskin	Investigation, Supervision, Validation, Funding acquisition, Writing - Review and editing
Nicholas Van Kuren	Data curation, Software
Matthew R. Lueder	Data curation
Noel Coleman	Data curation
Christopher Blackden	Resources

Author	Contributions
Jennifer L. Mason	Supervision
Saksham Phul	Data curation, Methodology, Formal analysis
Miguel A. Brown	Data curation, Methodology, Formal analysis
Alex Sickler	Methodology, Formal analysis
Kelsey Keith	Software, Writing - original draft, API, Formal Analysis, Data Curation, Visualization
Dave Hill	Software, Writing - original draft, API, Data Curation
Asif T Chinwalla	Project Administration, Supervision, Methodology, Investigation, Validation, API
Yuanchao Zhang	API, Software, Formal Analysis, Writing - original draft
Adam C. Resnick	Conceptualization, Funding acquisition, Resources, Supervision
Alvin Farrel	Supervision, Investigation, Visualization, Methodology, Funding acquisition
Deanne Taylor	Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Software, Supervision, Writing – Original draft, Writing - Review and editing
Jo Lynne Rokita^	Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Software, Supervision, Writing – Original draft, Writing - Review and editing

# **Declarations of Interest**

# **Figure Titles and Legends**

# **Table Titles and Legends**

# **OPENPEDCAN METHODS**

## **RESOURCE AVAILABILITY**

### **Lead contact**

Requests for access to OpenPedCan raw data and/or specimens may be directed to, and will be fulfilled by Jo Lynne Rokita (rokita@chop.edu).

# **Materials availability**

This study did not create new, unique reagents.

# Data and code availability

Within OpenPedCan (OPC), we harmonized, aggregated, and analyzed data from multiple sources. We harmonized data from the Therapeutically Applicable Research to Generate Effective Treatments (TARGET cohort) Initiative, an NCI-funded collection of disease-specific projects that seeks to identify the genomic changes of pediatric cancers [1]. We included already harmonized neuroblastoma samples from the Gabriella Miller Kids First (GMKF cohort) Pediatric Research Program, a large-scale effort to accelerate research and gene discovery in pediatric cancers and structural birth defects [2]. Additionally, we re-harmonized all samples from the Open Pediatric Brain Tumor Atlas (OpenPBTA, PBTA cohort), an open science initiative led by Alex's Lemonade Stand Foundation Childhood Cancer Data Lab and the Center for Data-Driven Discovery (D3B) at the Children's Hospital of Philadelphia (CHOP), which genomically characterized pediatric brain tumor data from the Children's Brain Tumor Network (CBTN), and the Pacific Pediatric Neuro-oncology Consortium (PNOC) [3,4]. Building on the work of OpenPBTA, OPC added the PBTA X01 data [5], the Chordoma Foundation data [6/], and the MI-ONCOSEQ Study [7], donated to CBTN by the University of Michigan, to the PBTA cohort. Finally, OPC includes the Children's Hospital of Philadelphia (CHOP) P30 Panel data generated by CHOP's Division of Genomic Diagnostics (DGD cohort) which includes targeted DNA and fusion panel data [3]. In addition to pediatric cancer data, OpenPedCan contains adult data from large science consortiums as references. For normal gene expression, GTEx [8] was used, and for comparison with adult cancers, The Cancer Genome Atlas (TCGA) [9] was included.

Merged summary files for the latest release of OpenPedCan are openly accessible in <u>CAVATICA</u> or via download-data.sh script in the <a href="https://github.com/PediatricOpenTargets/OpenPedCan-analysis">https://github.com/PediatricOpenTargets/OpenPedCan-analysis</a> repository. Cancer group summary data from release v11 are visible within the NCI's pediatric <a href="Molecular Targets Platform">Molecular Targets Platform</a>. Cohort, cancer group, and individual data are visible within <a href="PedcBioPortal">PedcBioPortal</a>

OpenPedCan analysis modules were developed within OpenPBTA [4], modified based on OpenPBTA, or newly created and can be found within the following publicly available repositories. OpenPBTA module analyses can be found at <a href="https://github.com/AlexsLemonade/OpenPBTA-analysis">https://github.com/AlexsLemonade/OpenPBTA-analysis</a>. OpenPedCan module analyses can be found at

https://github.com/PediatricOpenTargets/OpenPedCan-analysis. OpenPedCan api code can be found at https://github.com/PediatricOpenTargets/OpenPedCan-api.

All original code was developed within the following modules in the OpenPedCan analyses repository as listed below. Links to the modules are available here, and within each module is a detailed README that describes the purpose and intended usage of the scripts, along with pointers to the results from the data those scripts process.

List of OpenPedCan Analyses Modules
<u>chromosomal-instability</u>
<u>cnv-frequencies</u>
<u>collapse-rnaseq</u>
<u>compare-gistic</u>
copy number consensus call
<u>create-subset-files</u>
data-pre-release-qc
<u>efo-mondo-mapping</u>
<u>filter-mtp-tables</u>
focal-cn-file-preparation
<u>fusion-frequencies</u>

List of OpenPedCan Analyses Modules
<u>fusion-summary</u>
fusion filtering
gene-set-enrichment-analysis
gene match
<u>immune-deconv</u>
<u>independent-samples</u>
long-format-table-utils
methylation-preprocessing
methylation-summary
molecular-subtyping-ATRT
molecular-subtyping-CRANIO
molecular-subtyping-EPN
molecular-subtyping-EWS
molecular-subtyping-HGG
molecular-subtyping-LGAT
molecular-subtyping-MB
molecular-subtyping-NBL
molecular-subtyping-chordoma
molecular-subtyping-embryonal
molecular-subtyping-integrate
molecular-subtyping-neurocytoma
molecular-subtyping-pathology
<u>mtp-annotations</u>
mtp-tables-qc-checks
<u>mutational-signatures</u>
<u>pedcbio-cnv-prepare</u>
<u>pedcbio-sample-name</u>
<u>pedot-table-column-display-order-name</u>
<u>rna-seq-expression-summary-stats</u>
<u>rnaseq-batch-correct</u>
<u>run-gistic</u>
<u>snv-frequencies</u>
<u>tmb-calculation</u>
<u>tp53 nf1 score</u>
tumor-gtex-plots
<u>tumor-normal-differential-expression</u>

Software versions are documented in **Table XX**.

### **Data releases**

We maintained a data release folder on Amazon S3, downloadable directly from S3 or our open-access CAVATICA project, with merged files for each analysis (See data and code availability section). As we produced new results that we expected to be used across multiple analyses, or identified data issues, we created new data releases in a versioned manner.

### **METHOD DETAILS**

### Nucleic acids extraction and library preparation

For the PBTA X01 cohort, libraries were prepped using the Illumina TruSeq Strand-Specific Protocol to pull out poly-adenylated transcripts.

**cDNA Library Construction** Total RNA was quantified using the Quant-iT™ RiboGreen® RNA Assay Kit and normalized to 5ng/ul. Following plating, 2 uL of ERCC controls (using a 1:1000 dilution) were spiked into each sample. An aliquot of 325 ng for each sample was transferred into library preparation. The resultant 400bp cDNA went through dual-indexed library preparation: 'A' base addition, adapter ligation using P7 adapters, and PCR enrichment using P5 adapters. After enrichment, the libraries were quantified using Quant-iT PicoGreen (1:200 dilution). Samples were normalized to 5 ng/uL. The sample set was pooled and quantified using the KAPA Library Quantification Kit for Illumina Sequencing Platforms.

## **Data generation**

**PBTA X01 Illumina Sequencing** Pooled libraries were normalized to 2nM and denatured using 0.1 N NaOH prior to sequencing. Flowcell cluster amplification and sequencing were performed according to the manufacturer's protocols using the NovaSeq 6000. Each run was a 151bp paired-end with an eight-base index barcode read. Data was analyzed using the Broad Picard Pipeline which includes demultiplexing and data aggregation.

# **DNA WGS Alignment**

Please refer to the OpenPBTA manuscript for details [4].

# **Quality Control of Sequencing Data**

Please refer to the OpenPBTA manuscript for details [4]. We also ran somalier relate [10] to identify potential mismatched samples. We required that at least 20M total reads with 50% of RNA-Seq reads mapped to the human reference for samples to be included in analysis. We required at least 20X coverage for tumor DNA samples to be included in this analysis.

# SNP calling for B-allele Frequency (BAF) generation

Please refer to the OpenPBTA manuscript for details [4].

# **Somatic Mutation and INDEL Calling**

For matched tumor/normal samples, we used the same mutation calling methods as described in OpenPBTA manuscript for details [4]. For tumor only samples, we ran Mutect2 from GATK v4.2.2.0 using the following workflow.

### VCF annotation and MAF creation

Somatic variants were annotated by the Ensembl Variant Effect Predictor (VEP v105) [11]. From tumor only variant calls, we removed variants with  $alt_depth == 0$  or  $t_depth < 4$ .

### Consensus SNV Calling (tumor/normal only)

We adopted the consensus SNV calling method described in OpenPBTA manuscript with adjustment [4]. For SNV calling, we combined four consensus SNV calling algorithms: Strelka2[12], Mutect2[13], Lancet[14], and VarDict[doi? 10.1093/nar/gkw227].

Strelka2 outputs multi-nucleotide polymorphisms (MNPs) as consecutive single-nucleotide polymorphisms. In order preserve MNPs, we gather MNP calls from the other caller inputs, and search for evidence supporting these consecutive SNP calls as MNP candidates. Once found, the Strelka2 SNP calls supporting a MNP are converted to a single MNP call. This is done to preserve the predicted gene model as accurately as possible in our consensus calls. Consensus SNV from all four callers were collected and by default, calls that were detected in at least two calling algorithms or marked with "HotSpotAllele" were retained.

For all SNVs, potential non-hotspot germline variants were removed if they had a normal depth <= 7 and gnomAD allele frequency > 0.001. Final results were saved in MAF format.

# **Somatic Copy Number Variant (CNV) Calling**

We called copy number variants for tumor/normal samples using Control-FREEC [15,16] and CNVkit [17] as described in the OpenPBTA manuscript [4]. We used GATK [18] to call CNVs for matched tumor/normal WGS samples when there were at least 30 male and 30 female normals from the same sequencing platform available for panel of normal creation. For tumor only samples, we used Control-FREEC with the following modifications. Instead of the b-allele frequency germline input file, we used the dbSNP\_v153\_ucsc-compatible.converted.vt.decomp.norm.common\_snps.vcf.gz dbSNP common snps file and to avoid hard-to-call regions, utilized the hg38\_canonical\_150.mappability mappability file. Both are also linked in the public Kids First references CAVATICA project. The Control-FREEC tumor only workflow can be found here.

# Consensus CNV Calling (WGS samples only)

We adopted the consensus CNV calling described in OpenPBTA manuscript [4] with minor adjustments. For each caller and sample with WGS performed, we called CNVs based on consensus among Control-FREEC [15,16], CNVkit [17], and GATK [doi? 10.1101/gr.107524.110]. Sample and consensus caller files with more than 2,500 CNVs were removed to de-noise and increase data quality, based on cutoffs used in GISTIC [19]. For each sample, we included the following regions in the final consensus set: 1) regions with reciprocal overlap of 50% or more between at least two of the callers; 2) smaller CNV regions in which more than 90% of regions were covered by another caller. For GATK, if a panel of normal was not able to be created (required 30 male and 30 female with the same sequencing platform), consensus was run for that tumor using Control-FREEC, CNVkit, and MantaSV. We defined copy number as NA for any regions that had a neutral call for the samples included in the consensus file. We merged CNV regions within 10,000 bp of each other with the same direction of gain or loss into single region.

Any CNVs that overlapped 50% or more with immunoglobulin, telomeric, centromeric, segment duplicated regions, or that were shorter than 3000 bp were filtered out. The CNVKit calls for WXS samples were appended to the consensus CNV file.

## Somatic Structural Variant Calling (WGS samples only)

Please refer to the OpenPBTA manuscript for details [4].

# **Methylation Analysis**

## Methylation array preprocessing

We preprocessed raw Illumina 450K and EPIC 850K Infinium Human Methylation Bead Array intensities using the array preprocessing methods implemented in the minfi Bioconductor package [20]. We utilized either preprocessFunnorm when an array dataset had both tumor and normal samples or multiple OpenPedcan-defined cancer\_groups and preprocessQuantile when an array dataset had only tumor samples from a single OpenPedcan-defined cancer\_group to estimate usable methylation measurements (beta-values and m-values) and copy number (cn-values). Some Illumina Infinium array probes targeting CpG loci contain single-nucleotide polymorphisms (SNPs) near or within the probe [21], which could affect DNA methylation measurements [22]. As the minfi preprocessing workflow recommends, we dropped probes containing common SNPs in dbSNP (minor allele frequency > 1%) at the CpG interrogation or the single nucleotide extensions.

Details of methylation array preprocessing are available in the <u>OpenPedCan methylation-preprocessing module</u>.

# Methylation beta-values summaries

We comprehensively summarized gene-level and isoform-level metrics for the methylation betavalues estimated by array preprocessing to provide insight into the variations in overall genomic DNA methylation levels observed across different pediatric tumors by computing CpG probe-level summary metrics in each cancer group within a cohort, including 1) beta-values quantiles, 2) gene expression (TPM) and methylation (beta-values) correlation, 3) TPM median expression, and 4) transcript representation - a proxy for percent isoform expression in a gene. In addition, each CpG probe was annotated with a gene feature to identify the genomic regions likely involved in regulating gene expression.

Details of the analysis are available in the OpenPedCan methylation-summary module.

# Methylation sample classification

We ran the <a href="dkfz's brain classifier version 12.5">dkfz\_v12\_methylation\_mgmt\_status</a> (predicted methylation status), and dkfz\_v12\_methylation\_mgmt\_estimated (estimated MGMT methylation fraction).

### **Gene Expression**

The <a href="tumor-normal-differential-expression">tumor-normal-differential-expression</a> module perfoms differential expression analyses for all sets of Disease (cancer\_group) and Dataset (cohort) across all genes found in the gene-expression-rsem-tpm-collapsed.rds table. The purpose of this analysis is to highlight the correlation and understand the variability in gene expression in different cancer conditions across different histological tissues. For OpenPedCan v12 data release, this module performs expression analysis over 102 cancer groups across 52 histological tissues for all 54,346 genes found in the dataset. This analysis was performed on the Children's Hospital of Philadelphia HPC and was configured to use 96G of RAM per CPU, with one task (one iteration of expression analysis for each set of tissue and cancer group) per CPU (total 102\*52=5304 CPUs) using the <a href="R/DESeq2">R/DESeq2</a> package. Please refer to script run-tumor-normal-differential-expression.sh in the module for additional details on Slurm processing configuration. The same analysis can also be performed on CAVATICA, but requires further optimization. The module describes the steps for CAVATICA set up, and scripts to publish an application on the portal. The required data files are also available publicly on CAVATICA under the <a href="Open Pediatric Cancer">Open Pediatric Cancer</a> (OpenPedCan) Open Access. Refer to the module for detailed description and scripts.

### **Abundance Estimation**

Among the data sources used for OpenPedCan, GTEx and TCGA used GENCODE versions v26 and v36, respectively. Moreover, the gene symbols used in these different GENCODE versions also varied. Therefore, the gene symbols had to be harmonized for compatibility to map unique gene identifiers to their gene symbols. ENSG IDs from each data source were pulled and mapped to the GTF/GFF3 file from GENCODE v39 to extract unique gene symbols and remove duplicates. Additionally, the gene expression matrices had some instances where multiple Ensembl gene identifiers mapped to the same gene symbol. This was dealt with by filtering the expression matrix to only genes with [FPKM/TPM] > 0 and then selecting the instance of the gene symbol with the maximum mean [FPKM/TPM/Expected\_count] value across samples. This enabled many downstream modules that require RNA-seq data have gene symbols as unique gene identifiers. Refer to collapse-rnaseq module for scripts and details.

### **Gene Expression Summary Statistics**

We generated RNA-Seq gene expression (TPM) summary statistics for independent tumor samples from the combined OpenPedCan gene expression matrices, including cancers from pediatric cohorts (PBTA, GMKF, and TARGET) and adult cancers from the TCGA cohort. We grouped selected samples into two groups containing samples from a cancer group in either each cohort or all cohorts, and calculated TPM means, standard deviations, gene-wise z-scores, group-wise z-scores, and ranks for each group as described in the <a href="OpenPedCan rna-seq-expression-summary-stats module">OpenPedCan rna-seq-expression-summary-stats module</a> in detail. The resulting gene-wise and group-wise summary statistics tables were annotated with EFO and MONDO disease codes associated with the cancer groups.

### Gene fusion detection

# **QUANTIFICATION AND STATISTICAL ANALYSIS**

## Focal Copy Number Calling (focal-cn-file-preparation analysis module)

Please refer to the OpenPBTA manuscript for details on assignment of copy number status values to CNV segments, cytobands, and genes [4]. We applied criteria to resolve instances of multiple conflicting status calls for the same gene and sample, which are described in detail in the <u>focal-cn-file-</u>

<u>preparation</u> module. Briefly, we prioritized 1) non-neutral status calls, 2) calls made from dominant segments with respect to gene overlap, and 3) amplification and deep deletion status calls over gain and loss calls, respectively, when selecting a dominant status call per gene and sample. These methods resolved >99% of duplicated gene-level status calls.

### **Gene Set Variation Analysis (gene-set-enrichment-analysis analysis module)**

Please refer to the OpenPBTA manuscript for details [4].

### Fusion prioritization (fusion\_filtering analysis module)

The <u>fusion\_filtering</u> module filters artifacts and annotates fusion calls, with prioritization for oncogenic fusions, for the fusion calls from STAR-Fusion and Arriba. After artifact filtering, fusions were prioritized and annotated as "putative oncogenic fusions" when at least one gene was a known kinase, oncogene, tumor suppressor, curated transcription factor, on the COSMIC Cancer Gene Census List, or observed in TCGA. Fusions were retained in this module if they were called by both callers, recurrent or specific to a cancer group, or annotated as a putative oncogenic fusion. Please refer to the module linked above for more detailed documentation and scripts.

### **Splicing quantification**

To detect alternative splicing events, we utilized rMATS turbo (v. 4.1.0) with Ensembl/GENCODE v39 GFF annotations using the <u>Kids First RNA-Seq workflow</u>. We used \_-variable-read-length and \_ t paired options and applied an additional filter to include only splicing events with total junction read counts greater than 10.

## Mutational Signatures (mutational-signatures analysis module)

# Tumor Mutation Burden [TMB] (tmb-calculation analysis module)

Recent clinical studies have associated high TMB with improved patient response rates and survival benefit from immune checkpoint inhibitors [doi.org/10.1002/gcc.22733?].

The Tumor Mutation Burden (TMB) tmb-calculation module was adapted from the snv-callers module of the OpenPBTA project [4]. Here, we use mutations in the snv-consensus-plus-hotspots.maf.tsv.gz file which is generated using Kids First DRC Consensus Calling Workflow and is included in the OpenPedCan data download. The consensus MAF contains SNVs or MNVs called in at least 2 of the 4 callers (Mutect2, Strelka2, Lancet, and Vardict) plus hotspot mutations if called in 1 of the 4 callers. We calculated TMB for tumor samples sequenced with either WGS or WXS. Briefly, we split the SNV consensus MAF into SNVs and multinucleotide variants (MNVs). We split the MNV subset into SNV calls, merged those back with the SNVs subset, and then removed sample-specific redundant calls. The resulting merged and non-redundant SNV consensus calls were used as input for the TMB calculation. We tallied only nonsynonymous variants with classifications of high/moderate consequence ("Missense\_Mutation", "Frame\_Shift\_Del", "In\_Frame\_Ins", "Frame\_Shift\_Ins", "Splice\_Site", "Nonsense\_Mutation", "In\_Frame\_Del", "Nonstop\_Mutation", and "Translation Start Site") for the numerator.

**All mutation TMB** For WGS samples, we calculated the size of the genome covered as the intersection of Strelka2 and Mutect2's effectively surveyed areas, regions common to all variant callers, and used this as the denominator. WGS\_all\_mutations\_TMB = (total # mutations in consensus MAF) / intersection\_strelka\_mutect\_vardict\_genome\_size For WXS samples,

we used the size of the WXS bed region file as the denominator. WXS\_all\_mutations\_TMB =
(total # mutations in consensus MAF)) / wxs\_genome\_size

Coding only TMB We generated coding only TMB from the consensus MAF as well. We calculated the intersection for Strelka2 and Mutect2 surveyed regions using the coding sequence ranges in the GENCODE v39 gtf supplied in the OpenPedCan data download. We removed SNVs outside of these coding sequences prior to implementing the TMB calculation below: WGS\_coding\_only\_TMB = (total # coding mutations in consensus MAF) / intersection\_wgs\_strelka\_mutect\_vardict\_CDS\_genome\_size For WXS samples, we intersected each WXS bed region file with the GENCODE v39 coding sequence, sum only variants within this region for the numerator, and calculate the size of this region as the denominator. WXS\_coding\_only\_TMB = (total # coding mutations in consensus MAF) / intersection\_wxs\_CDS\_genome\_size

Finally, we include an option (nonsynfilter\_focr) to use specific nonsynonymous mutation variant classifications recommended from the <a href="https://example.com/memory.com/me

### **Clinical Data Harmonization**

### **WHO Classification of Disease Types**

### **Molecular Subtyping**

Here, we build upon the molecular subtyping performed in OpenPBTA [4].

### High-grade gliomas.

A new high-grade glioma entity called infant-type hemispheric gliomas (IHGs), characterized by distinct gene fusions enriched in receptor tyrosine kinase (RTK) genes including *ALK*, *NTRK1/2/3*, *ROS1* or *MET*, was identified in 2021 [doi? 10.1038/s41467-019-12187-5]. To identify IHG tumors, first, tumors which were classified as "IHG" by the DKFZ methylation classifier or diagnosed as "infant type hemispheric glioma" from pathology\_free\_text\_diagnosis were selected [23]. Then, the corresponding tumor RNA-seq data were utilized to seek the evidence for RTK gene fusion. Based on the specific RTK gene fusion present in the samples, IHGs were further classified as "IHG, ALK-altered", "IHG, NTRK-altered", "IHG, ROS1-altered", or "IHG, MET-altered". If no fusion was observed, the samples were identified as "IHG, To be classified".

### Atypical teratoid rhabdoid tumors.

Atypical teratoid rhabdoid tumors (ATRT) tumors were categorized into three subtypes: "ATRT, MYC", "ATRT, SHH", and "ATRT, TYR" [24]. In OpenPedCan, the molecular subtyping of ATRT was based solely on the DNA methylation data. Briefly, ATRT samples with a high confidence DKFZ methylation subclass score (>= 0.8) were selected and subtypes were assigned based on the DKFZ methylation subclass [doi10.1038/nature26000?]. Samples with low confidence DKFZ methylation subclass scores (< 0.8) were identified as "ATRT, To be classified".

#### Neuroblastoma tumors.

Neuroblastoma (NBL) tumors with a pathology diagnosis of neuroblastoma, ganglioneuroblastoma, or ganglioneuroma were subtyped based on their MYCN copy number status as either "NBL, MYCN amplified" or "NBL, MYCN non-amplified". If pathology\_free\_text\_diagnosis was "NBL, MYCN non-amplified" and the genetic data suggested MYCN amplification, the samples were subtyped as

"NBL, MYCN amplified". On the other hand, if pathology\_free\_text\_diagnosis was "NBL, MYCN amplified" and the genetic data suggested MYCN non-amplification, the RNA-Seq gene expression level of MYCN was used as a prediction indicator. In those cases, samples with MYCN gene expression above or below the cutoff (TPM >= 140.83 based on visual inspection of MYCN CNV status) were subtyped as "NBL, MYCN amplified" and "NBL, MYCN non-amplified", respectively. MYCN gene expression was also used to subtype samples without DNA sequencing data. If a sample did not fit none of these situations, it was denoted as "NBL, To be classified".

### Integration of brain tumor methylation classifications

## TP53 Alteration Annotation (tp53\_nf1\_score analysis module)

Please refer to the OpenPBTA manuscript for details [4].

# Prediction of participants' genetic sex

Please refer to the OpenPBTA manuscript for details [4].

# Selection of independent samples (independent-samples analysis module)

For analyses that require all input biospecimens to be independent, we use the OpenPedCan-analysis <u>independent-samples</u> module to select only one biospecimen from each input participant. For each input participant of an analysis, the independent biospecimen is selected based on the analysis-specific filters and preferences for the biospecimen metadata, such as experimental strategy, cancer group, and tumor descriptor.

# **Supplemental Information Titles and Legends**

### Consortia

# References

- 1. **dbGaP Study** <a href="https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\_id=phs000218.v23.p8">https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\_id=phs000218.v23.p8</a>
- 2. **dbGaP Study** <a href="https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\_id=phs001436.v1.p1">https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\_id=phs001436.v1.p1</a>
- 3. **dbGaP Study** <u>https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\_id=phs002517.v1.p1</u>
- 4. OpenPBTA: The Open Pediatric Brain Tumor Atlas

Joshua A Shapiro, Krutika S Gaonkar, Stephanie J Spielman, Candace L Savonen, Chante J Bethell, Run Jin, Komal S Rathi, Yuankun Zhu, Laura E Egolf, Bailey K Farrow, ... Jaclyn N Taroni *Cell Genomics* (2023-07) <a href="https://doi.org/gr92p6">https://doi.org/gr92p6</a>

DOI: 10.1016/j.xgen.2023.100340 · PMID: 37492101 · PMCID: PMC10363844

5. **2021 X01 Projects Abstracts | NIH Common Fund**<a href="https://commonfund.nih.gov/kidsfirst/2021X01projects#FY21">https://commonfund.nih.gov/kidsfirst/2021X01projects#FY21</a> Resnick

6. Home

Chordoma Foundation <a href="https://www.chordomafoundation.org/">https://www.chordomafoundation.org/</a>

- 7. Michigan Center for Translational Pathology <a href="https://mctp.med.umich.edu">https://mctp.med.umich.edu</a>
- 8. **dbGaP Study** <a href="https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\_id=phs000424.v9.p2">https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\_id=phs000424.v9.p2</a>
- 9. **The Cancer Genome Atlas Program (TCGA) NCI** (2022-05-13) <a href="https://www.cancer.gov/ccg/research/genome-sequencing/tcga">https://www.cancer.gov/ccg/research/genome-sequencing/tcga</a>
- 10. Somalier: rapid relatedness estimation for cancer and germline studies using efficient genome sketches

Brent S Pedersen, Preetida J Bhetariya, Joe Brown, Stephanie N Kravitz, Gabor Marth, Randy L Jensen, Mary P Bronner, Hunter R Underhill, Aaron R Quinlan

Genome Medicine (2020-07-14) https://doi.org/gtsm62

DOI: 10.1186/s13073-020-00761-2 · PMID: 32664994 · PMCID: PMC7362544

11. The Ensembl Variant Effect Predictor

William McLaren, Laurent Gil, Sarah E Hunt, Harpreet Singh Riat, Graham RS Ritchie, Anja Thormann, Paul Flicek, Fiona Cunningham

Genome Biology (2016-06-06) https://doi.org/gdz75c

DOI: <u>10.1186/s13059-016-0974-4</u> · PMID: <u>27268795</u> · PMCID: <u>PMC4893825</u>

12. Strelka2: fast and accurate calling of germline and somatic variants

Sangtae Kim, Konrad Scheffler, Aaron L Halpern, Mitchell A Bekritsky, Eunho Noh, Morten Källberg, Xiaoyu Chen, Yeonbin Kim, Doruk Beyter, Peter Krusche, Christopher T Saunders *Nature Methods* (2018-07-16) <a href="https://doi.org/gdwrp4">https://doi.org/gdwrp4</a>

DOI: 10.1038/s41592-018-0051-x · PMID: 30013048

13. Calling Somatic SNVs and Indels with Mutect2

David Benjamin, Takuto Sato, Kristian Cibulskis, Gad Getz, Chip Stewart, Lee Lichtenstein *Cold Spring Harbor Laboratory* (2019-12-02) <a href="https://doi.org/ggntwv">https://doi.org/ggntwv</a>

DOI: <u>10.1101/861054</u>

## 14. Genome-wide somatic variant calling using localized colored de Bruijn graphs

Giuseppe Narzisi, André Corvelo, Kanika Arora, Ewa A Bergmann, Minita Shah, Rajeeva Musunuri, Anne-Katrin Emde, Nicolas Robine, Vladimir Vacic, Michael C Zody *Communications Biology* (2018-03-22) https://doi.org/gfcfr8

DOI: 10.1038/s42003-018-0023-9 · PMID: 30271907 · PMCID: PMC6123722

### 15. Control-FREEC: a tool for assessing copy number and allelic content using nextgeneration sequencing data

Valentina Boeva, Tatiana Popova, Kevin Bleakley, Pierre Chiche, Julie Cappo, Gudrun Schleiermacher, Isabelle Janoueix-Lerosey, Olivier Delattre, Emmanuel Barillot *Bioinformatics* (2011-12-06) <a href="https://doi.org/ckt4vz">https://doi.org/ckt4vz</a>

DOI: <u>10.1093/bioinformatics/btr670</u> · PMID: <u>22155870</u> · PMCID: <u>PMC3268243</u>

# 16. Control-free calling of copy number alterations in deep-sequencing data using GC-content normalization

Valentina Boeva, Andrei Zinovyev, Kevin Bleakley, Jean-Philippe Vert, Isabelle Janoueix-Lerosey, Olivier Delattre, Emmanuel Barillot

Bioinformatics (2010-11-15) <a href="https://doi.org/c6bcps">https://doi.org/c6bcps</a>

DOI: <u>10.1093/bioinformatics/btg635</u> · PMID: <u>21081509</u> · PMCID: <u>PMC3018818</u>

# 17. CNVkit: Genome-Wide Copy Number Detection and Visualization from Targeted DNA Sequencing

Eric Talevich, AHunter Shain, Thomas Botton, Boris C Bastian *PLOS Computational Biology* (2016-04-21) <a href="https://doi.org/c9pd">https://doi.org/c9pd</a>

DOI: <u>10.1371/journal.pcbi.1004873</u> · PMID: <u>27100738</u> · PMCID: <u>PMC4839673</u>

# 18. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data

Aaron McKenna, Matthew Hanna, Eric Banks, Andrey Sivachenko, Kristian Cibulskis, Andrew Kernytsky, Kiran Garimella, David Altshuler, Stacey Gabriel, Mark Daly, Mark A DePristo *Genome Research* (2010-07-19) <a href="https://doi.org/bnzbn6">https://doi.org/bnzbn6</a>

DOI: 10.1101/gr.107524.110 · PMID: 20644199 · PMCID: PMC2928508

# 19. **GISTIC2.0** facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers

Craig H Mermel, Steven E Schumacher, Barbara Hill, Matthew L Meyerson, Rameen Beroukhim, Gad Getz

Genome Biology (2011-04-28) https://doi.org/10.1186/gb-2011-12-4-r41

DOI: 10.1186/gb-2011-12-4-r41

# 20. Preprocessing, normalization and integration of the Illumina HumanMethylationEPIC array with minfi

Jean-Philippe Fortin, Timothy J Triche Jr, Kasper D Hansen

Bioinformatics (2016-11-29) https://doi.org/f9x7kd

DOI: 10.1093/bioinformatics/btw691 · PMID: 28035024 · PMCID: PMC5408810

### 21. Review of processing and analysis methods for DNA methylation array data

CS Wilhelm-Benartzi, DC Koestler, MR Karagas, JM Flanagan, BC Christensen, KT Kelsey, CJ Marsit, EA Houseman, R Brown

British Journal of Cancer (2013-08-27) https://doi.org/gb9qvv

DOI: <u>10.1038/bjc.2013.496</u> · PMID: <u>23982603</u> · PMCID: <u>PMC3777004</u>

# 22. Impact of SNPs on methylation readouts by Illumina Infinium HumanMethylation450 BeadChip Array: implications for comparative population studies

Patrycja Daca-Roszak, Aleksandra Pfeifer, Jadwiga Żebracka-Gala, Dagmara Rusinek, Aleksandra Szybińska, Barbara Jarząb, Michał Witt, Ewa Ziętkiewicz

BMC Genomics (2015-11-25) https://doi.org/gb3h5r

DOI: <u>10.1186/s12864-015-2202-0</u> · PMID: <u>26607064</u> · PMCID: <u>PMC4659175</u>

### 23. DNA methylation-based classification of central nervous system tumours

David Capper, David TW Jones, Martin Sill, Volker Hovestadt, Daniel Schrimpf, Dominik Sturm, Christian Koelsche, Felix Sahm, Lukas Chavez, David E Reuss, ... Stefan M Pfister *Nature* (2018-03-14) <a href="https://doi.org/gc5t36">https://doi.org/gc5t36</a>

DOI: <u>10.1038/nature26000</u> · PMID: <u>29539639</u> · PMCID: <u>PMC6093218</u>

# 24. Molecular subgrouping of atypical teratoid/rhabdoid tumors—a reinvestigation and current consensus

Ben Ho, Pascal D Johann, Yura Grabovska, Mamy Jean De Dieu Andrianteranagna, Fupan Yao, Michael Frühwald, Martin Hasselblatt, Franck Bourdeaut, Daniel Williamson, Annie Huang, Marcel Kool

Neuro-Oncology (2019-12-31) https://doi.org/gn3kcm

DOI: <u>10.1093/neuonc/noz235</u> · PMID: <u>31889194</u> · PMCID: <u>PMC7229260</u>