

```

0;

function x = solvetridiagonal(b, d1, d2, d3)
    % n is the length of the center diagonal
    n = length(d2)

    % Scan down to reduce the lower diagonal to zero and the pivots to 1.
    for i = 1:n
        % Rescale the row so the pivot is 1.
        a = d2(i);
        d2(i) /= a;
        if i < n
            d3(i) /= a;
        end
        b(i) /= a;

        % Cancel out all non zero elements in the column except for the pivot.
        if i < n
            factor = -d1(i);
            d1(i) += factor;
            d2(i+1) += factor * d3(i);
            b(i+1) += factor * b(i);
        end
    end

    % Scan back upwards and reduce the upper diagonal to zero and thus solving for x.
    for i = flip(2:n)
        factor = -d3(i-1);
        d3(i-1) += factor;
        b(i-1) += factor * b(i);
    end

    % Finally set x = b
    x = b;
end

% 2.a

n = 10000
% construct the 3 diagonals
d1 = - (2:n);
d2 = 3 * (1:n);
d3 = - (1:n-1);
% pick b to be something
b = 1:n;

%x = solvetridiagonal(b, d1, d2, d3)

% 2.b

function gt = g(t)
    gt = (pi/2)^2*sin(pi/2 * t)
end

n = 100
h = 1/n

d1 = repmat(-1/h^2, 1, n-1)
d2 = repmat(2/h^2, 1, n)
d3 = repmat(-1/h^2, 1, n-1)
b = g((1:n) * h)

v = solvetridiagonal(b, d1, d2, d3)

u = sin(pi/2 * (1:n) * h)

norm(v-u, inf)

```