

# 1218 - NIRSpec-IFU Observations of a QSOs at z=6

Cycle: 1, Proposal Category: GTO


**INVESTIGATORS**

| *Name* | *Institution* | *E-Mail* |
|---|---|---|
| **Dr. Pierre Ferruit (PI) (ESA Member)** | **European Space Agency - ESTEC** | **pierre.ferruit@esa.int** |
| Dr. Roberto Maiolino (CoI) (ESA Member) (Contact) | University of Cambridge | r.maiolino@mrao.cam.ac.uk |
| Dr. Santiago Arribas (CoI) (ESA Member) | Consejo Superior de Investigaciones Cientificas | arribas@cab.inta-csic.es |


**OBSERVATIONS**

| *Folder* | *Observation* | *Label* | *Observing Template* | *Science Target* |
|---|---|---|---|---|
| Observation Folder | | | | |
| | 2 | NIRSpec IFU observation of J0100 | NIRSpec IFU Spectroscopy | (2) J0100+2802 |


**ABSTRACT**

Update June 2019: We have removed: J1148 +5251 and we have added R100 observations for J0100 +2802

---

Previous submision (STScI Edit number=5)

This proposal is associated with NIRSpec Observation IDs:

FERRUIT_3051

FERRUIT_3052

The goal of the observations is to map (redshifted) optical nebular lines in the host galaxy and circumgalactic region of two luminous quasars at z~6. The observations (obtained with the high resolution grating) will also be used to detect and characterize the quasar-driven outflows.

**OBSERVING DESCRIPTION**

Contact: Roberto Maiolino (r.maiolino@mrao.cam.ac.uk)

The NIRSpec IFU observations are done with the G395H grating and are aimed aimed primarily at mapping the strongest optical nebular lines (Hbeta, [OIII], Halpha, [NII]) in order to trace the distribution and kinematics of the medium in  the host galaxy and the circumgalactic region.

We are not using TA as Gaia GS are available for any orientation, hence ensuring a pointing accuracy adequate for our goals.

We use NRSIRS2RAPID in order to better identify and remove cosmic rays.

PA has been restricted to minimise MSA leakage effects

**Fixed Targets**

| # | Name | Target Coordinates | Targ. Coord. Corrections | Miscellaneous |
|---|---|---|---|---|
| (2) | J0100+2802 | RA: 01 00 13.0200 (15.0542500d)<br>Dec: +28 02 55.80 (28.04883d)<br>Equinox: J2000 | | |

*Comments:*
*Category=Galaxy*
*Description=[High-redshift galaxies, Quasars]*
*Extended=YES*

## Observation

**Proposal 1218, Observation 2: NIRSpec IFU observation of J0100** Mon Jun 24 18:02:42 GMT 2019

**Diagnostic Status: Warning**

Observing Template: NIRSpec IFU Spectroscopy

## Diagnostics

(Visit 2:1) Warning (Form): Overheads are provisional until the Visit Planner has been run.

## Fixed Targets

| # | Name | Target Coordinates | Targ. Coord. Corrections | Miscellaneous |
|---|---|---|---|---|
| (2) | J0100+2802 | RA: 01 00 13.0200 (15.0542500d)<br>Dec: +28 02 55.80 (28.04883d)<br>Equinox: J2000 | | |

*Comments:*
*Category=Galaxy*
*Description=[High-redshift galaxies, Quasars]*
*Extended=YES*

## Template

| TA Method |
|---|
| NONE |

## Dithers

| # | Dither Type | Size | Starting Point | Number of Points | Points |
|---|---|---|---|---|---|
| 1 | CYCLING | MEDIUM | 1 | 4 | |

## Spectral Elements

| # | Grating/Filter | Readout Pattern | Groups/Int | Integrations/Exp | Leakcal | Dither | Autocal | Total Dithers | Total Integrations | Total Exposure Time | ETC Wkbk.Calc ID |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | G395H/F290LP | NRSIRS2RAPID | 185 | 1 | false | true | NONE | 4 | 4 | 10854.134 | |
| 2 | PRISM/CLEAR | NRSIRS2RAPID | 62 | 1 | false | true | NONE | 4 | 4 | 3676.4 | |
| 3 | PRISM/CLEAR | NRSIRS2RAPID | 62 | 1 | true | false | NONE | 1 | 1 | 919.1 | |

## Special Requirements

Aperture PA Range 201.892975 to 212.892975 Degrees (V3 63.0 to 74.0)