

■ Scientific Justification

Actively accreting supermassive black holes, i.e. quasars, are known to exist at very high redshift, when the Universe was much less than a billion years old. However, we currently do not know how these first supermassive black holes (SMBHs) formed. From estimates of the black hole mass, it is seen that SMBHs with $> 10^9 M_{\odot}$ exist less than a billion years after the Big Bang. Current models (e.g. Thispaper (2012); Thispaper (2012), Thispaper (2012)) suggestion either direct collapse black holes (DCBHs) or Population III stars could be the progenitors of the very-high redshift SMBHs.

We propose to obtain MIRI imaging, coronagraphy and spectroscopy of all known $z \geq 6.8$ quasars. MIRI covers 4.9-28.8 μm thus granting access to the optical/near-IR 0.63-3.7 μm rest-frame at $z = 6.8$.

The Science Case includes:

- How did the first SMBHs form?
- What are the BH masses from multiple lines (including rest-frame Balmer)?
- What are the host galaxy properties of $z > 6.8$ quasars?
- What is the stellar content of the quasars host galaxy? Does the Magorrian relation hold at very high- z ?
- Is there evidence for (major) merging?
- What is the environment of the $z > 6.8$ quasars?
- Whats the optical/near-IR spectral slope for the highest redshift quasars?
- Are the $z > 6.8$ luminous quasars different from ‘regular AGN’ detected in the JWST Deep Fields?
- What are the details of the Reionization Epoch?

These questions directly address two of the four *JWST* science themes (The End of the Dark Ages: First Light and Reionization; and Assembly of Galaxies) No telescope or observatory in the midterm future will be able to access wavelengths longer than 5 μm to these sensitivities.

Building on the very high redshift quasar catalog in the infrared from Ross & Cross (2020, MNRAS accepted) and from Dr. Sarah Bosmans current list: there are 20 objects with $z \geq 6.80$.

MIRI coronagraphy is a capability unique to JWST.

Some other key things to note: The JWST GTO teams are already looking at $z > 6.70$ quasars. However, only quasar J1342+0928 at $z = 7.54$ is currently being observed by MIRI. Pretty obviously(!) I would strongly suspect other teams will be looking at the very high-redshifts quasars. A **key point** here is not to directly compete with those observations, but moreover, complement them. As such, our team has a *a zero proprietary period* making the data immediately accessible from a webpages (e.g. github.com/d80b2t/JWST_Cycle1) The GTO and ERS programs already have the targets and instruments and modes that they are going to be using. One could then start to build-up on that arguing that youll never have the lambda 5um space-based data again any time again soon, and Generating a sample for a

large number (all) of the currently known $z > 6.7$ quasars is exactly what JWST was built for, and this will tremendously legacy value. This could/would be suggested as a multi-cycle program (though not necessarily explicitly as that I dont think this category is implemented yet). Using the Cycle 1+GTO+(ERS?) data as type of Pilot program, observing maybe ~ 5 -10 objects in total first, and letting that guide observations in future cycles.

■ **Technical Justification**

■ **Special Requirements (if any)**

■ **Justify Coordinated Parallel Observations (if any)**

■ **Justify Duplications (if any)**

■ **Analysis Plan (AR only)**