JWST Proposal 1283 (Created: Friday, June 21, 2019 at 12:00:48 PM Eastern Standard Time) - Overview



# 1283 - The MIRI HUDF Deep Imaging Survey

Cycle: 1, Proposal Category: GTO

# **INVESTIGATORS**

Name	Institution	E-Mail
Dr. H.U. Norgaard-Nielsen (PI) (ESA Member)	Danish Space Research Institute	hunn@space.dtu.dk
Prof. Pablo G. Perez-Gonzalez (CoI) (ESA Member)	Universidad Complutense de Madrid	pgperez@ucm.es

### **OBSERVATIONS**

Folder	Observation	Label	Observing Template	Science Target		
Observa	ation Folder			_		
	1	MIRI XDF NIRCam F1 15W+F277W	MIRI Imaging	(1) MDS.MIRI-DEEP-SURVEY		
	2	MIRI XDF NIRCam F1 15W+F277W/F356W	MIRI Imaging	(2) MDS-MIRI-SURVEY-2		
	3	MIRI XDF NIRCam F1 15W+F356W	MIRI Imaging	(3) MDS-MIRI-SURVEY-3		
	4	MIRI XDF NIRISS JH- bands	MIRI Imaging	(4) MSD-MIRI-SURVEY-4		
	5	MIRI XDF NIRISS K- band	MIRI Imaging	(5) MDS-MIRI-SURVEY-5		

# **ABSTRACT**

A key theme of JWST is to provide unique information about the formation and evolution of galaxies in the early universe. A lot of effort will be performed to examine the details of how the reionization of the Universe was evolving from the recombination phase at  $z \sim 1100$  to  $z \sim 6$  (Epoch of Reionization, EoR). One of the fundamental unknowns in this transition is the source of the ionizing photons during EoR.

EC-MIRI is participating in the multi-instrument NIRCam – NIRSpec - MIRI coordinated imaging and spectroscopic study of the HUDF and surrounding field. Following on the spirit of this setup, and with the goal of strengthening the primary objectives of MIRI GTO high-z science, the

JWST Proposal 1283 (Created: Friday, June 21, 2019 at 12:00:48 PM Eastern Standard Time) - Overview

EC-MIRI team proposes coordinated parallel observations with NIRISS and NIRCam of a well selected field around HUDF, while executing the MIRI deep HUDF imaging survey for a total of 60 hours. The proposed coordinated parallels split the time into about 20 hours of NIRISS observations and about 40 hours of NIRCam imaging. The selected strategies complement those of NIRCam and NIRSpec in the same area of the sky in two very important aspects:

- 1) The proposed NIRCam coordinated parallels will supplement the NIRCam Survey by providing deeper images in an additional pointing with three filters selected to optimize the detection of z>10 candidates
- 2) The proposed NIRISS coordinated parallels will provide deep wide field slitless low resolution spectroscopy of all high-z sources in a field of 2.2 x 2.2 arcmin2 within the NIRSpec WIDE MOS survey

# **OBSERVING DESCRIPTION**

The MIRI HUDF Deep Imaging Survey

Scientific Justification

The whole Universe exploded about 13.8 billion years ago in the Big Bang. The detailed physics of the evolution of the Universe from a tiny fraction of a second after Big Bang to the recombination of hydrogen, when the Universe was 380.000 years old, are well understood, mainly because of very detailed temperature and polarization maps of the Cosmic Microwave Background (CMB) radiation (the latest provided by the Planck Mission). Our understanding of the next period from the recombination to the epoch when the Universe has been fully ionized again is much more uncertain. The end of this transition is established by observations of significant fluxes short ward of Ly in QSOs with  $z \sim 6$ , when the Universe was about 1 billion years old (Fan et al. 2001, Fan et al. 2006).

The fundamental question for the reionization epoch is that the sources responsible for emitting the ionizing photons have not been identified. AGN's are known to emit a lot of UV photons, but their number density is peaking around z = 2 - 3 and is decreasing dramatically up  $z \sim 6$ . This means that they cannot provide a significant contribution to the reionization of the Universes. The most likely source is intrinsically faint star forming galaxies, so faint that they have been below the detection limits of the available instruments.

An important constrain on the reionization scenario is the optical depth of Thomson scattering of the electrons by the CMB photons. The Planck Mission (Planck Collaboration XX 2015) finds =  $0.066\pm0.012$ , somewhat smaller than the value found by WMAP 9yr (Bennett et al. 2013),  $0.089\pm0.014$ . Robertson et al. (2013, 2015) show that it is much easier to explain this lower value from the known UV luminosity function (MUV > -17) but it is still necessary to extrapolate the luminosity functions below MUV < -13 (Robertson et al. 2015).

Recently, three investigations of the total number of high redshifts sources detected by HST and UltraVISTA have been published (Bouwens et al. 2015, Finkelstein 2015, Bowler et al. 2015). For z ~ 7 and z ~ 8 they discuss many hundreds of sources. For z ~ 10, only 6 sources have been identified. HST is covering wavelengths 1.6 µm. This means that at these redshifts HST is observing rest – frame wavelengths in the UV, 2100Å,

JWST Proposal 1283 (Created: Friday, June 21, 2019 at 12:00:48 PM Eastern Standard Time) - Overview

1900Å and 1500Å, for these 3 redshift intervals. This implies that from HST photometry of high redshift objects, mainly information about the content, ages etc. of young stars can be extracted.

With the unique sensitivity above 5 µm, MIRI will play an important role in studying the different phases of EoR. With MIRI photometry, it will be possible for the first time to get:

- 1) an unbiased estimate of the total stellar masses in objects with z > 4,
- 2) determine masses and ages of the young and old stellar populations in objects with z > 8.5, exploiting clean measurements of the Balmer Break. From these masses and ages, information about the star formation rate can be derived, and thereby estimate how these populations have contributed with ionizing photons during the EoR.

Furthermore, with the 5.6  $\mu$ m filter, MIRI will be the only instrument onboard JWST which will be able to study the H emission line for sources with  $z \sim 7.5$  and the H + [OIII] lines for sources with  $z \sim 10$ .

Other high priority scientific areas, where this program will provide new insight, are 'Mass assembly evolution', 'Morphology evolution' and 'Obscured AGN'.

The MIRI Deep Imaging Survey will be performed with the  $5.6 \,\mu m$  filter on the HST UDF area covering both the ALMA and the MUSE Deep Surveys, spending about  $60 \,h$  GTO time. With the estimated sensitivity of this filter given in ETC, the detection limit is  $28.3 \,AB$ , S/N = 4.0. With this detection limit a small, but significant, number of sources will be detected with z > 9.

EC-MIRI is participating in the multi-instrument NIRCam – NIRSpec - MIRI coordinated imaging and spectroscopic study of the HUDF and surrounding field. Following on the spirit of this effort, and with the goal of strengthening the primary objectives of MIRI GTO high-z science, the EC-MIRI team proposes coordinated parallel observations with NIRISS and NIRCam of a well selected field around HUDF, while executing the MIRI deep HUDF imaging survey for a total of 60 hours. The proposed coordinated parallels split the time into about 20 hours of NIRISS observations and about 40 hours of NIRCam imaging. The selected strategies complement those of NIRCam and NIRSpec in the same area of the sky in two very important aspects:

- 1) As part of their cosmological survey, NIRCam will be getting a deep pointing on the HUDF to a (almost) uniform depth of AB  $\sim$ 30 mag over the full wavelength coverage of NIRCam. The proposed NIRCam coordinated parallels will built upon this set of data by providing deeper images in an additional pointing with three filters selected to optimize the detection of z>10 candidates using a Lyman break technique.
- 2) The proposed NIRISS coordinated parallels will provide deep wide field slitless low resolution spectroscopy of all high-z sources in a field of 2.2 x 2.2 arcmin2 around the HUDF with some of the deepest ancillary HST data. This set of data will be taken in a subsection of the planned NIRSpec WIDE MOS survey, and therefore complement the NIRSpec survey by providing deeper spectroscopy without pre-selection of targets.

  The MIRI EC Team is collaborating with the MIRI US, the NIRCam and the NIRSpec Teams on all the observations collected in HUDF within the

JWST Proposal 1283 (Created: Friday, June 21, 2019 at 12:00:48 PM Eastern Standard Time) - Overview JWST GTO program

Proposal 1283 - Targets - The MIRI HUDF Deep Imaging Survey

#	Name	Target Coordinates	Targ. Coord. Corrections	Miscellaneous	
(1)	MDS.MIRI-DEEP-SURVEY	RA: 03 32 38.1090 (53.1587875d)			
		Dec: -27 46 41.86 (-27.77829d)			
		Equinox: J2000			
Comments					
	=Unidentified on=[Blank field]				
(2)	MDS-MIRI-SURVEY-2	RA: 03 32 38.3160 (53.1596500d)			
		Dec: -27 46 41.86 (-27.77829d)			
		Equinox: J2000			
Comments					
_	=Unidentified on=[Blank field]				
(3)	MDS-MIRI-SURVEY-3	RA: 03 32 38.1090 (53.1587875d)			
(3)		Dec: -27 46 39.11 (-27.77753d)			
-		Equinox: J2000			
Comments Category=					
Category=	=Unidentified on=[Blank field]				
(4)	MSD-MIRI-SURVEY-4	RA: 03 32 37.9018 (53.1579242d)			
		Dec: -27 46 41.86 (-27.77829d)			
		Equinox: J2000			
Comments					
Category= Description	=Unidentified on=[Blank field]				
(5)	MDS-MIRI-SURVEY-5	RA: 03 32 38.1090 (53.1587875d)			
		Dec: -27 46 44.61 (-27.77906d)			
		Equinox: J2000			
Comments					
Category=	=Unidentified on=[Blank field]				

Pro	posal 1283	3 - Observation	1 - The MIRI	HUDF Deep I	maging Surv	ey				
ے	Proposal 1283, 0	Observation 1: MIRI X	DF NIRCam F115W	F277W					Fri Jun 21	17:00:48 GMT 2019
.፬	Diagnostic Statu	s: Warning								
Va	Observing Temp	late: MIRI Imaging								
ē	Coordinated Para	illel Template(s): NIRC	am Imaging							
Observation	with other JWST	to the uncertainty on the observations of HUDF ate the APT once the ov		_			o update the specificat	ions on the ranges of	PA angles, to assure a	n optimal overlab
8		ng (Form): Data volume								
Diagnostics	` /	ng (Form): Overheads a				36000.0 MB.				
osi	(VISIL 1.1) Walli	ng (Form). Overneaus a	ne provisional until the	Visit Flamet has been	Tuii.					
l ĝ										
<u>ä</u>										
-	# N:	ame	Target Coordin	ates		Targ. Coord. Correct	tions	Miscella	neous	
Fixed Targets	(1) M	DS.MIRI-DEEP-SURV		90 (53.1587875d)		8				
l ž	,		Dec: -27 46 41.8							
ΙË			Equinox: J2000	, , , , , ,						
l g	Comments:		1.							
Ιĉ	Category=Unide									
	Description=[Blo	апк пеіа [				NIDC I				
ate	MIRI Imaging					NIRCam Imaging				
Ιē	Subarray: FULL					Module: ALL				
Template						Subarray: FULL				
_	#	Dither Type	Starting Point	Number of Points	Points	Starting Set	Number of Sets	Optimized For	Direction	Pattern Size
ē	1	4-POINT-MIRI-	1	5		1	1			DEFAULT
Dithers		F1000W-WITH- NIRCam								
Elements	MIRI Imaging	Filter F	Readout Pattern Gro		ations/Exp Exposu	res/Dith Dither	Total Dither	rs Total Integrations	Total Exposure Time	ETC Wkbk.Calc ID
ΙĔ	1	F560W F	FAST 95	36	1	Dither 1	4	144	37962.547	
I≗										
<u></u>										
Spectral										
Ιĕ										
S										
Spectral Elements	NIRCam Imagir	ng Short Filter	Long Filter	Readout Pattern	Groups/Int	Integrations/Exp	<b>Total Integrations</b>	Total Dithers	Total Exposure Time	ETC Wkbk.Calc ID
ΙĔ	1	F115W	F277W	DEEP8	6	8	32	4	37406.907	
<u></u>										
ΙË										
ĕ										
Š										
								•		

<u>P</u>	Proposal 1283 - Observation 1 - The MIRI HUDF Deep Imaging Survey
	Aperture PA Range 294.38 to 334.59 Degrees (V3 289.930295 to 330.140295) Aperture PA Range 355.81 to 42.89 Degrees (V3 351.360295 to 38.440295) No Parallel Background Limited. Background no more than 10% above minimum
	හි   හි
	Special Specia
8	$\frac{\mathbf{a}}{\mathbf{b}}$

Pi	rOI	nosal 1283	- Ohservatio	on 2 - The M	IRI HLIDE	Deen	Imaging	ı Surv	ev					
Г				XDF NIRCam F1			magnic	Cuiv	Су				Fri Jun 21	17:00:48 GMT 2019
Observation	5	Diagnostic Status:		11221 1 1221 0 1111 1 2 2	277 1772								11104112	17.001.0 0.111 2019
1 2	3	Observing Templat	-											
	5		el Template(s): NIR	Cam Imaging										
4	2	Comments: Due to	the uncertainty on t	he date of the launci	h of JWST, we exp	ect, once	a reliable la	unch date	e has been	established	to update the specific	ations on the ranges o	of PA angles, to assure	an optimal overlab
lc			bservations of HUD e the APT once the i	F overheads for parall	el observations h	we heen i	mnlemented	in the AP	T softwar	,				
7	_			me for this visit 567										
Diagnostics	3			s are provisional unt										
	<u> </u>		, , , , , , , , , , , , , , , , , , , ,											
֓֞֝֞֝֞֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֡֡֓֓֓֓֓֓֡֡֝֓֡֓֡֡֓֡	ָּהָּ <u> </u>													
ڠ	5													
7	,	# Nan	ne	Target Co	ordinates			ŗ	Targ. Coo	rd. Correc	tions	Miscella	neous	
Targete	<u> </u>	(2) MD	S-MIRI-SURVEY-	2 RA: 03 32	38.3160 (53.1596	500d)								
1.6	3			Dec: -27 46	5 41.86 (-27.7782	9d)								
-	5			Equinox: J2	2000									
E Ve	2	Comments:												
Įü	-	Category=Unident Description=[Blan	ifiea k field]											
9	3	MIRI Imaging							NIRCar	n Imaging				
	2	Subarray: FULL							Module:	ALL				
Template	ξl								Subarray	: FULL				
Ľ														
Į,	5	#	Dither Type	Starting Point	-	f Points	Points		Starting	Set	Number of Sets	Optimized For	Direction	Pattern Size
Dithere		1	4-POINT-MIRI- F1000W-WITH- NIRCam		5									DEFAULT
Flomonte	3	MIRI Imaging	Filter	Readout Pattern	Groups/Int	Integr	ations/Exp	Exposur	res/Dith	Dither	Total Dither	rs Total Integrations	Total Exposure Time	ETC Wkbk.Calc ID
Į		1	F560W	FAST	95	18		1		Dither 1	4	72	18981.274	
1 🖁	}	2	F560W	FAST	95	18		1		Dither 1	4	72	18981.274	
7	3													
{	5													
Spectral	3													
$\vdash$	_		GI . TIII	T 7711			G 7		<b>.</b> .			m . 1511	T . 17	
4		NIRCam Imaging	Short Filter	Long Filter	Readout I	'attern	Groups/In	ıt	Integra	ions/Exp	<b>Total Integrations</b>	Total Dithers	Total Exposure Time	ETC Wkbk.Calc ID
	2	1	F115W	F277W	DEEP8		6		4		16	4	18681.98	
1 -	2	2	F115W	F356W	DEEP8		6		4		16	4	18681.98	
Spectral Fleme	5													
}	5													
] 2	2													
Ű	)													

<u>P</u>	roposal 1283 - Observation 2 - The MIRI HUDF Deep Imaging Survey
Pedilip	Aperture PA Range 294.38 to 334.59 Degrees (V3 289.930295 to 330.140295) Aperture PA Range 355.81 to 42.89 Degrees (V3 351.360295 to 38.440295) No Parallel Background Limited. Background no more than 10% above minimum
Special	

Pro	posal 128	3 - Observatio	on 3 - The M	IRI HUD	)F Deep	Imaging	Surve	ev					
ے	Proposal 1283,	Observation 3: MIRI	XDF NIRCam F1	15W+F356W	V							Fri Jun 21	17:00:48 GMT 2019
Į. <u>ē</u>	Diagnostic Stat	us: Warning											
Za	Observing Temp	plate: MIRI Imaging											
ē	Coordinated Par	allel Template(s): NIR	Cam Imaging										
Observation	overlab with oth	er JWST observations	of HUDF		_				blished,	to update the specific	ations on the ranges	of the PA angles, to ass	sure an optimal
<del>                                     </del>		late the APT, once the											
Diagnostics	` /	ning (Form): Data volum					owed of 5	8000.0 MB.					
St	(Visit 3:1) Warn	ning (Form): Overheads	s are provisional unt	il the Visit Pl	lanner has bee	n run.							
١Ĕ													
<u>ja</u> č													
-	# N	Name	Target Co	ordinatos			т	arg. Coord.	Correct	ions	Miscella	noons	
Fixed Targets		MDS-MIRI-SURVEY-		38.1090 (53.	1587875d)		1.	arg. Coord.	COLLECT	ions	Miscella	ncous	
ΙĒ	(3)	ADS-MIKI-SOK VL 1 -		5 39.11 (-27.7	, in the second								
⊢			Equinox: J	,	77733 <b>d</b> )								
e	Comments:		Equilion. 3.	2000									
۱ <u></u>	Category=Unide												
—	Description=[Bi	lank field]											
ag I	MIRI Imaging							NIRCam In					
٦	Subarray: FULL							Module: ALl					
Template								Subarray: FU	LL				
_	#	Dither Type	Starting Point	Numl	ber of Points	Points		Starting Set		Number of Sets	Optimized For	Direction	Pattern Size
je	1	4-POINT-MIRI-	-	5	oci oi i oines	Tomes		Starting Sec		Transer of Sets	Optimizeu 1 01	Direction	DEFAULT
Dithers	•	F1000W-WITH- NIRCam		J									DEFREE
Elements	MIRI Imaging	Filter	Readout Pattern	Groups/Int	Integ	rations/Exp	Exposure	es/Dith Di	her	Total Dither	Total Integrations	Total Exposure Time	ETC Wkbk.Calc ID
ΙĔ	1	F560W	FAST	95	36		1	Di	her 1	4	144	37962.547	
<u>=</u> [													
Spectral													
ĕ													
S													
Spectral Elements	NIRCam Imagi	ing Short Filter	Long Filter	Reado	out Pattern	Groups/Int	t .	Integrations	/Ехр	<b>Total Integrations</b>	<b>Total Dithers</b>	Total Exposure Time	ETC Wkbk.Calc ID
ΙĔ	1	F115W	F356W	DEEP	28	6		8		32	4	37406.907	
<u>=</u> [													
1 # 5													
l e													
ပ													

<u> </u>	oposai 1283 - Observation 3 - The Mirki HUDF Deep Imaging Survey
	Aperture PA Range 294.38 to 334.59 Degrees (V3 289.930295 to 330.140295) Aperture PA Range 355.81 to 42.89 Degrees (V3 351.360295 to 38.440295) No Parallel Background Limited. Background no more than 10% above minimum
Special	

Proposal 1283 - Observation 4 - The MIRI HUDF Deep Imaging Survey

Proposal 1283, Observation 4: MIRI XDF NIRISS JH-bands

Diagnostic Status: Warning

Observing Template: MIRI Imaging

Coordinated Parallel Template(s): NIRISS Wide Field Slitless Spectroscopy

Observation Comments: Due to the uncertainty in the date of the launch of JWST, we expect, once a reliable launch date has been established, to update the spefifications on the ranges of the PA angles, to assure an optinal overlab with other JWST observations of HUDF

Fri Jun 21 17:00:48 GMT 2019

Miscellaneous

We expect to update the APT, once the overheads on parallel observations have been implemented in the APT softare

(Visit 4:1) Warning (Form): Data volume for this visit 48183.79 MB exceeds half the maximum allowed of 58000.0 MB.

(Visit 4:1) Warning (Form): Overheads are provisional until the Visit Planner has been run.

Fixed Targets Name **Target Coordinates** Targ. Coord. Corrections MSD-MIRI-SURVEY-4 RA: 03 32 37.9018 (53.1579242d)

Dec: -27 46 41.86 (-27.77829d)

Equinox: J2000

Comments:

Diagnostics

Category=Unidentified Description=[Blank field]

MIRI Imaging NIRISS Wide Field Slitless Spectroscopy

Subarray: FULL

Dither Direct Images Primes: NO\_DITHERING

	Г	rs	#	Dither Type	Starting Point	t Number	of Points	Points	Starting	g Set	Number of Sets	Optimized For	Direction	Pattern Size
MIRI Imaging Filter Readout Pattern Groups/Int Integrations/Exp Exposures/Dith Dither Total Dithers Total Total Exposure ETC Wkbk.Cale	L	he	1						1		1			SMALL
	Ž	ă												
			MIRI Imaging	Filter	Readout Pattern	Groups/Int	Integr	ations/Exp	Exposures/Dith	Dither	Total Ditl			ETC Wkbk.Calc

ı		MIKI Imaging	Filter	Readout Pattern	Groups/Int	Integrations/Exp	Exposures/Dith	Dither	Total Dithers	Integrations	Total Exposure Time	ID WKbk.Calc
ı		1	F560W	FAST	92	1	1	None	1	1	255.304	
ı	Ŋ	2	F560W	FAST	97	9	1	Dither 1	4	36	9690.44	
ı	ments	3	F560W	FAST	92	1	1	None	1	1	255.304	
ı	Ĕ	4	F560W	FAST	92	1	1	None	1	1	255.304	
ı	Ele	5	F560W	FAST	97	9	1	Dither 1	4	36	9690.44	
ı		6	F560W	FAST	92	1	1	None	1	1	255.304	
ı	ctral	7	F560W	FAST	92	1	1	None	1	1	255.304	
ı	Spe	8	F560W	FAST	97	9	1	Dither 1	4	36	9690.44	
ı	$\overline{\mathbf{v}}$	9	F560W	FAST	92	1	1	None	1	1	255.304	
ı		10	F560W	FAST	92	1	1	None	1	1	255.304	
ı		11	F560W	FAST	97	9	1	Dither 1	4	36	9690.44	
1		12	F560W	FAST	92	1	1	None	1	1	255.304	

Proposal 1283 - Observation 4 - The MIRI HUDF Deep Imaging Survey

	NIRISS Wide Field Slitless Spectroscopy	Exposure Type	Filter	Grism	Readout Pattern	Groups/Int	Integrations/Exp	Total Dithers	Total Integrations	Total Exposure Time	ETC Wkbk.Calc ID
l	1	DIRECT	F115W		NIS	6	1	1	1	268.419	
ts	2	GRISM	F115W	GR150C	NIS	6	9	4	36	9663.093	
l e	3	DIRECT	F115W		NIS	6	1	1	1	268.419	
Elements	4	DIRECT	F115W		NIS	6	1	1	1	268.419	
	5	GRISM	F115W	GR150R	NIS	6	9	4	36	9663.093	
व्य	6	DIRECT	F115W		NIS	6	1	1	1	268.419	
Spectral	7	DIRECT	F150W		NIS	6	1	1	1	268.419	
၂ ရွိ	8	GRISM	F150W	GR150R	NIS	6	9	4	36	9663.093	
ľ	9	DIRECT	F150W		NIS	6	1	1	1	268.419	
l	10	DIRECT	F150W		NIS	6	1	1	1	268.419	
	11	GRISM	F150W	GR150C	NIS	6	9	4	36	9663.093	
	12	DIRECT	F150W		NIS	6	1	1	1	268.419	

Aperture PA Range 294.38 to 334.59 Degrees (V3 289.930295 to 330.140295) Aperture PA Range 355.81 to 42.89 Degrees (V3 351.360295 to 38.440295) No Parallel Special Requirements

Proposal 1283, Observation 5: MIRI XDF NIRISS K-band Fri Jun 21 17:00:48 GMT 2019 Observation **Diagnostic Status: Warning** Observing Template: MIRI Imaging Coordinated Parallel Template(s): NIRISS Wide Field Slitless Spectroscopy Comments: Due to the uncertainty on the date of the launch of JWST, we expect, once a reliable launch date has been established, to update the specifications on the ranges of PA angles, to assure an optimal overlab with other JWST observations We expect to update the APT once the overheads for parallel observations have been implemented in the APT software **Diagnostics** (Visit 5:1) Warning (Form): Overheads are provisional until the Visit Planner has been run. **Targ. Coord. Corrections** Miscellaneous Fixed Targets Name **Target Coordinates** (5) MDS-MIRI-SURVEY-5 RA: 03 32 38.1090 (53.1587875d) Dec: -27 46 44.61 (-27.77906d) Equinox: J2000 Comments: Category=Unidentified Description=[Blank field] Template MIRI Imaging **NIRISS Wide Field Slitless Spectroscopy** Subarray: FULL Dither Direct Images Primes: NO\_DITHERING Dithers Dither Type **Starting Point** Number of Points Points **Starting Set** Number of Sets **Optimized For** Direction Pattern Size 2 4-POINT-MIRI-**SMALL** 1 F1000W-WITH-NIRISS | N | 1 | 2 | MIRI Imaging Filter Readout Pattern Groups/Int Integrations/Exp Exposures/Dith Dither **Total Dithers** Total Total Exposure ETC Wkbk.Calc Integrations Time ID F560W 92 255.304 **FAST** 1 1 None 1 F560W **FAST** 90 9 1 Dither 1 4 36 8991.13

L	ı 1~		1 300 11	17101	70	/	•	Dittier 1	•	50	0771.13	
			F560W	FAST	92	1	1	None	1	1	255.304	
Į	$\begin{bmatrix} 3 \\ 4 \end{bmatrix}$		F560W	FAST	77	1	1	None	1	1	213.678	
	<b>D</b> 5		F560W	FAST	90	9	1	Dither 1	4	36	8991.13	
Ú	6	I	F560W	FAST	77	1	1	None	1	1	213.678	
1 3	≝  F	VIRISS Wide Field Slitless Spectroscopy	Exposure Type	Filter	Grism	Readout Pattern	Groups/Int	Integrations/Exp	<b>Total Dithers</b>	Total Integrations	Total Exposure Time	ETC Wkbk.Calc ID
}	S   S   1   2		DIRECT	F200W		NIS	6	1	1	1	268.419	
1	2		GRISM	F200W	GR150R	NIS	5	10	4	40	9018.887	
3	<b>ਰ</b> 3		DIRECT	F200W		NIS	6	1	1	1	268.419	
13	ة  4		DIRECT	F200W		NIS	5	1	1	1	225.472	
	5 S		GRISM	F200W	GR150C	NIS	5	10	4	40	9018.887	
Ľ	6	I	DIRECT	F200W		NIS	5	1	1	1	225.472	

<u>Pro</u>	oposal 1283 - Observation 5 - The MIRI HUDF Deep Imaging Survey
Requirements	Aperture PA Range 294.38 to 334.59 Degrees (V3 289.930295 to 330.140295) Aperture PA Range 355.81 to 42.89 Degrees (V3 351.360295 to 38.440295) No Parallel Background Limited. Background no more than 10% above minimum
Special	