



WISE - the Wide-field Infrared Survey Explorer

Ned Wright (UCLA)

Project Overview




Science

- Sensitive all sky survey with 8X redundancy
 - Find the most luminous galaxies in the universe
 - Find the closest stars to the sun
 - Provide an important catalog for JWST
 - Provide lasting research legacy


Salient Features

- 4 imaging channels covering 3 - 25 microns wavelength
- 40 cm telescope operating at <17K
- Two stage solid hydrogen cryostat
- Delta launch from WTR: 14 Dec 2009
- Sun-synchronous 6am 530km orbit
- Scan mirror provides efficient mapping
- Expected life: 10 months, actual 7.7-9.5
- 4 TDRSS tracks per day

Wide Field Infrared Survey Explorer



Infrared



- Optical
- Reflected light


Near-IR
different colors

Thermal-IR
emitted radiation



Why Space?


“Ground-based infrared astronomy is like observing stars in broad daylight with a telescope made out of fluorescent lights” — George Rieke.



40 cm WISE
telescope in
space equals
six thousand
8-meter
telescopes on
the ground!



Animated Scan Mirror Icon





WISE Science Heritage

- NIRAS SMEX proposal in 1988.
CJL & ELW as co-I's.
- Review panel suggested ground-based survey which became 2MASS.
- NIRST mission concept study in 1994. PRME & ELW as co-PI's.
- NGSS proposal in 1998.
- NGSS Phase A study in 1999.
- NGSS proposal in 2001.
- NGSS Phase A study in 2002.
- Renamed WISE in 2003

PROPOSAL TO
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
FOR A
NEAR-INFRARED ASTRONOMY SATELLITE

P1976-9-88

For the period 1 July 1989 through 30 September 1993

Total Estimated Cost: \$15,457,092
(not including costs from NASA Centers)

Volume I - Investigation and Technical Plan

Principal Investigator

Dr. Giovanni G. Fazio
Smithsonian Astrophysical Observatory
60 Garden Street
Cambridge MA 02138
(617) 495-7458

Associate Director for
Optical and Infrared Astronomy
Dr. David W. Latham

Co-Investigators

Smithsonian Astrophysical Observatory
Dr. John P. Huchra
Dr. David G. Koch
Dr. Gary J. Melnick
Dr. Steven P. Willner

University of California
at Los Angeles
Dr. Edward L. Wright

California Institute of Technology
Infrared Processing and Analysis Center
Dr. Carol Lonsdale, Lead Scientist
Dr. Charles A. Beichman
Dr. Baruch T. Soifer

Space Telescope
Science Institute
Dr. Richard Burg, Lead Scientist
Dr. Brian McLean
Dr. Michael Shara

California Institute of Technology
Dr. Jeremy R. Mould
Dr. I. Neill Reid

Jet Propulsion Laboratory
Dr. Richard Capps

NASA/Ames Research Center
Dr. Craig R. McCreight

Air Force Geophysics Laboratory
Dr. Stephan D. Price

September 1988

Dr. Irwin I. Shapiro
Director

Smithsonian Institution
Astrophysical Observatory
Cambridge, Massachusetts 02138



NIRAS 1988

- All sky survey.
- Big arrays (58x62).
- Continuous slew.
- Scan mirror to freeze image on array.
- Exposures every 4.5 seconds.




Figure 3-3. Optical Diagram of the Telescope

A scan mirror is used to freeze the sky field-of-view on the detectors for a period of 4.3 seconds as the spacecraft rotates at the orbital rate. The field-of-view is then advanced within 0.2 seconds.


Table 3-1. NIRAS Sensitivity

WAVELENGTH (microns)	BANDWIDTH (microns)	S/N	NUMBER OF PASSES	FLUX DENSITY		
				POINT SOURCE (mag)	DIFFUSE (mJy)	DIFFUSE (μ Jy/arcsec 2)
1.87	1.14	1	1	14.6	1.2	1.8
		5	1	12.8	6.0	
		1	4	15.3	0.6	0.9
		5	4	13.6	3.0	
3.52	2.16	1	1	13.4	1.2	1.7
		5	1	11.7	6.0	
		1	4	14.2	0.6	0.9
		5	4	12.4	3.0	




WISE Survey Strategy Provides Minimum of 8 Exposures Per Position

- Scan mirror enables efficient surveying
 - 8.8-s exposure/11-s duty cycle
- 10% frame to frame overlap
- 90% orbit to orbit overlap
- Sky covered in 6 months observing




- Single observing mode
- Minimum 8, median 14 exposures/position after losses to Moon and SAA






Asteroids Move

- Four frames of data taken on 2010 Jan. 8 during in-orbit checkout.
- Blue = 3.6um; green = 4.6um; red = 12um
- Circled asteroids are (L to R in the first frame, diameters in km):
17818 MBA D~12.4
153204 MBA D~2.8
22006 MBA D~11.5
87355 MBA D~4.3
80590 MBA D~4.1



Field of view = 34 x 25 arcmin (whole WISE FOV is 47 x 47 arcmin)


2010 AB₇₈





Results

- Diameter 1.3 km
- Anomalously faint in July, giving rotation pole at $(\alpha,\delta) = (185 \pm 26^\circ, -35 \pm 13^\circ)$
 - Obliquity of pole is $90 \pm 17^\circ$ with respect to the orbit
- Bond albedo $A=0.02$, $p_V=0.05$
- $\Theta = 0.55$ at 1 AU from Sun
- $\sqrt{\kappa\rho C} = 220 \pm 110 \text{ J/K/m}^2/\sqrt{\text{s}}$






National Aeronautics and Space
Administration
Jet Propulsion Laboratory
California Institute of Technology

Wide-field Infrared Survey Explorer (WISE)



NEO albedo distribution model



UCLA

ELW - 12
04 Apr 13



Most Hazardous WISE Discovery

- 2010 MU₁₁₂ - recovered in Feb 2013 by David Tholen
- Minimum Orbit Intersection Distance = 0.0011 AU
- Closest approach in next hundred years, 12 Dec 2082 at 0.007 AU
- Diameter 600 m, Albedo = 3%, estimated mass 200 megatons
- $a = 1.756 \text{ AU}$, $e = 0.54$, $i = 48^\circ$
- $v_\infty = 29.5 \text{ km/sec}$
- Impact energy in TNT equivalent = $\text{Mass} * (v_\infty^2 + 11^2) / 2.9^2$

–24 billion tons of TNT




National Aeronautics and Space
Administration
Jet Propulsion Laboratory
California Institute of Technology

Wide-field Infrared Survey Explorer (WISE)





Inhabitants of WISE Color Space





SDSS Classifications:

- Galaxies
 - $z \sim 0.4$ LIRGs
 - Local LIRGs
 - Local ULIRGs
 - QSOs
-
- Blackbodies
 - Power Laws




- Distance 2 pc! Proper motion 2.8"/yr
- Luhman (2013) trawled through the WISE single frame detections and found this lurking nearby.

Wide-field Infrared Survey Explorer (WISE)
WISE Color Space
A Very Red...Blue Compact Galaxy



WISE image




SDSS image

- $z=0.0425$
- $Z \sim 1/12 Z_{\odot}$



BCDs & Green Peas

- WISE colors of BCDs and green peas
- From C-W Tsai et al poster 333.11 at the Jan 2011 AAS meeting
- Griffith et al, 2011 ApJL, 736 L22 (arXiv:1106.4844)



A semi-WISE team result



A variable mid-infrared synchrotron break associated with the compact jet in GX 339–4



P. Gandhi¹, A.W. Blain², D.M. Russell³, P. Casella⁴, J. Malzac^{5,6}, S. Corbel⁷, P. D'Avanzo⁸, F.W. Lewis⁹, S. Markoff³, M. Cadolle Bel¹⁰, P. Goldoni^{11,12}, S. Wachter¹³, D. Khangulyan¹ and A. Mainzer¹⁴

ABSTRACT

Many X-ray binaries remain undetected in the mid-infrared, a regime where emission from their compact jets is likely to dominate. Here, we report the detection of the black hole binary GX 339–4 with the Wide-field Infrared Survey Explorer (WISE) during a very bright, hard accretion state in 2010. Combined with a rich contemporaneous multiwavelength dataset, clear spectral curvature is found in the infrared, associated with the peak flux density expected from



Flaring black hole GX339-4



- WISE observations occurred during a very active period




National Aeronautics and Space
Administration
Jet Propulsion Laboratory
California Institute of Technology

Wide-field Infrared Survey Explorer (WISE)



The far-off Universe




UCLA



L^* at $z=0.33$, $z=6.4$ QSO, $z=3$ ULIRG: FSC15307 x 3

ELW - 20
04 Apr 13

WISE Band 1 and 2 Dropouts




- $W1 > 17.4$ and $W2 > 15.9$ and ($W3 < 10.6$ or $W4 < 7.7$)
- W1814+34 (Eisenhardt et al 2012, ApJ 755, 173; Bridge et al 2012)
- $z=2.452$
- Extended Lyman alpha emission (~ 40 kpc)



SED of W1814+34




- AGN with $A_V = 50$
- Starburst
- Spiral Galaxy
- Warm Spitzer data to get 3.6 & 4.5 μm since WISE did not detect it at 3.4 & 4.6 μm .
- SHARC II (CSO) at 350 μm
- VLA radio data
- Peak $vL_v = 10^{13.38} L_\odot$






Warm Spitzer Followup

- Objects not detected by WISE at 3.4 & 4.6 μm can be measured using warm Spitzer
 - bigger mirror
 - longer integration times
- Synergy between surveys and great observatories







Herschel Followup Program



- Example: W2207+19
- Warm Spitzer at 3.6 & 4.5 μm
- WISE at 12 & 22 μm
- Herschel at 70, 160, 250, 350 & 500 μm
- Peak vF_v at 22 μm
- Peak $vL_v = 10^{13.13} L_\odot$








Many W12 drops


- About 1000/sky
- High percentage with high z 's: see histogram
- Spitzer followup usually picks up 3.6 and 4.5 μm flux
- Herschel followup usually detects far-IR flux






AGN Selection

- Stern et al
2012, ApJ,
753, 30
- Density
70/sq.deg
- 60% have
published
z's in
COSMOS
field







Z-distribution




ULAS 1120+0641




- $W1-W2 \approx 1.17 \pm 0.31$
- $\approx 43 \pm 8 \mu\text{Jy}$ at $3.4 \mu\text{m}$
- $z = 7.085$
- Mortlock et al, 2011, Nature, 474, 616, arXiv:1106.6088








Blain et al 2013





W1 image of SPT z=1.13 cluster




20 60 100 140


WISE $z \sim 1.3$ Galaxy Cluster Candidate


WISE W1

WISE $z \sim 1.3$ Galaxy Cluster Candidate



SDSS r





r J K (Subaru)
WISE z ~ ~~1.3~~
0.99 Galaxy Cluster Candidate




Blind Survey of WISE Sources

- Lake et al, arXiv:1111.0341
- 762 DEIMOS (on Keck) spectra of “all” WISE sources in FoV’s centered on 10 ULIRG candidates
- Three different levels of W1 flux limits
 - $W1 > 120 \mu\text{Jy}$, the required sensitivity
 - $W1 > 80 \mu\text{Jy}$, the all-sky achieved sensitivity
 - No limit on $W1$, with many sources pulled in by the other bands
- For $W1 > 120 \mu\text{Jy}$, 60% of all high-latitude sources are galaxies with median redshift 0.3
- Stars at high $|b|$ are mainly M dwarfs




Stellar Type Histogram





Redshift Histogram





NGC 628



NGC 1398



NGC 1566



NGC 2403




M 81



M 51



M 83



M 101



NGC 5907



NGC 6822




NGC 6946




IC 342



Warming Telescope




- Data from 8/7/10 to 9/30/10 released in June 2012.
- AllWISE coaddition of data from 10/1/10 to 2/1/11 will give proper motions and deeper 3.4 & 4.6 μ m catalogs.





Proper Motion Sensitivity






National Aeronautics and Space
Administration
Jet Propulsion Laboratory
California Institute of Technology

Wide-field Infrared Survey Explorer (WISE)



Coverage Released 14 Mar 2012

Actual Coverage Achieved for W4




UCLA

ELW - 46
04 Apr 13



More Coverage upto 9/30/10


1884474 frames thru 10-273.0; 68.0% to 16x+





Final 2 band coverage

2784184 frames thru end of mission





Thus WISE has

- Discovered many new NEOs and potentially hazardous asteroids and gave radiometric diameters for nearly 160,000 objects.
- Searched for the $\frac{1}{2}$ to $\frac{2}{3}$ of the stars in the solar neighborhood that have not yet been seen, including the closest stars to the Sun.
- Surveyed star formation in the Milky Way and in massive Ultra-Luminous Infrared Galaxies.

- Or at least we have the data now: 10 trillion pixels worth. We have lots of work left analyzing this treasure trove of information.



WISE Summary

- Launched 14 Dec 2009
- Band centers 3.4, 4.6, 12 & 22 microns
- Sensitivity better than 0.08, 0.11, 1 & 6 mJy
- Saturation at 0.3, 0.5, 0.7 & 10 Jy point sources
- Angular Resolution 6, 6, 6 & 12 arc-seconds
- Position accuracy about 0.15 arc-seconds 1σ 1-axis for high SNR
- Completed all-sky survey 17 July, big tank ran out hydrogen 5 Aug, little tank empty on 29 Sep, two-band survey for asteroids continued until 1 Feb 2011.
- Data releases:
 - Preliminary release of 57% of the sky on 14 April 2011
 - All-sky Release 14 March 2012
 - Three band data release 27 June 2012
 - Two band single image release 31 July 2012
- Data products include image atlas and source catalog



Two Proposals Under Review

- WITS (Wide-field Infrared Time-Domain Survey) submitted to NASA Astrophysics Explorer MoO call
 - TURN WISE back ON
 - Scan wider range of solar elongations to get observations over two week period every six months for two years
 - Measure variability of MW Cepheids, Miras, QSOs
 - Better SNR on $z=1$ galaxies
- Re-Animation of NEOWISE – submitted to the Human Exploration & Operations Mission Directorate (HEOMD)
 - TURN WISE back ON
 - Resume WISE scan pattern to find more NEOs
 - Obviously also gets variability of Miras, QSOs and better SNR on $z=1$ galaxies



JWST is coming

- JWST covers the WISE bands.
- Mining the WISE data to find targets for JWST will pay off dramatically in 5 years.

