

# Near and Mid-infrared properties of known $z \geq 5$ Quasars

Nicholas P. Ross<sup>1\*</sup> and Nicholas J. G. Cross<sup>1</sup>

<sup>1</sup>*Institute for Astronomy, University of Edinburgh, Royal Observatory, Edinburgh, EH9 3HJ, United Kingdom*

28 May 2019

## ABSTRACT

We assemble a catalogue of 463 spectroscopically confirmed very high ( $z \geq 5.00$ ) redshift quasars and report their near ( $zZyYJHK_s$  and  $K$ ) infrared and mid-infrared (WISE) properties. Using archival WFCAM/UKIRT and VIRCAM/VISTA data we check for photometric variability in the near-infrared that might be expected from super-Eddington accretion and find *blah*. Extrapolating the known quasar luminosity function we suggest that  $x\%$  of the possibly detected  $z \geq 5$  quasars in the current datasets have been discovered. All the data, analysis codes and plots used and generated here can be found at: [github.com/d80b2t/VHzQ](https://github.com/d80b2t/VHzQ).

**Key words:** Astronomical data bases: surveys – Quasars: general – galaxies: evolution – galaxies: infrared.

## 1 INTRODUCTION

Very high redshift quasars (VHzQ; defined here to have redshifts  $z \geq 5.00$ ) are excellent probes of the early Universe. This includes studies of the Epoch of Reionization for hydrogen (see e.g. Fan et al. 2006; Mortlock 2016, for reviews), the formation and build-up of supermassive black holes (e.g., Rees 1984; Wyithe & Loeb 2003; Volonteri 2010; Agarwal et al. 2016; Valiante et al. 2018; Latif et al. 2018; Wise et al. 2019) and early metal enrichment (see e.g., Simcoe et al. 2012; Chen et al. 2017; Bosman et al. 2017).

Super-critical accretion, where  $\dot{M} > \dot{M}_{\text{Edd}}$ , is a viable mechanism to explain the high, potentially super-Eddington, luminosity and rapid growth of supermassive black holes in the early universe (e.g., Alexander & Natarajan 2014; Madau et al. 2014; Volonteri et al. 2015; Pezzulli et al. 2016; Lupi et al. 2016; Pezzulli et al. 2017; Takeo et al. 2018). Thus, one could expect VHzQs to potentially vary in luminosity as they go through phases of super-critical accretion. These signatures of photometric variability should be looked for, noting the rest-frame optical emission is redshifted into the observed near-infrared (NIR) at redshifts  $z > 5$ . Fortunately, data are now in place from deep, wide-field NIR instruments and surveys such as the Wide Field Camera (WFCAM) instrument on the United Kingdom Infra-Red Telescope (UKIRT) in the Northern Hemisphere and the VISTA InfraRed CAMera (VIRCAM) on the Visible and Infrared Survey Telescope for Astronomy (VISTA) in the Southern Hemisphere, that are necessary for identifying VHzQs.

Quasars are known to be prodigious emitters of infrared

emission, thought to be from the thermal emission of dust grains heated by continuum emission from the accretion disc (e.g., Richards et al. 2006; Leipski et al. 2014; Hill et al. 2014; Hickox et al. 2017). Observations in the mid-infrared, e.g.  $\sim 3\text{--}30\mu\text{m}$  allow discrimination between AGN<sup>1</sup> and passive galaxies due to the  $1.6\mu\text{m}$  “bump” entering the MIR at  $z \approx 0.8\text{--}0.9$  (e.g., Wright et al. 1994; Sawicki 2002; Lacy et al. 2004; Stern et al. 2005; Richards et al. 2006; Timlin et al. 2016) as well as between AGN and star-forming galaxies due to the presence of Polycyclic Aromatic Hydrocarbon (PAHs) at  $\lambda > 3\mu\text{m}$  (e.g., Yan et al. 2007; Tielens 2008).

Jiang et al. (2006) and Jiang et al. (2010) report on the discovery of a quasar without hot-dust emission in a sample of 21  $z \approx 6$  quasars. Such apparently hot-dust-free quasars have no counterparts at low redshift. Moreover, those authors demonstrate that the hot-dust abundance in the 21 quasars builds up in tandem with the growth of the central black hole. But understanding how dust first forms and appears in the central engine remains an open question (Wang et al. 2008, 2011).

WISE mapped the sky in 4 passbands, in bands centered at wavelengths of 3.4, 4.6, 12, and  $23\mu\text{m}$ . The all sky ‘ALLWISE’ catalogue release, contains nearly 750 million detections at high-significance<sup>2</sup>, of which over 4.5M AGN candidates have been identified with 90% reliability (Assef

<sup>1</sup> Historically, “quasars” and “Active Galactic Nuclei (AGN)” have described different luminosity/classes of objects. In recognition of the fact that both terms describe accreting supermassive black holes, we use these terms interchangeably, with a preference for quasar, since we are generally in the higher- $L$  regime (e.g. Haardt et al. 2016).

<sup>2</sup> [wise2.ipac.caltech.edu/docs/release/allwise/expsup/sec2.1.html](https://wise2.ipac.caltech.edu/docs/release/allwise/expsup/sec2.1.html)

\* Corresponding Author: [npross@roe.ac.uk](mailto:npross@roe.ac.uk)


et al. 2018). Blain et al. (2013) presented WISE mid-infrared (MIR) detections of 17 (55%) of the then known 31 quasars at  $z > 6$ . However, Blain et al. (2013) was compiled with the WISE ‘All-Sky’ data release, as opposed to the superior AllWISE catalogues. That sample only examined the 31 known  $z > 6$  quasars; our sample has 170 objects with redshift  $z \geq 6.00$  (with 108 detected in WISE). Bañados et al. (2016) reports WISE W1, W2, W3 and W4 magnitudes for the Panoramic Survey Telescope and Rapid Response System 1 (Pan-STARRS1, PS1; Kaiser et al. 2002, 2010), but with no further investigation into the reddest WISE waveband for the VHzQs.

Critically, we now have available to us new W1 and W2 photometry from the ‘unWISE Source Catalog’ (Schlafly & Meisner 2018), a WISE-selected catalogue that is based on significantly deeper imaging and has a more extensive modeling of crowded regions than the ALLWISE release. For the first time in a catalogue, unWISE takes advantage of the ongoing mid-IR Near-Earth Object Wide-Field Infrared Survey Explorer Reactivation mission (NEOWISE-R; Mainzer et al. 2014), and achieves depths  $\sim 0.7$  mag deeper than ALLWISE (in W1/2). This additional depth is a significant advantage in the detection and study of VHzQs in the 3–5 micron regime.

Here we present for the first time the combined near-infrared properities (from UKIRT and VIRCAM) and the new mid-infrared unWISE for all the spectroscopically known  $z \geq 5.00$  quasars. Our motivations are numerous and include: (i) establishing the first complete catalogue of  $z > 5.00$  quasars since the pioneering work from SDSS; (ii) utilizing all the WFCAM and VISTA near-infrared photometry available for the quasars; (iii) making the first study of near- and mid-IR variability of the VHzQ population and (iv) establishing the photometric properties for upcoming surveys and telescopes including the Large Synoptic Survey Telescope (LSST)<sup>3</sup>, ESA *Euclid*<sup>4</sup> and the *James Webb Space Telescope* (JWST)<sup>5,6,7,8</sup>. We chose redshift  $z = 5.00$  as our lower redshift limit due to a combination of garnishing a large sample, adequately spanning physical properties (e.g. luminosity, age of the Universe) and to highlight the parts of  $L - z$  parameter space where  $z > 5$  quasars still wait to be discovered.

This paper is organized as follows. In Section 2, we present the assembled list of the 463  $z \geq 5.00$  VHzQs that we have compiled. We then give a high-level overview of the photometric surveys and datasets we use and present the photometry of the VHzQs. In Section 3 we ... and In Section 4 we ... We conclude in Section 5 and present all the necessary details to obtain our dataset in the Appendices.

We present all our photometry and magnitudes on the AB zero-point system (Oke & Gunn 1983; Fukugita et al. 1996). This includes the near-infrared, as well as the mid-infrared magnitudes. Appendix ?? gives the AB to Vega transforms for a wide range of optical, NIR and MIR filters. We use a flat  $\Lambda$ CDM cosmology with  $H0 = 67.7$  km s<sup>−1</sup>



**Figure 1.** The spectral bands used by different survey telescopes and that are relevant here. The *grizy* filters are from the Pan-STARRS survey. The *JHK* are from UKIRT/WFCAM, while *K<sub>s</sub>* is a VISTA/VIRCAM filter. The narrow W-band centered at  $\lambda \approx 14,500\text{\AA}$  is a CFHT/Wircam filter. [NJC: Do we use the W-band anywhere?] The WISE passbands W1–4 are also presented. The quasar spectrum is a composite based on Vanden Berk et al. (2001) and Bañados et al. (2016). The L and T dwarf spectra are from Cushing et al. (2006).

Mpc1,  $\Omega_M = 0.307$ , and  $\Omega_\Lambda = 0.693$  (Planck Collaboration et al. 2016) to be consistent with Bañados et al. (2016) and all logarithms are to the base 10.

## 2 METHOD AND DATA

Quasars are generally identified by photometric selection followed by spectroscopic confirmation. Here, we reverse this method obtaining first a list of spectroscopic quasars and then obtain photometric information.

We have compiled a list of 463 quasars with redshifts  $z \geq 5.00$ . We use all the  $z \geq 5.00$  quasars that have been discovered, spectroscopically confirmed and published as of 2018 December 31 (MJD 58483). We then obtain optical, near-infrared and mid-infrared photometry for the spectral dataset. The optical data comes from the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS) survey (Chambers et al. 2016). The near-infrared data comes from two sources: first, the WFCAM (Casali et al. 2007) on the UKIRT, primarily, but not exclusively, as part of the UKIRT Infrared Deep Sky Survey (UKIDSS; Lawrence et al. 2007). And second, data from the VIRCAM on the VISTA (Emerson et al. 2006; Dalton et al. 2006). The mid-infrared,  $\lambda = 3 - 30\mu\text{m}$  wavelength data is from the the Wide-Field Infrared Survey Explorer (WISE; Wright et al. 2010; Cutri 2013) mission. For reference, Figure 1 displays the wavelength and normalised transmission of the filters in question.

### 2.1 Spectroscopy

We compile the list of all known, spectroscopically confirmed quasars from the literature. This list was compiled from a range of surveys and papers. Specifically, we use data from: Bañados et al. (2014, 2016, 2018), Becker et al. (2015), Calura et al. (2014), Carilli et al. (2007, 2010), Carnall et al. (2015), Cool et al. (2006), De Rosa et al. (2011), Fan et al. (2000, 2001, 2003, 2004, 2006, 2018), Goto (2006), Ikeda et al. (2017), Jiang et al. (2008, 2009, 2015); ?, Kashikawa et al. (2015), Koptelova et al. (2017), Kim et al. (2015),

<sup>3</sup> lsst.org

<sup>4</sup> sci.esa.int/euclid/

<sup>5</sup> jwst.nasa.gov;

<sup>6</sup> sci.esa.int/jwst;

<sup>7</sup> www.asc-csa.gc.ca/eng/satellites/jwst;

<sup>8</sup> jwst.stsci.edu

Survey	# VHzQs	(%)	Survey reference
ATLAS	4	(0.86)	Shanks et al. (2015)
CFHQS	20	(4.32)	Willott et al. (2007)
DELS	16	(3.46)	Dey et al. (2018)
ELAIS	1	(0.22)	Väisänen et al. (2000)
FIRST	1	(0.22)	Becker et al. (1995)
HSC	8	(1.73)	Miyazaki et al. (2018)
IMS	5	(1.08)	Kim et al. (2015)
MMT	12	(2.59)	McGreer et al. (2013)
NDWFS	1	(0.22)	Jannuzi & Dey (1999)
PSO	83	(17.93)	Kaiser et al. (2002, 2010)
RD	1	(0.22)	Mahabal et al. (2005)
SDSS	170	(36.72)	Stoughton et al. (2002)
SDWISE <sup>b</sup>	27	(5.83)	Wang et al. (2016)
SHELLQs	55	(11.88)	Matsuoka et al. (2016)
SUV <sup>c</sup>	20	(4.32)	Yang et al. (2017)
UHS	1	(0.22)	Wang et al. (2017)
ULAS	10	(2.16)	Lawrence et al. (2007)
VDES <sup>d</sup>	17	(3.67)	Reed et al. (2017)
VHS	1	(0.22)	Wang et al. (2018a)
VIK	9	(1.94)	Edge et al. (2013)
VIMOS	1	(0.22)	Le Fèvre et al. (2003)


**Table 1.** The source and number of the VHzQ, with the key survey reference also given. Recent survey name and acronyms include: <sup>a</sup>DESI Legacy Imaging Survey; <sup>b</sup>SDWISE = SDSS+WISE; <sup>c</sup>SUV = SDSS-ULAS/VHS; <sup>d</sup>VDES = VHS/VIKING+DES;

2018), Kurk et al. (2007, 2009), Leipski et al. (2014), Mahabal et al. (2005), Matsuoka et al. (2016, 2018a,b), Mazzuchelli et al. (2017), Morganson et al. (2012), Mortlock et al. (2009, 2011), McGreer et al. (2006, 2013), Reed et al. (2015, 2017), Stern et al. (2007), Tang et al. (2017), Venemans et al. (2007, 2012, 2013, 2015b,a, 2016), Wang et al. (2016, 2017, 2018b,a), Willott et al. (2007, 2009, 2010, 2013, 2015), Wu et al. (2015) Yang et al. (2018b,a) and Zeimann et al. (2011).

Most of these objects are easily identified by their broad Ly $\alpha$  emission line, Nv emission and characteristic shape blueward of 1215Å in the rest-frame. As we shall see, some of the recently discovered objects are close to the galaxy luminosity function characteristic luminosity  $M^*$ , and some have relatively weak or maybe even completely absorbed Ly $\alpha$  (e.g. Figures 7 and 10 in Bañados et al. 2016). We leave aside detailed investigation and discussion into spectral features and line strengths, and take as given the published spectra and redshift identifications.

The breakdown of how many VHzQ each survey reports is given in Table 1. The Sloan Digital Sky Survey (SDSS) and the Pan-STARRS1 (PS1; PSO in Table 1) survey and alone identified over half (54.6%) of the VHzQ population. Data from the Hyper Suprime-Cam (HSC) on the Subaru telescope is responsible for 13.6% of our dataset (HSC+SHELLQs in Table 1). The combination of surveys is also vital for identifying VHzQs. The UKIDSS Large Area Survey (ULAS) on its own, or in combination with other surveys is responsible for 6.5% of the sample (SUV+ULAS) including the highest- $z$  object. Where more than one survey is used for the high-redshift identification (e.g. via shorter-band veto and longer wavelength detection) we follow the discovery paper naming convention.

The redshifts for the VHzQs generally come from the measurement of broad UV/optical emission lines. There are



**Figure 2.** The redshift distribution  $N(z)$  of the VHzQ sample. The bins are  $\delta - z = 0.075$  in width.

$z \geq$	Age / Myr	No. of objects
7.50	700	1
7.00	767	4
6.78	800	14
6.50	845	40
6.19	900	86
6.00	937	170
5.70	1000	267
5.00	1180	463

**Table 2.** The number of objects at or above a given redshift. The age of the Universe in megayears is also given.

far infra-red emission lines e.g. CII 158 micron available for several objects, but at the level of our current analysis broadband redshifts are sufficient.

The number of objects at or above various redshifts, along with the corresponding age of the Universe is given in Table 2.

The  $N(z)$  redshift histogram is given for the sample in Figure 2. We split the contribution up by survey. For clarity we show the individual surveys of SDSS, PS1, HSC, the ULAS detection, and tally the remaining surveys together (“various”). In Section ?? we discuss the form of the  $N(z)$  as well as the coverage of the luminosity-redshift  $L - z$  plane.

## 2.2 Optical Photometry

We query the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS)<sup>9</sup> Data Release 1 (DR1) Catalog Archive Server Jobs System (CasJobs) service at [mas-tweb.stsci.edu/ps1casjobs/](https://web.stsci.edu/ps1casjobs/). The Pan-STARRS1 (PS1) survey observed the 30,000 deg<sup>2</sup> of sky north of declination  $\delta = -30$  degrees in the five *grizy* filters. PS1 is the first part of Pan-STARRS to be completed and is the basis for the DR1. Chambers et al. (2016), Magnier et al. (2016c), Waters et al. (2016), Magnier et al. (2016a), Magnier et al. (2016b) and Flewelling et al. (2016) describe the instrument, survey, and data analyses. The principal science product of the PS1 survey is the catalog accessible through the CasJobs interface.

The PS1 Casjobs SQL Server is located at [mast-](https://mast-)

<sup>9</sup> <https://outerspace.stsci.edu/display/PANSTARRS>

[web.stsci.edu/ps1casjobs](http://web.stsci.edu/ps1casjobs). The top level documentation is given [here](#) while the description of tables is given [here](#).

The main tables are the `objectThin` and `meanObject` tables. We query and return the mean PSF magnitudes from the *grizy* filters (`MeanPSFMag`) which are in the AB system for our 463 VHzQ sample. Our SQL and links to the main PS1 DR1 tables are given at Appendix ??.

### 2.3 Near-infrared photometry

The near-infrared data in this paper comes from the Wide Field Astronomy Unit’s (WFAU) Science Archives for UKIRT-WFCAM, the WFCAM Science Archive (WSA; [Hambly et al. 2008](#)) and VISTA-VIRCAM, the VISTA Science Archive (VSA; [Cross et al. 2012](#)). These archives were developed for the VISTA Data Flow System (VDFS [Emerson et al. 2004](#)).

We access both the WSA and the VSA and include all non-proprietary WFCAM data, which covers all public surveys and PI projects from Semester 05A to 2017-Jan-01, and all non-proprietary VISTA data, which covers all public surveys and PI projects from science verification on 2009-Oct-15 to 2016-Apr-01.

Here we are not just quering the WSA or VSA data tables. We are taking a list of objects (positions) are performing matched aperture (“forced”) photometry on the NIR imaging data. As such, we generate a set of tables that are different in subtle ways to the regular “Detection” tables. The two most important tables for our needs are the `[w/v]serv1000MapRemeasurement` and `[w/v]serv1000MapRemeasAver`.

We produce and provide a two new databases with all the necessary quantities and measurements to fully reproduce our tables, figures and results herein. Moreover, these databases report considerably more information than we report here. Full documentation can be found at the [WSA Schema Browser](#) and the [VSA Schema Browser](#).

#### 2.3.1 Averaging matched photometry

The data was processed using a matched-aperture photometry method where flux is measured at the spectroscopic position of the quasar, without necessarily knowing if there is a formal detection in the NIR photometry beforehand. The matched-aperture pipeline is discussed in [Cross et al. \(2013\)](#) and with fuller details to appear in a forthcoming paper (Cross et al., 2019, in prep).

We query the WSA and VSA performing matched-aperture photometry at the positions of our 463 VHzQs. This database is world-readable and we give the full recipe and relevant SQL queries for accessing both databases in Appendix B as well as online.

The photometry in a single epoch image often has low signal-to-noise. The advantage of matched aperture photometry on quasars is that co-adding is relatively simple if each epoch is taken in the same aperture and the aperture photometry has been corrected to total. Indeed, the standard aperture corrections work well for point sources. Coadding using the matched-aperture photometry, where the individual epochs are taken from multiple projects with different pointings and orientations, should help with issues such as scattered light, pixel distortion and aperture corrections.

We average the aperture corrected calibrated fluxes (e.g. `aperJky3`), and then convert to magnitudes. Since we do not have a deep image for each set of averages, we cannot calculate non-aperture corrected values, so the photometry is only appropriate for point-sources.

$$\bar{F} = \frac{\sum_i^N (w_i F_i)}{\sum_i^N w_i} \quad (1)$$

where  $F_i$  is the  $i^{th}$  epoch measurement of a parameter to be averaged such as the aperture corrected calibrated flux in a  $1''$  aperture (`aperJky3`) and  $\bar{F}$  is the weighted mean average of this parameter. The weight for each epoch  $w_i = 1/(\sigma_F)^2$  if the epoch is included and  $w_i = 0$  if an epoch is excluded for quality control purposes.

We calculate a set of averaged catalogues, for each pointing and filter, based on the requirements in `RequiredMapAverages`, in these cases over time spans of 7, 14, 30, 91, 183 days, 365 days, 730 days, over 10 epochs and over all epochs. The averaging process starts at the first epoch and works onwards from there. Again, we present these measurements in the new SQL tables.

We detect 359 unique quasars in the WFCAM WSA database, 220 quasars are detected in the VISTA VSA database with 130 objects in common with both WFCAM and VISTA data. We give the necessary SQL queries syntax in Appendix B.

### 2.4 MIR data

The MIR data for this study comes from the Wide-field Infrared Survey Explorer (WISE) mission, and we utilize data from the WISE cryogenic and NEOWISE (Mainzer et al. 2011 ApJ, 731, 53) post-cryogenic survey phases.

We use data from the the beginning of the WISE mission (2010 January; [Wright et al. 2010](#)) through the fourth-year of NEOWISE-R operations (Mainzer et al. 2011, 2017 December;). More specifically, we utilise the recently released “unWISE Catalog”<sup>10</sup>. The unWISE effort<sup>10</sup> is the unblurred coadds of the WISE imaging using the AllWISE and NEOWISE-R stacked data (Lang 2014; Meisner et al. 2018a,b).

All fluxes in the unWISE catalog are reported there are in “Vega nanoMaggies”, with the Vega magnitude of a source is given by  $m_{\text{Vega}} = 22.52.5 \log(f)$ , where  $f$  is the source flux. The absolute calibration for unWISE is ultimately inherited from AllWISE through the calibration of ?. This inheritance depends on details of the PSF normalization at large radii, which is uncertain. Subtracting 4 millimag from the unWISE W1, and 32 millimag from unWISE W2 fluxes improves the agreement between unWISE and AllWISE fluxes.

Thus to convert unWISE Vega magnitudes onto the AB system, we have:

$$\begin{aligned} W1_{\text{AB,unWISE}} &= 22.5 - 2.5 \log(f_{W1}) - 0.004 + 2.699 \\ W2_{\text{AB,unWISE}} &= 22.5 - 2.5 \log(f_{W2}) - 0.032 + 3.339. \end{aligned}$$

For the our MIR variability investigations, we do not

<sup>10</sup> <http://unwise.me/>

use the unWISE coadds, but instead use the [AllWISE](#) catalogue and the [NEOWISE 2018 Data Release](#). NEOWISE 2018 makes available the 3.4 and 4.6  $\mu$ m (W1 and W2) single-exposure images and extracted source information that was acquired between 2016 December 13 and 2017 December 13 UTC, which was the fourth year of survey operations of the Near-Earth Object Wide-field Infrared Survey Explorer Re-activation Mission (NEOWISE; Mainzer et al. 2014, ApJ, 792, 30). The fourth year NEOWISE data products are concatenated with those from the first three years into a single archive.

The WISE scan pattern leads to coverage of the full-sky approximately once every six months (a “sky pass”), but the satellite was placed in hibernation in 2011 February and then reactivated in 2013 October. Hence, our light curves have a cadence of 6 months with a 32 month sampling gap.

Tables ??, ?? and ??, represent the culmination of this effort, and we now describe the assembly of their contents in more detail.

survey	qsoName	ra	dec	redshift	Z	Y	J	H	K	S	W1
PSO	J000.3401+26.8358	0.34011	26.83588	5.75	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$19.28 \pm 0.062$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$16.280 \pm 0.02$
SDSS	J0002+2550	0.66412	25.84304	5.82	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$19.37 \pm 0.069$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$16.250 \pm 0.02$
SDWISE	J0008+3616	2.21429	36.27041	5.17	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$19.33 \pm 0.063$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$16.018 \pm 0.02$
PSO	J002.3786+32.8702	2.37870	32.87026	6.10	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$20.99 \pm 0.249$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$17.951 \pm 0.10$
SDSS	J0012+3632	3.13700	36.53781	5.44	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$19.01 \pm 0.049$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$15.821 \pm 0.03$
PSO	J004.3936+17.0862	4.39361	17.08630	5.80	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$20.56 \pm 0.202$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$17.834 \pm 0.10$
PSO	J006.1240+39.2219	6.12404	39.22193	6.62	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$21.28 \pm 0.422$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$17.364 \pm 0.06$
SDWISE	J0025-0145	6.36183	-1.75903	5.07	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$17.74 \pm 0.004$	$-1000.00 \pm -1000.000$	$-1000.00 \pm -1000.000$	$14.851 \pm 0.00$
PSO	J007.0273+04.9571	7.02733	4.95712	6.00	$-1000.00 \pm -1000.000$	$20.33 \pm 0.056$	$20.23 \pm 0.074$	$20.29 \pm 0.108$	$20.19 \pm 0.105$	$-1000.00 \pm -1000.000$	$17.178 \pm 0.06$
SDWISE	J0031+0710	7.85775	7.17692	5.33	$-1000.00 \pm -1000.000$	$20.03 \pm 0.082$	$20.20 \pm 0.146$	$19.49 \pm 0.106$	$19.61 \pm 0.123$	$-1000.00 \pm -1000.000$	$16.658 \pm 0.03$

**Table 3.** The caption without a number



Selection	number detected (%)
Any band ( $ZYJHK/K_s$ )	449 (97.0)
Z-band	75 (16.1)
Y-band	273 (59.0)
J-band	447 (96.5)
H-band	269 (58.1)
K or $K_s$ -band	322 (69.5)

**Table 4.** Detection rate of VH $z$ Qs in the near-infrared.

### 3 RESULTS

Having collated the sample of 463 VH $z$ Qs, and obtained their optical, near- and mid-infrared photometry we report here the various photometric properties of the quasars.

First, we will concentrate on detection rate in the infrared, go on to report on the color-redshift and color-color properties of our sample and then report on how the current sample populates the luminosity-redshift  $Lz$ -plane.

#### 3.1 Detection Rates in the NIR

Table 4 gives the detection rates for the VH $z$ Qs in the NIR  $YJHK/K_s$ -bands. The first thing to note is that the coverage of the NIR surveys for example from the UKIDSS LAS and VISTA VHS, does not overlap the full area for where the VH $z$ Qs are detected.

There are 14 objects that have no NIR detections. 3 of these (PSOJ053.9605-15.7956, PSO J056.7168-16.4769 and DELSJ0411-0907) have been observed (by VHS) but are out of our queried time range. 6 objects have not been observed (or at least the data is not in the WSA/VSA archives yet) and 5 objects at  $\geq +60$  deg<sup>2</sup> are too far north for UKIRT and cannot be observed.

##### 3.1.1 Comparing WFCAM and VISTA

There are 130 overlapping QSOs between WFCAM and VISTA. Using the *VegaToAB* value<sup>11</sup> to put these objects on the same AB system, and for each object compared the two measurements. First, the calculated weighted average (calibrated flux) in each filter of both and calculated the ratio and difference between each measurement and the average. Then for each filter we calculated the weighted average of the differences (in mag) for each instrument to see if there were significant offsets. The results are given in Table 5. The only filter with a significant offset is the Y-band. All of the VISTA averages are negative and all of the WFCAM ones are positive. The  $K_s$  versus  $K$  band may be slightly dodgy, given the different shapes of the filters.


We investigate the VH $z$ Qs in the UltraVISTA field. UltraVISTA is... One detail to note is that depths quoted in the UltraVISTA Data Releases are for more realistic 1.8 arc-sec diameter apertures (J. Dunlop, priv. comm.).

#### 3.2 Detection Rates in the MIR

Unlike the NIR coverage, the WISE satellite and mission performed an all-sky survey, so the location of every VH $z$ Q

abs(VIRCAM - WFCAM)	millimag	no. of objects
Z	23.2	3
Y	57.3	53
J	2.1	106
H	45.8	96
$K_s/K$	25.2	110

**Table 5.** Comparing the magnitudes in different WFCAM/UKIRT and VIRCAM/VISTA near-infrared bands.



**Figure 3.** WISE W1/2/3/4 magnitude against signal-to-noise, colour coded by *wxcov* the mean coverage depth, in each corresponding band.

in our dataset is covered. However, the depth of the WISE ALLWISE survey depends heavily on sky location, with locations near the Ecliptic Poles having the highest number of exposures.

Before reporting on the detection rates, we investigate this effect. Figure 4 shows the WISE magnitude versus signal-to-noise, colour coded by *wxcov* the mean coverage depth, in each corresponding band. In the two shorter bands W1/2 we see the clear and expected trend for brighter objects to have larger SNR, and also for the higher signal to noise for objects with more exposures at a given magnitude. The behaviour for the W3/4 bands is different, with two populations clearly evident in W3 and although a bit more mixed, also in W4. With the suggested split at  $\text{SNR} > 2$ , and no obvious R.A./Declination dependence seen, this behaviour is explained by the fact that there are non-detections in W3/4 for objects (with high W1/2 SNR) that are reported in the ALLWISE catalogue.

For the 278 VH $z$ Q with coverage detections, the mean number of exposures for the W1/2 bands is 32.0 and 31.5, respectively, with a minimum number of exposures 17 and 12, and the maximum number of exposures being 114 (for both bands). For the W3/4 filters, the corresponding mean, minimum and maximum exposure are 17.4 and 17.5, 5.8 and 6.8 and 69 (for both bands). These values are directly from the *wxcov* entries in the [WISE ALLWISE catalogue](#).

<sup>11</sup> What is this exactly??

Selection	number detected (%)
W1 SNR > 2.0	275 (64.9)
W2 SNR > 2.0	255 (60.1)
W1 $\wedge$ W2 SNR > 2.0	
W3 SNR > 2.0	99 (23.3)
W4 SNR > 2.0	29 (6.8)
Any W1/2/3/4 SNR > 2.0	
W1/2 SNR < 2.0 $\wedge$ W3 SNR > 2.0	

**Table 6.**

Selection	number detected (% of full spectra)
From “Source”, “Rejects”,	245, 40 (67.2)
W1 SNR > 2.0	279 (65.8)
W2 SNR > 2.0	258 (60.8)
W1 $\wedge$ W2 SNR > 2.0	253 (59.7)
W3 SNR > 2.0	97 (22.9)
W4 SNR > 2.0	33 (7.8)
W1/2 SNR < 2.0 $\wedge$ W3 SNR > 2.0	3 (0.7)



**Table 7.** Data from the AllWISE Source Catalog and AllWISE Reject Table, from the [NASA/IPAC Infrared Science Archive](#)**Figure 4.** WISE signal-to-noise measures for the four bands, as well as for (W1-W2) colour. The points are colour coded by redshift.

Table 7 gives the detection rates for the VH $z$ Qs in the MIR WISE W1-4 bands.

[Blain et al. \(2013\)](#)

Recently, [Assef et al. \(2018\)](#) released two large catalogues of AGN candidates identified across 30,000 deg<sup>2</sup> of extragalactic sky from the WISE AllWISE Data Release. The “R90” catalogue, is contains 4.5M AGN candidates at 90% reliability (and  $\approx 150$  AGN candidates per deg<sup>2</sup>) while the “C75” catalog consists of 20.9M AGN candidates at 75% completeness (and  $\approx 700$  AGN candidates per deg<sup>2</sup>). Cross-matching out catalogue of 463 VH $z$ Qs with these catalogues, produces 42 matches with the R90 sample and 98 matches with the C75 sample. Both catalogues unsurprisingly match to the ultraluminous quasar SDSS J0100+2802 ([Wu et al. 2015](#)) while the C75, but not the R90 catalogue matches to ULAS J1120+0641 ([Mortlock et al. 2011](#)). Neither catalogue matches J1342+0928 ([Bañados et al. 2018](#)).

Very High- $z$  Quasars Detected in WISE W3 and W4.

**Figure 5.** Histogram showing the number of NEOWISE-R epochs and detections there are for each VH $z$ Q.

### 3.3 Variability

VH $z$ Qs, if accreting at, or above the Eddington Limit, might well have large values of changing mass accretion rate,  $\dot{m}_{\text{accr}}$ . A consequence of this would be that these quasar exhibit signs of variability, most likely showing up in their UV/optical rest-frame spectra. We look for evidence of this variability signature in the NIR and MIR light-curves of the VH $z$ Qs. As a guide, C IV enters the  $Y$ -band at redshift  $z=5.32$  and exits at  $z=5.99$ , and enters the  $J$ -band at redshift  $z=6.55$  and exits at  $z=7.57$ . Mg II enters the  $H$ -band at redshift  $z=4.33$  and exits at  $z=5.37$  and enters the  $K$ -band at redshift  $z=6.25$  and exits at 7.50.

Using the extended datasets described in Section 2.3 and ??, we

Figure 5 gives the number of NEOWISE-R epochs and detections there are for each VH $z$ Q, while Figure ?? presents three examples of the MIR lightcurves and associated colour changes. Here we show J0100+2802 ([Wu et al. 2015](#)), J0224-4711 and J1626+2751. **NJC: What about NIR light-curves / combined light-curves**

The two with large differences are:

SDSSJ0349+0034, in K/Ks band: VSA =  $18.36 \pm 0.10$ , WSA =  $19.13 \pm 0.24$  and


SDSSJ2220-0101, in J band: VSA =  $19.38 \pm 0.04$ , WSA =  $22.23 \pm 0.15$

## 4 SURFACE DENSITY OF VH $z$ Qs

## 5 DISCUSSION AND CONCLUSIONS

In this study, we have, for the first time, ompiled the list of all  $z > 5$  spectroscopically confirmed quasars. We have assemble the NIR ( $y/Y, J, H, K/K_s$ ) and MIR (WISE W1/2/3/4) photometry for these objects, given their detection rates and SEDs. We find that:





**Figure 6.** Here we show the MIR NEOWISE-R for J0100+2802 (Wu et al. 2015), J0224-4711 and J1626+2751. Red points are the W1 band; cyan points the W2 band.

We can gain a good appreciation for what these missions will discover by collating the datasets we currently have.

- Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam porta sodales est, vel cursus risus porta non. Vivamus vel pretium velit. Sed fringilla suscipit felis, nec iaculis lacus convallis ac.
- Fusce pellentesque condimentum dolor, quis vehicula tortor hendrerit sed. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos himenaeos. Etiam interdum tristique diam eu blandit. Donec in lacinia libero.
- Sed elit massa, eleifend non sodales a, commodo ut felis. Sed id pretium felis. Vestibulum et turpis vitae quam aliquam convallis. Sed id ligula eu nulla ultrices tempus. Phasellus mattis erat quis metus dignissim malesuada. Nulla tincidunt quam volutpat nibh facilisis euismod. Cras vel auctor neque. Nam quis diam risus.

Nunc lacus nibh, convallis ac lobortis ut, tempus ac lectus. Maecenas eu elit massa. Nulla vel lacus lorem. Proin et lobortis tortor. Phasellus ultrices nisl non enim porttitor dictum. Curabitur nec nunc ac nibh ornare elementum. Nunc ultrices hendrerit ultricies. Aliquam dapibus semper est et gravida. Etiam cursus, massa eget tempor elementum, lectus urna feugiat nisi, eget sagittis.

### Author Contributions

N.P.R. initiated the project, compiled the list of  $z > 5.00$  quasars, wrote most of the analysis code, developed the the plotting scripts, and developed and wrote the initial and

subsequent drafts of the manuscript. N.J.G.C. supplied the critical near-infrared expertise and database for which the bulk of the project relies. N.J.G.C. also contributed directly to the writing of the manuscript.

### Availability of Data and computer analysis codes

All materials, databases, data tables and code are fully available at: <https://github.com/d80b2t/VHzQ>

### ACKNOWLEDGEMENTS

NPR acknowledges support from the STFC and the Ernest Rutherford Fellowship scheme.

We thank Mike Read at the ROE WFAU for help with the WFCAM Science Archiv (WSA), and also the VISTA Science Archive (VSA). We thank Bernie Shiao at STScI for help with the Pan-STARRS1 DR1 CasJobs interface.

This paper heavily used TOPCAT (v4.4) (Taylor 2005, 2011). This research made use of Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013; The Astropy Collaboration et al. 2018).

The Pan-STARRS1 Surveys (PS1) and the PS1 public science archive have been made possible through contributions by the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max-Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, the Queen's University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under Grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation Grant No. AST-1238877, the University of Maryland, Eotvos Lorand University (ELTE), the Los Alamos National Laboratory, and the Gordon and Betty Moore Foundation.

This project used data obtained with the Dark Energy Camera (DECam) and the NOAO Data Lab, The Data Lab is operated by the National Optical Astronomy Observatory, the national center for ground-based nighttime astronomy in the United States operated by the Association of Universities for Research in Astronomy (AURA) under cooperative agreement with the National Science Foundation.

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, and NEOWISE, which is a project of the Jet Propulsion Laboratory/California Institute of Technology. WISE and NEOWISE are funded by the National Aeronautics and Space Administration.

CasJobs was originally developed by the Johns Hopkins University/ Sloan Digital Sky Survey (JHU/SDSS) team. With their permission, MAST used version 3.5.16 to con-

struct CasJobs-based tools for GALEX, Kepler, the Hubble Source Catalog, and PanSTARRS.

This research has made use of the SVO Filter Profile Service (<http://svo2.cab.inta-csic.es/theory/fps/>) supported from the Spanish MINECO through grant AyA2014-55216. The SVO Filter Profile Service<sup>12</sup> describes the Spanish VO Filter Profile Service. The Filter Profile Service Access Protocol. Rodrigo, C., Solano, E. <http://ivoa.net/documents/Notes/SVOFPSDAL/index.html>

## APPENDIX A: NEAR-INFRARED WFCAM SCIENCE ARCHIVE SQL QUERIES

Here we give the recipe and SQL that returned the near-infrared photometry for the VHzQs from the WFCAM Science Archive.

The data are on the WFCAM Science Archive: [wsa.roe.ac.uk](http://wsa.roe.ac.uk). Access the User Login form [wsa.roe.ac.uk/login.html](http://wsa.roe.ac.uk/login.html) with these credentials::

- Username: **WSERV1000**
- password: **highzqso**
- community: **nonsurvey**

Then going to the [Free Form SQL Query](#) page the Database release **WSERV1000v20180716** can be accessed which contains all the data we use here.

We *nota bene* a few things. First, the quantity **aperJky3** and **aperJky3Err** are found in the **wserv1000MapRemeasAver** and **wserv1000MapRemeasurement**, so care has to be taken to return unique column names (otherwise e.g. [astropy.io.fits](#) will crash). As such, we alias **aver.aperJky3** to **aperJky3Aver** and likewise for the error quantity. Aliases will be necessary in some cases anyway, because some queries can be done sensibly on multiple instances of the same table. Other times, one may join tables on quantities such as **catalogueID** or **apertureID**, where you are meaning the same thing, but aliases would again be sensible.

Second, the **RA** and **DEC** values returned by the WSA are in radians, if used directly. To return values in degrees, use a selection with an alias, e.g. **RA as RADeg** and **DEC as DECDeg**.

<sup>12</sup> Rodrigo, C., Solano, E., Bayo, A. <http://ivoa.net/documents/Notes/SVOFPS/index.html>

Then the following SQL will return the values in Table ??.

```

1  SELECT
2  qso.qsoName, qso.ra as raJ2000, qso.dec as decJ2000,
3  aver.apertureID, aver.aperJky3 as aperJky3Aver,
4  aver.aperJky3Err as aperJky3AverErr, aver.sumWeight,
5  aver.ppErrBits as ppErrBitsAver, m.mjdObs,
6  m.filterID, remeas.aperJky3,
7  remeas.aperJky3Err,
8  w.weight, remeas.ppErrBits,
9  m.project
10
11 FROM
12 finalQsoCatalogue as qso,
13 MapApertureIDshighzQsoMap as ma,
14 wserv1000MapRemeasAver as aver,
15 wserv1000MapRemeasurement as remeas,
16 MapProvenance as v,
17 wserv1000MapAverageWeights as w,
18 MapFrameStatus as mfs,
19 Multiframe as m
20
21 WHERE
22 qso.qsoID=ma.objectID and
23 ma.apertureID=aver.apertureID and
24 aver.apertureID=remeas.apertureID and
25 aver.catalogueID=v.combicatID and
26 v.avSetupID=1 and
27 v.catalogueID=remeas.catalogueID and
28 w.combicatID=v.combicatID and
29 w.catalogueID=v.catalogueID and
30 w.apertureID=aver.apertureID and
31 mfs.catalogueID=remeas.catalogueID and
32 m.multiframeID=mfs.multiframeID and
33 mfs.programmeID=10999 and
34 mfs.mapID=1
35 order by v.combicatID, m.mjdObs

```

## APPENDIX B: NEAR-INFRARED VISTA SCIENCE ARCHIVE SQL QUERIES

In a very similar manner to the WSA, we give here the details on how to access the VISTA Science Archive (VSA)

At the [VSA Login](#), enter with these credentials::

- Username: **VSERV1000**
- password: **highzqso**
- community: **proprietary**

Then head to the [Freeform SQL Query](#) page where the database release to use is **VSERV1000v20180716**.

## REFERENCES

- Agarwal B., Smith B., Glover S., Natarajan P., Khochfar S., 2016, *MNRAS*, 459, 4209
- Alexander T., Natarajan P., 2014, *Science*, 345, 1330
- Assef R. J., Stern D., Noirot G., Jun H. D., Cutri R. M., Eisenhardt P. R. M., 2018, *ApJS*, 234, 23
- Astropy Collaboration et al., 2013, *Astron. & Astrophys.*, 558, A33
- Bañados E., et al., 2014, *AJ*, 148, 14
- Bañados E., et al., 2016, *ApJS*, 227, 11
- Bañados E., et al., 2018, *Nat*, 553, 473
- Becker G. D., Bolton J. S., Lidz A., 2015, *PASA*, 32, 45
- Becker R. H., White R. L., Helfand D. J., 1995, *ApJ*, 450, 559
- Blain A., et al., 2013, *ArXiv e-prints*
- Bosman S. E. I., et al., 2017, *MNRAS*, 470, 1919
- Calura F., Gilli R., Vignali C., Pozzi F., Pipino A., Matteucci F., 2014, *MNRAS*, 438, 2765
- Carilli C. L., et al., 2007, *ApJ Lett.*, 666, L9
- Carilli C. L., et al., 2010, *ApJ*, 714, 834
- Carnall A. C., Shanks T., Chehade B., Fumagalli M., Rauch M., Irwin M. J., Gonzalez-Solares E., Findlay J. R., Metcalfe N., 2015, *MNRAS*, 451, L16
- Casali M., et al., 2007, *Astron. & Astrophys.*, 467, 777
- Chambers K. C., et al., 2016, *arXiv:1612.05560v3*
- Chen S.-F. S., et al., 2017, *ApJ*, 850, 188
- Cool R. J., et al., 2006, *AJ*, 132, 823
- Cross N., Hambly N., Collins R., Sutorius E., Read M., Blake R., 2013, in Adamson A., Davies J., Robson I., eds, *Thirty Years of Astronomical Discovery with UKIRT Vol. 37 of Astrophysics and Space Science Proceedings, Discovery of Variables in WFCAM and VISTA Data*. p. 193
- Cross N. J. G., et al., 2012, *Astron. & Astrophys.*, 548, A119
- Cushing M. C., et al., 2006, *ApJ*, 648, 614
- Cutri R. M. o., 2013, *Technical report, Explanatory Supplement to the AllWISE Data Release Products*
- Dalton G. B., et al., 2006, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 6269 of Proc. SPIE, The VISTA infrared camera*. p. 62690X
- De Rosa G., Decarli R., Walter F., Fan X., Jiang L., Kurk J., Pasquali A., Rix H. W., 2011, *ApJ*, 739, 56
- Dey A., et al., 2018, *ArXiv e-prints*
- Edge A., Sutherland W., Kuijken K., Driver S., McMahon R., Eales S., Emerson J. P., 2013, *The Messenger*, 154, 32
- Emerson J., McPherson A., Sutherland W., 2006, *The Messenger*, 126, 41
- Emerson J. P., Irwin M. J., Lewis J., Hodgkin S., Evans D., Bunclark P., McMahon R., Hambly N. C., Mann R. G., Bond I., Sutorius E., Read M., Williams P., Lawrence A., Stewart M., 2004, in P. J. Quinn & A. Bridger ed., *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 5493 of Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, VISTA data flow system: overview*. pp 401–410
- Fan X., Carilli C. L., Keating B., 2006, *ARA&A*, 44, 415
- Fan X., et al., 2000, *AJ*, 119, 1
- Fan X., et al., 2004, *AJ*, 128, 515
- Fan X., et al., 2006, *AJ*, 132, 117
- Fan X., et al., 2018, *ArXiv e-prints*
- Fan X., Narayanan V. K., Lupton R. H., Strauss M. A., Knapp G. R., Becker R. H., White R. L., Pentericci L., et al., 2001, *AJ*, 122, 2833
- Fan X., Strauss M. A., Schneider D. P., Becker R. H., White R. L., Haiman Z., Gregg M., Pentericci L., et al., 2003, *AJ*, 125, 1649
- Flewelling H. A., et al., 2016, *arXiv:1612.05243v2*
- Fukugita M., Ichikawa T., Gunn J. E., Doi M., Shimasaku K., Schneider D. P., 1996, *AJ*, 111, 1748
- Goto T., 2006, *MNRAS*, 371, 769
- Haardt F., Gorini V., Moschella U., Treves A., Colpi M., eds, 2016, *Astrophysical Black Holes Vol. 905 of Lecture Notes in Physics*, Berlin Springer Verlag
- Hambly N. C., et al., 2008, *MNRAS*, 384, 637
- Hickox R. C., Myers A. D., Greene J. E., Hainline K. N., Zakamska N. L., DiPompeo M. A., 2017, *ApJ*, 849, 53
- Hill A. R., Gallagher S. C., Deo R. P., Peeters E., Richards G. T., 2014, *MNRAS*, 438, 2317
- Ikedda H., Nagao T., Matsuoka K., Kawakatu N., Kajisawa M., Akiyama M., Miyaji T., Morokuma T., 2017, *ApJ*, 846, 57
- Jannuzi B. T., Dey A., 1999, in Weymann R., et al. eds, *ASP Conf. Ser. 191: Photometric Redshifts and the Detection of High Redshift Galaxies* p. 111
- Jiang L., et al., 2006, *AJ*, 132, 2127
- Jiang L., et al., 2008, *AJ*, 135, 1057
- Jiang L., et al., 2009, *AJ*, 138, 305
- Jiang L., et al., 2010, *Nat*, 464, 380
- Jiang L., McGreer I. D., Fan X., Bian F., Cai Z., Clément B., Wang R., Fan Z., 2015, *AJ*, 149, 188
- Kaiser N., et al., 2002, in J. A. Tyson & S. Wolff ed., *Society of Photo-Optical Instrumentation Engineers (SPIE) Vol. 4836, Pan-STARRS: A Large Synoptic Survey Telescope Array*. pp 154–164
- Kaiser N., et al., 2010, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Vol. 7733, The Pan-STARRS wide-field optical/NIR imaging survey*. p. 0
- Kashikawa N., Ishizaki Y., Willott C. J., Onoue M., Im M., Furusawa H., Toshikawa J., Ishikawa S., Niino Y., Shimasaku K., Ouchi M., Hibon P., 2015, *ApJ*, 798, 28
- Kim Y., et al., 2015, *ApJ Lett.*, 813, L35
- Kim Y., et al., 2018, *ArXiv e-prints*
- Koptelova E., Hwang C.-Y., Yu P.-C., Chen W.-P., Guo J.-K., 2017, *Scientific Reports*, 7, 41617
- Kurk J. D., et al., 2007, *ApJ*, 669, 32
- Kurk J. D., Walter F., Fan X., Jiang L., Jester S., Rix H.-W., Riechers D. A., 2009, *ApJ*, 702, 833
- Lacy M., et al., 2004, *ApJS*, 154, 166
- Lang D., 2014, *AJ*, 147, 108
- Latif M. A., Volonteri M., Wise J. H., 2018, *arXiv:1801.07685v1*
- Lawrence A., et al., 2007, *MNRAS*, 379, 1599
- Le Fèvre O., et al., 2003, in Iye M., Moorwood A. F. M., eds, *Instrument Design and Performance for Optical/Infrared Ground-based Telescopes Vol. 4841 of Proc. SPIE, Commissioning and performances of the VLT-VIMOS instrument*. pp 1670–1681
- Leipski C., et al., 2014, *ApJ*, 785, 154
- Lupi A., Haardt F., Dotti M., Fiacconi D., Mayer L., Madau P., 2016, *MNRAS*, 456, 2993
- Madau P., Haardt F., Dotti M., 2014, *ApJ Lett.*, 784, L38
- Magnier E. A., et al., 2016a, *arXiv:1612.05242v2*
- Magnier E. A., et al., 2016b, *arXiv:1612.05244v2*

- Magnier E. A., et al., 2016c, arXiv:1612.05240v2
- Mahabal A., Stern D., Bogosavljević M., Djorgovski S. G., Thompson D., 2005, *ApJ Lett.*, 634, L9
- Mainzer A., et al., 2011, *ApJ*, 731, 53
- Mainzer A., et al., 2014, *ApJ*, 792, 30
- Matsuoka Y., et al., 2016, *ApJ*, 828, 26
- Matsuoka Y., et al., 2018a, *PASJ*, 70, S35
- Matsuoka Y., et al., 2018b, *ApJS*, 237, 5
- Mazzucchelli C., et al., 2017, *ApJ*, 849, 91
- McGreer I. D., Becker R. H., Helfand D. J., White R. L., 2006, *ApJ*, 652, 157
- McGreer I. D., et al., 2013, *ApJ*, 768, 105
- Meisner A. M., Lang D., Schlegel D. J., 2018a, *Research Notes of the American Astronomical Society*, 2, 1
- Meisner A. M., Lang D. A., Schlegel D. J., 2018b, *Research Notes of the American Astronomical Society*, 2, 202
- Miyazaki S., et al., 2018, *PASJ*, 70, S1
- Morganson E., et al., 2012, *AJ*, 143, 142
- Mortlock D., 2016, in Mesinger A., ed., *Understanding the Epoch of Cosmic Reionization: Challenges and Progress* Vol. 423 of *Astrophysics and Space Science Library*, Quasars as Probes of Cosmological Reionization. p. 187
- Mortlock D. J., et al., 2009, *Astron. & Astrophys.*, 505, 97
- Mortlock D. J., et al., 2011, *Nat*, 474, 616
- Oke J. B., Gunn J. E., 1983, *ApJ*, 266, 713
- Pezzulli E., Valiante R., Schneider R., 2016, *MNRAS*, 458, 3047
- Pezzulli E., Volonteri M., Schneider R., Valiante R., 2017, *MNRAS*, 471, 589
- Reed S. L., et al., 2015, *MNRAS*, 454, 3952
- Reed S. L., et al., 2017, *MNRAS*, 468, 4702
- Rees M. J., 1984, *ARA&A*, 22, 471
- Richards G. T., et al., 2006, *ApJS*, 166, 470
- Sawicki M., 2002, *AJ*, 124, 3050
- Schlaflly E. F., Meisner A. M., 2018, *ApJS*
- Shanks T., et al., 2015, *MNRAS*, 451, 4238
- Simcoe R. A., Sullivan P. W., Cooksey K. L., Kao M. M., Matejek M. S., Burgasser A. J., 2012, *Nat*, 492, 79
- Stern D., et al., 2005, *ApJ*, 631, 163
- Stern D., et al., 2007, *ApJ*, 663, 677
- Stoughton C., et al., 2002, *AJ*, 123, 485
- Takeo E., Inayoshi K., Ohsuga K., Takahashi H. R., Mineshige S., 2018, *MNRAS*, 476, 673
- Tang J.-J., et al., 2017, *MNRAS*, 466, 4568
- Taylor M., , 2011, *TOPCAT: Tool for OPERations on Catalogues And Tables*, *Astrophysics Source Code Library*
- Taylor M. B., 2005, in Shopbell P., Britton M., Ebert R., eds, *Astronomical Data Analysis Software and Systems XIV* Vol. 347 of *Astronomical Society of the Pacific Conference Series*, *TOPCAT & STIL: Starlink Table/VOTable Processing Software*. p. 29
- The Astropy Collaboration et al., 2018, *ArXiv e-prints*
- Tielens A. G. G. M., 2008, *ARA&A*, 46, 289
- Timlin J. D., Ross N. P., et al., 2016, *ApJS*, 225, 1
- Väisänen P., Tollestrup E. V., Willner S. P., Cohen M., 2000, *ApJ*, 540, 593
- Valiante R., Schneider R., Graziani L., Zappacosta L., 2018, *MNRAS*, 474, 3825
- Vanden Berk D. E., et al., 2001, *AJ*, 122, 549
- Venemans B. P., et al., 2012, *ApJ Lett.*, 751, L25
- Venemans B. P., et al., 2013, *ApJ*, 779, 24
- Venemans B. P., et al., 2015a, *MNRAS*, 453, 2259
- Venemans B. P., et al., 2015b, *ApJ Lett.*, 801, L11
- Venemans B. P., McMahon R. G., Warren S. J., Gonzalez-Solares E. A., Hewett P. C., Mortlock D. J., Dye S., Sharp R. G., 2007, *MNRAS*, 376, L76
- Venemans B. P., Walter F., Zschaechner L., Decarli R., De Rosa G., Findlay J. R., McMahon R. G., Sutherland W. J., 2016, *ApJ*, 816, 37
- Volonteri M., 2010, *A&ARv*, 18, 279
- Volonteri M., Silk J., Dubus G., 2015, *ApJ*, 804, 148
- Wang F., et al., 2016, *ApJ*, 819, 24
- Wang F., et al., 2017, *ApJ*, 839, 27
- Wang F., et al., 2018a, arXiv:1810.11926v1
- Wang F., et al., 2018b, arXiv:1810.11925v1
- Wang R., et al., 2008, *ApJ*, 687, 848
- Wang R., et al., 2011, *ApJ Lett.*, 739, L34
- Waters C. Z., et al., 2016, arXiv:1612.05245v4
- Willott C. J., Bergeron J., Omont A., 2015, *ApJ*, 801, 123
- Willott C. J., et al., 2007, *AJ*, 134, 2435
- Willott C. J., et al., 2009, *AJ*, 137, 3541
- Willott C. J., et al., 2010, *AJ*, 139, 906
- Willott C. J., Omont A., Bergeron J., 2013, *ApJ*, 770, 13
- Wise J. H., Regan J. A., O'Shea B. W., Norman M. L., Downes T. P., Xu H., 2019, arXiv:1901.07563v1
- Wright E. L., Eisenhardt P. E., Fazio G. G., 1994, *ArXiv Astrophysics e-prints*
- Wright E. L., et al., 2010, *AJ*, 140, 1868
- Wu X.-B., et al., 2015, *Nat*, 518, 512
- Wyithe J. S. B., Loeb A., 2003, *ApJ*, 586, 693
- Yan L., Sajina A., Fadda D., Choi P., Armus L., Helou G., Teplitz H., Frayer D., Surace J., 2007, *ApJ*, 658, 778
- Yang J., et al., 2017, *AJ*, 153, 184
- Yang J., et al., 2018a, arXiv:1811.11915v1
- Yang J., et al., 2018b, arXiv:1810.11927v1
- Zeimann G. R., White R. L., Becker R. H., Hodge J. A., Stanford S. A., Richards G. T., 2011, *ApJ*, 736, 57