

# **AGN Feedback: Simulations of Black Hole**

Ramesh Narayan

# MIND THE OTHER GAP

8 - 12 July, 2013 at the Institute of Astronomy & Kavli Institute for Cosmology (KICC)  
University of Cambridge, UK. Website: <http://www.ast.cam.ac.uk/meetings/2013/MindTheGap>

Gas Accretion on SMBH

AGN

Jets, Winds, Radiation

Lucio Mayer  
Norman Murray  
Thorsten Naab  
Ramesh Narayan  
Eve Ostriker  
Elia Quataert

Local ISM

Host galaxy

Universe

Virginia Bennett  
Cathie Clarke  
Tiago Costa  
Mike Curtis  
Martin Haehnelt  
Simon Karlsson

# Galactic Nucleus




Image credit: Lincoln Greenhill, Jim Moran

# Three Accretion Regimes

## ADAF/Slim Disk


Super-Eddington, radiation trapped  
(Begelman '79; Abramowicz et al. '89)

## Thin Accretion Disk

(Pringle & Rees '72; Shakura & Sunyaev '73; Novikov & Thorne '73)  
Quasars, XRBs in high soft state

## ADAF/RIAF

Radiatively inefficient  
(Ichimaru '77; Rees et al. '82; Narayan & Yi '94, '95; Abramowicz et al. '95)



# Accretion and Outflows

- Analytic disk theory (**1D**) is okay for understanding basic physics of accretion
- **Jets** and **outflows** involve **2D** motions and are beyond analytical theory
- We need numerical simulations:
  - GR (black hole – Kerr metric!) ✓
  - MHD (magnetic fields essential) ✓
  - Radiation (tough problem) ?✓

# Numerical Simulations

- Simulations of varying degrees of complexity have been done over the years
  - Pseudo-Newtonian hydrodynamics
  - Pseudo-N magnetohydrodynamics (MHD)
  - General Relativistic MHD (GRMHD) \*\*
  - Radiation hydro/MHD → GRRMHD
- **Good news:** GRMHD simulations
  - Produce jets and winds from “generic” initial conditions
  - Provide new insights on accretion/jet physics
  - Provide useful information for AGN feedback




# Feedback in Radio Mode/ Maintenance Mode

## ADAF/RIAF

Radiatively inefficient


(Ichimaru '77; Rees et al. '82; Narayan  
& Yi '94, '95; Abramowicz et al. '95)



# GRMHD Simulations of ADAFs

- ADAF/RIAF is the easiest of the three accretion modes to simulate
- We can safely ignore radiation
- Geometrically thick: everything goes fast
- Simulations reach steady state out to fairly large radii
  - $\sim \text{few} \times 100M$  in the best cases (still  $\ll R_B$ )
- ADAFs readily form jets and outflows






Sądowski et al. (2013)

# Two Important Parameters


- BH Jet is powered by BH spin energy
- Jet power is sensitive to BH Spin and Magnetic Flux:

$$P_{\text{jet}} \approx \Phi_{\text{mag}}^2 \Omega_{\text{H}}^2 / c$$


- For a given  $\dot{M}$ , there is a limit to how much Magnetic Flux  $\Phi_{\text{mag}}$  can be pushed into the BH
- System at this limit: Magnetically Arrested Disk (MAD)
- GRMHD simulations of ADAFs readily achieve the MAD limit if sufficient coherent magnetic flux is available
- Jets are highly collimated: feedback efficiency low?



Tchekhovskoy



Sądowski et al. (2013)



BH Jet in MAD state can have a large efficiency:  $\eta_{\text{jet}} = P_{\text{jet}} / \dot{M} c^2$  can even exceed 100% (Tchekhovskoy et al. 2012)

Strong dependence of  $\eta_{\text{jet}}$  on spin parameter  $a_*$

# Disk Wind

- The Disk Wind is more boring:
  - At best only mildly relativistic:  
 $\sim 0.1-0.2 c$
  - Power source is primarily the Disk
  - Power depends modestly on BH spin
  - Power depends modestly on BH Mag Flux
- Large solid angle:  $\sim 2\pi$
- Low power in comparison to jet
- Likely to be efficient source of feedback

v



# Maintenance Mode Feedback Efficiency

Feedback efficiency depends on 3 parameters:

$\dot{M}/\dot{M}_{\text{Edd}}$  ( $\dot{M}/M$ )

$\Omega_{\text{H}}$  (range: 0—1)

$\Phi_{\text{mag}}$  (range: 0— $\Phi_{\text{max}}$ )

Perhaps  $\Phi_{\text{mag}} \rightarrow \Phi_{\text{max}}$   
(MAD)


Still, we need  $\dot{M}$ ,  $\Omega_{\text{H}}$   
before we can “predict” how  
much energy or momentum  
feedback occurs

$$\begin{aligned}\dot{E}_{\text{jet}} &\approx 0.5 \left( \frac{\Phi}{\Phi_{\text{max}}} \right)^2 \left( \frac{\Omega_{\text{H}}}{0.2} \right)^2 \dot{M} c^2 \\ \dot{E}_{\text{wind}} &\approx 0.005 \left[ 1 + 3 \left( \frac{\Phi}{\Phi_{\text{max}}} \right)^2 \left( \frac{\Omega_{\text{H}}}{0.2} \right)^2 \right] \dot{M} c^2 \\ \dot{P}_{\text{jet}} &\approx 0.5 \left( \frac{\Phi}{\Phi_{\text{max}}} \right)^2 \left( \frac{\Omega_{\text{H}}}{0.2} \right)^2 \dot{M} c \\ \dot{P}_{\text{wind}} &\approx 0.1 \dot{M} c\end{aligned}$$

Sądowski et al. (2013)

Available in principle in





# Major Caveat

- We do not have very good information on mass loss in the wind
- Serious limitation for feedback estimates

Unless we figure out the mapping between  $\dot{M}_B$  and  $\dot{M}_{BH}$ , it will be hard to come up with a predictive prescription for AGN energy/mmtm feedback in the maintenance mode

$$\dot{M}_{\text{wind}} \approx \dot{M}_{BH} \left( \frac{r}{r_{\text{in}}} \right)^s$$
$$\frac{\dot{M}_{BH}}{\dot{M}_B} \approx \left( \frac{r_{\text{in}}}{r_B} \right)^s \ll 1 ??$$

# MIND THE OTHER GAP

8 - 12 July, 2013 at the Institute of Astronomy & Kavli Institute for Cosmology (KICC)  
University of Cambridge, UK. Website: <http://www.ast.cam.ac.uk/meetings/2013/MindTheGap>

Gas Accretion on SMBH

AGN

Jets, Winds, Radiation

Lucio Mayer  
Norman Murray  
Thorsten Naab  
Ramesh Narayan  
Eve Ostriker  
Elia Quataert

??

✓✓

Local ISM

Host galaxy

Universe


Virginia Bennett  
Cathie Clarke  
Tiago Costa  
Mike Curtis  
Martin Haehnelt  
Simon Karlsson

# Quasar Mode : I

## Thin Accretion Disk

(Pringle & Rees '72; Shakura & Sunyaev '73; Novikov & Thorne '73)

Quasars, XRBs in high soft state



# Thin Accretion Disk Model

- Makes robust predictions for the radiative luminosity  $L_{\text{disk}}$  (no  $a$  dependence)
- Radiative feedback is straightforward
  - $\eta_{\text{disk}}(a_*)$
- How about mechanical feedback via jets and winds?
- GRMHD simulations have become feasible in recent years, so we can check



# No Jets in Simulations of Thin Accretion Disks

- Thin disk simulations do not show anything that looks like a jet
- However:
  - thin disks are hard to simulate
  - models are converged only to  $R \sim 20M$
  - No jet or wind out to  $20M$
- XRBs in the Thermal-Dominant State (thin disk regime) do not have jets



# Quasar Mode : II

## ADAF/Slim Disk


Super-Eddington, radiation trapped  
(Begelman '79; Abramowicz et al. '89)

Radiative luminosity should be limited  
to at most  $\sim \text{few } L_{\text{Edd}}$

What if  $\dot{M} \gg \dot{M}_{\text{Edd}}$ ? What  
happens to all the energy?


How much energy comes out via a jet  
or a wind?

Need 3D Radiation GRMHD simulations



# Numerical Simulations of Super-Eddington Accretion

- The field has been dominated by Ohsuga (2003...): Radiation hydro/MHD
- Important results on winds
- However, no GR or even SR
- Recent developments:
  - GR+Hydro+Rad(M1) (Sadowski+ '13)
  - GR+MHD+Rad(M1) (McKinney+ '13)
- First results will be out soon



# Very Preliminary

$$\dot{M}_{\text{BH}} = 78 L_{\text{Edd}} / c^2$$

$$L_{\text{radiation}} = 1.1 L_{\text{Edd}}, \eta_{\text{rad}} = 0.015$$

$$L_{\text{Poynting}} = 9.2 L_{\text{Edd}}$$

$$L_{\text{matter}} = 2.3 L_{\text{Edd}}$$

$$L_{\text{total}} = 12.6 L_{\text{Edd}}, \eta_{\text{total}} = 0.16$$

# Summary

- Given  $M_{\text{BH}}$ ,  $\dot{M}_{\text{BH}}$ ,  $\Omega_{\text{H}}$ ,  $\Phi_{\text{mag}}$  ( $= \Phi_{\text{max}}?$ ),  
BH simulators are able to estimate  
 $\dot{E}_{\text{jet}}$ ,  $\dot{P}_{\text{jet}}$ ,  $\dot{E}_{\text{wind}}$ ,  $\dot{P}_{\text{wind}}$
- What other quantities would you like?
  - Angular distribution of energy/mmtm?
  - Lorentz factor/velocity?
  - SMBH spinup/spindown?
- But major uncertainty:  $\dot{M}_{\text{B}}$  vs  $\dot{M}_{\text{BH}}$ 
  - Prognosis is uncertain



# MIND THE OTHER GAP

8 - 12 July, 2013 at the Institute of Astronomy & Kavli Institute for Cosmology (KICC)  
University of Cambridge, UK. Website: <http://www.ast.cam.ac.uk/meetings/2013/MindTheGap>

Gas Accretion on SMBH

AGN

Jets, Winds, Radiation

Lucio Mayer  
Norman Murray  
Thorsten Naab  
Ramesh Narayan  
Eve Ostriker  
Elia Quataert

??

✓✓

Local ISM

Host galaxy

Universe

Virginia Bennett  
Cathie Clarke  
Tiago Costa  
Mike Curtis  
Martin Haehnelt  
Simon Karlsson