

1 Useful background for torques at the ISCO

From e.g. Zimmerman et al. (2005), we can write the maximum accretion disk temperature as

$$T_{\max} = f \left(\frac{3GM\dot{M}}{8\pi R_{in}^3 \sigma} \right)^{1/4} \quad (1)$$

where M is the central mass, \dot{M} is the accretion rate, R_{in} is the innermost radius of the disk and f is a parameter ($O(1)$) that approximates a spectral hardening modification from pure black body ($f = 1$). We can parameterize the innermost (maximum) disk temperature for a generic thin disk as

$$T_{\max} \approx 5.6 \times 10^5 \text{K} \left(\frac{M_{\text{BH}}}{10^8 M_{\odot}} \right)^{-1/4} \left(\frac{\dot{M}}{\dot{M}_{\text{Edd}}} \right)^{1/4} \left(\frac{\eta}{0.1} \right)^{-1/4} \left(\frac{R_{in}}{6r_g} \right)^{-1/4} \quad (2)$$

where M_{BH} is the black hole mass, \dot{M} is the accretion rate (in units of \dot{M}_{Edd}), the Eddington accretion rate, $\eta \sim 0.1$ is the standard accretion efficiency, and $R_{in} = 6r_g$, the ISCO for a Schwarzschild BH. For comparison, keeping all the parameters the same but changing $M_{\text{BH}} = 10M_{\odot}$ yields $T_{\max} \sim 3.1 \times 10^7 \text{K}$.

If the disk is thin, we expect there to be zero-torques at R_{in} , as the material plunges in free-fall at $r < R_{in}$. The resulting zero-torque temperature (T_{ZT}) profile is given by (Zimmermann et al. 2005)

$$T_{ZT} = T_{\max} \left(\frac{r}{R_{in}} \right)^{-3/4} \left[1 - \left(\frac{r}{R_{in}} \right)^{-1/2} \right]^{1/4} \quad (3)$$

and is given by the red curve in Fig. 1. However, magnetic torques (Gammie 1999; Krolik & Agol 2000) or a puffed-up disk (Narayan et al. 1997; Ashford & Paczynski 2003) can have a finite or quite large torque at the inner edge. In this case, the disk temperature profile looks like

$$T_{NZT} = T_{\max} \left(\frac{r}{R_{in}} \right)^{-3/4} \quad (4)$$

or the black curve in Fig. 1.

Integrate the disk over the temperature profiles above (Zimmermann et al. 2005) find

$$L_{\text{disk}}[ZT, NZT] = [1, 3] \frac{GM\dot{M}}{2R_{in}} \quad (5)$$




Figure 1: Comparison of the disk temperature profile due to a change in the boundary condition at the inner edge (R_{in}). Red=standard zero torque at ISCO assumption. Black= non-zero torque at ISCO.

or the disk is $\times 3$ more luminous due to this extra torquing at R_{in} .

2 The heartbeat in GRS195+105

Now if we look at the 'heartbeat' state in GRS 1915 + 105, we have $M_{bh} = 12.4^{+2}_{-1.8} M_{\odot}$ and $d = 8.6^{+2.0}_{-1.6}$ kpc. The period of the oscillation is 50s. Translating this to a $M_{bh} = 10^8 M_{\odot}$, we get an equivalent light-crossing period of 50×10^7 s or ~ 16 years. (Interesting: I wonder if e.g. PG 1302-105 is in a heartbeat state!?)

From Nielsen et al. 2011, their model (section 6 in their paper, Fig. 14, Table 3) for the heartbeat goes as:

1. a wave of excess material (from \dot{M}) originating between $20 - 30r_g$ and propagates radially inward and outward.
2. Disk responds by increasing R_{in} at constant temperature.
3. Disk luminosity increases quickly. At max. R_{in} drops sharply, temp. spikes, disk becomes unstable
4. Disk ejects material. Collides with corona. Hard X-ray pulse.
5. Disk relaxes, density wave subsides.
6. Intense X-ray wind from outer disk due to X-rays
7. Short lived jet

Relevant disk timescales around a $M_{bh} = 10^8 M_{\odot}$ at $R \sim 25r_g$ as:

$$t_{orb} \sim 2\text{days} \left(\frac{R}{25r_g} \right)^{3/2} \frac{r_g}{c} \quad (6)$$

$$t_{th} \sim 2.5\text{months} \left(\frac{\alpha}{0.03} \right)^{-1} \left(\frac{R}{25r_g} \right)^{3/2} \frac{r_g}{c} \quad (7)$$

$$t_{front} \sim 4\text{yr} \left(\frac{h/R}{0.05} \right)^{-1} \left(\frac{\alpha}{0.03} \right)^{-1} \left(\frac{R}{25r_g} \right)^{3/2} \frac{r_g}{c} \quad (8)$$

$$t_{\nu} \sim 82\text{yr} \left(\frac{h/R}{0.05} \right)^{-2} \left(\frac{\alpha}{0.03} \right)^{-1} \left(\frac{R}{25r_g} \right)^{3/2} \frac{r_g}{c}. \quad (9)$$

3 So is the heartbeat relevant?

From t_ν above, it seems like we'd need a slow rise (80-100years) for the first (long) part of the cycle. But that's not consistent with a $\sim 16\text{yr}$ period for the oscillation that you might expect from a simple scaling of light-crossing time ($10M_\odot \rightarrow 10^8M_\odot$). At $10r_g$, t_ν above is 20yrs, so maybe. But why is the disk behaving like this at $10r_g$ (or $25r_g$ for that matter). Could predict observables based on the sequence above from Nielsen et al. (2011).

Nielsen et al. talk about a R_{in} getting closer to the BH accounting for the increase in L_{disk} but they don't explore a state-change as in Fig. 1. That should give us a different prediction and (shorter) timescales in general.