

A new interpretation of optical and infrared variability in quasars

Nicholas P. Ross¹ et al.

¹*Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, United Kingdom*

Changing-look quasars are a recently identified class of active galaxies in which the strong UV continuum and/or broad optical hydrogen emission lines associated with unobscured quasars either appear or disappear on timescales of months to years [1, 2, 3, 4, 5]. The physical processes responsible for this behaviour are still debated, but changes in the black hole accretion rate or accretion disk structure appear more likely than changes in obscuration [6, 7]. Here we report on three epochs of spectroscopy of SDSS J110057.70-005304.5, a quasar at a redshift of $z = 0.378$ whose UV continuum and broad hydrogen emission lines have dramatically faded over the past ≈ 20 years. The change in this quasar was initially seen in the infrared, and an archival spectrum from 2010 shows an intermediate phase of the transition during which the flux below rest-frame 3400\AA has collapsed. This combination is unique compared to previously published examples of changing-look quasars, and is best explained by dramatic changes in the innermost regions of the accretion disk. The optical continuum has been rising again since mid-2016, leading to a prediction of a rise in hydrogen emission line flux in the next year. If our model is confirmed, the physics of changing-look quasars are governed by processes at the innermost stable circular orbit (ISCO) around the black hole, and the structure of the innermost disk. The easily identifiable and monitored changing-look quasars would then provide a new probe of the strong gravity regime.

The changing-look quasar phenomenon, where the dramatic disappearance, or appearance, of the strong UV continuum and/or the prominent broad optical emission lines is seen on month-to-year timescales, is now widely observed [1, 2, 3, 4, 5, 8, 9] yet poorly understood. Changes in obscuration are generally disfavoured due to the timescales currently observed [6, 7], and it is clear that changing-look quasars are a key laboratory for understanding accretion physics and active galactic nuclei (AGN).

The Shakura-Sunyaev α -disk model [10] has long been used to (over-)simply describe the basic properties of optically thick, geometrically thin accretion disks. Indeed we utilize the alpha-disk model in this paper, however, the model is ad hoc, only parameterizing disk viscosity, and does not permit predictions of global changes to the disk [11]. Real AGN disks seem to be cooler [e.g., 12] and larger [e.g., 13, 14, 15, 16] than the alpha-disk model predicts, and viscosity seems likely due to magnetic fields [17] with additional contributions to turbulence from the effects of embedded objects in the disk [e.g., 18].

Changing-look quasars have traditionally been discovered by looking for large, $|\Delta m| > 1$ magnitude changes in optical light curves of quasars or galaxies (e.g. across $3720\text{\AA} < \lambda < 5680\text{\AA}$, in the g -band). However, we have taken advantage of the ongoing Near-Earth Object WISE Reactivation mission [NEOWISE-R; 19, 20, 21], as well as the Dark Energy Camera Legacy Survey (DECaLS¹) in order to discover new changing-look quasars. While previous efforts have used the 1-year baseline of the WISE mission to identify CLQs [22, e.g., []], our team is the first to extend . Our team is the first to extend this selection to the infrared using NEOWISE-R mission data, and we have identified

¹legacysurvey.org/decamls/




Figure 1 Multi-wavelength light curve of J1100-0053, including optical data from LINEAR, CRTS, SDSS, PanSTARRS and DECaLS, and mid-IR data from the WISE satellite. The three vertical lines illustrate the epochs of the three optical spectra presented in Figure 2. J1100-0053 was flagged for further study due to the extreme IR fading observed by WISE. Note that the optical emission has been recovering over the past few years; we predict the IR emission will similarly recover over the next few years.

a sample of SDSS quasars that show dramatic decreases in their IR flux over the course of a few years. These changes are on timescales too short to be due to changes in obscuration, so a different explanation is needed.

In this article we present the $z = 0.378$ quasar SDSS J110057.70-005304.5 (hereafter J1100-0053) for which we have spectral observations showing a transition in the blue-continuum slope traditionally associated with the black-body spectrum of an object with broad hydrogen emission lines, into a ‘dim state’ where the rest-frame UV flux is suppressed, and then returning to a blue-continuum sloped quasar. We present a model that invokes changes at the ISCO to be the triggering event for the change in the accretion disk, which along with the changes in the broad emission lines, explains a major change to the disk interior to $150r_g$ (where r_g is the gravitational radius; $r_g = \frac{GM}{c^2}$) as well as the IR light curves. Critically, our model makes predictions to the future behaviour of J1100-0053.

1 Target Selection and Observations

We started by matching the SDSS-III BOSS Data Release 12 Quasar catalog [DR12Q; 23] to the NEOWISE-R IR data (WISE W1 at $3.4\mu\text{m}$, WISE W2 at $4.6\mu\text{m}$). We found ~ 200 objects identified by a factor of 2 or more change in




Figure 2 Optical spectra of J1100-0053 obtained on MJD 51908 (blue; SDSS), 55302, (red; BOSS) and 57809 (black; Palomar/DBSP). *Left::* The full optical spectra; *Right (top)::* Zoom in on the $H\beta$ -[O III] complex; *Right (bottom)::* Zoom in on the $H\alpha$ -N II complex.

the observed WISE W1 and W2 bands over the course of typically three or four years [see 21, and the Supplemental Material for the detailed NEOWISE-R selection]. Scanning these 200 objects, we also examined the change in optical colour using the SDSS and DECaLS imaging surveys in order to identify changes suggestive of changing-look quasars. From this inspection, a priority list of ≈ 70 quasar targets was derived and we obtained new optical spectroscopy from the Palomar 5m telescope. J1100-0053 was one of these 70 objects, but critically, had spectra from both SDSS and BOSS and was thus a priority target.

Figure 1 presents the light curve of J1100-0053. Along with WISE IR data, optical data from the SDSS, Catalina Real-time Transient Survey [CRTS; 24, 25], LINEAR [26] and PanSTARRS [27, 28, 29, 30] are available. Figure 2 shows the three optical spectra of J1100-0053 from the SDSS, BOSS and Palomar observations taken on MJD 51908 (UT 2000 December 30), 55302 (UT 2010 April 16) and 57809 (UT 2017 February 25), respectively. The first-epoch SDSS spectrum shows a typical blue quasar, but blue continuum then collapses in the second epoch BOSS spectrum taken 10 years later. However, the blue continuum has then returned in the third epoch spectrum 7 years later, albeit at a diminished flux from the initial spectrum. The Supplemental Material gives further observation details including an achival ROSAT detection.

2 Discussion

Our model of thermal emission from a multicolour disk implies changes in the region from the ISCO to \sim few tens-100 r_g are required to suppress flux into the observed g -band. In particular, we suggest a physical collapse of the disk scale height due to a cooling front propagating outward from the ISCO.

For J1100-0053 we apply our model as follows. We start with an inflated (slim) disk. We assume a non-zero torque at the ISCO and $h/r \sim 0.2$ inside of $r \sim 100r_g$. This is the initial state circa 2000 (MJD 51900). A non-zero torque at the ISCO implies that matter in the plunging region is connected (however weakly) to matter outside the ISCO, probably by magnetic fields [e.g., 31, 32]. A non-zero torque at the ISCO maintains a hotter innermost disk than a condition of zero torque at the ISCO, and an assumption of non-zero torque is particularly appropriate if disk viscosity and accretion are indeed driven by magnetic fields. In order to explain data subsequent to 2007, we assume a cooling front propagates out from the ISCO over a timescale t_{front} . The simplest explanation is that the non-zero torque condition at the ISCO changes to a (more nearly) zero-torque condition, leading to a dramatically cooler, thinner disk near the ISCO. As the cooling front propagates, the drop in temperature leads to a drop in flux. Our model requires cooler regions behind the front to emit 10% the flux of the initial hotter disk, and the disk height drops by a factor ~ 2 . The dimming of the inner disk causes a drop in the ionizing photon flux (L_{ion}), which will cause: the Balmer lines to drop in flux after a light travel time of months and the IR from the outer disk/torus to drop in flux after a light travel time of ~ 3 years.

If the inner accretion disk is usually inflated [see e.g., 33, 34, 35], such a cooling front will naturally produce: 1) a collapse in the scale height of the disk; 2) a decrease in flux moving from UV to longer optical wavelengths; 3) a temporarily thicker scattering atmosphere, further decreasing flux at short wavelengths. This model implies changes to the optical emission moving from shorter to longer wavelengths (as the radius of the cooling front increases), on months-to-years-long timescales. It also predicts a longer time to recover the original flux (compared to the initial




Figure 3 A cartoon illustration of our model explaining the unusual spectral evolution of J1100-0053. In 2000, corresponding to the SDSS spectral epoch, the quasar has a standard modified α -disk, with a geometrically thin, optically thick disk that is inflated near the ISCO, i.e., the inner edge of the accretion disk. Circa 2007, an event occurs that deflates the inner disk, leaving some scattering clouds, and causing a cooling front to propagate inwards. Circa 2012, the front reflects back, re-heating the inner accretion disk. We predict that in the next year, the quasar should roughly return to its initial state.

collapse) as a front will move more slowly in a thinner disk (see Fig. 2). A decrease in the UV flux would be expected to cause a decrease in IR flux, as the heating of the IR-emitting dusty torus is reduced; however, there should be a delay due to light travel time as well [e.g., 36].

Since we assume the disk starts in a puffed-up state ($h/r \sim 0.2$) and since the front cooling time (equation 5 in the Supplemental Material) is inversely proportional to h/r and α , the front propagates faster in a puffed-up disk than in a thinner disk. By 2010 (MJD 55300) the front has reached $r \sim 50r_g$. During that time, the collapsing disk height increases the number density of scatterers and the temporary cold phase formed during cooling produces the remarkable blue downturn in the 2010 spectrum. The cooling front continues to propagate radially outward but cools less efficiently at larger disk radii. Eventually, a heating front propagates back inwards, analogous to the well-known accretion disk limit cycle mechanism in models of dwarf novae outbursts [e.g., 37]. The returning heating front travels more slowly because the disk is thinner (and t_{front} is inversely proportional to h/r), and will re-inflate the disk as it propagates inwards towards the SMBH. This means the return to normal will be asymmetric in time, as observed, and the g -band bottoms out first, because that wavelength is dominated by emission coming from $r \sim 100r_g$ (see discussion in the Supplemental Material).

Using [38] and [33], Figure 3 shows a model for a $M_{\text{BH}} = 3 \times 10^8 M_\odot$, radiative efficiency of $\epsilon = 0.1$, accretion rate in units of Eddington accretion, $\dot{M} = 0.032$, inner disk radius of $6r_g$ and outer disk radius of $10,000r_g$. The resulting model spectra can be seen in Figure 3. We expect the front to return to the ISCO in about 2018. That means the broad Balmer lines will come back a few months later, but the WISE IR flux should not come back until about 2021.

[39] observed a similar event to J1100-0053 with the source SDSS J231742.60+000535.1. However, their object provided an ambiguous case, as the IR brightness of their source did not decline. This is consistent with our model, as their cooling event is relatively brief. We discuss this object and the [39] result further in the Supplemental Material.

In this letter, we have shown that a simple phenomenological model with a propagating cooling front is capable of describing the gross spectral and temporal variations in a changing looking quasar. Our model makes a prediction for this source, testable over the next few years and implies that changing looking quasars as a class are driven by changes near the ISCO, close to the SMBH. By monitoring changing look quasars we introduce new tests of models of accretion disk physics and a new probe of the strong gravity regime.

1. LaMassa, S. M. *et al.* The Discovery of the First “Changing Look” Quasar: New Insights Into the Physics and Phenomenology of Active Galactic Nucleus. *ApJ* **800**, 144 (2015). 1412.2136.
2. Runnoe, J. C. *et al.* Now you see it, now you don’t: the disappearing central engine of the quasar J1011+5442. *MNRAS* **455**, 1691–1701 (2016). 1509.03640.
3. MacLeod, C. L. *et al.* A systematic search for changing-look quasars in SDSS. *MNRAS* **457**, 389–404 (2016). 1509.08393.
4. Ruan, J. J. *et al.* Toward an Understanding of Changing-look Quasars: An Archival Spectroscopic Search in SDSS. *ApJ* **826**, 188 (2016). 1509.03634.

5. Yang, Q. *et al.* Discovery of 21 New Changing-look AGNs in Northern Sky. *ArXiv e-prints* (2017). 1711.08122v1.
6. Hutsemékers, D., Agís González, B., Sluse, D., Ramos Almeida, C. & Acosta Pulido, J.-A. Polarization of the changing-look quasar J1011+5442. *A&A* **604**, L3 (2017). 1707.05540.
7. Sheng, Z. *et al.* Mid-infrared Variability of Changing-look AGNs. *ApJ* **846**, L7 (2017). 1707.02686.
8. Gezari, S. *et al.* iPTF Discovery of the Rapid “Turn-on” of a Luminous Quasar. *ApJ* **835**, 144 (2017). 1612.04830.
9. Rumbaugh, N. *et al.* Extreme variability quasars from the Sloan Digital Sky Survey and the Dark Energy Survey. *ArXiv e-prints* (2017). 1706.07875.
10. Shakura, N. I. & Sunyaev, R. A. Black holes in binary systems. Observational appearance. *A&A* **24**, 337 (1973).
11. King, A. Accretion disc theory since Shakura and Sunyaev. *Mem. Soc. Astron. Italiana* **83**, 466 (2012). 1201.2060.
12. Lawrence, A. The UV peak in active galactic nuclei: a false continuum from blurred reflection? *MNRAS* **423**, 451–463 (2012). 1110.0854.
13. Pooley, D., Blackburne, J. A., Rappaport, S. & Schechter, P. L. X-Ray and Optical Flux Ratio Anomalies in Quadruply Lensed Quasars. I. Zooming in on Quasar Emission Regions. *ApJ* **661**, 19–29 (2007). astro-ph/0607655.
14. Morgan, C. W., Kochanek, C. S., Morgan, N. D. & Falco, E. E. The Quasar Accretion Disk Size-Black Hole Mass Relation. *ApJ* **712**, 1129–1136 (2010). 1002.4160.
15. Morgan, C. W. *et al.* Further Evidence that Quasar X-Ray Emitting Regions are Compact: X-Ray and Optical Microlensing in the Lensed Quasar Q J0158-4325. *ApJ* **756**, 52 (2012). 1205.4727.
16. Mosquera, A. M. & Kochanek, C. S. The Microlensing Properties of a Sample of 87 Lensed Quasars. *ApJ* **738**, 96 (2011). 1104.2356.
17. Balbus, S. A. & Hawley, J. F. A powerful local shear instability in weakly magnetized disks. I - Linear analysis. II - Nonlinear evolution. *ApJ* **376**, 214–233 (1991).
18. McKernan, B., Ford, K. E. S., Kocsis, B., Lyra, W. & Winter, L. M. Intermediate-mass black holes in AGN discs - II. Model predictions and observational constraints. *MNRAS* **441**, 900–909 (2014). 1403.6433.
19. Mainzer, A. *et al.* Initial Performance of the NEOWISE Reactivation Mission. *ApJ* **792**, 30 (2014). 1406.6025.
20. Meisner, A. M., Lang, D. & Schlegel, D. J. Deep Full-sky Coadds from Three Years of WISE and NEOWISE Observations. *AJ* **154**, 161 (2017). 1705.06746.
21. Meisner, A. M. *et al.* Searching for Planet Nine with Coadded WISE and NEOWISE-Reactivation Images. *AJ* **153**, 65 (2017). 1611.00015.
22. Assef, R. J. *et al.* The WISE AGN Catalog. *1706.09901v1* (2017). 1706.09901.

23. Pâris, I., Petitjean, P., Ross, N. P. *et al.* The Sloan Digital Sky Survey Quasar Catalog: Twelfth data release. *A&A* **597**, A79 (2017). 1608.06483.
24. Drake, A. J. *et al.* First Results from the Catalina Real-Time Transient Survey. *ApJ* **696**, 870–884 (2009). 0809.1394.
25. Mahabal, A. A. *et al.* Discovery, classification, and scientific exploration of transient events from the Catalina Real-time Transient Survey. *Bulletin of the Astronomical Society of India* **39**, 387–408 (2011). 1111.0313.
26. Sesar, B. *et al.* Exploring the Variable Sky with LINEAR. I. Photometric Recalibration with the Sloan Digital Sky Survey. *AJ* **142**, 190 (2011). 1109.5227.
27. Kaiser, N. *et al.* The Pan-STARRS wide-field optical/NIR imaging survey. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 7733 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 0 (2010).
28. Stubbs, C. W. *et al.* Precise Throughput Determination of the PanSTARRS Telescope and the Gigapixel Imager Using a Calibrated Silicon Photodiode and a Tunable Laser: Initial Results. *ApJS* **191**, 376–388 (2010). 1003.3465.
29. Tonry, J. L. *et al.* The Pan-STARRS1 Photometric System. *ApJ* **750**, 99 (2012). 1203.0297.
30. Magnier, E. A. *et al.* The Pan-STARRS 1 Photometric Reference Ladder, Release 12.01. *ApJS* **205**, 20 (2013). 1303.3634.
31. Gammie, C. F. Efficiency of Magnetized Thin Accretion Disks in the Kerr Metric. *ApJ* **522**, L57–L60 (1999). astro-ph/9906223.
32. Agol, E. & Krolik, J. H. Magnetic Stress at the Marginally Stable Orbit: Altered Disk Structure, Radiation, and Black Hole Spin Evolution. *ApJ* **528**, 161–170 (2000). astro-ph/9908049.
33. Sirko, E. & Goodman, J. Spectral energy distributions of marginally self-gravitating quasi-stellar object discs. *MNRAS* **341**, 501–508 (2003). astro-ph/0209469.
34. Thompson, T. A., Quataert, E. & Murray, N. Radiation Pressure-supported Starburst Disks and Active Galactic Nucleus Fueling. *ApJ* **630**, 167–185 (2005). astro-ph/0503027.
35. Hopkins, P. F. & Quataert, E. An analytic model of angular momentum transport by gravitational torques: from galaxies to massive black holes. *MNRAS* **415**, 1027–1050 (2011). 1007.2647.
36. Jun, H. D. *et al.* Infrared Time Lags for the Periodic Quasar PG 1302-102. *ApJ* **814**, L12 (2015). 1511.01515.
37. Cannizzo, J. K. On the $M_V(\text{peak})$ versus Orbital Period Relation for Dwarf Nova Outbursts. *ApJ* **493**, 426–430 (1998). astro-ph/9712210.
38. Ford, K. E. S. *et al.* *in prep.* (2018).
39. Guo, H. *et al.* The Optical Variability of SDSS Quasars from Multi-epoch Spectroscopy. III. A Sudden UV Cutoff in Quasar SDSS J2317+0005. *ApJ* **826**, 186 (2016). 1605.07301.