

A new interpretation of optical and infrared variability in Quasars

Nicholas P. Ross¹ et al.

¹*Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, United Kingdom*

Changing-look quasars are a class of recently identified object in which the strong UV continuum and broad optical hydrogen emission lines associated with unobscured quasars either appear or disappear on timescales of months to years (1; 2; 3; 4). The physical processes responsible for this behaviour are still debated, but changes in the black hole accretion rate or accretion disk structure appear more likely than changes in obscuration (5; 6). Here we report on three epochs of spectroscopy of SDSS J110057.70-005304.5, a quasar whose UV continuum and broad hydrogen emission lines have dramatically faded over the past 20 years. An archival spectrum of this quasar from 2010 shows an intermediate phase of the transition during which the flux below rest-frame 340nm has collapsed. This is unique compared to previously published examples of changing-look quasars, and is best explained by dramatic changes in the innermost regions of the accretion disk. The optical continuum has been rising again since mid-2016, leading to a prediction of a rise in hydrogen emission line flux in the next few months. If our model is confirmed, the physics of ‘changing look’ quasars are governed by processes at the innermost stable circular orbit (ISCO) around the black hole, and the structure of the innermost disk. Thus, the easily identifiably and monitored Changing Look Quasars would then provide a new probe of the strong gravity regime.

The “Changing-Look” quasar phenomenon, where the dramatic disappearance, or appearance, of prominent broad optical emission lines is seen on year timescales, is now widely observed, (1; 2; 3; 4; 7; 8). yet poorly understood. Changes in obscuration are generally disfavoured, (5; 6), and it is clear that the CLQs are a key laboratory into understanding accretion physics and the nature of the AGN broad line region (BLR). Zooming in on the AGN accretion disk, the famous α -disk model (9) for a optically thick, geometrically thin disk ($h/R \ll 1$, where h is the vertical scale height of the disk) is known to have serious short-comings e.g. (10; 11; 12) AGN seem to be cooler than they ought to be (e.g., 12) with the SEDs of AGN showing a universal near-UV shape, reaching a maximum in νS_ν around 1100Å, Such a peak suggests a characteristic temperature of $T \sim 30\,000\text{K}$, whereas for a thermal model, the characteristic temperature should be roughly $T \sim 100\,000\text{K}$. Moreover, constraints from microlensing observations for the size of the optical emission region (13; 14; 15; 16, e.g.,) suggestion this region is larger than the one predicted by the standard Shakura-Sunyaev disk.

CLQs have traditionally been discovered by looking for large, $|\Delta m| > 1$ magnitude changes in the optical light curves (e.g. in the g -band). However, we have taken advantage of the ongoing Near-Earth Object WISE Reactivation mission (NEOWISE-R)(17; 18; 19), as well as the Dark Energy Camera Legacy Survey (DECaLS¹) in order to discover new CLQs. Our team is the first to extend this selection to the infrared using NEOWISE-R mission data. Indeed, we have found a sample of SDSS quasars that show *dramatic decreases in their IR flux over the course of a few years*. These changes are on timescales too short to be considered due to changes in obscuration, so a new explanation is needed.

¹legacysurvey.org/decamls/




Figure 1 The light curve of J110057. SDSS, DECaLS and PanSTARRS give the optical photometry. The WISE IR light curves are shown and their dramatic decrease led to the identification of J110057. The three spectral epochs are shown by the vertical lines.

In this article we present the $z = 0.378$ quasar SDSS J110057.70-005304.5 that we have observed transitioning from a blue continuum sloped object to become a regular galaxy. However, along with the changes in the BELs, we see a major change to the disk interior to $150R_g$.

1 Results

Matching the SDSS/BOSS Data Release 12 Quasar catalog (Paris17) to the NEOWISE-R IR data (W1 is $3.4\mu\text{m}$, W2 is $4.6\mu\text{m}$) our team found ≈ 200 objects with fading light IR light curves. These objects were identified by a factor of 2 or more drop in the observed WISE W1 and W2 bands.

Figure 1 gives the optical light curve of J110057. Figure 2 shows the three optical spectra of J110057.

Checking the data archives we found there was no source within 30 arcsec in the VLA FIRST, i.e., at 21 cm radio frequencies. None of the *Hubble Space Telescope*, the *Spitzer Space Telescope* or the *Kepler Mission* has observed J110057 patch of sky. It is also not in the HSC DR1 footprint. There is a detection in ROSAT, using the 2nd all-sky survey (2RXS; Boller et al. 2016, A&A, 588, 103) as 2RXS J110058.1-005259 with 27.00 counts (count error 6.14), count rate = 0.06 ± 0.01 and an exposure time of = 431.95 seconds. The NED gives J110057 as $1.27 \pm 0.28 \times 10^{-12}$ erg/cm²/s in the 0.1-2.4 keV range (unabsorbed flux). J110057 Neither *Chandra* or *XMM-Newton*.

Figure 2 shows the optical spectra for J110057 from the SDSS, the BOSS and Palomar observations taken on MJD 51908, 55302, and 57809. The MJD 51908 SDSS spectrum is of a typical blue quasar, the blue continuum then collapses




Figure 2 Optical spectra of J110057 obtained on MJD 51908 (blue; SDSS), 55302, (red; BOSS) and 57809 (black; Palomar/DBSP). *Left::* The full optical spectra; *Right (top)::* Zoom in on the $H\beta$ -[O III] complex; *Right (bottom)::* Zoom in on the $H\alpha$ -N II complex.

2 Discussion

Our preliminary model of emission from a multicolor disk (Fig. 3) imply changes from the innermost stable circular orbit (ISCO) to \sim few tens-100 R_g are required to suppress flux into the observed g -band. In particular, we suggest a physical collapse of the disk scale height due to a cooling front propagating outward from the ISCO. If the inner accretion disk is usually inflated (see e.g. Sirko & Goodman 2003, Thompson et al. 2005, Hopkins & Quataert 2011), such a cooling front will naturally produce: 1) a collapse in the scale height of the disk; 2) a decrease in flux moving from UV to longer optical wavelengths; 3) a temporarily thicker scattering atmosphere, further decreasing flux at short wavelengths. This model implies changes to the optical emission moving from shorter to longer wavelengths (as the radius of the cooling front increases), on months-to-years-long timescales. It also predicts a longer time to recover the original flux (compared to the initial collapse, as a front will move more slowly in a thinner disk (see Fig. 2). A decrease in the UV flux would also be expected to cause a decrease in IR flux, as the heating of the IR-emitting dusty torus is reduced; however, there should also be a delay due to light travel time.

For J110057 we apply our model as follows. We start with an inflated disk, with non-zero torque at the ISCO, and $h/R \sim 0.2$ inside of $R \sim 100R_g$. This is the initial state in circa 2000 (\approx MJD 51900). Around 2007, there is a switch to a zero torque at ISCO state. A cooling front is set-up, which propagates out from the ISCO at the timescale, t_{front} . Regions behind the front emit flux at $\sim 0.1L$ of what they did prior to the passage of the front (**NPR** comment: due to drop in T ? $T \downarrow \times 1.78 \Rightarrow L \downarrow \times 10$), and the temperature decrease leads the height to drop by a factor of 2. (**NPR** comment: just due to less kinetic energy??). L_{ion} starts decreasing due to the drop in ionizing photons, which in turn causes the Balmer lines to also decrease in flux.

Since the disk starts puffed-up, the cooling front time is not long, and by 2010 (MJD 55300), the front has




Figure 3 PLACE HOLDER OF OBSERVED AND MODEL FOR J110057.

reached $R \sim 50R_g$. During that time, the collapsing disk height increases the number density of scatterers, which in turn causes Rayleigh scattering producing the blue downturn in the 2010 spectrum. The cooling front keeps going, until it hits the part of the disk where it is normally thin, around $R = 100R_g$, arriving around 2012. This sets up another (heating) front, which will travel *back in* towards the SMBH, and re-inflate the disk. This ‘returning’ front travels more slowly because the disk is thinner. It also means the return to normal will be asymmetric in time, as observed, and the *g*-band bottoms out first because that is coming from $R \sim 100R_g$.

Using Ford et al and Sirko & Goodman 2003, Figure 3 shows a model for a $M_{\text{BH}} = 3 \times 10^8 M_\odot$, radiative efficiency of $\epsilon = 0.1$, accretion rate in units of Eddington accretion, $\dot{M} = 0.032$, inner and outer disk radii in units of r_g of SMBH of $\text{radius}_{\text{in}}=6.0$, $\text{radius}_{\text{out}}=1.0 \times 10^4$. The resulting model spectra can be seen in Figure 3.

We expect the front to return to the ISCO in about 2018. That means the H lines will come back a few months later, but the WISE IR flux shouldn’t come back until about 2021.

(20) observed a similar event to J110057, with SDSS J231742.60 +000535.1. However, their object provided an ambiguous case, as the IR brightness of their source did not decline. However this is consistent with our model, as their cooling event is relatively brief.

1. LaMassa, S. M. *et al.* The Discovery of the First “Changing Look” Quasar: New Insights Into the Physics and Phenomenology of Active Galactic Nucleus. *ApJ* **800**, 144 (2015). 1412.2136.
2. Runnoe, J. C. *et al.* Now you see it, now you don’t: the disappearing central engine of the quasar J1011+5442. *MNRAS* **455**, 1691–1701 (2016). 1509.03640.
3. MacLeod, C. L. *et al.* A systematic search for changing-look quasars in SDSS. *MNRAS* **457**, 389–404 (2016). 1509.08393.

4. Ruan, J. J. *et al.* Toward an Understanding of Changing-look Quasars: An Archival Spectroscopic Search in SDSS. *ApJ* **826**, 188 (2016). 1509.03634.
5. Hutsemékers, D., Agís González, B., Sluse, D., Ramos Almeida, C. & Acosta Pulido, J.-A. Polarization of the changing-look quasar J1011+5442. *Astron. & Astrophys.* **604**, L3 (2017). 1707.05540.
6. Sheng, Z. *et al.* Mid-infrared Variability of Changing-look AGNs. *ApJ Lett.* **846**, L7 (2017). 1707.02686.
7. Gezari, S. *et al.* iPTF Discovery of the Rapid “Turn-on” of a Luminous Quasar. *ApJ* **835**, 144 (2017). 1612.04830.
8. Rumbaugh, N. *et al.* Extreme variability quasars from the Sloan Digital Sky Survey and the Dark Energy Survey. *ArXiv e-prints* (2017). 1706.07875.
9. Shakura, N. I. & Sunyaev, R. A. Black holes in binary systems. Observational appearance. *Astron. & Astrophys.* **24**, 337 (1973).
10. Antonucci, R. Constraints on Disks Models of The Big Blue Bump from UV/Optical/IR Observations. In Poutanen, J. & Svensson, R. (eds.) *High Energy Processes in Accreting Black Holes*, vol. 161 of *Astronomical Society of the Pacific Conference Series*, 193 (1999). astro-ph/9810067.
11. Koratkar, A. & Blaes, O. The Ultraviolet and Optical Continuum Emission in Active Galactic Nuclei: The Status of Accretion Disks. *PASP* **111**, 1–30 (1999).
12. Lawrence, A. The UV peak in active galactic nuclei: a false continuum from blurred reflection? *MNRAS* **423**, 451–463 (2012). 1110.0854.
13. Pooley, D., Blackburne, J. A., Rappaport, S. & Schechter, P. L. X-Ray and Optical Flux Ratio Anomalies in Quadruply Lensed Quasars. I. Zooming in on Quasar Emission Regions. *ApJ* **661**, 19–29 (2007). astro-ph/0607655.
14. Morgan, C. W., Kochanek, C. S., Morgan, N. D. & Falco, E. E. The Quasar Accretion Disk Size-Black Hole Mass Relation. *ApJ* **712**, 1129–1136 (2010). 1002.4160.
15. Morgan, C. W. *et al.* Further Evidence that Quasar X-Ray Emitting Regions are Compact: X-Ray and Optical Microlensing in the Lensed Quasar Q J0158-4325. *ApJ* **756**, 52 (2012). 1205.4727.
16. Mosquera, A. M. & Kochanek, C. S. The Microlensing Properties of a Sample of 87 Lensed Quasars. *ApJ* **738**, 96 (2011). 1104.2356.
17. Mainzer, A. *et al.* Initial Performance of the NEOWISE Reactivation Mission. *ApJ* **792**, 30 (2014). 1406.6025.
18. Meisner, A. M., Lang, D. & Schlegel, D. J. Deep Full-sky Coadds from Three Years of WISE and NEOWISE Observations. *AJ* **154**, 161 (2017). 1705.06746.
19. Meisner, A. M. *et al.* Searching for Planet Nine with Coadded WISE and NEOWISE-Reactivation Images. *AJ* **153**, 65 (2017). 1611.00015.
20. , H. *et al.* The Optical Variability of SDSS Quasars from Multi-epoch Spectroscopy. III. A Sudden UV Cutoff in Quasar SDSS J2317+0005. *ApJ* **826**, 186 (2016). 1605.07301.