

185: Introduction to Complex Analysis

Nir Elber

Spring 2022

CONTENTS

1	Introducing Complex Numbers	3
1.1	January 19	3
1.2	January 21	5
1.3	January 24	10
1.4	January 26	15
1.5	January 28	21
1.6	January 31	26
1.7	February 2	32
1.8	February 4	36
1.9	February 7	42

THEME 1: INTRODUCING COMPLEX NUMBERS

1.1 January 19

It is reportedly close enough to start.

1.1.1 Logistics

We are online for the first two weeks, as with the rest of Berkeley. We will be using bCourses a lot, so check it frequently. There is also a website. There is a homework due on Friday, but do not worry about it.

Here are some syllabus things.

- Our main text is *Complex Variables and Applications, 8th Edition* because it is the version that Professor Morrow used. There is a free copy online.
- Homeworks are readings (for each course day) and weekly problem sets. Late homeworks are never accepted.
- Lowest two homework scores are dropped.
- There are two midterms and a final. The final is cumulative, as usual. The final can replace one midterm if the score is higher.
- Regrade requests can be made in GradeScope within one week of being graded.
- The class is curved but usually only curved at the end. The average on exams is expected to be 80%–83%.

1.1.2 Complex Numbers

Welcome to complex analysis. What does that mean?




Idea 1.1. In complex analysis, we study functions $f : \mathbb{C} \rightarrow \mathbb{C}$, usually analytic to some extent.

There are two pieces here: we should study \mathbb{C} in themselves and then we will study the functions.

Complex
numbers

Definition 1.2 (Complex numbers). The set of complex numbers \mathbb{C} is $\{a + bi : a, b \in \mathbb{R}\}$, where $i^2 = -1$.

Hopefully \mathbb{R} is familiar from real analysis. As an aside, we see $\mathbb{R} \subseteq \mathbb{C}$ because $a = a + 0i \in \mathbb{C}$ for each $a \in \mathbb{R}$. The complex numbers have an inherent geometry as a two-dimensional plane.



The point is that \mathbb{C} looks like the real plane \mathbb{R}^2 . More precisely, $\mathbb{C} \cong \mathbb{R}^2$ as an \mathbb{R} -vector space, where our basis is $\{1, i\}$.

We would like to understand \mathbb{C} geometrically, “as a space.” The first step here is to create a notion of size.

Norm on \mathbb{C}

Definition 1.3 (Norm on \mathbb{C}). We define the **norm map** $|\cdot| : \mathbb{C} \rightarrow \mathbb{R}_{\geq 0}$ by $|z| := \sqrt{z\bar{z}}$. In other words,

$$|a + bi| := \sqrt{a^2 + b^2}.$$

Note that this agrees with the absolute value on \mathbb{R} : for $a \in \mathbb{R}$, we have $\sqrt{a^2} = |a|$.

Norm functions, as in the real case, give us a notion of distance.

Metric on \mathbb{C}

Definition 1.4 (Metric on \mathbb{C}). We define the *metric on \mathbb{C}* to be $d_{\mathbb{C}}(z_1, z_2) := |z_1 - z_2|$.

One can check that this is in fact a metric, but we will not do so here.

Remark 1.5. The distance in \mathbb{C} is defined to match the distance in \mathbb{R}^2 under the basis $\{1, i\}$.

Again as we discussed in real analysis, having a metric gives us a metric topology by open balls. Lastly it is this topology that our geometry will follow from: we have turned \mathbb{C} into a topological space.

1.1.3 Complex Functions

There are lots of functions on \mathbb{C} , and lots of them are terrible. So we would like to focus on functions with some structure. We’ll start with *continuous functions*, which are more or less the functions that respect topology.

Then from continuous functions, we will be able to define *holomorphic functions*, which are complex differentiable. This intended to be similar to being real differentiable, but complex differentiable turns out to be a very strong condition. Nevertheless, everyone’s favorite functions are holomorphic.

Example 1.6. Polynomials, \exp , \sin , and \cos are all holomorphic.

To make concrete that complex differentiability is stronger than real differentiability, the Cauchy–Riemann equations which provides a partial differential equation to test complex differentiability.

From here we define *analytic functions*, which essentially are defined as taking the form

$$f(z) := \sum_{k=0}^{\infty} a_k z^k.$$

Analytic functions are super nice in that we have an ability to physically write them down, so the following theorem is amazing.

Theorem 1.7. Holomorphic functions on \mathbb{C} are analytic.

To prove this, we will need the following result, which is what Professor Morrow calls the most fundamental result in complex analysis, the *Cauchy integral formula*.

In short, the Cauchy integral formula lets us talk about the value of holomorphic functions (and derivatives) at a point in terms of integrals around the point. This will essentially let us build the power series for a holomorphic function by hand. But as described, we will need a notion of complex (path) integration to even be able to talk about the Cauchy integral formula.

The Cauchy integral formula has lots of applications; for example, *Liouville's theorem* on holomorphic functions and the *Fundamental theorem of algebra*.

Remark 1.8. It is very hard to spell Liouville.

Additionally, we remark that our study of holomorphic functions, via the Cauchy integral formula, will boil down to a study of complex path integrals. So we will finish out our story with the *Residue theorem*, which provides a very convenient way to compute such integrals.

Then as a fun addendum, we talk about automorphisms of the complex numbers.

Automor-
phisms of \mathbb{C}

Definition 1.9 (Automorphisms of \mathbb{C}). A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is an *automorphism of \mathbb{C}* if it is bijective and both f and f^{-1} are holomorphic.

What is amazing is that all of these functions have a concrete description in terms of *Möbius transformations*.

1.1.4 Why Care?

Whenever taking a class, it is appropriate to ask why one should care. Here are some reasons to care.

- Algebraic geometry in its study of complex analytic spaces uses complex analysis.
- Analytic number theory (e.g., the Prime number theorem) makes heavy use of complex analysis.
- Combinatorics via generating functions can use complex analysis.
- Physics uses complex analysis.

The first two Professor Morrow is more familiar with, the last two less so.

1.2 January 21

We're reviewing set theory today.

1.2.1 Set Theory Notation

We have the following definitions.

- \emptyset means the empty set.
- $a \in X$ means that a is an element of the set X .
- $A \subseteq B$ means that A is a subset of B .
- $A \subsetneq B$ means that A is a proper subset of B .

- $A \cup B$ consists of the elements which are in at least one of A or B .
- $A \cap B$ consists of the elements which are in both A and B .
- $A \setminus B$ consists of the elements of A which are not in B .
- Two sets A and B are *disjoint* if and only if $A \cap B = \emptyset$.
- Given a set X , we define $\mathcal{P}(X)$ to be the set of all subsets of X .
- $|X| = \#X$ is the cardinality of X , or (roughly speaking) the number of elements of X .

As an example of unwinding notation, we have the following.

Proposition 1.10 (De Morgan's Laws). Fix $\mathcal{S} \subseteq \mathcal{P}(X)$ a collection of subsets of a set X . Then

$$X \setminus \bigcap_{S \in \mathcal{S}} S = \bigcup_{S \in \mathcal{S}} (X \setminus S) \quad \text{and} \quad X \setminus \bigcup_{S \in \mathcal{S}} S = \bigcap_{S \in \mathcal{S}} (X \setminus S).$$

Proof. We take these one at a time.

- Note $a \in X \setminus \bigcap_{S \in \mathcal{S}} S$ if and only if $a \in X$ and $a \notin \bigcap_{S \in \mathcal{S}} S$. However, $a \notin \bigcap_{S \in \mathcal{S}} S$ is merely saying that a is not in all of the sets $S \in \mathcal{S}$, which is equivalent to saying $a \notin S$ for one of the $S \in \mathcal{S}$.
Thus, this is equivalent to saying $a \in X$ while $a \notin S$ for some $S \in \mathcal{S}$, which is equivalent to $a \in \bigcup_{S \in \mathcal{S}} (X \setminus S)$.
- Note $a \in X \setminus \bigcup_{S \in \mathcal{S}} S$ if and only if $a \in X$ and $a \notin \bigcup_{S \in \mathcal{S}} S$. However, $a \notin \bigcup_{S \in \mathcal{S}} S$ is merely saying that a is not in any of the sets $S \in \mathcal{S}$, which is equivalent to saying $a \notin S$ for each of the $S \in \mathcal{S}$.
Thus, this is equivalent to saying $a \in X$ while $a \notin S$ for each $S \in \mathcal{S}$, which is equivalent to $a \in \bigcap_{S \in \mathcal{S}} (X \setminus S)$. ■

1.2.2 Some Conventions

In this class, we take the following names of standard sets.

- $\mathbb{N} = \{0, 1, 2, \dots\}$ is the set of natural numbers. Importantly, $0 \in \mathbb{N}$.
- $\mathbb{N}^+ = \{1, 2, 3, \dots\}$ is the set of positive integers.
- $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$ is the set of integers.
- $\mathbb{Q} = \{p/q : p, q \in \mathbb{Z} \text{ and } q \neq 0\}$ is the set of rationals.
- \mathbb{R} is the set of real numbers. We will not specify a construction here; see any real analysis class.
- $\mathbb{R}^\times = \{x \in \mathbb{R} : x \neq 0\}$ is the nonzero real numbers.
- $\mathbb{R}^+ = \{x \in \mathbb{R} : x > 0\}$ is the positive real numbers.
- $\mathbb{R}_{\geq 0} = \{x \in \mathbb{R} : x \geq 0\}$ is the nonnegative real numbers.
- $\mathbb{R}_{\leq 0} = \{x \in \mathbb{R} : x \leq 0\}$ is the nonpositive real numbers.
- \mathbb{C} is the complex numbers.
- $\mathbb{C}^\times = \{z \in \mathbb{C} : z \neq 0\}$ is the set of nonzero complex numbers.

1.2.3 Relations

Let's review some set theory definitions.

Cartesian
product

Definition 1.11 (Cartesian product). Given two sets A and B , we define the *Cartesian product* $A \times B$ to be the set of ordered pairs (a, b) such that $a \in A$ and $b \in B$.

Binary
relation

Definition 1.12 (Binary relation). A *binary relation* on A is any subset $R \subseteq A^2 := A \times A$. We may sometimes notate $(x, y) \in R$ by xRy , read as " x is related to y ."

Example 1.13. Equality is a binary relation on any set A ; namely, it is the subset $\{(a, a) : a \in A\}$.

The best relations are equivalence relations.

Equivalence
relation

Definition 1.14 (Equivalence relation). An *equivalence relation* on A is a binary relation R satisfying the following three conditions.

- Reflexive: each $x \in A$ has $(x, x) \in R$.
- Symmetric: each $x, y \in A$ has $(x, y) \in R$ implies $(y, x) \in R$.
- Transitive: each $x, y, z \in A$ has $(x, y) \in R$ and $(y, z) \in R$ implies $(x, z) \in R$.

Equivalence relations are nice because they allow us to partition the set into "equivalence classes."

Equivalence
class

Definition 1.15 (Equivalence class). Fix A a set and $R \subseteq A^2$ an equivalence relation. Then, for given $x \in A$, we define

$$[x]_R := \{y \in A : (x, y) \in R\}$$

to be the *equivalence class* of x .

The hope is that equivalence classes partition the set. What is a partition?

Partition

Definition 1.16 (Partition). A *partition* of a set A is a collection of nonempty subsets $S \subseteq \mathcal{P}(A)$ of A such that any two distinct $S_1, S_2 \in \mathcal{S}$ are disjoint while $A = \bigcup_{S \in \mathcal{S}} S$.

And now let's manifest our hope.

Lemma 1.17. Equivalence relations are in one-to-one correspondence with partitions of A .

Proof. Given an equivalence relation R , we define the collection

$$\mathcal{S}(R) = \{[x]_R : x \in A\}.$$

We claim that $R \mapsto \mathcal{S}(R)$ is our needed bijection. We have the following checks.

- Well-defined: observe that $\mathcal{S}(R)$ does partition A : if we have $[x]_R, [y]_R \in \mathcal{S}$, then $[x]_R \cap [y]_R \neq \emptyset$ implies there is some z with $(x, z) \in R$ and $(z, y) \in R$, so $x \in [y]_R$ and then $[x]_R \subseteq [y]_R$ follows. So by symmetry, $[y]_R \subseteq [x]_R$ as well, so we finish the disjointness check.

Further, we see that

$$A = \bigcup_{x \in A} \{x\} \subseteq \bigcup_{x \in A} [x]_R \subseteq A$$

because $x \in [x]_R$, so indeed the equivalence classes cover A .

- **Injective:** suppose R_1 and R_2 have $\mathcal{S}(R_1) = \mathcal{S}(R_2)$. We show that $R_1 \subseteq R_2$, and $R_2 \subseteq R_1$ will follow by symmetry, finishing.

We notice that, for any \mathcal{S} partitioning A , being a partition, will have exactly one subset which contains x . But for $\mathcal{S}(R)$ for an equivalence relation R , we see $x \in [x]_R \in \mathcal{S}(R)$, so this equivalence class must be the one.

So because $[x]_{R_1}$ and $[x]_{R_2}$ are the only subsets of $\mathcal{S}(R_1)$ and $\mathcal{S}(R_2)$ containing x (respectively), we must have $[x]_{R_1} = [x]_{R_2}$. Thus, $(x, y) \in R_1$ implies $y \in [x]_{R_1} = [x]_{R_2}$ implies $(x, y) \in R_2$.

- **Surjective:** suppose that \mathcal{S} is a partition of A . As noted above, each $x \in A$ is a member of exactly one set $S \in \mathcal{S}$, which we call $[x]$. Then we define $R \subseteq A^2$ by $(x, y) \in R$ if and only if $y \in [x]$. One can check that this is an equivalence relation, which we will not do here in detail.¹

The point is that

$$[x]_R = \{y : (x, y) \in R\} = \{y : y \in [x]\} = [x],$$

so $\mathcal{S}(R) = \mathcal{S}$. So our mapping is surjective. ■

We continue our discussion.

Quotient set

Definition 1.18 (Quotient set). Given an equivalence relation $R \subseteq A^2$, we define the *quotient set* A/R is the set of equivalence classes of R . In other words,

$$A/R = \{[x]_R : x \in A\}.$$

Intuitively, the quotient set is the set where we have gone ahead and identified the elements which are "similar" or "related."

We would like a more concrete way to talk about equivalence classes, for which we have the following.

Representatives

Definition 1.19 (Representatives). Given an equivalence relation $R \subseteq A^2$, we say that $C \subseteq A$ is a *set of representatives of R -equivalence classes of A* if and only if C consists of exactly one element from each equivalence class in A/R .

1.2.4 Functions

To finish off, we discuss functions.

Functions

Definition 1.20 (Functions). A *function* $f : X \rightarrow Y$ is a relation $f \subseteq X \times Y$ satisfying the following.

- For each $x \in X$, there is some $y \in Y$ such that $(x, y) \in f$. Intuitively, each $x \in X$ goes somewhere.
- For each $x \in X$ and given some $y_1, y_2 \in Y$ such that $(x, y_1), (x, y_2) \in f$, then $y_1 = y_2$. Intuitively, each $x \in X$ goes to at most one place.

We will write $f(x) = y$ as notational sugar for $(x, y) \in f$. Note this equality is legal because the value y with $(x, y) \in f$ is uniquely given.

We would like to create new functions from old. Here are two ways to do this.

Restriction

Definition 1.21 (Restriction). Given a function $f : X \rightarrow Y$ and a subset $A \subseteq X$, we define

$$f|_A = \{(x, y) \in f : x \in A\} \subseteq A \times Y$$

to be a function $f|_A : A \rightarrow Y$.

¹ Note $x \in [x]$ by definition of $[x]$. If $y \in [x]$, then note $y \in [y]$ as well, so $[x] = [y]$ is forced by uniqueness, so $x \in [y]$. If $y \in [x]$ and $z \in [y]$, then again by uniqueness $[x] = [y] = [z]$, so $z \in [x]$ follows.

We will not check that $f|_A$ is actually a function; it is, roughly speaking inherited from f .

Definition 1.22. Given two functions $f : X \rightarrow Y$ and $g : Y \rightarrow Z$, we define the *composition* of f and g to be some function $g \circ f : X \rightarrow Z$ defined by

$$(g \circ f)(x) := g(f(x)).$$

Again, we will not check that this makes a function; it is.

Functions can also help create new sets.

Image

Definition 1.23 (Image). Given a function $f : X \rightarrow Y$, we define the *image* of f to be

$$\text{im } f = f(X) := \{y \in Y : \text{there is } x \in X \text{ such that } f(x) = y\}.$$

Namely, $\text{im } f$ consists of all elements hit by someone in X hit by f .

Fiber,
pre-image

Definition 1.24 (Fiber, pre-image). Given a function $f : X \rightarrow Y$ and some $y \in Y$, we define the *fiber* of f over y to be

$$f^{-1}(y) = \{x \in X : f(x) = y\} \subseteq X.$$

In general, we define the *pre-image* of a subset $A \subseteq Y$ to be

$$f^{-1}(A) := \{x \in X : f(x) \in A\} = \bigcup_{a \in A} \{x \in X : f(x) = a\} = \bigcup_{a \in A} f^{-1}(a).$$

Some functions have nicer properties than others.

Inj-, sur-,
bijective

Definition 1.25 (Inj-, sur-, bijective). Fix a function $f : X \rightarrow Y$. We have the following.

- Then f is *injective* or *one-to-one* if and only if, given $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies $x_1 = x_2$.
- Then f is *surjective* or *onto* if and only if $\text{im } f = Y$. In other words, for each $y \in Y$, there exists $x \in X$ with $f(x) = y$.
- Then f is *bijective* if and only if it is both injective and surjective.

Here is an example.

Identity

Definition 1.26 (Identity). For a given set X , the function $\text{id}_X : X \rightarrow X$ defined by $\text{id}_X(x) := x$ is called the *identity function*.

For completeness, here are the checks that id_X is bijective.

- Injective: given $x_1, x_2 \in X$, we see $\text{id}_X(x_1) = \text{id}_X(x_2)$ implies $x_1 = \text{id}_X(x_1) = \text{id}_X(x_2) = x_2$.
- Surjective: given $x \in X$, we see that $x \in \text{im } \text{id}_X$ because $x = \text{id}_X(x)$.

We leave with some lemmas, to be proven once in one's life.

Lemma 1.27. Fix a finite sets X and Y such that $\#X = \#Y$. Then a function $f : X \rightarrow Y$ is bijective if and only if it is injective or surjective.

Proof. Certainly if f is bijective, then it is both injective and surjective, so there is nothing to say.

The reverse direction is harder. We proceed by induction on $\#X = \#Y$. If $\#X = \#Y = 0$, then $X = Y = \emptyset$, and all functions $f : \emptyset \rightarrow \emptyset$ are vacuously bijective: for injective, note that any $x_1, x_2 \in \emptyset$ have $x_1 = x_2$; for surjective, note that any $x \in \emptyset$ has $f(x) = x$.

Otherwise $\#X = \#Y > 0$. We have two cases.

- Take f injective; we show f is surjective. In this case, $\#X > 0$, so choose some $a \in X$. Note that $x \in X$ with $x \neq a$ will have $f(x) \neq f(a)$ by injectivity, so we may define the restriction

$$f|_{X \setminus \{a\}} : X \setminus \{a\} \rightarrow Y \setminus \{f(a)\}.$$

Observe that $f|_{X \setminus \{a\}}$ is injective because f is: if $x_1, x_2 \in X \setminus \{a\}$ have

$$f(x_1) = f|_{X \setminus \{a\}}(x_1) = f|_{X \setminus \{a\}}(x_2) = f(x_2),$$

then $x_1 = x_2$ follows.

Now, $\#(X \setminus \{a\}) = \#(Y \setminus \{f(a)\}) = \#X - 1$, so by induction $f|_{X \setminus \{a\}}$ will be bijective because it is injective. In particular, f by way of $f|_{X \setminus \{a\}}$ fully hits $Y \setminus \{f(a)\}$ in its image, so because $f(a) \in \text{im } f$ as well, we conclude $\text{im } f = Y$. So f is surjective.

- Take f surjective; we show f is injective. Define a function $g : Y \rightarrow X$ as follows: for each $y \in Y$, the surjectivity of f promises some $x \in X$ such that $f(x) = y$, so choose any such x and define $g(y) := x$.² Observe that $f(g(y)) = y$ by construction.

Now, we notice that g is injective: if $y_1, y_2 \in Y$ have $g(y_1) = g(y_2)$, then $y_1 = f(g(y_1)) = f(g(y_2)) = y_2$. So the previous case tells us that g is in fact bijective.

So now choose any $x_1, x_2 \in X$ such that $f(x_1) = f(x_2)$. The surjectivity of f promises some $y_1, y_2 \in Y$ such that $g(y_1) = x_1$ and $g(y_2) = x_2$, so we see that

$$x_1 = g(y_1) = g(f(g(y_1))) = g(f(x_1)) = g(f(x_2)) = g(f(g(y_2))) = g(y_2) = x_2,$$

proving our injectivity. ■

Lemma 1.28. Fix $f : X \rightarrow Y$ a bijective function. Then there is a unique function $g : Y \rightarrow X$ such that $f \circ g = \text{id}_Y$ and $g \circ f = \text{id}_X$.

Proof. We show existence and uniqueness separately.

- We show existence. Note that, because $f : X \rightarrow Y$ is surjective, each $y \in Y$ has some $x \in X$ such that $f(x) = y$. In fact, this $x \in X$ is uniquely defined because $f(x_1) = f(x_2)$ implies $x_1 = x_2$, so we may define $g(y)$ as the value x for which $f(x) = y$.

By construction, $f(g(y)) = y$, so $f \circ g = \text{id}_Y$. Additionally, we note that, given any $x \in X$, the value x_0 for which $f(x) = f(x_0)$ is $x = x_0$ by the injectivity, so $g(f(x)) = x$. Thus, $g \circ f = \text{id}_X$, as claimed.

- We show uniqueness. Suppose that we have two functions $g_1, g_2 : Y \rightarrow X$ which satisfy

$$f \circ g_1 = f \circ g_2 = \text{id}_Y \quad \text{and} \quad g_1 \circ f = g_2 \circ f = \text{id}_X.$$

Then we see that

$$g_1 = g_1 \circ \text{id}_Y = g_1 \circ (f \circ g_2) = (g_1 \circ f) \circ g_2 = \text{id}_X \circ g_2 = g_2,$$

where we have used the fact that function composition associates. This finishes. ■

1.3 January 24

Good morning everyone.

² Technically we are using the Axiom of Choice here. One can remove this with an induction because all sets are finite, but I won't bother.

1.3.1 Algebraic Structure

Today we are reviewing the complex numbers (reportedly, “some basics”). Or at least it is hopefully mostly review. Here is our main character this semester.

Complex
numbers

Definition 1.29 (Complex numbers). The set \mathbb{C} of *complex numbers* is

$$\mathbb{C} := \{a + bi : a, b \in \mathbb{R}\}.$$

Here i is some symbol such that $i^2 = -1$ formally.

In particular, two complex numbers $a_1 + b_1i$ and $a_2 + b_2i$ are equal if and only if $a_1 = a_2$ and $b_1 = b_2$.
The complex numbers also have some algebraic structure.

+ and \times in \mathbb{C}

Definition 1.30 (+ and \times in \mathbb{C}). Given complex numbers $a_1 + b_1i, a_2 + b_2i \in \mathbb{C}$, we define

$$(a_1 + b_1i) + (a_2 + b_2i) = (a_1 + a_2) + (b_1 + b_2)i,$$

and

$$(a_1 + b_1i)(a_2 + b_2i) = (a_1a_2 - b_1b_2) + (a_1b_2 + a_2b_1)i,$$

defined essentially by direct expansion, upon recalling $i^2 = -1$.

Here is the corresponding algebraic structure.

Proposition 1.31. The set \mathbb{C} with the above operations is a two-dimensional \mathbb{R} -vector space with basis $\{1, i\}$.

Proof. The elements $\{1, i\}$ span \mathbb{C} because all complex numbers in \mathbb{C} can be written as $a + bi = a \cdot 1 + b \cdot i$ by definition.

To see that these elements are linearly independent, suppose $a + bi = 0$. If $b = 0$, then $a = 0$ follows, and we are done. Otherwise, take $b \neq 0$, but then we see $(-a/b) = i$, so

$$(-a/b)^2 = -1 < 0,$$

which does not make sense for real numbers. This finishes. ■

Proposition 1.32. The set \mathbb{C} with the above operations is a field.

Proof. We have the following checks.

- The element $0 + 0i$ is our additive identity. Indeed, one can check that $(0 + 0i) + (a + bi) = (a + bi) + (0 + 0i) = a + bi$.
- The element $1 + 0i$ is our multiplicative identity. Indeed, one can check that $(1 + 0i)(a + bi) = (a + bi)(1 + 0i) = a + bi$.
- Commutativity of addition and multiplication follow from by expansion.
- The distributive laws can again be checked by expansion.
- The additive inverse of $a + bi$ is $(-a) + (-b)i$.
- The multiplicative inverse of $a + bi$ can be found by wishing really hard and writing

$$\frac{1}{a + bi} = \frac{1}{a + bi} \cdot \frac{a - bi}{a - bi} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Then one can check this works. ■

Sometimes we would like to extract our coefficients from our basis.

Re and Im

Definition 1.33 (Re and Im). Given $z := a + bi \in \mathbb{C}$, we define the operations

$$\operatorname{Re} z := a \quad \text{and} \quad \operatorname{Im} z := b.$$

Importantly, $\operatorname{Re} : \mathbb{C} \rightarrow \mathbb{R}$ and $\operatorname{Im} : \mathbb{C} \rightarrow \mathbb{R}$.

Because we are merely doing basis extraction, it makes sense that these operations will preserve some (additive) structure.

Proposition 1.34. Fix $z = a + bi$ and $w = c + di$. Then the following.

- (a) $\operatorname{Re}(z + w) = \operatorname{Re} z + \operatorname{Re} w$.
- (b) $\operatorname{Im}(z + w) = \operatorname{Im} z + \operatorname{Im} w$.

Proof. We proceed by direct expansion. Observe

$$\operatorname{Re}(z + w) = \operatorname{Re}((a + c) + (b + d)i) = a + c = \operatorname{Re} z + \operatorname{Re} w,$$

and

$$\operatorname{Im}(z + w) = \operatorname{Im}((a + c) + (b + d)i) = b + d = \operatorname{Im} z + \operatorname{Im} w.$$

This finishes. ■

It also turns out that the complex numbers have a very special transformation.

Conjugate

Definition 1.35 (Conjugate). Given $z := a + bi \in \mathbb{C}$, we define the *complex conjugate* to be $\bar{z} := a - bi \in \mathbb{C}$.

We promised conjugation would be special, so here are some special things.

Proposition 1.36. Fix $z = a + bi \in \mathbb{C}$. Then the following.

- (a) $z + \bar{z} = 2 \operatorname{Re} z$.
- (b) $z - \bar{z} = 2i \operatorname{Im} z$.
- (c) $\overline{\bar{z}} = z$.

Proof. We take these one at a time.

(a) Write $a + bi + \overline{a + bi} = a + bi + a - bi = 2a$.

(b) Write $a + bi - \overline{a + bi} = a + bi - (a - bi) = 2bi$.

(c) Write $\overline{\overline{a + bi}} = \overline{a - bi} = a + bi$. ■

In fact, more is true.

Proposition 1.37. Fix $z = a + bi \in \mathbb{C}$ and $w = c + di \in \mathbb{C}$. Then the following.

- (a) $\overline{z + w} = \bar{z} + \bar{w}$.
- (b) $\overline{z\bar{w}} = \bar{z} \cdot \bar{w}$.

Proof. We take these one at a time.

- Write

$$\overline{z + w} = (a + c) - (b + d)i = (a - bi) + (c - di) = \bar{z} + \bar{w}.$$

- Write

$$\begin{aligned}\bar{z} \cdot \bar{w} &= (a - bi)(c - di) \\ &= (ac - bd) - (ad + bc)i \\ &= \overline{(ac - bd) + (ad + bc)i} \\ &= \overline{zw}.\end{aligned}$$

This finishes. ■

1.3.2 Defining Distance

Complex conjugation actually gives rise to a notion of size.

Norm on \mathbb{C}

Definition 1.38 (Norm on \mathbb{C}). Given $z := a + bi$, we define the *norm function on \mathbb{C}* by

$$|z| := \sqrt{a^2 + b^2}.$$


Size actually gives distance.

Distance on \mathbb{C}

Definition 1.39 (Distance on \mathbb{C}). Given complex numbers $z = a + bi$ and $w = c + di$, we define the *distance between z and w* to be

$$|z - w| = \sqrt{(a - c)^2 + (b - d)^2}.$$

Here are some examples.



One can ask what is the distance between $0 + 0i$ and $1 + 1i$, and we can compute directly that this is $\sqrt{1 + 1} = \sqrt{2}$. Similarly, the distance between $1 + 2i$ and $1 + i$ is $|(1 + 2i) - (1 + i)| = |i| = 1$. It should agree with our geometric intuition.

We mentioned complex conjugation is involved here, so we have the following lemma.

Lemma 1.40. Fix $z, w \in \mathbb{C}$. The following are true.

- (a) $|z|^2 = z\bar{z}$.
- (b) $|\operatorname{Re} z| \leq |z|$ and $|\operatorname{Im} z| \leq |z|$.
- (c) $|z| = |\bar{z}| = |-z|$.
- (d) $|z| = 0$ if and only if $z = 0$.
- (e) $|zw| = |z| \cdot |w|$.

Proof. We take these one at a time. Set $z = a + bi$.

(a) We have

$$|z|^2 = a^2 + b^2 = (a + bi)(a - bi) = z\bar{z}.$$

Here we have used subtraction of two squares, which one can see when writing $a^2 + b^2 = a^2 - (ib)^2$.

(b) We have $a^2 \leq a^2 + b^2$ and $b^2 \leq a^2 + b^2$ by the Trivial inequality, so

$$|\operatorname{Re} z| = |a| \leq \sqrt{a^2 + b^2} = |z|,$$

and similarly,

$$|\operatorname{Im} z| = |b| \leq \sqrt{a^2 + b^2} = |z|.$$

(c) Note

$$|\bar{z}| = |a - bi| = \sqrt{a^2 + (-b)^2} = \sqrt{a^2 + b^2} = |z|,$$

and

$$|-z| = |-a - bi| = \sqrt{(-a)^2 + (-b)^2} = \sqrt{a^2 + b^2} = |z|.$$

(d) From (b), we know that $|\operatorname{Re} z|, |\operatorname{Im} z| \leq |z|$, but $|z| = 0$ then forces $\operatorname{Re} z = \operatorname{Im} z = 0$, so $z = 0$.

(e) From (a), we can write $|zw|^2 = zw \cdot \overline{zw}$, which will expand out into

$$z \cdot w \cdot \bar{z} \cdot \bar{w}.$$

We can collect this into $z\bar{z} \cdot w\bar{w} = |z|^2 |w|^2$. Thus, by (a) again, $|zw|^2 = |z|^2 |w|^2$. But because all norms must be nonnegative real numbers, we may take square roots to conclude $|zw| = |z| \cdot |w|$. ■

Remark 1.41. Norms are actually more general constructions. For example, the requirement $|zw| = |z| \cdot |w|$ makes $|\cdot|$ into a “multiplicative” norm.

To finish off, we actually show that our distance function is good: we show the triangle inequality.

Lemma 1.42 (Triangle inequality). For every $x, y, z \in \mathbb{C}$, we claim

$$|z - x| \leq |z - y| + |y - x|.$$

This claim should be familiar from real analysis. Intuitively, it means that travelling between z and x cannot be made into a shorter trip by taking a detour to some other point y first.

Proof. Let $a := z - y$ and $b := y - x$ so that $a + b = z - x$. Thus, we are showing that

$$|a + b| \stackrel{?}{\leq} |a| + |b|,$$

which is nicer because it only has two letters. For this, because everything is a nonnegative real numbers, it suffices to show the square of this requirement; i.e., we show

$$(|a| + |b|)^2 - |a + b|^2 \stackrel{?}{\geq} 0.$$

Fully expanding, it suffices to show

$$|a|^2 + |b|^2 + 2|a| \cdot |b| - |a + b|^2 \stackrel{?}{\geq} 0.$$

Expanding out $|w|^2 = w\bar{w}$ for $w \in \mathbb{C}$, we are showing

$$a\bar{a} + b\bar{b} + 2|a| \cdot |b| - (a + b)(\bar{a} + \bar{b}) \stackrel{?}{\geq} 0.$$

This is nice because the expansion of the rightmost term will induce some cancellation: it expands into $a\bar{a} + a\bar{b} + \bar{a}b + b\bar{b}$, so we are left with showing

$$2|a| \cdot |b| - (a\bar{b} + b\bar{a}) \stackrel{?}{\geq} 0.$$

Note that $\bar{a}b = \overline{ab}$, so we can collect the final term as $2\operatorname{Re}(a\bar{b})$. Similarly, we can write $|a| \cdot |b| = |a| \cdot |\bar{b}| = |a\bar{b}|$, so we are showing

$$2|a\bar{b}| - 2\operatorname{Re}(a\bar{b}) \geq 0,$$

which is true because the real part does not exceed the norm. This finishes. ■

1.4 January 26

In-person class should start on Monday. Homework #2 will be released on Friday.

1.4.1 Geometry on \mathbb{C}


So let's try to build a topology on \mathbb{C} today. We pick up the following definition.

Convex


Definition 1.43 (Convex). A subset $X \subseteq \mathbb{C}$ is *convex* if and only if, for every $z, w \in X$ and $t \in [0, 1]$, we have that $w + t(z - w) \in X$.

Intuitively, "convex" means that X contains the line segment of any two points in X .

Example 1.44. The circle is convex: any line with endpoints in the circle lives in the circle.



Non-Example 1.45. The star-shape is not convex: the given line goes outside the star.



To define our open sets, we will define balls first.

Open ball

Definition 1.46 (Open ball). Given some $z_0 \in \mathbb{C}$, then *open ball* centered at z_0 with radius $r > 0$ is

$$B(z_0, r) := \{z \in \mathbb{C} : |z - z_0| < r\}.$$

Observe $z_0 \in B(z_0, r)$.

Open balls let us define all sorts of properties.

Isolated

Definition 1.47 (Isolated). Fix $X \subseteq \mathbb{C}$. A point $z \in X$ is isolated in X if and only if there exists $r > 0$ such that

$$B(z, r) \cap X = \{z\}.$$


Discrete

Definition 1.48 (Discrete). A subset $X \subseteq \mathbb{C}$ is *discrete* if and only if every point is isolated.**Example 1.49.** Any finite subset of $X \subseteq \mathbb{C}$ is discrete. Namely, any point $z \in X$ can take

$$r = \frac{1}{2} \min_{w \in X \setminus \{z\}} |z - w|.$$

Example 1.50. The subset $\mathbb{Z} \subseteq \mathbb{C}$ is isolated. Namely, take $r = \frac{1}{2}$ for any given point.

Bounded

Definition 1.51 (Bounded). A subset $X \subseteq \mathbb{C}$ is *bounded* if and only if there is an M such that $X \subseteq B(0, M)$.**Example 1.52.** The star from earlier fits into a large circle and is therefore bounded.

And here is our fundamental definition for our topology.

Open

Definition 1.53 (Open). A subset $X \subseteq \mathbb{C}$ is *open* if and only if, for each $z \in X$, there exists $r > 0$ such that $B(z, r) \subseteq X$.**Remark 1.54** (Nir). We should probably show that open balls are open; let $B(z, r)$ be an open ball. Well, for any $w \in B(z, r)$, set $r_w := r - |z - w|$, which is positive because $w \in B(z, r)$ requires $|z - w| < r$. Now, $w' \in B(w, r_w)$ implies that $|w - w'| < r - |z - w|$, so by the triangle inequality,

$$|z - w'| \leq |z - w| + |w - w'| < r,$$

so $w' \in B(z, r)$ follows. So indeed, each $w \in B(z, r)$ has $B(w, r_w) \subseteq B(z, r)$.

Open lets us define closed.

Closed

Definition 1.55 (Closed). A subset $X \subseteq \mathbb{C}$ is *closed* if and only if $\mathbb{C} \setminus X$ is open.



Warning 1.56. Sets are not doors: a set can be both open and closed.

1.4.2 Unions and Intesections

Here are some basic properties of our topology.

Lemma 1.57. The subsets \emptyset and \mathbb{C} are open and closed in \mathbb{C} .

Proof. It suffices to show that \emptyset and \mathbb{C} are both open, by definition of closed. That \emptyset is open holds vacuously because one cannot find any $z \in \emptyset$ anyways. That \mathbb{C} is open holds because open balls are subsets of \mathbb{C} , so any $z \in \mathbb{C}$ can take $r = 1$ so that

$$B(z, r) \subseteq \mathbb{C}.$$

So we are done. ■

Lemma 1.58. Fixing some $z \in \mathbb{C}$, the set $\{z\}$ is closed.

Proof. We show that $U := \mathbb{C} \setminus \{z\}$ is open. Well, fix any $w \in U$, and because $w \neq z$, we note $|z - w| > 0$, so we set $r := \frac{1}{2}|z - w|$. It follows that

$$z \notin B(w, r)$$

because $|z - w| > r$. But this is equivalent to $B(w, r) \subseteq \mathbb{C} \setminus \{z\} = U$, so we are done. ■

We would like to make new open and closed subsets from old ones. Here is one way to do so.

Lemma 1.59. The following are true.

- (a) Arbitrary union: if \mathcal{U} is any collection of open subsets of \mathbb{C} , then the union $\bigcup_{U \in \mathcal{U}} U$ is also open.
- (b) Arbitrary intersection: if \mathcal{V} is any collection of closed subsets of \mathbb{C} , then intersection $\bigcap_{V \in \mathcal{V}} V$ is also closed.

Proof. We take these one at a time.

- (a) Fix $z \in \bigcup_{U \in \mathcal{U}} U$. We need to show there is some $r > 0$ such that

$$B(z, r) \stackrel{?}{\subseteq} \bigcup_{U \in \mathcal{U}} U.$$

Well, we know there must be some $U_z \in \mathcal{U}$ such that $z \in U_z$ by definition of the union. But now U_z is open, and therefore we are promised an $r > 0$ such that

$$B(z, r) \subseteq U_z \subseteq \bigcup_{U \in \mathcal{U}} U,$$

so we are done.

(b) Fix \mathcal{V} a collection of closed subsets of \mathbb{C} . We want to show that

$$\mathbb{C} \setminus \bigcap_{V \in \mathcal{V}} V$$

is open, which by de Morgan's law is equivalent to

$$\bigcup_{V \in \mathcal{V}} (\mathbb{C} \setminus V)$$

being open. However, each $V \in \mathcal{V}$ is closed, so $\mathbb{C} \setminus V$ will be open, so we are done by (a). ■

Lemma 1.60. The following are true.

- (a) Finite intersection: if $\{U_k\}_{k=1}^n$ is a finite collection of open subsets of \mathbb{C} , then the intersection $\bigcap_{k=1}^n U_k$ is also open.
- (b) Finite union: if $\{V_k\}_{k=1}^n$ is a finite collection of closed subsets of \mathbb{C} , then $\bigcup_{k=1}^n V_k$ is also closed.

Proof. We take these one at a time.

(a) Fix $z \in \bigcap_{k=1}^n U_k$ so that we need to find $r > 0$ such that

$$B(z, r) \subseteq \bigcup_{k=1}^n U_k.$$

Well, $z \in U_k$ for each k , and each U_k is open, so there is an $r_k > 0$ such that $B(z, r_k) \subseteq U_k$. Thus, we set $r := \min_k \{r_k\}$; because there are only finitely many r_k , we are assured that $r > 0$. Now, we observe that

$$B(z, r) \subseteq B(z, r_k) \subseteq U_k.$$

(Explicitly, $|w - z| < r$ implies $|w - z| < r_k$ because $r \leq r_k$.) Thus, it follows that

$$B(z, r) \subseteq \bigcap_{k=1}^n U_k,$$

as desired.

(b) We use de Morgan's laws. We want to show that

$$\mathbb{C} \setminus \bigcup_{k=1}^n V_k$$

is open, which by de Morgan's laws is the same thing as showing that

$$\bigcap_{k=1}^n (\mathbb{C} \setminus V_k)$$

is open. However, each $\mathbb{C} \setminus V_k$ is open by hypothesis on the V_k , so the full intersection is open by (a). This finishes. ■

Remark 1.61. The finiteness is in fact necessary. For example,

$$\bigcap_{n \in \mathbb{N}} B(0, 1/n) = \{0\}.$$

Then one can check that each open ball is open while singletons in \mathbb{C} are not.

1.4.3 Interior, Closure

Let's see more definitions.

Interior

Definition 1.62 (Interior). Given a subset $X \subseteq \mathbb{C}$, we define the *interior* X° of X to be the union of all open sets contained in X (which will be open by Lemma 1.59).

Remark 1.63. In fact, X° is the largest open subset of X , for any open subset $U_0 \subseteq \mathbb{C}$ contained in X will have

$$U_0 \subseteq \bigcup_{\text{open } U \subseteq X} U = X^\circ.$$

It follows X is open if and only if $X = X^\circ$: if $X = X^\circ$, then X is open because X° is open; if X is open, then X is the largest open subset of \mathbb{C} contained in X , so $X = X^\circ$.

Closure

Definition 1.64 (Closure). Given a subset $X \subseteq \mathbb{C}$, we define the *closure* \overline{X} of X to be the intersection of all closed sets containing X (which will be closed by Lemma 1.59).

Remark 1.65. In fact, \overline{X} is the smallest closed set containing X , for any closed subset $V_0 \subseteq \mathbb{C}$ containing X will have

$$V_0 \supseteq \bigcap_{\text{open } V \supseteq X} V = \overline{X}.$$

It follows X is closed if and only if $X = \overline{X}$: if $X = \overline{X}$, then X is closed because \overline{X} is closed; if X is closed, then X is the smallest closed subset of \mathbb{C} containing X , so $X = \overline{X}$.

By the above definitions, it is not too hard to see that $X^\circ \subseteq X \subseteq \overline{X}$.

The interior and closure also let us define the boundary.

Frontier,
boundary

Definition 1.66 (Frontier, boundary). Given a subset $X \subseteq \mathbb{C}$, we define the *frontier* or *boundary* ∂X of X to be $\overline{X} \setminus X^\circ$.

1.4.4 Connectivity

Discon-
nected

Definition 1.67 (Disconnected). A subset $X \subseteq \mathbb{C}$ is *disconnected* if and only if there exists nonempty disjoint open subsets U_1 and U_2 such that $X \subseteq U_1 \cup U_2$ and $X \cap U_1, X \cap U_2 \neq \emptyset$. (In other words, the subspace of $X \subseteq \mathbb{C}$ is (topologically) disconnected.) In this case, we say that U_1 and U_2 *disconnect* X .

Lastly, we say X is *connected* if and only if it is not disconnected.

Example 1.68. The set \emptyset is connected because it is impossible for $U \cap \emptyset \neq \emptyset$ for any open set U of \mathbb{C} .

Example 1.69. Any singleton $\{z\}$ is connected. In fact, one cannot decompose $\{x\}$ into two disjoint sets at all, much less into disjoint sets of the form $U \cap \{x\}$ with U open.

Example 1.70. Any open ball $B(z, r)$ is connected. This is surprisingly annoying to check.

Example 1.71. The set $\{1, 2\}$ is disconnected by $U_1 = B(1, 1/2)$ and $U_2 = B(2, 1/2)$.

Connectivity plays nicely with the rest of our definitions as well.

Lemma 1.72. A given subset $X \subseteq \mathbb{C}$ is connected if and only if the only subsets of X which are both open and closed (in the subspace topology) are \emptyset and X .

Proof. We take the directions independently. For the forwards direction, take X connected, and suppose that $U \subseteq X$ is open and closed. In this subspace topology, we get that $X \setminus U$ will also be open, and then the subsets U and $X \setminus U$ are both open, disjoint and have

$$X = U \cup (X \setminus U).$$

Thus, we require $U = \emptyset$ or $X \setminus U = \emptyset$, so $U \in \{\emptyset, X\}$.

We leave the reverse direction as an exercise. Suppose that X is disconnected, and we show that there is a nonempty proper closed and open subset of X . Well, because X is disconnected, we have disjoint open sets U_1 and U_2 of \mathbb{C} such that $X \cap U_1, X \cap U_2 \neq \emptyset$ and $X \subseteq U_1 \cup U_2$. It follows that

$$X = (U_1 \cap X) \cup (U_2 \cap X). \quad (*)$$

However, now consider the open subset $U := U_1 \cap X$ of X . We note that $(U_1 \cap X) \cap (U_2 \cap X) = \emptyset$, so by $(*)$ we see that $U_1 \cap X = X \setminus (U_2 \cap X)$, so $U_1 \cap X$ is closed as well.

To finish, we note that $U \neq \emptyset$ is nonempty, and its complement is $X \setminus U = U_2 \cap X$ is also nonempty, so $U \neq X$ is proper. Thus, U is a proper nonempty closed and open subset of X . This finishes. ■

Remark 1.73 (Nir). It is actually important that the open subsets in the above lemma are in the subspace topology and are not required to be \mathbb{C} -open. For example, $X = \{1, 2\}$ is disconnected, but it has no nonempty \mathbb{C} -open subsets to witness this.

Lemma 1.74. Fix \mathcal{S} a collection of connected subsets of \mathbb{C} . If $\bigcap_{S \in \mathcal{S}} S$ is nonempty, then $\bigcup_{S \in \mathcal{S}} S$ will be connected.

Proof. Suppose $\bigcup_{S \in \mathcal{S}} S$ is contained in the disjoint open subsets U_1 and U_2 of \mathbb{C} ; we claim $U_1 \cap (\bigcup_{S \in \mathcal{S}} S) = \emptyset$ or $U_2 \cap (\bigcup_{S \in \mathcal{S}} S) = \emptyset$, which will finish.

Pick up some

$$z \in \bigcap_{S \in \mathcal{S}} S,$$

which exists because the intersection is nonempty. Without loss of generality, we may assume that $z \in U_1$.

Now, $z \in S$ for each $S \in \mathcal{S}$, so we see $U_1 \cap S \neq \emptyset$, so because $(U_1 \cap S) \cup (U_2 \cap S) = S$, we see that $U_2 \cap S = \emptyset$ by hypothesis on S 's connectivity. Thus, unioning over the $U_2 \cap S = \emptyset$,

$$U_2 \cap \left(\bigcup_{S \in \mathcal{S}} S \right) = \emptyset,$$

which finishes the proof. ■

Remark 1.75. The condition with nonempty intersection is necessary: $\{0\}$ and $\{1\}$ are connected, but $\{0\} \cup \{1\}$ is not.

1.5 January 28

Hopefully we'll be in-person on Monday. Homework 2 will be released later today, due next Friday.

1.5.1 Sequences

Today we're talking about sequences, building towards a theory of sequences and series. Next week we will begin studying holomorphic functions and actually doing complex analysis.

Anyways, here is a series of definitions.

Sequence

Definition 1.76 (Sequence). A sequence of complex numbers is a function $f : \mathbb{N} \rightarrow \mathbb{C}$. Often we will notate this by $\{z_n\}_{n \in \mathbb{N}}$ where $z_n := f(n)$.

By convention, all of our sequences will be sequences of complex numbers unless otherwise stated.

Subsequence

Definition 1.77 (Subsequence). A sequence $\{w_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ is a *subsequence* of a sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ if and only if there is some strictly increasing function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $w_n = z_{g(n)}$.

Bounded

Definition 1.78 (Bounded). A sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ is *bounded* if and only if there exists a positive real number $M > 0$ such that

$$\{z_n\}_{n \in \mathbb{N}} \subseteq B(0, M).$$

In other words, $|z_n| < M$ for each $n \in \mathbb{N}$.

We are in particular interested in convergence in analysis.

Converges

Definition 1.79 (Converges). A sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ *converges* to some $z \in \mathbb{C}$ if and only if, for each $\varepsilon > 0$, there exists some N such that $n > N$ implies

$$|z - z_n| < \varepsilon.$$

We will notate this by $z_n \rightarrow z$ or $\lim_{n \rightarrow \infty} z_n = z$.

Note that the definition of the limit above says that

$$\lim_{n \rightarrow \infty} z_n = z \iff \lim_{n \rightarrow \infty} |z_n - z| = 0.$$

Intuitively, the distance between the z_n and the z has to "narrow in" on z .

We would like some real-analytic tools for our complex analysis. Here is a convergence lemma.

Lemma 1.80. Fix $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ a sequence. Then, letting $z_n := x_n + y_n i$, we have that $z_n \rightarrow z$ where $z = x + yi$ if and only if $x_n \rightarrow x$ and $y_n \rightarrow y$.

Proof. This is essentially by definition of the metric on \mathbb{C} . We take the directions one at a time.

- Suppose that $z_n \rightarrow z$ in \mathbb{C} . Then we claim that $\operatorname{Re} z_n \rightarrow \operatorname{Re} z$ and $\operatorname{Im} z_n \rightarrow \operatorname{Im} z$ in \mathbb{R} . Indeed, for any $\varepsilon > 0$, there is N such that

$$n > N \implies |z - z_n| < \varepsilon.$$

But now we see that $|\operatorname{Re} z_n - \operatorname{Re} z|, |\operatorname{Im} z_n - \operatorname{Im} z| \leq \sqrt{(\operatorname{Re} z_n - \operatorname{Re} z)^2 + (\operatorname{Im} z_n - \operatorname{Im} z)^2}$, so it follows

$$n > N \implies |\operatorname{Re} z_n - \operatorname{Re} z|, |\operatorname{Im} z_n - \operatorname{Im} z| < \varepsilon,$$

finishing.

- Suppose that $\operatorname{Re} z_n \rightarrow x$ and $\operatorname{Im} z_n \rightarrow y$. We claim that $z_n \rightarrow x + yi$. Indeed, for any $\varepsilon > 0$, there exists N_x such that

$$n > N_x \implies |\operatorname{Re} z_n - x| < \varepsilon/2$$

and N_y such that

$$n > N_y \implies |\operatorname{Im} z_n - y| < \varepsilon/2.$$

It follows that

$$n > \max\{N_x, N_y\} \implies |z_n - (x + yi)| = \sqrt{|\operatorname{Re} z_n - x|^2 + |\operatorname{Im} z_n - y|^2} \leq \sqrt{\left(\frac{\varepsilon}{2}\right)^2 + \left(\frac{\varepsilon}{2}\right)^2} < \varepsilon.$$

This finishes. ■

Essentially, this means that checking convergence of complex numbers is the same as checking real and imaginary parts individually, so we can turn convergence questions into ones from real analysis.

We also have the following basic properties about convergence.

Proposition 1.81. Fix $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ a convergent sequence. The following are true.

- (a) $\{z_n\}_{n \in \mathbb{N}}$ is bounded.
- (b) The limit of $\{z_n\}_{n \in \mathbb{N}}$ is unique.
- (c) Every subsequence of $\{z_n\}_{n \in \mathbb{N}}$ converges to z .

Proof. We take the claims one at a time. Let $z \in \mathbb{C}$ be so that $z_n \rightarrow z$.

- (a) Fix $\varepsilon = 1$ so that there exists N so that $n > N$ implies $|z_n - z| < 1$. Now set

$$M := \max(\{|z_n| + 1 : n \leq N\} \cup \{|z| + 1\}).$$

We claim that $|z_n| < M$ for each $n \in \mathbb{N}$. We have two cases.

- If $n \leq N$, then $|z_n| < |z_n| + 1 \leq M$.
- Otherwise $n > N$ so that

$$|z_n| \leq |z_n - z| + |z| < |z| + 1 \leq M,$$

so we are done.

- (b) Suppose that $z_n \rightarrow z'$ for some $z' \in \mathbb{C}$, and we show $z = z'$. Indeed, if $z = z'$, then we are done, so suppose that $z \neq z'$ so that $|z - z'| \neq 0$. Then we set $\varepsilon := \frac{1}{2}|z - z'| > 0$, and we are promised some N, N' such that

$$n > N \implies |z - z_n| < \frac{\varepsilon}{2} \quad \text{and} \quad n > N' \implies |z' - z_n| < \frac{\varepsilon}{2}.$$

In particular, we see that, for $n > \max\{N, N'\}$, we have

$$|z - z'| \leq |z - z_n| + |z_n - z'| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon = \frac{1}{2}|z - z'|.$$

But because $0 \leq |z - z'|$, we see that this forces $|z - z'| = 0$, so $z = z'$ follows. (Technically we have hit contradiction, but we do not need to use this.)

- (c) Note that subsequences can be characterized by choosing a strictly increasing function $f : \mathbb{N} \rightarrow \mathbb{N}$ so that we want to show $z_{f(n)} \rightarrow z$. Indeed, for any $\varepsilon > 0$, we are promised some N so that

$$n > N \implies |z - z_n| < \varepsilon.$$

Now, for each $n \in \mathbb{N}$, we have³ $f(n) \geq n$, so we see that

$$n > N \implies f(n) > N \implies |z - z_{f(n)}| < \varepsilon,$$

which finishes. ■

Sequences themselves have an arithmetic.

³ We can show this by induction on n , for $f(0) \geq 0$ and $f(n+1) > f(n)$ forces $f(n+1) \geq f(n) + 1$.

Proposition 1.82. Fix $\{z_n\}_{n \in \mathbb{N}}, \{w_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ sequences such that $z_n \rightarrow z$ and $w_n \rightarrow w$. Then the following hold.

- (a) $z_n + w_n \rightarrow z + w$.
- (b) $z_n w_n \rightarrow zw$.
- (c) If $w \neq 0$ and $w_n \neq 0$ for each $n \in \mathbb{N}$, then $\frac{1}{w_n} \rightarrow \frac{1}{w}$.

Proof. We take these one at a time, essentially borrowing the proof from metric spaces.

- (a) Fix some $\varepsilon > 0$. We can find some N_z such that

$$n > N_z \implies |z - z_n| < \varepsilon/2$$

and some N_w such that

$$n > N_w \implies |w - w_n| < \varepsilon/2.$$

Now, taking $N := \max\{N_z, N_w\}$ so that the triangle inequality gives

$$n > N \implies |(z + w) - (z_n + w_n)| \leq |z - z_n| + |w - w_n| < \varepsilon,$$

which finishes.

- (b) We have to use the fact that the sequences are bounded here. Because $w_n \rightarrow w$, the sequence is bounded, so there is an M so that $|w_n| < M$ for each $n \in \mathbb{N}$. Now, the key inequality is that

$$|z_n w_n - zw| \leq |z_n w_n - zw_n| + |zw_n - zw| \leq M|z_n - z| + |z| \cdot |w_n - w|. \quad (*)$$

So with this in mind, fix any $\varepsilon > 0$, and we see that we are promised N_z such that

$$n > N_z \implies |z_n - z| < \frac{\varepsilon}{2M}$$

and some N_w such that

$$n > N_w \implies |w_n - w| < \frac{\varepsilon}{2|z|}$$

so that (*) implies

$$n > \max\{N_z, N_w\} \implies |z_n w_n - zw| < \varepsilon,$$

finishing.

- (c) We begin with some motivating arithmetic. Observe that

$$\left| \frac{1}{w} - \frac{1}{w_n} \right| = \frac{|w_n - w|}{|w w_n|}.$$

We can upper-bound the numerator without tears, so we see the main difficulty is lower-bounding the denominator. Well, because $w \neq 0$, we can set $\varepsilon = |w|/2$ so that there exists N_0 such that

$$n > N_0 \implies |w_n - w| < |w|/2.$$

In particular, it follows that $|w_n| \geq |w| - |w - w_n| = |w|/2$ for $n > N_0$.

With this in mind, fix any $\varepsilon > 0$. Then we are promised some N_1 such that

$$n > N_1 \implies |w_n - w| < |w|^2 \varepsilon / 2$$

so that we see

$$n > \max\{N_0, N_1\} \implies \left| \frac{1}{w} - \frac{1}{w_n} \right| = \frac{|w_n - w|}{|w| \cdot |w_n|} \leq \frac{|w|^2 \varepsilon / 2}{|w| \cdot |w|/2} = \varepsilon,$$

finishing. ■

1.5.2 Limit Points

Here is our main character.

Limit point

Definition 1.83 (Limit point). Fix $X \subseteq \mathbb{C}$ and some $z \in \mathbb{C}$. Then we say that z is a *limit point* if and only if there exists some sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ such that $z_n \rightarrow z$.

Accumulation point

Definition 1.84 (Accumulation point). Fix $X \subseteq \mathbb{C}$ and some $z \in \mathbb{C}$. Then we say that z is an *accumulation point* if and only if there exists some sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq X \setminus \{z\}$ such that $z_n \rightarrow z$.

Essentially accumulation points do not allow isolated points while limit points do.

The above essentially gives a more directly topological definition of “closed set.” It also gives us a more directly topological definition of the closure.

Lemma 1.85. Fix $X \subseteq \mathbb{C}$ and $z \in \mathbb{C}$. The following are equivalent.

- (a) We have that $z \in \overline{X}$.
- (b) For all $\varepsilon > 0$, we have $B(z, \varepsilon) \cap X \neq \emptyset$.
- (c) There exists $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ such that $z_n \rightarrow z$.

Proof. We show our directions one at a time.

- We show (a) implies (b). Suppose $z \in \overline{X}$, and for the sake of contradiction suppose we have $\varepsilon > 0$ such that $B(z, \varepsilon) \cap X = \emptyset$. In particular, $z \notin X$.

Now, $z \in \overline{X}$ implies that z is contained in every closed set containing X by definition of \overline{X} . But because $B(z, \varepsilon)$ is open and is disjoint from X , we see

$$z \in \overline{X} \subseteq \mathbb{C} \setminus B(z, \varepsilon),$$

which is a contradiction.

- We show (b) implies (c). For each $n \in \mathbb{N}$, we know that $B(z, 1/n) \cap X \neq \emptyset$, so we find some $z_n \in B(z, 1/n)$. Now, for any $\varepsilon > 0$, choose $N := 1/\varepsilon$ so that

$$n > N \implies |z_n - z| < \frac{1}{n} < \frac{1}{N} = \varepsilon,$$

so indeed $z_n \rightarrow z$.

- We show (b) implies (a). We proceed by contraposition. Suppose that $z \notin \overline{X}$. It follows that $z \in \mathbb{C} \setminus \overline{X}$, which is open, so there exists an $r > 0$ such that

$$B(z, r) \subseteq \mathbb{C} \setminus \overline{X} \subseteq \mathbb{C} \setminus X.$$

It follows that $B(z, r) \cap X = \emptyset$.

- We show (c) implies (b). Suppose $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ has $z_n \rightarrow z$ for some $z \in \mathbb{C}$. For any $\varepsilon > 0$, there exists N such that

$$n > N \implies |z_n - z| < \varepsilon,$$

so in particular, choosing any $n := \lceil N \rceil + 1$ has $z_n \in B(z, \varepsilon) \cap X$, so $B(z, \varepsilon) \cap X \neq \emptyset$. ■

The above discussion can give us a more directly topological definition of “closed.”

Lemma 1.86. A subset $X \subseteq \mathbb{C}$ is closed in \mathbb{C} if and only if X contains all of its limit points.

Proof. By the previous lemma, we see that $z \in \overline{X}$ if and only if z is a limit point of X , so \overline{X} is the set of limit points of X . Now, X is closed if and only if $X = \overline{X}$, so X is closed if and only if all limit points of X are in fact points of X . (Note that all points of X are automatically limit points essentially because $X \subseteq \overline{X}$ for free.) ■

While we're here, we can pick up a nice topological result.

Lemma 1.87. Fix $X \subseteq \mathbb{C}$ a connected subset. Then \overline{X} is also connected.

Proof. This argument is purely topological. We proceed by contraposition: suppose \overline{X} is disconnected by $U_1, U_2 \subseteq \mathbb{C}$. We claim that U_1, U_2 disconnect X . Well, we already know that $A \subseteq \overline{A} \subseteq U_1 \cup U_2$, and we already know that U_1 and U_2 are disjoint.

We claim that, for $U \subseteq \mathbb{C}$ an open subsets, if $U \cap \overline{X} \neq \emptyset$, then $U \cap X \neq \emptyset$ as well. Indeed, we proceed by contraposition: if $U \cap X = \emptyset$, then $X \subseteq \mathbb{C} \setminus U$, but $\mathbb{C} \setminus U$ is closed, so

$$\overline{X} \subseteq \mathbb{C} \setminus U,$$

so $\overline{X} \cap U = \emptyset$.

Thus, it follows from $U_1 \cap \overline{X}, U_2 \cap \overline{X} \neq \emptyset$ that $U_1 \cap X, U_2 \cap X \neq \emptyset$. This finishes the proof that U_1 and U_2 disconnect X . Indeed, ■

1.5.3 Cauchy Sequences

Just like in real analysis, we will be interested in Cauchy sequences.

Cauchy
sequence

Definition 1.88 (Cauchy sequence). A sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ is a *Cauchy sequence* if and only if, for each $\varepsilon > 0$, there exists an N such that

$$n, m > N \implies |z_n - z_m| < \varepsilon.$$

We have the following results on Cauchy sequences.

Proposition 1.89. Fix $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ a sequence. If $\{z_n\}_{n \in \mathbb{N}}$ converges, it is Cauchy.

Proof. This proof uses no special properties of \mathbb{C} . If $z_n \rightarrow z$, then for a given $\varepsilon > 0$, there exists N such that

$$n > N \implies |z_n - z| < \varepsilon/2.$$

It follows that

$$n, m > N \implies |z_n - z_m| < |z_n - z| + |z_m - z| < \varepsilon,$$

finishing. ■

Proposition 1.90. Every Cauchy sequence in \mathbb{C} converges.

Proof. If $\{z_n\}_{n \in \mathbb{N}}$ is Cauchy, then we claim $\{\operatorname{Re} z_n\}_{n \in \mathbb{N}}$ and $\{\operatorname{Im} z_n\}_{n \in \mathbb{N}}$ are Cauchy sequences. Indeed, for any $\varepsilon > 0$, there exists N so that

$$n, m > N \implies |z_n - z_m| < \varepsilon,$$

but then $|\operatorname{Re} z_n - \operatorname{Re} z_m| < |z_n - z_m|$ and $|\operatorname{Im} z_n - \operatorname{Im} z_m| < |z_n - z_m|$, so the same N witnesses that $\{\operatorname{Re} z_n\}_{n \in \mathbb{N}}$ and $\{\operatorname{Im} z_n\}_{n \in \mathbb{N}}$ are Cauchy in \mathbb{R} .

Now, Cauchy sequences in \mathbb{R} converge, so there are reals $x, y \in \mathbb{R}$ such that $\operatorname{Re} z_n \rightarrow x$ and $\operatorname{Im} z_n \rightarrow y$. It follows that $z_n \rightarrow x + yi$, finishing. ■

1.5.4 A Little More Topology

We close with one more topological definition.

Sequentially
compact

Definition 1.91 (Sequentially compact). A subset $X \subseteq \mathbb{C}$ is *sequentially compact* if and only if every $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ has a convergent subsequence which converges in X .

Remark 1.92. This happens to be equivalent to X is compact because $\mathbb{C} \cong \mathbb{R}^2$ satisfies the fact that all compact sets are closed and bounded.

Example 1.93. Every finite set is compact.

And here is a last definition.

Tends to
infinity

Definition 1.94 (Tends to infinity). A sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ *tends to infinity* (notated $z_n \rightarrow \infty$) if and only if each $M > 0$ has some $N \in \mathbb{N}$ such that

$$n > N \implies |z_n| > M.$$

Essentially the points of $\{z_n\}_{n \in \mathbb{N}}$ wander infinitely away.

1.6 January 31

So we are lecturing in-person today. Good morning everyone.

Quote 1.95. If I don't fall off the stage, I will consider it a major accomplishment.

Homework 2 is due Friday, the 4th of February. Office hours will occur at the usual times, but they will now occur in-person at Evans 749.

1.6.1 Series

Today we're mostly talking about series, and on Friday we'll talk about continuous functions.

Series

Definition 1.96 (Series). An *infinite series* over \mathbb{C} is an infinite sum

$$S := \sum_{n=1}^{\infty} z_n$$

where $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ is a sequence of complex numbers.

With respect to series, we really want to know when various series converge so that the series is well-defined.

Converge,
diverge

Definition 1.97 (Converge, diverge). Fix a sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ of complex numbers, we define the m th partial sum to be

$$S_m := \sum_{n=0}^m z_n.$$

Then we say that the infinite series *converges* if and only if the sequence $\{S_m\}$ of partial sums converges. Otherwise, we say that S is *divergent*.

As usual, we start with some basic examples.

Exercise 1.98. Fix some $z \in \mathbb{C}$ with $|z| < 1$, we define $z_n := z^n$. Then we have

$$S = \sum_{k=0}^{\infty} z^k = \frac{1}{1-z}.$$

Proof. Fix some partial sum

$$S_N := \sum_{k=0}^N z^k = 1 + z + z^2 + \cdots + z^N.$$

Multiplying by z , we see that

$$zS_N = z + z^2 + \cdots + z^N + z^{N+1}.$$

It follows that

$$S_N - zS_N = 1 - z^{N+1}.$$

Because $|z| < 1$, we have $z \neq 1$, so we may write

$$S_N = \frac{1}{1-z} - \frac{z^{N+1}}{1-z}.$$

However, we may note that as $N \rightarrow \infty$, the bad term z^{N+1} will have size

$$|z^{N+1}| = |z|^{N+1},$$

which goes to 0 (because $|z| < 1$).⁴ It follows that

$$\lim_{N \rightarrow \infty} S_N = \frac{1}{1-z},$$

which is what we wanted. ■

Anyways, here are some basic lemmas.

Lemma 1.99 (Divergence test). Suppose that $\{z_n\}_{n \in \mathbb{N}}$ is a sequence of complex numbers such that $\sum z_n$ converges. Then $z_n \rightarrow 0$ as $n \rightarrow \infty$.

Proof. Let S_n be the n th partial sum so that we are given $S_n \rightarrow L$ for some $L \in \mathbb{C}$. But now we see that

$$z_{n+1} = \left(\sum_{k=0}^{N+1} z_k \right) - \left(\sum_{k=0}^N z_k \right) = S_{n+1} - S_n.$$

Using limit laws, we see that

$$\lim_{n \rightarrow \infty} z_{n+1} = \lim_{n \rightarrow \infty} S_{n+1} - \lim_{n \rightarrow \infty} S_n = L - L = 0.$$

Shifting the indices back gives $z_n \rightarrow 0$ as $N \rightarrow \infty$. ■

Here is an important example of a divergent series.

⁴ This is surprisingly annoying to rigorize with a ε - δ proof, so we won't do so here. The interested can try to use induction to manually bound $|z|^n$ by $\frac{c}{n}$ for some c .

Exercise 1.100. We claim that

$$S = \sum_{k=1}^{\infty} \frac{1}{k}$$

does not converge.

Proof. We will show that the sequence of partial sums $\{S_n\}_{n=1}^{\infty}$ is not Cauchy, which will show that the series diverges. Well, observe that

$$S_{2^{n+1}} - S_{2^n} = \sum_{k=2^n+1}^{2^{n+1}} \frac{1}{k}$$

after cancelling out all of our terms. However, each term in the sum is at least $\frac{1}{2^{n+1}}$, so we may bound

$$S_{2^{n+1}} - S_{2^n} \geq \frac{1}{2^{n+1}} (2^{n+1} - 2^n) = \frac{1}{2}.$$

We now show that the partial sums are not Cauchy. Fix ε . Supposing for the sake of contradiction that the sequence is Cauchy, there exists N so that $n, m > N$ has

$$|S_n - S_m| < \frac{1}{2}.$$

However, we can find some power of 2 named 2^r which exceeds N , in which case we find $2^{r+1}, 2^r > N$ and

$$|S_{2^{r+1}} - S_{2^r}| \geq \frac{1}{2},$$

which is our contradiction. ■

Remark 1.101. Because a sequence will converge if and only if its real and imaginary parts do, we can turn a convergene test into a real-number test by taking the real and imaginary parts of the sum.

1.6.2 The Comparison Test

Recall the comparison test in \mathbb{R} .

Theorem 1.102 (Comparison test). Fix $\{x_n\}_{n \in \mathbb{N}}, \{y_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$ sequences of real numbers. Further, suppose that we there exists a positive constatatn $c > 0$ such that $0 \leq x_n \leq cy_n$. Then the following hold.

- If $\sum y_n$ converges, then $\sum x_n$ converges as well.
- If $\sum x_n$ diverges, then $\sum y_n$ diverges as well.

Proof. We appeal to real analysis. The interested can see Theorem 2.1.21 in Eterović. The main point is to use the Monotone sequence theorem. ■

Here is an example.

Exercise 1.103. Fix $s > 1$ an integer. Then the series

$$S = \sum_{k=1}^{\infty} \frac{1}{k^s}$$

converges.

Proof. Because s is an integer, we have $s \geq 2$. Namely, $\frac{1}{k^s} \leq \frac{1}{k^2}$, so by the comparison test it suffices to just show the convergence of

$$S' := \sum_{k=1}^{\infty} \frac{1}{k^2}.$$

For this, we apply some trickery. In particular, for $k > 1$, we can bound

$$\frac{1}{k^2} < \frac{1}{k(k-1)} = \frac{1}{k-1} - \frac{1}{k}.$$

In particular,

$$S' = 1 + \sum_{k=2}^{\infty} \frac{1}{k^2} < 1 + \sum_{k=2}^{\infty} \left(\frac{1}{k-1} - \frac{1}{k} \right).$$

Thus, by the comparison test, it suffices to show the convergence of

$$T := \sum_{k=2}^{\infty} \left(\frac{1}{k-1} - \frac{1}{k} \right).$$

But the n th partial sum will telescope, giving

$$T_n := \sum_{k=2}^n \left(\frac{1}{k-1} - \frac{1}{k} \right) = 1 - \frac{1}{n},$$

so $T_n \rightarrow 1$ as $n \rightarrow \infty$, and $T = 1$. It follows that S' is upper-bounded by $1 + T \leq 2$. ■

1.6.3 Absolute Convergence

The following kind of convergence is nontrivially stronger, but that makes it better.

Absolute
convergence

Definition 1.104 (Absolute convergence). Fix a sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ of complex numbers. Then the sum $S := \sum z_n$ converges *absolutely* if and only if the series

$$\sum_{n=0}^{\infty} |z_n|$$

also converges. In other words, the partial sums of the above series converges.

We have the following quick lemma to justify naming this “convergence.”

Lemma 1.105. If a series converges absolutely, then the series also converges.

Proof. The idea is to use the triangle inequality. Fix our series

$$S := \sum_{n=0}^{\infty} z_n$$

for which

$$T := \sum_{n=0}^{\infty} |z_n|$$

converges. Let S_n be the n th partial sum of S and T_n the n th partial sum of T .

Our goal is to show that $\{S_n\}_{n \in \mathbb{N}}$ is Cauchy. Observe $\{T_n\}_{n \in \mathbb{N}}$ is an increasing sequence of real numbers because $|z| \geq 0$ always. To start off our arithmetic, we note that, for $n, m \in \mathbb{N}$ with $n > m$, we have

$$|S_n - S_m| = \left| \sum_{k=m+1}^n z_k \right|,$$

which by the triangle inequality can be bounded by

$$|S_n - S_m| \leq \sum_{k=m+1}^n |z_k| = T_n - T_m.$$

But now we can use the fact that $\{T_n\}_{n \in \mathbb{N}}$ must be Cauchy to finish: for any $\varepsilon > 0$, there exists some N such that $n > m > N$ implies $T_m - T_n < \varepsilon$. But then this same N promises $n > m > N$ implies

$$|S_n - S_m| < T_m - T_n < \varepsilon,$$

which is what we wanted. ■

Here is a surprise tool that will help us later.

Lemma 1.106. Fix a sequence $\{a_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ of nonzero complex numbers. Further suppose that the sequence $\{a_n\}_{n \in \mathbb{N}}$ tends to infinity (i.e., $|a_n| \rightarrow \infty$ as $n \rightarrow \infty$), then for any positive real number $r \in \mathbb{R}^+$, the series

$$\sum_{k=0}^{\infty} \left(\frac{r}{|a_k|} \right)^k$$

converges.

Proof. We need the a_n to be nonzero in order to allow division, so the real puzzle is to determine how to use the fact $|a_n| \rightarrow \infty$. Well, there exists some N such that $n > N$ has

$$|a_n| > 2r.$$

But then $\frac{r}{|a_n|} < \frac{1}{2}$ for each $n > N$, so we can use the comparison test as follows: observe that

$$\sum_{k=0}^{\infty} \frac{1}{2^k}$$

will converge, and there will exist some $c > 1$ so that

$$\frac{r}{|a_k|} < \frac{c}{2^k}$$

for $0 \leq k \leq N$; and then for $n > N$, we get the above inequality anyways as discussed earlier (observe we took $c > 1$). ■

Quote 1.107. I can't break math on the first day of class. I can do it later on.

Lemma 1.108. Suppose that we have two series $S := \sum_{k \in \mathbb{N}} z_k$ and $T := \sum_{k \in \mathbb{N}} w_k$ are both absolutely convergent. Then the sum

$$P := \sum_{k=0}^{\infty} \left(\sum_{i+j=k} z_i w_j \right)$$

is absolutely convergent as well. In fact, P will converge to ST .

Proof. We sketch the result, and the remaining details are in Eterović. As usual, consider the partial sums

$$A_n := \sum_{k=0}^n |z_k| \quad \text{and} \quad B_n = \sum_{k=0}^n |w_k|,$$

both of which will converge as $n \rightarrow \infty$. Brazenly multiplying these together, we see that

$$A_n B_n = \sum_{i,j=0}^n |z_i w_j| = \sum_{k=0}^n \sum_{\substack{i+j=k \\ 0 \leq i,j \leq n}} |z_i w_j| + \sum_{k>n} \sum_{\substack{i+j=k \\ 0 \leq i,j \leq n}} |z_i w_j|.$$

In the first sum, observe that any time $i + j = k$, we will automatically have $i, j \leq k \leq n$. It follows that

$$A_n B_n = \sum_{k=0}^n \left(\sum_{i+j=k} z_i w_j \right) + \underbrace{\sum_{\substack{i+j>n \\ 0 \leq i,j \leq n}} |z_i w_j|}_{R_n}.$$

The key claim is that $R_n \rightarrow 0$. The main idea is that $i + j > n$ implies that $i \geq n/2$ or $j \geq n/2$, so we can write

$$|R_n| \leq \sum_{i=0}^n \sum_{j=n/2}^n |z_i w_j| + \sum_{i=n/2}^n \sum_{j=0}^n |z_i w_j| = \left(\sum_{i=0}^n |z_i| \right) \left(\sum_{j=n/2}^n |w_j| \right) + \left(\sum_{i=n/2}^n |z_i| \right) \left(\sum_{j=0}^n |w_j| \right).$$

Now, fix any $\varepsilon > 0$, and we show there exists X so that $n > X$ has $|R_n| < \varepsilon$. Note $A := \sum |z_k|$ and $B := \sum |w_k|$ both converge and hence have Cauchy partial sums. Because the partial sums are increasing, we may bound

$$|R_n| \leq A \left(\sum_{j=n/2}^n |w_j| \right) + B \left(\sum_{i=n/2}^n |z_i| \right)$$

So there exists N such that $n > m > N$ has

$$\sum_{i=m+1}^n |z_i| < \frac{\varepsilon}{2B}$$

Similarly there exists M so that $n > m > M$ has

$$\sum_{j=m+1}^n |w_j| < \frac{\varepsilon}{2A},$$

from which it follows that $n > n/2 > \max\{N, M\}$ will have

$$|R_n| \leq A \cdot \frac{\varepsilon}{2A} + B \cdot \frac{\varepsilon}{2B} = \varepsilon,$$

which finishes.

Now, because $R_n \rightarrow 0$, we see

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \left(\sum_{i+j=k} |z_i| \cdot |w_j| \right) = \lim_{n \rightarrow \infty} A_n B_n - \lim_{n \rightarrow \infty} R_n,$$

which does indeed converge, so indeed the series

$$\sum_{k=0}^{\infty} \left(\sum_{i+j=k} |z_i| \cdot |w_j| \right)$$

will converge. By the comparison test (using the triangle inequality), it follows that

$$P = \sum_{k=0}^{\infty} \left(\sum_{i+j=k} z_i w_j \right)$$

will also absolutely converge.

To show that P converges to ST , we observe that the difference of the n th partial sum is

$$P_n - S_n T_n = \sum_{k=0}^n \left(\sum_{i+j=k} z_i w_j \right) - \sum_{i,j=0}^n z_i w_j = \sum_{k=0}^n \left(\sum_{i+j=k} z_i w_j \right) - \sum_{k=0}^n \left(\sum_{i+j=k}^n z_i w_j \right) + \sum_{\substack{0 \leq i,j \leq n \\ i+j < n}} z_i w_j,$$

so

$$P_n - S_n T_n = \sum_{\substack{0 \leq i,j \leq n \\ i+j < n}} z_i w_j.$$

But by the triangle inequality, we see $|P_n - S_n T_n| \leq R_n$, so $P_n - S_n T_n \rightarrow 0$ as $n \rightarrow \infty$. It follows P_n and $S_n T_n$ have the same limit as $n \rightarrow \infty$ (which exists because S_n and T_n have a limit). So indeed, $P = ST$. ■

1.7 February 2

Good morning everyone. Here is some house-keeping.

- Homework #2 is due at Friday at 11:59, on GradeScope. The assignment has just been added.
- There are office hours to talk about the homework. Please come if you have questions.

1.7.1 Summation Review

Today we finish our discussion of series, for now. We quickly recall the definitions.

Series

Definition 1.109 (Series). An *infinite series* over \mathbb{C} is an infinite sum

$$S := \sum_{n=1}^{\infty} z_n$$

where $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ is a sequence of complex numbers.

Converge,
diverge

Definition 1.110 (Converge, diverge). Fix a sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ of complex numbers, we define the m th partial sum to be

$$S_m := \sum_{n=0}^m z_n.$$

Then we say that the infinite series *converges* if and only if the sequence $\{S_m\}$ of partial sums converges. Otherwise, we say that S is *divergent*.

Today we are building towards proving Dirichlet's convergence theorem. We pick up the following lemmas.

Lemma 1.111. Fix sequences $\{z_{k,\ell}\}_{k,\ell \in \mathbb{N}}$ a collection of complex numbers satisfying the following conditions.

- Fixing any k , the sum $\sum_{\ell=0}^{\infty} |z_{k,\ell}|$ converges.
- The sum $\sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}|$ converges.

Then the following are true.

- (a) Fix any ℓ , the sum $\sum_{k=0}^{\infty} |z_{k,\ell}|$ converges; i.e., the terms in the left sum below are well-defined.
- (b) We have that

$$\sum_{\ell=0}^{\infty} \sum_{k=0}^{\infty} z_{k,\ell} = \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} z_{k,\ell},$$

and both sums converge.

Intuitively, the first condition is requiring that the series horizontally does not grow too fast. The second condition is requiring an absolute convergence condition. Then the first conclusion says we can sum vertically instead, and the second conclusion says that we can move around the order of summation.

Proof. We will sketch this proof; we prove (a) and (b) more or less simultaneously. To turn the infinite double sum into something we can consider finite partial sums of, we set, for each natural N ,

$$S_n := \sum_{k=0}^n \sum_{\ell=0}^n |z_{k,\ell}|.$$

The main claim is that

$$\lim_{n \rightarrow \infty} S_n \stackrel{?}{=} \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}|.$$

Indeed, fix any $\varepsilon > 0$. Because the latter sum converges, there exists some natural A such that

$$\sum_{k>A} \sum_{\ell=0}^{\infty} |z_{k,\ell}| < \frac{\varepsilon}{2}.$$

Further, there exists some natural B_k such that

$$\sum_{\ell>B_k} |z_{k,\ell}| < \frac{\varepsilon}{2A}$$

for each $k \in \mathbb{N}$. Take $B := \max_{0 \leq k < A} B_k$. Now, we set $N := \max\{A, B\}$. To start off our inequalities, we note that

$$S_n = \sum_{k=0}^n \sum_{\ell=0}^n |z_{k,\ell}| \leq \sum_{k=0}^n \sum_{\ell=0}^{\infty} |z_{k,\ell}| \leq \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}|,$$

so we know the sign of our difference. In particular, for any $n > N$, we see that

$$S_n = \sum_{k=0}^N \sum_{\ell=0}^N |z_{k,\ell}| \geq \sum_{k=0}^K \sum_{\ell=0}^L |z_{k,\ell}|.$$

Thus,

$$0 \leq \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - S_n \leq \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - \sum_{k=0}^K \sum_{\ell=0}^L |z_{k,\ell}| = \sum_{k>K} \sum_{\ell=0}^{\infty} |z_{k,\ell}| + \sum_{k=0}^K \sum_{\ell>L} |z_{k,\ell}|$$

after some cancellation. But we can upper-bound the last quantity by $\frac{\varepsilon}{2} + K \cdot \frac{\varepsilon}{2K} = \varepsilon$, so we are done.

The main point of the above lemma is that we are told each $\varepsilon > 0$ has some N so that $n > N$ implies

$$\sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - S_n = \sum_{\substack{(k,\ell) \in \mathbb{Z}^2 \\ k > n \text{ or } \ell > n}} |z_{k,\ell}| < \varepsilon.$$

We now take the two parts in sequence.

- (a) Fix an index ℓ' ; we show absolute convergence by showing that the partial sums of $\sum_{k=0}^{\infty} |z_{k,\ell'}|$ are Cauchy. Indeed, fix some $\varepsilon > 0$, and we know there exists N so that each $n > N$ has

$$\sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - S_n < \varepsilon.$$

Now, we see that any $n > m > N$ will have

$$\sum_{k=m+1}^n |z_{k,\ell'}| \leq \sum_{k=m+1}^N \sum_{\ell=0}^{\infty} |z_{k,\ell}| \leq \sum_{k=N+1}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| + \sum_{k=0}^N \sum_{\ell=N+1}^{\infty} |z_{k,\ell}| < \varepsilon,$$

so we are done.

- (b) As above, fix some $\varepsilon > 0$, and we are promised N so that

$$\sum_{\substack{(k,\ell) \in \mathbb{Z}^2 \\ k > N \text{ or } \ell > N}} |z_{k,\ell}| < \varepsilon/2.$$

Observe, for $K, L > N$, we have by the triangle inequality tht

$$\left| \sum_{\ell=0}^L \sum_{k=0}^K z_{k,\ell} - S_N \right| < \varepsilon/2.$$

This bounds holds for any K , so we can send K arbitrarily large; that inner sum will converge, so in fact we can send K to ∞ without ill effect. (Formally, the inner term is an increasing sequence bounded above, so it will converge as $K \rightarrow \infty$.) This gives

$$\left| \sum_{\ell=0}^L \sum_{k=0}^{\infty} z_{k,\ell} - S_N \right| \leq \varepsilon/2.$$

Again, the inner term is an increasing sequence as $L \rightarrow \infty$ but still bounded above as $\varepsilon/2$, so the inner sum will converge as $L \rightarrow \infty$ and still give the inequality

$$\left| \sum_{\ell=0}^{\infty} \sum_{k=0}^{\infty} z_{k,\ell} - S_N \right| < \varepsilon.$$

Now as we send $\varepsilon \rightarrow 0$, we see that $\lim_{N \rightarrow \infty} S_N = \sum_{\ell=0}^{\infty} \sum_{k=0}^{\infty} z_{k,\ell}$, which finishes. ■

1.7.2 Dirichlet Test

We now go directly for the Dirichlet test for convergence.

Lemma 1.112 (Summation by parts). Fix sequences $\{a_n\}_{n \in \mathbb{N}}$ and $\{b_n\}_{n \in \mathbb{N}}$ sequences of complex numbers. Then we define

$$B_n := \sum_{k=0}^n b_k,$$

and $B_{-1} = 0$ to be the empty sum. It follows that, for any $n, m \in \mathbb{N}$ with $n > m$,

$$\sum_{k=m}^n a_k b_k = a_n B_n - a_m B_{m-1} + \sum_{k=m}^{n-1} B_k (a_k - a_{k+1}).$$

Proof. This comes down to a direct computation. The main point is that $b_k = B_k - B_{k-1}$, which even works with $k = 0$. Indeed,

$$\begin{aligned}
 \sum_{k=m}^n a_k b_k &= \sum_{k=m}^n a_k (B_k - B_{k-1}) \\
 &= \sum_{k=m}^n a_k B_k - \sum_{k=m}^n a_k B_{k-1} \\
 &\stackrel{*}{=} a_n B_n + \sum_{k=m}^{n-1} a_k B_k - a_m B_{m-1} - \sum_{k=m}^n a_{k+1} B_k \\
 &= a_n B_n - a_m B_{m-1} + \sum_{k=m}^{n-1} B_k (a_k - a_{k+1}),
 \end{aligned}$$

which is what we wanted. The important step to pay attention to is the rearrangement we did in $\stackrel{*}{=}$ in order to massage the sums together. ■

And here is our theorem.

Theorem 1.113 (Dirichlet's test). Fix $\{a_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$ a sequence of real numbers and $\{b_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$ a sequence of complex numbers satisfying the following conditions.

- The sequence $\{a_n\}_{n \in \mathbb{N}}$ is decreasing.
- We have $a_n \rightarrow 0$ as $n \rightarrow \infty$.
- Bounded partial sums: there exists a positive real number M such that

$$\left| \sum_{k=0}^n b_k \right| < M$$

for each $n \in \mathbb{N}$.

Then we claim that

$$\sum_{k=0}^{\infty} a_k b_k$$

converges.

Proof. As usual, fix our partial sums

$$S_n := \sum_{k=0}^n a_k b_k \quad \text{and} \quad B_n := \sum_{k=0}^n b_k.$$

We are given that the B_k are bounded, so we are going to want to use Lemma 1.112, which tells us that

$$S_n = a_n B_n + \sum_{k=0}^{n-1} B_k (a_k - a_{k+1}).$$

We examine the convergence of these terms individually.

- We will show that the summation term absolutely converges. We are given that the partial sums B_n are bounded by M , so we note $|B_k (a_k - a_{k+1})| < M |a_k - a_{k+1}|$, so it suffices to show that

$$M \sum_{k=0}^{n-1} |a_k - a_{k+1}|$$

converges as $n \rightarrow \infty$. It would be great if this would telescope, and indeed it does! Because the a_k are decreasing,

$$\sum_{k=0}^{\infty} |a_k - a_{k+1}| = \sum_{k=0}^{\infty} (a_k - a_{k+1}) = a_0 - a_{n+1}.$$

Because $a_n \rightarrow 0$ as $n \rightarrow \infty$, we see that this sum will converge to a_0 . It follows that

$$\sum_{k=0}^{\infty} |B_k(a_k - a_{k+1})|$$

will converge by the Comparison test, so

$$\sum_{k=0}^{\infty} B_k(a_k - a_{k+1})$$

converges by absolute convergence.

- Note that the B_n are bounded in norm by M , so $|a_n B_n| \leq M|a_n|$, but $|a_n| \rightarrow 0$ as $n \rightarrow \infty$, so $|a_n B_n| \rightarrow 0$. ■

Eterović has lots of different convergence tests in his notes, but we don't care about most of them. Here is one that we do care about.

Theorem 1.114 (Integral test). Fix a decreasing function $f : [1, \infty) \rightarrow \mathbb{R}^+$ and for which

$$\int_k^{k+1} f(x) dx$$

always exists. Then the sequence of integrals $I_n := \int_1^n f(x) dx$ converges if and only if the summation

$$\sum_{k=1}^{\infty} f(k)$$

converges.

Proof. We omit this proof; it's a reasonably standard real-analytic test. ■

1.8 February 4

Today we are talking about continuity.



Warning 1.115. The first half of this lecture was transcribed from Professor Morrow's notes because

1.8.1 Limits

Before defining continuity, we have the following definitions.

Limit

Definition 1.116 (Limit). Fix $f : X \rightarrow \mathbb{C}$ a function and $z_0 \in \overline{X}$. Then we say *the limit of $f(z)$ as z approaches z_0 equals w* , denoted

$$\lim_{z \rightarrow z_0} f(z) = w,$$

if and only if, for each $\varepsilon > 0$, there exists $\delta > 0$ such that

$$|z - z_0| < \delta \implies |f(z) - w| < \varepsilon$$

for $z \in X$.

This is the standard ε - δ definition.

We also pick up the following convention as a surprise tool that may help us later.

Infinite limits

Definition 1.117 (Infinite limits). Fix $f : X \rightarrow \mathbb{C}$ a function. Then we say *the limit of $f(z)$ as z tends to infinity equals w* , denoted

$$\lim_{z \rightarrow \infty} f(z) = w,$$

if and only if, for each $\varepsilon > 0$, there exists $N > 0$ such that

$$|z| > N \implies |f(z) - w| < \varepsilon$$

for $z \in X$.

As in real analysis, the ε - δ definition of a limit can be translated to a statement about sequences.

Proposition 1.118. Fix $\alpha \in \overline{X}$. Then $\lim_{z \rightarrow \alpha} f(z) = w$ if and only if, for each $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ such that $z_n \rightarrow \alpha$ as $n \rightarrow \infty$, we have $f(z_n) \rightarrow w$ as $n \rightarrow \infty$.

Proof. In the forwards direction, fix $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ such that $z_n \rightarrow \alpha$, and we show that $f(z_n) \rightarrow w$. Well, for any $\varepsilon > 0$, there exists $\delta > 0$ such that

$$|z - \alpha| < \delta \implies |f(z) - f(\alpha)| < \varepsilon,$$

where $z \in X$. But for this $\delta > 0$, there exists N such that

$$n > N \implies |z_n - \alpha| < \delta \implies |f(z_n) - f(\alpha)| < \varepsilon.$$

So indeed, $f(z_n) \rightarrow f(\alpha)$.

In the reverse direction, suppose that $f(z)$ does not approach w as $z \rightarrow \alpha$. Then, there exists $\varepsilon_0 > 0$ such that, for any $\delta > 0$, there is $z \in X$ such that $|z - \alpha| < \delta$ while $|f(z) - w| > \varepsilon_0$. Well, for any $n \in \mathbb{N}$, taking $\delta = 1/(n+1)$, we are promised $z_n \in X$ such that

$$|z_n - \alpha| < \frac{1}{n+1} \text{ and } |f(z_n) - w| > \varepsilon_0.$$

So to finish, we claim that $z_n \rightarrow \alpha$ as $n \rightarrow \infty$, but $f(z_n)$ does not approach w as $n \rightarrow \infty$.

- For any $\varepsilon > 0$, we note that $N := 1/\varepsilon$ has $n > N$ implies

$$|z_n - \alpha| < \frac{1}{n+1} < \frac{1}{N+1} < \frac{1}{N} = \varepsilon,$$

so indeed $z_n \rightarrow \alpha$ as $n \rightarrow \infty$.

- We note that $\varepsilon_0 > 0$ satisfies that

$$|f(z_n) - w| > \varepsilon_0$$

for any $n \in \mathbb{N}$, so no N will have $n > N$ implies $|f(z_n) - w| < \varepsilon_0$. Thus, $f(z_n)$ does not approach w as $n \rightarrow \infty$.

The sequence $\{z_n\}_{n \in \mathbb{N}}$ now completes the proof by showing the reverse direction by contraposition. ■

While we're here, we pick up the following definitions.

Bounded

Definition 1.119 (Bounded). A function $f : X \rightarrow \mathbb{C}$ is *bounded* if there exists $R > 0$ such that $\text{im } f \subseteq B(0, R)$.

Bounded
near

Definition 1.120 (Bounded near). Fix a nonempty open subset $\Omega \subseteq \mathbb{C}$ and $z_0 \in \Omega$. Then $f : \Omega \setminus \{z_0\} \rightarrow \mathbb{C}$ is *bounded near* z_0 if and only if

$$\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0.$$

1.8.2 Continuity

And here is our central definition for today.

Continuous

Definition 1.121 (Continuous). A function $f : X \rightarrow \mathbb{C}$ is *continuous at* $z_0 \in X$ if and only if, for each $\varepsilon > 0$, there exists $\delta > 0$ such that

$$|z - z_0| < \delta \implies |f(z) - f(z_0)| < \varepsilon,$$

where $z \in X$. Further, f is *continuous on* X if and only if f is continuous at each $z_0 \in X$.

We have the following lemma of equivalent definitions.

Lemma 1.122. Suppose that $f : X \rightarrow \mathbb{C}$.

- (a) Then f is continuous at w if and only if every sequence $\{z_n\} \subseteq X$ such that $z_n \rightarrow z$ implies $f(z_n) \rightarrow f(z)$.
- (b) We have that f is continuous on X if and only if every open set $U \subseteq \mathbb{C}$ has $f^{-1}(U)$ open in X .
- (c) We have that f is continuous on X if and only if each closed set $V \subseteq X$ has $f^{-1}(V)$ closed in X .
- (d) We have that f is continuous at if and only if, for each $\varepsilon > 0$ and $z \in \mathbb{C}$, we have that $f^{-1}(B(z, \varepsilon))$ is open in X .

Proof. We take the parts one at a time.

- (a) We could use Proposition 1.118, but we will just do this by hand. For the forwards direction, suppose that $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ converges to some w . Then let $\varepsilon > 0$. By assumption, there exists some $\delta > 0$ such that

$$|z - w| < \delta \implies |f(z) - f(w)| < \varepsilon.$$

It follows from $z_n \rightarrow w$ that there exists some N such that

$$n > N \implies |z_n - w| < \delta \implies |f(z_n) - f(w)| < \varepsilon,$$

so it follows that $f(z_n) \rightarrow f(w)$.

In the reverse direction, take f not continuous at w , so there exists $\varepsilon > 0$ so that for all $n \in \mathbb{N}$, there exists some chosen z_n with

$$|z_n - w| < \delta \implies |f(z_n) - f(w)| > \varepsilon.$$

But as $z_n \rightarrow w$, we see that $f(z_n)$ does not approach $f(w)$, so we are done.

- (b) In the forwards direction, suppose that $U \subseteq \mathbb{C}$ is open, and we show that $f^{-1}(U)$ is open in X . Well, suppose that $z \in f^{-1}(U)$, and we will find $\delta > 0$ such that $B(z, \delta) \subseteq f^{-1}(U)$.

Well, $f(z) \in U$, so there exists $\varepsilon > 0$ such that $B(f(z), \varepsilon) \subseteq U$. Thus, continuity of f requires some $\delta > 0$ such that

$$|w - z| < \delta \implies |f(w) - f(z)| < \varepsilon,$$

which implies $f(w) \in B(f(z), \varepsilon) \subseteq U$ implies $w \in f^{-1}(U)$. So indeed, $B(z, \delta) \subseteq f^{-1}(U)$.

In the reverse direction, suppose that each open $U \subseteq \mathbb{C}$ has $f^{-1}(U)$ is open. Now fix any $z \in X$ and $\varepsilon > 0$. The set $B(f(z), \varepsilon)$ is open, so

$$f^{-1}(B(f(z), \varepsilon))$$

is open. But $z \in f^{-1}(B(f(z), \varepsilon))$, so we can find $\delta > 0$ such that $B(z, \delta) \subseteq f^{-1}(B(f(z), \varepsilon))$. Thus, $|w - z| < \delta$ implies $w \in f^{-1}(B(f(z), \varepsilon))$ implies $f(w) \in B(f(z), \varepsilon)$ implies $|f(w) - f(z)| < \varepsilon$, finishing.

- (c) In the forwards direction, suppose f is continuous so that $U \subseteq \mathbb{C}$ open implies $f^{-1}(U) \subseteq X$ is open. But then, if $V \subseteq \mathbb{C}$ is closed, then $\mathbb{C} \setminus V$ is open, so⁵

$$f^{-1}(\mathbb{C} \setminus V) = f^{-1}(\mathbb{C}) \setminus f^{-1}(V) = X \setminus f^{-1}(V)$$

is open, so $f^{-1}(V)$ is closed.

In the backwards direction, suppose f^{-1} preserves closed sets. Then, if $U \subseteq \mathbb{C}$ is open, $\mathbb{C} \setminus U$ is closed, so

$$f^{-1}(\mathbb{C} \setminus U) = f^{-1}(\mathbb{C}) \setminus f^{-1}(U) = X \setminus f^{-1}(U)$$

is closed, so $f^{-1}(U)$ is open. Thus, f^{-1} preserves open sets, so f must be continuous.

- (d) In the forwards direction, fix $\varepsilon > 0$ and $z \in \mathbb{C}$, so $B(z, \varepsilon)$ is open, so $f^{-1}(B(z, \varepsilon))$ is open in X , finishing.

In the other direction fix $\varepsilon > 0$ and $z \in \mathbb{C}$ to consider $B(f(z), \varepsilon) \subseteq U$. Thus, continuity of f requires some $\delta > 0$ such that

$$|w - z| < \delta \implies |f(w) - f(z)| < \varepsilon,$$

which implies $f(w) \in B(f(z), \varepsilon) \subseteq U$ implies $w \in f^{-1}(U)$. So indeed, $B(z, \delta) \subseteq f^{-1}(U)$.

In the reverse direction, fix U open, and we show that $f^{-1}(U)$ is open. Well, each $z \in U$ has some ε_z such that $B(z, \varepsilon_z) \subseteq U$. But $f^{-1}(B(z, \varepsilon_z))$ is open by hypothesis, so

$$f^{-1}(U) = f^{-1}\left(\bigcup_{z \in U} B(z, \varepsilon_z)\right) = \bigcup_{z \in U} f^{-1}(B(z, \varepsilon_z))$$

is an arbitrary union of open sets and hence open. ■

And here are some special examples.

Example 1.123. Fix some $z_0 \in \mathbb{C}$. Then $f(z) := |z - z_0|$ is continuous on \mathbb{C} . Indeed, fix any $w \in \mathbb{C}$. Then for any $\varepsilon > 0$, we set $\delta := \varepsilon$ so that $|z - w| < \delta$ implies

$$|f(z) - f(w)| = \left| |z - z_0| - |w - z_0| \right| \leq |z - w| < \delta = \varepsilon.$$

Example 1.124. The functions Re and Im is continuous. Indeed, fix any $w \in \mathbb{C}$. Then, for any $\varepsilon > 0$, take $\delta := \varepsilon$ so that $|z - w| < \delta$ implies

$$|\operatorname{Re} z - \operatorname{Re} w| = |\operatorname{Re}(z - w)| \leq |z - w| < \delta = \varepsilon,$$

and similarly,

$$|\operatorname{Im} z - \operatorname{Im} w| = |\operatorname{Im}(z - w)| \leq |z - w| < \delta = \varepsilon,$$

Continuous functions also have some arithmetic.

⁵ To see $f^{-1}(A \setminus B) = f^{-1}(A) \setminus f^{-1}(B)$, note that $x \in f^{-1}(A \setminus B)$ if and only if $f(x) \in A \setminus B$ if and only if $f(x) \in A$ but $f(x) \notin B$ if and only if $x \in f^{-1}(A)$ but $x \notin f^{-1}(B)$.

Proposition 1.125. Fix $f, g : X \rightarrow \mathbb{C}$ to functions continuous at $z_0 \in X$. Then $f + g, f \cdot g$ are both continuous at $z_0 \in X$, and f/g is continuous at z_0 provided $g(z_0) \neq 0$.

Proof. The point is to appeal to the corresponding results on convergence of sequences. In particular, we use the idea that f is continuous at z_0 if and only if each sequence $z_n \rightarrow z_0$ in X has $f(z_n) \rightarrow f(z_0)$. We omit the details because they are essentially the same as in a real analysis class. ■

Corollary 1.126. Every polynomial in one variable is a continuous function $X \rightarrow \mathbb{C}$ for any $X \subseteq \mathbb{C}$.

Proof. Note that $x \mapsto x$ is continuous, so by induction $x \mapsto x^n$ is continuous for each $n \in \mathbb{N}$. Taking a \mathbb{C} -linear combination gives arbitrary polynomials. ■

Here is another sort of arithmetic.

Lemma 1.127. The composition of two continuous functions is continuous.

Proof. Omitted. ■

1.8.3 Connectedness

We want to build towards a particular type of continuous function.

Proposition 1.128. Fix $X \subseteq \mathbb{C}$ a connected subset. Then a continuous function $f : X \rightarrow \mathbb{C}$ has connected image $f(X)$.


Proof. The main point is to use the topological characterization of continuity. In particular, suppose that $f(X)$ is disconnected, and we show that X is disconnected. In particular, suppose that U_1 and U_2 disconnect $f(X)$, and we have that $f^{-1}(U_1)$ and $f^{-1}(U_2)$ disconnect X . We will not run all the checks here; the main point is that $f^{-1}(U_1)$ and $f^{-1}(U_2)$ are open because f is continuous. ■

Path **Definition 1.129 (Path).** A path in \mathbb{C} is a continuous function $\gamma : [a, b] \rightarrow \mathbb{C}$ where $a < b$ are real numbers.


Definition 1.130. We say that a path γ is *closed* if and only if $\gamma(a) = \gamma(b)$. We say that γ is *simple* if and only if $\gamma : (a, b) \rightarrow \mathbb{C}$ is injective.

Remark 1.131. We restrict γ to the closed interval at the end so that closed, simple paths are allowed to exist.


Example 1.132. Here is a path.




Example 1.133. Here is a closed path.



Example 1.134. Here is a simple path.



Example 1.135. Here is a closed, simple path, also called a loop.




Concena-
tion

Definition 1.136 (Concatenation). Fix $\gamma_1 : [a, b] \rightarrow \mathbb{C}$ and $\gamma_2 : [c, d] \rightarrow \mathbb{C}$ paths in \mathbb{C} such that $\gamma_1(b) = \gamma_2(c)$. Then we define the *concatenation* of γ_1 and γ_2 to be

$$(\gamma_1 * \gamma_2)(t) := \begin{cases} \gamma_1(t) & t \in [a, b], \\ \gamma_2(t) & t \in [b, d + c - b]. \end{cases}$$

The main point is that we are doing one path after the other.

Example 1.137. The following shows an example concatenation of $\gamma_1 * \gamma_2$, where $\gamma_1, \gamma_2 : [0, 1] \rightarrow \mathbb{C}$.



The entire path is $\gamma_1 * \gamma_2$.

Paths give us the following notion.

Path-
connected

Definition 1.138 (Path-connected). A subset $X \subseteq \mathbb{C}$ is *path connected* if and only if, for any two $x_0, x_1 \in X$, there exists a path $\gamma : [0, 1] \rightarrow X$ such that $\gamma(0) = x_0$ and $\gamma(1) = x_1$.

Lemma 1.139. The open ball $B(z, r)$ and closed ball $\overline{B(z, r)}$ are both path-connected.

Proof. The point is that $B(z, r)$ and $\overline{B(z, r)}$ are both convex, so the path

$$\gamma(t) := z_0 + t(z_1 - z_0)$$

will work. ■

Here is the basic result.

Proposition 1.140. A space X is path-connected implies that X is connected. If X is open and connected, then X is path-connected.

Proof. We will show this next class. ■

1.9 February 7

Good morning everyone. A few announcements.

- Homework #3 is due on Friday.
- There will be no in-person class on Wednesday or Friday.
- Office hours this week are today (1:00PM–2:30PM) and tomorrow (2:00PM–3:30PM).

1.9.1 Connectedness

Today we're going to talk more about continuous functions.

Last time we ended with the following proposition.


Proposition 1.141. A space X is path-connected implies that X is connected. If X is open and connected, then X is path-connected.

Proof. We do these separately.

- Suppose that $X = U_1 \sqcup U_2$ is disconnected, and we show that X is not path-connected. Namely, we have $U_1, U_2 \subseteq X$ open subsets (in X) which are disjoint and nonempty. Because U_1 and U_2 are nonempty, find $x_1 \in U_1$ and $x_2 \in U_2$.

However, we claim there is no continuous path $\gamma : [0, 1] \rightarrow X$ with $\gamma(0) = x_1$ and $\gamma(1) = x_2$. Indeed, the image of $\gamma([0, 1])$ must be connected, but then we can disconnect $\gamma([0, 1])$ by U_1 and U_2 : $\gamma([0, 1]) \subseteq U_1 \cup U_2$ and $x_\bullet \in U_\bullet \cap \gamma([0, 1])$ and $U_1 \cap U_2 = \emptyset$.

At a high level, here is the image that a disconnected X cannot have a path between any two pair points: there is no possible red path below which stays in the gray region.




- Suppose we have a point $z \in X$, and we set

$$C(z) := \{w \in X : \text{there is a path from } z \text{ to } w\}.$$

We claim that $C(z)$ is closed and open in X , which will force $C(z) = X$ because X is connected and $C(z)$ is nonempty ($z \in C(z)$ by the trivial path $\gamma : t \mapsto z$).

We start by showing $C(z)$ is open: because X is open, there exists $r > 0$ such that $B(w, r) \subseteq X$. But with $w \in C(z)$, there will be a path between any point in $p \in B(z, w)$ and w , so there is a path from z to w to p . Here is the image.



Now we show that $C(z)$ is closed. Suppose that $w \in X \setminus C(z)$, and we have to show that there is an open ball around w in $X \setminus C(z)$. To see this, fix an open ball $B(w, r) \subseteq X$ for $r > 0$, but now there can be no path from z to anywhere in $B(w, r)$, for then we could just run the above argument again to show that $w \in C(z)$. ■

Remark 1.142. The proof for the second part merely needs X to be locally path-connected, not a metric space.

Corollary 1.143. We have that \mathbb{C} is path-connected and therefore connected.

Proof. Given any two points $z, w \in \mathbb{C}$, we choose the path $\gamma : [0, 1] \rightarrow \mathbb{C}$ by

$$\gamma(t) = tz + (1 - t)w.$$

Indeed, $\gamma(0) = w$ and $\gamma(1) = z$, and γ is somewhat clearly continuous by, say, checking componentwise. ■

1.9.2 Compactness

Let's do compactness better this time.

Lemma 1.144. Fix $X \subseteq \mathbb{C}$ (sequentially) compact. Then X is both closed and bounded.

Proof. We start by showing X is closed. For this, we show that X contains all of its limit points.

Well, suppose that $z \in X$ is a limit point so that we have a sequence $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ such that $z_n \rightarrow z$. But by the (sequential) compactness of X , this sequence has a convergent subsequence $\{z_{\sigma n}\}_{n \in \mathbb{N}}$ which does converge in X . But any subsequence will converge to the same limit (!), so $z_{\sigma n} \rightarrow z$ as well, so $z \in X$ is forced.

We now show that X is bounded. We proceed by contraposition: if X is unbounded, then for any $n \in \mathbb{N}$, then we can find some $z_n \in X \setminus B(0, n)$. But then we can check that $\{z_n\}_{n \in \mathbb{N}}$ has no convergent subsequence, essentially because it tends off to infinity. ■

Our goal for the rest of class is to prove the following two results.

Proposition 1.145. A subset $X \subseteq \mathbb{C}$ is (sequentially) compact if and only if it is closed and bounded.

Theorem 1.146 (Heine–Borel). A subset $X \subseteq \mathbb{C}$ is (sequentially) compact if and only if every open cover of X has a finite subcover.

On the homework, we showed the backward direction of Theorem 1.146.

Remark 1.147. Our hope is to have lots of equivalent characterizations of compactness so that we can have easier proofs of statements about compact sets.

To start off, here are some lemmas we will need.

Lemma 1.148. Fix $X \subseteq \mathbb{C}$ (sequentially) compact. For any $\varepsilon > 0$, there exist only finitely many points $z_1, \dots, z_n \in X$ such that

$$X \subseteq \bigcup_{k=1}^n B(z_k, \varepsilon).$$

Proof. The point is to build some inductive argument: one fixes an $\varepsilon > 0$ and then continues choosing random points out of X until we cover X . Indeed, if the process does not terminate, then the sequence we generate has no convergent subsequence.

Rigorously, if X is empty, then just choose no points at all and be done. Otherwise, we can find some $z_1 \in X$. Inductively, suppose we have a sequence $\{z_1, \dots, z_m\}$. If

$$X \subseteq \bigcup_{k=1}^m B(z_k, \varepsilon),$$

then we are done. Otherwise, we can find $z_{m+1} \in X \setminus \bigcup_{k=1}^m B(z_k, \varepsilon)$.

If the above inductive process terminates, then we get the result. Otherwise, there is a sequence $\{z_n\}_{n \in \mathbb{N}}$ such that

$$z_{n+1} \in X \setminus \bigcup_{k=1}^n B(z_k, \varepsilon).$$

We claim that $\{z_n\}_{n \in \mathbb{N}}$ has subsequence converging in X . Indeed, suppose for the sake of contradiction that $z_{\sigma n} \rightarrow z$ for some strictly increasing σ and $z \in X$. Then there exists N such that $n > N$ implies

$$|z_{\sigma n} - z| < \varepsilon/2.$$

But then, finding some $n+1, n > N$, we have

$$|z_{\sigma(n+1)} - z_{\sigma n}| < |z_{\sigma(n+1)} - z| + |z_{\sigma n} - z| < \varepsilon,$$

so

$$z_{\sigma(n+1)} \in \bigcup_{k=1}^{\sigma(n+1)-1} B(z_k, \varepsilon),$$

which is our contradiction to the construction of z_n . ■

Lemma 1.149. Fix $X \subseteq \mathbb{C}$ (sequentially) compact with some open cover \mathcal{U} of X . Then there is an $\varepsilon > 0$ such that, for every $z \in X$, there is $U \in \mathcal{U}$ such that $B(z, \varepsilon) \subseteq U$.

Proof. Suppose that, for all $\varepsilon > 0$, there exists some $z \in X$ such that no $U \in \mathcal{U}$ has $B(z, \varepsilon) \subseteq U$. We construct a sequence in X with no subsequence converging in X . Indeed, for any $n \in \mathbb{N}$, we find $z_n \in X$ such that no $U \in \mathcal{U}$ has $B(z_n, 1/n) \subseteq U$. We claim that $\{z_n\}_{n \in \mathbb{N}}$ has no subsequence converging in X .

Indeed, suppose that we have $z \in X$ and strictly increasing $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $z_{\sigma n} \rightarrow z$. We will then be able to find some z_n such that $B(z_n, 1/n) \subseteq U$ for some $U \in \mathcal{U}$, which will be a contradiction. Indeed, $z \in X$, and \mathcal{U} covers z , so there is some $U \in \mathcal{U}$ with $z \in U$. In fact, U is open, so there is an $\varepsilon > 0$ such that

$$B(z, \varepsilon) \subseteq U.$$

Now, there is N such that for $n > N$, we can guarantee that $|z - z_n| < \varepsilon/2$. Further, for $n > 2/\varepsilon$, we have $1/n < \varepsilon/2$. So $n > \max\{N, 2/\varepsilon\}$ will have $\sigma n > \max\{N, 2/\varepsilon\}$, implying

$$|w - z_n| < 1/n < \varepsilon/2 \implies |w - z| < |w - z_n| + |z - z_n| = \varepsilon \implies w \in B(z, \varepsilon) \subseteq U,$$

so $B(z_n, 1/n) \subseteq U$. This contradiction finishes. ■

This is saying that there is a universal ε that we can find for our open cover.

Lemma 1.150. Fix X a bounded set. Then, for any $\varepsilon > 0$, there exist finitely many points z_1, \dots, z_n such that

$$X \subseteq \bigcup_{k=1}^n B(z_k, \varepsilon).$$

Proof. The point is to reduce this to the case of $[-M, M]^2$ which can cover X by boundedness, and then we can create the cover for X by hand. ■

Now let's attack one of our equivalent conditions for compactness.

Proposition 1.145. A subset $X \subseteq \mathbb{C}$ is (sequentially) compact if and only if it is closed and bounded.

Proof. The forwards direction we have already done.

In the backwards direction, suppose that $\{z_n\}_{n \in \mathbb{N}} \subseteq X$ is some sequence. Our main goal is to construct a convergent subsequence. By boundedness, we can choose $w_{1,1}, \dots, w_{1,\ell_1}$ such that

$$X \subseteq \bigcup_{k=1}^{\ell_1} B(w_{1,k}, 1/2).$$

Now, because $\{z_n\}_{n \in \mathbb{N}}$ is infinite, there must be some index $w_1 := w_{1,k_1}$ such that

$$L_1 = \{n \in \mathbb{N} : z_n \in B(w_{1,k_1}, 1/2)\}$$

is infinite. The important point is that $\{z_n\}_{n \in L_1}$ has formed a subsequence which lives inside of a ball of radius $1/2$. We can continue this process: by boundedness, we can find some $w_{2,1}, \dots, w_{2,\ell_2} \in B(w_{1,k_1}, 1/2)$ such that

$$B(w_{1,k_1}, 1/2) \subseteq \bigcup_{k=1}^{\ell_2} B(w_{2,k}, 1/4).$$

Then we can choose L_2 from here by choosing one of the $w_{2,k}$ with infinitely many indices. Continuing this process forces our sequence to converge.

To more explicitly appeal to choice, we note that we can always find some sequence $\{w_{k,i}\} \subseteq X$ such that

$$X \subseteq \bigcup_{k=1}^{\ell_n} B(w_{k,i}, 1/2^k),$$

but L_{i-1} is infinite, so there is a specific $w_k := w_{k,i}$ such that

$$L_i := \{n \in L_{i-1} : |z_n - w_k| < 2^{-k}\}$$

is infinite. To actually construct our sequence from these infinite subsets, we define a choice function over our indices: define $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ such that $\varphi(n+1)$ is the smallest number exceeding $\varphi(n)$ with $\varphi(n+1) \in L_{n+1}$. Then we know that

$$|z_{\varphi(n)} - w_k| < 2^{-n}$$

for each $1 \leq k \leq n$. Thus, for $n \geq m > N$, we have

$$|z_{\varphi(n)} - z_{\varphi(m)}| \leq |z_{\varphi(n)} - w_m| + |z_{\varphi(m)} - w_m| < 2 \cdot 2^{-m} < 2^{-N+1},$$

so for any $\varepsilon > 0$, we can choose $N := 1 - \log_2 \varepsilon$ sufficiently large so that $n, m > N$ implies

$$|z_{\varphi(n)} - z_{\varphi(m)}| < 2^{-N+1} = \varepsilon.$$

It follows that the subsequence defined by φ is Cauchy and hence converges. But because X is closed, any convergent sequence in X will be in X , so our sequence in X has a convergent subsequence. ■