

# 185: Introduction to Complex Analysis

Nir Elber

Spring 2022

---

# CONTENTS

---

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	January 19 . . . . .	3
1.2	January 21 . . . . .	5
<b>2</b>	<b>Complex Numbers and Their Topology</b>	<b>11</b>
2.1	January 24 . . . . .	11
2.2	January 26 . . . . .	15
2.3	January 28 . . . . .	21
2.4	January 31 . . . . .	27
2.5	February 2 . . . . .	33
2.6	February 4 . . . . .	37
2.7	February 7 . . . . .	43
2.8	February 9 . . . . .	47
<b>3</b>	<b>Holomorphic Functions</b>	<b>53</b>
3.1	February 11 . . . . .	53
3.2	February 14 . . . . .	57
3.3	February 16 . . . . .	61
3.4	February 18 . . . . .	63

---

# THEME 1: INTRODUCTION

---

*Our reality isn't about what's real, it's about what we pay attention to.*

—Hank Green

## 1.1 January 19

It is reportedly close enough to start.

### 1.1.1 Logistics

We are online for the first two weeks, as with the rest of Berkeley. We will be using `bCourses` a lot, so check it frequently. There is also a website. There is a homework due on Friday, but do not worry about it.

Here are some syllabus things.

- Our main text is *Complex Variables and Applications, 8th Edition* because it is the version that Professor Morrow used. There is a free copy online.
- The homework consists of readings (for each course day) and weekly problem sets. Late homework is never accepted.
- Lowest two homework scores are dropped.
- There are two midterms and a final. The final is cumulative, as usual. The final can replace one midterm if the score is higher.
- Regrade requests can be made in GradeScope within one week of being graded.
- The class is curved but usually only curved at the end. The average on exams is expected to be 80%–83%.

### 1.1.2 Complex Numbers

Welcome to complex analysis. What does that mean?




**Idea 1.1.** In complex analysis, we study functions  $f : \mathbb{C} \rightarrow \mathbb{C}$ , usually analytic to some extent.

There are two pieces here: we should study  $\mathbb{C}$  in themselves, and then we will study the functions.

**Definition 1.2** (Complex numbers). The set of complex numbers  $\mathbb{C}$  is  $\{a + bi : a, b \in \mathbb{R}\}$ , where  $i^2 = -1$ .

Hopefully  $\mathbb{R}$  is familiar from real analysis. As an aside, we see  $\mathbb{R} \subseteq \mathbb{C}$  because  $a = a + 0i \in \mathbb{C}$  for each  $a \in \mathbb{R}$ . The complex numbers have an inherent geometry as a two-dimensional plane.



The point is that  $\mathbb{C}$  looks like the real plane  $\mathbb{R}^2$ . More precisely,  $\mathbb{C} \cong \mathbb{R}^2$  as an  $\mathbb{R}$ -vector space, where our basis is  $\{1, i\}$ .

We would like to understand  $\mathbb{C}$  geometrically, “as a space.” The first step here is to create a notion of size.

**Definition 1.3** (Norm on  $\mathbb{C}$ ). We define the **norm map**  $|\cdot| : \mathbb{C} \rightarrow \mathbb{R}_{\geq 0}$  by  $|z| := \sqrt{z\bar{z}}$ . In other words,

$$|a + bi| := \sqrt{a^2 + b^2}.$$

Note that this agrees with the absolute value on  $\mathbb{R}$ : for  $a \in \mathbb{R}$ , we have  $\sqrt{a^2} = |a|$ .

Norm functions, as in the real case, give us a notion of distance.

**Definition 1.4** (Metric on  $\mathbb{C}$ ). We define the *metric on  $\mathbb{C}$*  to be  $d_{\mathbb{C}}(z_1, z_2) := |z_1 - z_2|$ .

One can check that this is in fact a metric, but we will not do so here.

**Remark 1.5.** The distance in  $\mathbb{C}$  is defined to match the distance in  $\mathbb{R}^2$  under the basis  $\{1, i\}$ .

Again as we discussed in real analysis, having a metric gives us a metric topology by open balls. Lastly it is this topology that our geometry will follow from: we have turned  $\mathbb{C}$  into a topological space.

### 1.1.3 Complex Functions

There are lots of functions on  $\mathbb{C}$ , and lots of them are terrible. So we would like to focus on functions with some structure. We’ll start with *continuous functions*, which are more or less the functions that respect topology.

Then from continuous functions, we will be able to define *holomorphic functions*, which are complex differentiable. This intended to be similar to being real differentiable, but complex differentiable turns out to be a very strong condition. Nevertheless, everyone’s favorite functions are holomorphic.

**Example 1.6.** Polynomials,  $\exp$ ,  $\sin$ , and  $\cos$  are all holomorphic.

To make concrete that complex differentiability is stronger than real differentiability, the Cauchy–Riemann equations which provides a partial differential equation to test complex differentiability.

From here we define *analytic functions*, which essentially are defined as taking the form

$$f(z) := \sum_{k=0}^{\infty} a_k z^k.$$

Analytic functions are super nice in that we have an ability to physically write them down, so the following theorem is amazing.

**Theorem 1.7.** Holomorphic functions on  $\mathbb{C}$  are analytic.

To prove this, we will need the following result, which is what Professor Morrow calls the most fundamental result in complex analysis, the *Cauchy integral formula*.

In short, the Cauchy integral formula lets us talk about the value of holomorphic functions (and derivatives) at a point in terms of integrals around the point. This will essentially let us build the power series for a holomorphic function by hand. But as described, we will need a notion of complex (path) integration to even be able to talk about the Cauchy integral formula.

The Cauchy integral formula has lots of applications; for example, *Liouville's theorem* on holomorphic functions and the *Fundamental theorem of algebra*.

**Remark 1.8.** It is very hard to spell Liouville.

Additionally, we remark that our study of holomorphic functions, via the Cauchy integral formula, will boil down to a study of complex path integrals. So we will finish out our story with the *Residue theorem*, which provides a very convenient way to compute such integrals.

Then as a fun addendum, we talk about automorphisms of the complex numbers.

**Definition 1.9** (Automorphisms of  $\mathbb{C}$ ). A function  $f : \mathbb{C} \rightarrow \mathbb{C}$  is an *automorphism of  $\mathbb{C}$*  if it is bijective and both  $f$  and  $f^{-1}$  are holomorphic.

What is amazing is that all of these functions have a concrete description in terms of *Möbius transformations*.

### 1.1.4 Why Care?

Whenever taking a class, it is appropriate to ask why one should care. Here are some reasons to care.

- Algebraic geometry in its study of complex analytic spaces uses complex analysis.
- Analytic number theory (e.g., the Prime number theorem) makes heavy use of complex analysis.
- Combinatorics via generating functions can use complex analysis.
- Physics uses complex analysis.

The first two Professor Morrow is more familiar with, the last two less so.

## 1.2 January 21

We're reviewing set theory today.

### 1.2.1 Set Theory Notation

We have the following definitions.

- $\emptyset$  means the empty set.
- $a \in X$  means that  $a$  is an element of the set  $X$ .
- $A \subseteq B$  means that  $A$  is a subset of  $B$ .
- $A \subsetneq B$  means that  $A$  is a proper subset of  $B$ .

- $A \cup B$  consists of the elements which are in at least one of  $A$  or  $B$ .
- $A \cap B$  consists of the elements which are in both  $A$  and  $B$ .
- $A \setminus B$  consists of the elements of  $A$  which are not in  $B$ .
- Two sets  $A$  and  $B$  are *disjoint* if and only if  $A \cap B = \emptyset$ .
- Given a set  $X$ , we define  $\mathcal{P}(X)$  to be the set of all subsets of  $X$ .
- $|X| = \#X$  is the cardinality of  $X$ , or (roughly speaking) the number of elements of  $X$ .

As an example of unwinding notation, we have the following.

**Proposition 1.10** (De Morgan's Laws). Fix  $\mathcal{S} \subseteq \mathcal{P}(X)$  a collection of subsets of a set  $X$ . Then

$$X \setminus \bigcap_{S \in \mathcal{S}} S = \bigcup_{S \in \mathcal{S}} (X \setminus S) \quad \text{and} \quad X \setminus \bigcup_{S \in \mathcal{S}} S = \bigcap_{S \in \mathcal{S}} (X \setminus S).$$

*Proof.* We take these one at a time.

- Note  $a \in X \setminus \bigcap_{S \in \mathcal{S}} S$  if and only if  $a \in X$  and  $a \notin \bigcap_{S \in \mathcal{S}} S$ . However,  $a \notin \bigcap_{S \in \mathcal{S}} S$  is merely saying that  $a$  is not in all the sets  $S \in \mathcal{S}$ , which is equivalent to saying  $a \notin S$  for one of the  $S \in \mathcal{S}$ .  
Thus, this is equivalent to saying  $a \in X$  while  $a \notin S$  for some  $S \in \mathcal{S}$ , which is equivalent to  $a \in \bigcup_{S \in \mathcal{S}} (X \setminus S)$ .
- Note  $a \in X \setminus \bigcup_{S \in \mathcal{S}} S$  if and only if  $a \in X$  and  $a \notin \bigcup_{S \in \mathcal{S}} S$ . However,  $a \notin \bigcup_{S \in \mathcal{S}} S$  is merely saying that  $a$  is not in any of the sets  $S \in \mathcal{S}$ , which is equivalent to saying  $a \notin S$  for each of the  $S \in \mathcal{S}$ .  
Thus, this is equivalent to saying  $a \in X$  while  $a \notin S$  for each  $S \in \mathcal{S}$ , which is equivalent to  $a \in \bigcap_{S \in \mathcal{S}} (X \setminus S)$ . ■

### 1.2.2 Some Conventions

In this class, we take the following names of standard sets.

- $\mathbb{N} = \{0, 1, 2, \dots\}$  is the set of natural numbers. Importantly,  $0 \in \mathbb{N}$ .
- $\mathbb{N}^+ = \{1, 2, 3, \dots\}$  is the set of positive integers.
- $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  is the set of integers.
- $\mathbb{Q} = \{p/q : p, q \in \mathbb{Z} \text{ and } q \neq 0\}$  is the set of rationals.
- $\mathbb{R}$  is the set of real numbers. We will not specify a construction here; see any real analysis class.
- $\mathbb{R}^\times = \{x \in \mathbb{R} : x \neq 0\}$  is the nonzero real numbers.
- $\mathbb{R}^+ = \{x \in \mathbb{R} : x > 0\}$  is the positive real numbers.
- $\mathbb{R}_{\geq 0} = \{x \in \mathbb{R} : x \geq 0\}$  is the nonnegative real numbers.
- $\mathbb{R}_{\leq 0} = \{x \in \mathbb{R} : x \leq 0\}$  is the nonpositive real numbers.
- $\mathbb{C}$  is the complex numbers.
- $\mathbb{C}^\times = \{z \in \mathbb{C} : z \neq 0\}$  is the set of nonzero complex numbers.

### 1.2.3 Relations

Let's review some set theory definitions.

**Definition 1.11** (Cartesian product). Given two sets  $A$  and  $B$ , we define the *Cartesian product*  $A \times B$  to be the set of ordered pairs  $(a, b)$  such that  $a \in A$  and  $b \in B$ .

**Definition 1.12** (Binary relation). A *binary relation* on  $A$  is any subset  $R \subseteq A^2 := A \times A$ . We may sometimes notate  $(x, y) \in R$  by  $xRy$ , read as " $x$  is related to  $y$ ."

**Example 1.13.** Equality is a binary relation on any set  $A$ ; namely, it is the subset  $\{(a, a) : a \in A\}$ .

The best relations are equivalence relations.

**Definition 1.14** (Equivalence relation). An *equivalence relation* on  $A$  is a binary relation  $R$  satisfying the following three conditions.

- Reflexive: each  $x \in A$  has  $(x, x) \in R$ .
- Symmetric: each  $x, y \in A$  has  $(x, y) \in R$  implies  $(y, x) \in R$ .
- Transitive: each  $x, y, z \in A$  has  $(x, y) \in R$  and  $(y, z) \in R$  implies  $(x, z) \in R$ .

Equivalence relations are nice because they allow us to partition the set into "equivalence classes."

**Definition 1.15** (Equivalence class). Fix  $A$  a set and  $R \subseteq A^2$  an equivalence relation. Then, for given  $x \in A$ , we define

$$[x]_R := \{y \in A : (x, y) \in R\}$$

to be the *equivalence class* of  $x$ .

The hope is that equivalence classes partition the set. What is a partition?

**Definition 1.16** (Partition). A *partition* of a set  $A$  is a collection of nonempty subsets  $S \subseteq \mathcal{P}(A)$  of  $A$  such that any two distinct  $S_1, S_2 \in S$  are disjoint while  $A = \bigcup_{S \in S} S$ .

And now let's manifest our hope.

**Lemma 1.17.** Equivalence relations are in one-to-one correspondence with partitions of  $A$ .

*Proof.* Given an equivalence relation  $R$ , we define the collection

$$\mathcal{S}(R) = \{[x]_R : x \in A\}.$$

We claim that  $R \mapsto \mathcal{S}(R)$  is our needed bijection. We have the following checks.

- Well-defined: observe that  $\mathcal{S}(R)$  does partition  $A$ : if we have  $[x]_R, [y]_R \in \mathcal{S}$ , then  $[x]_R \cap [y]_R \neq \emptyset$  implies there is some  $z$  with  $(x, z) \in R$  and  $(z, y) \in R$ , so  $x \in [y]_R$  and then  $[x]_R \subseteq [y]_R$  follows. So by symmetry,  $[y]_R \subseteq [x]_R$  as well, so we finish the disjointness check.

Further, we see that

$$A = \bigcup_{x \in A} \{x\} \subseteq \bigcup_{x \in A} [x]_R \subseteq A$$

because  $x \in [x]_R$ , so indeed the equivalence classes cover  $A$ .

- **Injective:** suppose  $R_1$  and  $R_2$  have  $\mathcal{S}(R_1) = \mathcal{S}(R_2)$ . We show that  $R_1 \subseteq R_2$ , and  $R_2 \subseteq R_1$  will follow by symmetry, finishing.

We notice that, for any  $\mathcal{S}$  partitioning  $A$ , being a partition, will have exactly one subset which contains  $x$ . But for  $\mathcal{S}(R)$  for an equivalence relation  $R$ , we see  $x \in [x]_R \in \mathcal{S}(R)$ , so this equivalence class must be the one.

So because  $[x]_{R_1}$  and  $[x]_{R_2}$  are the only subsets of  $\mathcal{S}(R_1)$  and  $\mathcal{S}(R_2)$  containing  $x$  (respectively), we must have  $[x]_{R_1} = [x]_{R_2}$ . Thus,  $(x, y) \in R_1$  implies  $y \in [x]_{R_1} = [x]_{R_2}$  implies  $(x, y) \in R_2$ .

- **Surjective:** suppose that  $\mathcal{S}$  is a partition of  $A$ . As noted above, each  $x \in A$  is a member of exactly one set  $S \in \mathcal{S}$ , which we call  $[x]$ . Then we define  $R \subseteq A^2$  by  $(x, y) \in R$  if and only if  $y \in [x]$ . One can check that this is an equivalence relation, which we will not do here in detail.<sup>1</sup>

The point is that

$$[x]_R = \{y : (x, y) \in R\} = \{y : y \in [x]\} = [x],$$

so  $\mathcal{S}(R) = \mathcal{S}$ . So our mapping is surjective. ■

We continue our discussion.

**Definition 1.18** (Quotient set). Given an equivalence relation  $R \subseteq A^2$ , we define the *quotient set*  $A/R$  is the set of equivalence classes of  $R$ . In other words,

$$A/R = \{[x]_R : x \in A\}.$$

Intuitively, the quotient set is the set where we have gone ahead and identified the elements which are “similar” or “related.”

We would like a more concrete way to talk about equivalence classes, for which we have the following.

**Definition 1.19** (Representatives). Given an equivalence relation  $R \subseteq A^2$ , we say that  $C \subseteq A$  is a set of *representatives of  $R$ -equivalence classes of  $A$*  if and only if  $C$  consists of exactly one element from each equivalence class in  $A/R$ .

## 1.2.4 Functions

To finish off, we discuss functions.

**Definition 1.20** (Functions). A *function*  $f : X \rightarrow Y$  is a relation  $f \subseteq X \times Y$  satisfying the following.

- For each  $x \in X$ , there is some  $y \in Y$  such that  $(x, y) \in f$ . Intuitively, each  $x \in X$  goes somewhere.
- For each  $x \in X$  and given some  $y_1, y_2 \in Y$  such that  $(x, y_1), (x, y_2) \in f$ , then  $y_1 = y_2$ . Intuitively, each  $x \in X$  goes to at most one place.

We will write  $f(x) = y$  as notational sugar for  $(x, y) \in f$ . Note this equality is legal because the value  $y$  with  $(x, y) \in f$  is uniquely given.

We would like to create new functions from old. Here are two ways to do this.

**Definition 1.21** (Restriction). Given a function  $f : X \rightarrow Y$  and a subset  $A \subseteq X$ , we define

$$f|_A = \{(x, y) \in f : x \in A\} \subseteq A \times Y$$

to be a function  $f|_A : A \rightarrow Y$ .

<sup>1</sup> Note  $x \in [x]$  by definition of  $[x]$ . If  $y \in [x]$ , then note  $y \in [y]$  as well, so  $[x] = [y]$  is forced by uniqueness, so  $x \in [y]$ . If  $y \in [x]$  and  $z \in [y]$ , then again by uniqueness  $[x] = [y] = [z]$ , so  $z \in [x]$  follows.



We will not check that  $f|_A$  is actually a function; it is, roughly speaking inherited from  $f$ .

**Definition 1.22.** Given two functions  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ , we define the *composition* of  $f$  and  $g$  to be some function  $g \circ f : X \rightarrow Z$  defined by

$$(g \circ f)(x) := g(f(x)).$$

Again, we will not check that this makes a function; it is.

Functions can also help create new sets.

**Definition 1.23** (Image). Given a function  $f : X \rightarrow Y$ , we define the *image* of  $f$  to be

$$\text{im } f = f(X) := \{y \in Y : \text{there is } x \in X \text{ such that } f(x) = y\}.$$

Namely,  $\text{im } f$  consists of all elements hit by someone in  $X$  hit by  $f$ .

**Definition 1.24** (Fiber, pre-image). Given a function  $f : X \rightarrow Y$  and some  $y \in Y$ , we define the *fiber* of  $f$  over  $y$  to be

$$f^{-1}(y) = \{x \in X : f(x) = y\} \subseteq X.$$

In general, we define the *pre-image* of a subset  $A \subseteq Y$  to be

$$f^{-1}(A) := \{x \in X : f(x) \in A\} = \bigcup_{a \in A} \{x \in X : f(x) = a\} = \bigcup_{a \in A} f^{-1}(a).$$

Some functions have nicer properties than others.

**Definition 1.25** (Inj-, sur-, bijective). Fix a function  $f : X \rightarrow Y$ . We have the following.

- Then  $f$  is *injective* or *one-to-one* if and only if, given  $x_1, x_2 \in X$ ,  $f(x_1) = f(x_2)$  implies  $x_1 = x_2$ .
- Then  $f$  is *surjective* or *onto* if and only if  $\text{im } f = Y$ . In other words, for each  $y \in Y$ , there exists  $x \in X$  with  $f(x) = y$ .
- Then  $f$  is *bijective* if and only if it is both injective and surjective.

Here is an example.

**Definition 1.26** (Identity). For a given set  $X$ , the function  $\text{id}_X : X \rightarrow X$  defined by  $\text{id}_X(x) := x$  is called the *identity function*.

For completeness, here are the checks that  $\text{id}_X$  is bijective.

- Injective: given  $x_1, x_2 \in X$ , we see  $\text{id}_X(x_1) = \text{id}_X(x_2)$  implies  $x_1 = \text{id}_X(x_1) = \text{id}_X(x_2) = x_2$ .
- Surjective: given  $x \in X$ , we see that  $x \in \text{im } \text{id}_X$  because  $x = \text{id}_X(x)$ .

We leave with some lemmas, to be proven once in one's life.

**Lemma 1.27.** Fix finite sets  $X$  and  $Y$  such that  $\#X = \#Y$ . Then a function  $f : X \rightarrow Y$  is bijective if and only if it is injective or surjective.

*Proof.* Certainly if  $f$  is bijective, then it is both injective and surjective, so there is nothing to say.

The reverse direction is harder. We proceed by induction on  $\#X = \#Y$ . If  $\#X = \#Y = 0$ , then  $X = Y = \emptyset$ , and all functions  $f : \emptyset \rightarrow \emptyset$  are vacuously bijective: for injective, note that any  $x_1, x_2 \in \emptyset$  have  $x_1 = x_2$ ; for surjective, note that any  $x \in \emptyset$  has  $f(x) = x$ .

Otherwise,  $\#X = \#Y > 0$ . We have two cases.

- Take  $f$  injective; we show  $f$  is surjective. In this case,  $\#X > 0$ , so choose some  $a \in X$ . Note that  $x \in X$  with  $x \neq a$  will have  $f(x) \neq f(a)$  by injectivity, so we may define the restriction

$$f|_{X \setminus \{a\}} : X \setminus \{a\} \rightarrow Y \setminus \{f(a)\}.$$

Observe that  $f|_{X \setminus \{a\}}$  is injective because  $f$  is: if  $x_1, x_2 \in X \setminus \{a\}$  have

$$f(x_1) = f|_{X \setminus \{a\}}(x_1) = f|_{X \setminus \{a\}}(x_2) = f(x_2),$$

then  $x_1 = x_2$  follows.

Now,  $\#(X \setminus \{a\}) = \#(Y \setminus \{f(a)\}) = \#X - 1$ , so by induction  $f|_{X \setminus \{a\}}$  will be bijective because it is injective. In particular,  $f$  by way of  $f|_{X \setminus \{a\}}$  fully hits  $Y \setminus \{f(a)\}$  in its image, so because  $f(a) \in \text{im } f$  as well, we conclude  $\text{im } f = Y$ . So  $f$  is surjective.

- Take  $f$  surjective; we show  $f$  is injective. Define a function  $g : Y \rightarrow X$  as follows: for each  $y \in Y$ , the surjectivity of  $f$  promises some  $x \in X$  such that  $f(x) = y$ , so choose any such  $x$  and define  $g(y) := x$ .<sup>2</sup> Observe that  $f(g(y)) = y$  by construction.

Now, we notice that  $g$  is injective: if  $y_1, y_2 \in Y$  have  $g(y_1) = g(y_2)$ , then  $y_1 = f(g(y_1)) = f(g(y_2)) = y_2$ . So the previous case tells us that  $g$  is in fact bijective.

So now choose any  $x_1, x_2 \in X$  such that  $f(x_1) = f(x_2)$ . The surjectivity of  $f$  promises some  $y_1, y_2 \in Y$  such that  $g(y_1) = x_1$  and  $g(y_2) = x_2$ , so we see that

$$x_1 = g(y_1) = g(f(g(y_1))) = g(f(x_1)) = g(f(x_2)) = g(f(g(y_2))) = g(y_2) = x_2,$$

proving our injectivity. ■

**Lemma 1.28.** Fix  $f : X \rightarrow Y$  a bijective function. Then there is a unique function  $g : Y \rightarrow X$  such that  $f \circ g = \text{id}_Y$  and  $g \circ f = \text{id}_X$ .

*Proof.* We show existence and uniqueness separately.

- We show existence. Note that, because  $f : X \rightarrow Y$  is surjective, each  $y \in Y$  has some  $x \in X$  such that  $f(x) = y$ . In fact, this  $x \in X$  is uniquely defined because  $f(x_1) = f(x_2)$  implies  $x_1 = x_2$ , so we may define  $g(y)$  as the value  $x$  for which  $f(x) = y$ .

By construction,  $f(g(y)) = y$ , so  $f \circ g = \text{id}_Y$ . Additionally, we note that, given any  $x \in X$ , the value  $x_0$  for which  $f(x) = f(x_0)$  is  $x = x_0$  by the injectivity, so  $g(f(x)) = x$ . Thus,  $g \circ f = \text{id}_X$ , as claimed.

- We show uniqueness. Suppose that we have two functions  $g_1, g_2 : Y \rightarrow X$  which satisfy

$$f \circ g_1 = f \circ g_2 = \text{id}_Y \quad \text{and} \quad g_1 \circ f = g_2 \circ f = \text{id}_X.$$

Then we see that

$$g_1 = g_1 \circ \text{id}_Y = g_1 \circ (f \circ g_2) = (g_1 \circ f) \circ g_2 = \text{id}_X \circ g_2 = g_2,$$

where we have used the fact that function composition associates. This finishes. ■

<sup>2</sup> Technically we are using the Axiom of Choice here. One can remove this with an induction because all sets are finite, but I won't bother.

---

## THEME 2: COMPLEX NUMBERS AND THEIR TOPOLOGY

---

*This somewhat laborious proof could have been avoided if one had defined a complex analytic structure*

— Jean-Pierre Serre

### 2.1 January 24

Good morning everyone.

#### 2.1.1 Algebraic Structure

Today we are reviewing the complex numbers (reportedly, "some basics"). Or at least it is hopefully mostly review. Here is our main character this semester.

**Definition 2.1** (Complex numbers). The set  $\mathbb{C}$  of *complex numbers* is

$$\mathbb{C} := \{a + bi : a, b \in \mathbb{R}\}.$$

Here  $i$  is some symbol such that  $i^2 = -1$  formally.

In particular, two complex numbers  $a_1 + b_1i$  and  $a_2 + b_2i$  are equal if and only if  $a_1 = a_2$  and  $b_1 = b_2$ .  
The complex numbers also have some algebraic structure.

**Definition 2.2** (+ and  $\times$  in  $\mathbb{C}$ ). Given complex numbers  $a_1 + b_1i, a_2 + b_2i \in \mathbb{C}$ , we define

$$(a_1 + b_1i) + (a_2 + b_2i) = (a_1 + a_2) + (b_1 + b_2)i,$$

and

$$(a_1 + b_1i)(a_2 + b_2i) = (a_1a_2 - b_1b_2) + (a_1b_2 + a_2b_1)i,$$

defined essentially by direct expansion, upon recalling  $i^2 = -1$ .

Here is the corresponding algebraic structure.

**Proposition 2.3.** The set  $\mathbb{C}$  with the above operations is a two-dimensional  $\mathbb{R}$ -vector space with basis  $\{1, i\}$ .

*Proof.* The elements  $\{1, i\}$  span  $\mathbb{C}$  because all complex numbers in  $\mathbb{C}$  can be written as  $a + bi = a \cdot 1 + b \cdot i$  by definition.

To see that these elements are linearly independent, suppose  $a + bi = 0$ . If  $b = 0$ , then  $a = 0$  follows, and we are done. Otherwise, take  $b \neq 0$ , but then we see  $(-a/b) = i$ , so

$$(-a/b)^2 = -1 < 0,$$

which does not make sense for real numbers. This finishes. ■

**Proposition 2.4.** The set  $\mathbb{C}$  with the above operations is a field.

*Proof.* We have the following checks.

- The element  $0 + 0i$  is our additive identity. Indeed, one can check that  $(0 + 0i) + (a + bi) = (a + bi) + (0 + 0i) = a + bi$ .
- The element  $1 + 0i$  is our multiplicative identity. Indeed, one can check that  $(1 + 0i)(a + bi) = (a + bi)(1 + 0i) = a + bi$ .
- Commutativity of addition and multiplication follow from by expansion.
- The distributive laws can again be checked by expansion.
- The additive inverse of  $a + bi$  is  $(-a) + (-b)i$ .
- The multiplicative inverse of  $a + bi$  can be found by wishing really hard and writing

$$\frac{1}{a + bi} = \frac{1}{a + bi} \cdot \frac{a - bi}{a - bi} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Then one can check this works. ■

Sometimes we would like to extract our coefficients from our basis.

**Definition 2.5** (Re and Im). Given  $z := a + bi \in \mathbb{C}$ , we define the operations

$$\operatorname{Re} z := a \quad \text{and} \quad \operatorname{Im} z := b.$$

Importantly,  $\operatorname{Re} : \mathbb{C} \rightarrow \mathbb{R}$  and  $\operatorname{Im} : \mathbb{C} \rightarrow \mathbb{R}$ .

Because we are merely doing basis extraction, it makes sense that these operations will preserve some (additive) structure.

**Proposition 2.6.** Fix  $z = a + bi$  and  $w = c + di$ . Then the following.

- $\operatorname{Re}(z + w) = \operatorname{Re} z + \operatorname{Re} w$ .
- $\operatorname{Im}(z + w) = \operatorname{Im} z + \operatorname{Im} w$ .

*Proof.* We proceed by direct expansion. Observe

$$\operatorname{Re}(z + w) = \operatorname{Re}((a + c) + (b + d)i) = a + c = \operatorname{Re} z + \operatorname{Re} w,$$

and

$$\operatorname{Im}(z + w) = \operatorname{Im}((a + c) + (b + d)i) = b + d = \operatorname{Im} z + \operatorname{Im} w.$$

This finishes. ■

It also turns out that the complex numbers have a very special transformation.

**Definition 2.7** (Conjugate). Given  $z := a + bi \in \mathbb{C}$ , we define the *complex conjugate* to be  $\bar{z} := a - bi \in \mathbb{C}$ .

We promised conjugation would be special, so here are some special things.

**Proposition 2.8.** Fix  $z = a + bi \in \mathbb{C}$ . Then the following.

- (a)  $z + \bar{z} = 2 \operatorname{Re} z$ .
- (b)  $z - \bar{z} = 2i \operatorname{Im} z$ .
- (c)  $\overline{\bar{z}} = z$ .

*Proof.* We take these one at a time.

- (a) Write  $a + bi + \overline{a + bi} = a + bi + a - bi = 2a$ .
- (b) Write  $a + bi - \overline{a + bi} = a + bi - (a - bi) = 2bi$ .
- (c) Write  $\overline{\overline{a + bi}} = \overline{a - bi} = a + bi$ . ■

In fact, more is true.

**Proposition 2.9.** Fix  $z = a + bi \in \mathbb{C}$  and  $w = c + di \in \mathbb{C}$ . Then the following.

- (a)  $\overline{z + w} = \bar{z} + \bar{w}$ .
- (b)  $\overline{zw} = \bar{z} \cdot \bar{w}$ .

*Proof.* We take these one at a time.

- Write

$$\overline{z + w} = (a + c) - (b + d)i = (a - bi) + (c - di) = \bar{z} + \bar{w}.$$

- Write

$$\begin{aligned} \bar{z} \cdot \bar{w} &= (a - bi)(c - di) \\ &= (ac - bd) - (ad + bc)i \\ &= \overline{(ac - bd) + (ad + bc)i} \\ &= \overline{zw}. \end{aligned}$$

This finishes. ■

### 2.1.2 Defining Distance

Complex conjugation actually gives rise to a notion of size.

**Definition 2.10** (Norm on  $\mathbb{C}$ ). Given  $z := a + bi$ , we define the *norm function* on  $\mathbb{C}$  by


$$|z| := \sqrt{a^2 + b^2}.$$

Size actually gives distance.

**Definition 2.11** (Distance on  $\mathbb{C}$ ). Given complex numbers  $z = a + bi$  and  $w = c + di$ , we define the *distance* between  $z$  and  $w$  to be

$$|z - w| = \sqrt{(a - c)^2 + (b - d)^2}.$$

Here are some examples.



One can ask what is the distance between  $0+0i$  and  $1+i$ , and we can compute directly that this is  $\sqrt{1+1} = \sqrt{2}$ . Similarly, the distance between  $1+2i$  and  $1+i$  is  $|(1+2i) - (1+i)| = |i| = 1$ . It should agree with our geometric intuition.

We mentioned complex conjugation is involved here, so we have the following lemma.

**Lemma 2.12.** Fix  $z, w \in \mathbb{C}$ . The following are true.

- (a)  $|z|^2 = z\bar{z}$ .
- (b)  $|\operatorname{Re} z| \leq |z|$  and  $|\operatorname{Im} z| \leq |z|$ .
- (c)  $|z| = |\bar{z}| = |-z|$ .
- (d)  $|z| = 0$  if and only if  $z = 0$ .
- (e)  $|zw| = |z| \cdot |w|$ .

*Proof.* We take these one at a time. Set  $z = a + bi$ .

(a) We have

$$|z|^2 = a^2 + b^2 = (a + bi)(a - bi) = z\bar{z}.$$

Here we have used subtraction of two squares, which one can see when writing  $a^2 + b^2 = a^2 - (ib)^2$ .

(b) We have  $a^2 \leq a^2 + b^2$  and  $b^2 \leq a^2 + b^2$  by the Trivial inequality, so

$$|\operatorname{Re} z| = |a| \leq \sqrt{a^2 + b^2} = |z|,$$

and similarly,

$$|\operatorname{Im} z| = |b| \leq \sqrt{a^2 + b^2} = |z|.$$

(c) Note

$$|\bar{z}| = |a - bi| = \sqrt{a^2 + (-b)^2} = \sqrt{a^2 + b^2} = |z|,$$

and

$$|-z| = |-a - bi| = \sqrt{(-a)^2 + (-b)^2} = \sqrt{a^2 + b^2} = |z|.$$

(d) From (b), we know that  $|\operatorname{Re} z|, |\operatorname{Im} z| \leq |z|$ , but  $|z| = 0$  then forces  $\operatorname{Re} z = \operatorname{Im} z = 0$ , so  $z = 0$ .

(e) From (a), we can write  $|zw|^2 = zw \cdot \overline{zw}$ , which will expand out into

$$z \cdot w \cdot \bar{z} \cdot \bar{w}.$$

We can collect this into  $z\bar{z} \cdot w\bar{w} = |z|^2 |w|^2$ . Thus, by (a) again,  $|zw|^2 = |z|^2 |w|^2$ . But because all norms must be nonnegative real numbers, we may take square roots to conclude  $|zw| = |z| \cdot |w|$ . ■

**Remark 2.13.** Norms are actually more general constructions. For example, the requirement  $|zw| = |z| \cdot |w|$  makes  $|\cdot|$  into a “multiplicative” norm.

To finish off, we actually show that our distance function is good: we show the triangle inequality.

**Lemma 2.14** (Triangle inequality). For every  $x, y, z \in \mathbb{C}$ , we claim

$$|z - x| \leq |z - y| + |y - x|.$$

This claim should be familiar from real analysis. Intuitively, it means that travelling between  $z$  and  $x$  cannot be made into a shorter trip by taking a detour to some other point  $y$  first.

*Proof.* Let  $a := z - y$  and  $b := y - x$  so that  $a + b = z - x$ . Thus, we are showing that

$$|a + b| \stackrel{?}{\leq} |a| + |b|,$$

which is nicer because it only has two letters. For this, because everything is a nonnegative real numbers, it suffices to show the square of this requirement; i.e., we show

$$(|a| + |b|)^2 - |a + b|^2 \stackrel{?}{\geq} 0.$$

Fully expanding, it suffices to show

$$|a|^2 + |b|^2 + 2|a| \cdot |b| - |a + b|^2 \stackrel{?}{\geq} 0.$$

Expanding out  $|w|^2 = w\bar{w}$  for  $w \in \mathbb{C}$ , we are showing

$$a\bar{a} + b\bar{b} + 2|a| \cdot |b| - (a + b)(\bar{a} + \bar{b}) \stackrel{?}{\geq} 0.$$

This is nice because the expansion of the rightmost term will induce some cancellation: it expands into  $a\bar{a} + a\bar{b} + \bar{a}b + b\bar{b}$ , so we are left with showing

$$2|a| \cdot |b| - (a\bar{b} + b\bar{a}) \stackrel{?}{\geq} 0.$$

Note that  $\overline{ab} = \bar{a}\bar{b}$ , so we can collect the final term as  $2\operatorname{Re}(a\bar{b})$ . Similarly, we can write  $|a| \cdot |b| = |a| \cdot |\bar{b}| = |a\bar{b}|$ , so we are showing

$$2|a\bar{b}| - 2\operatorname{Re}(a\bar{b}) \geq 0,$$

which is true because the real part does not exceed the norm. This finishes. ■

## 2.2 January 26

In-person class should start on Monday. Homework #2 will be released on Friday.


### 2.2.1 Geometry on $\mathbb{C}$

So let's try to build a topology on  $\mathbb{C}$  today. We pick up the following definition.


**Definition 2.15** (Convex). A subset  $X \subseteq \mathbb{C}$  is *convex* if and only if, for every  $z, w \in X$  and  $t \in [0, 1]$ , we have that  $w + t(z - w) \in X$ .

Intuitively, “convex” means that  $X$  contains the line segment of any two points in  $X$ .

**Example 2.16.** The circle is convex: any line with endpoints in the circle lives in the circle.



**Non-Example 2.17.** The star-shape is not convex: the given line goes outside the star.



To define our open sets, we will define balls first.

**Definition 2.18** (Open ball). Given some  $z_0 \in \mathbb{C}$ , then *open ball* centered at  $z_0$  with radius  $r > 0$  is

$$B(z_0, r) := \{z \in \mathbb{C} : |z - z_0| < r\}.$$

Observe  $z_0 \in B(z_0, r)$ .

Open balls let us define all sorts of properties.

**Definition 2.19** (Isolated). Fix  $X \subseteq \mathbb{C}$ . A point  $z \in X$  is *isolated* in  $X$  if and only if there exists  $r > 0$  such that

$$B(z, r) \cap X = \{z\}.$$

**Definition 2.20** (Discrete). A subset  $X \subseteq \mathbb{C}$  is *discrete* if and only if every point is isolated.

**Example 2.21.** Any finite subset of  $X \subseteq \mathbb{C}$  is discrete. Namely, any point  $z \in X$  can take

$$r = \frac{1}{2} \min_{w \in X \setminus \{z\}} |z - w|.$$

**Example 2.22.** The subset  $\mathbb{Z} \subseteq \mathbb{C}$  is isolated. Namely, take  $r = \frac{1}{2}$  for any given point.

**Definition 2.23** (Bounded). A subset  $X \subseteq \mathbb{C}$  is *bounded* if and only if there is an  $M$  such that  $X \subseteq B(0, M)$ .



**Example 2.24.** The star from earlier fits into a large circle and is therefore bounded.



And here is our fundamental definition for our topology.

**Definition 2.25** (Open). A subset  $X \subseteq \mathbb{C}$  is *open* if and only if, for each  $z \in X$ , there exists  $r > 0$  such that  $B(z, r) \subseteq X$ .

**Remark 2.26** (Nir). We should probably show that open balls are open; let  $B(z, r)$  be an open ball. Well, for any  $w \in B(z, r)$ , set  $r_w := r - |z - w|$ , which is positive because  $w \in B(z, r)$  requires  $|z - w| < r$ . Now,  $w' \in B(w, r_w)$  implies that  $|w - w'| < r - |z - w|$ , so by the triangle inequality,

$$|z - w'| \leq |z - w| + |w - w'| < r,$$

so  $w' \in B(z, r)$  follows. So indeed, each  $w \in B(z, r)$  has  $B(w, r_w) \subseteq B(z, r)$ .

Open lets us define closed.

**Definition 2.27** (Closed). A subset  $X \subseteq \mathbb{C}$  is *closed* if and only if  $\mathbb{C} \setminus X$  is open.



**Warning 2.28.** Sets are not doors: a set can be both open and closed.

## 2.2.2 Unions and Intersections

Here are some basic properties of our topology.

**Lemma 2.29.** The subsets  $\emptyset$  and  $\mathbb{C}$  are open and closed in  $\mathbb{C}$ .

*Proof.* It suffices to show that  $\emptyset$  and  $\mathbb{C}$  are both open, by definition of closed. That  $\emptyset$  is open holds vacuously because one cannot find any  $z \in \emptyset$  anyways. That  $\mathbb{C}$  is open holds because open balls are subsets of  $\mathbb{C}$ , so any  $z \in \mathbb{C}$  can take  $r = 1$  so that

$$B(z, r) \subseteq \mathbb{C}.$$

So we are done. ■

**Lemma 2.30.** Fixing some  $z \in \mathbb{C}$ , the set  $\{z\}$  is closed.

*Proof.* We show that  $U := \mathbb{C} \setminus \{z\}$  is open. Well, fix any  $w \in U$ , and because  $w \neq z$ , we note  $|z - w| > 0$ , so we set  $r := \frac{1}{2}|z - w|$ . It follows that

$$z \notin B(w, r)$$

because  $|z - w| > r$ . But this is equivalent to  $B(w, r) \subseteq \mathbb{C} \setminus \{z\} = U$ , so we are done. ■

We would like to make new open and closed subsets from old ones. Here is one way to do so.

**Lemma 2.31.** The following are true.

- (a) Arbitrary union: if  $\mathcal{U}$  is any collection of open subsets of  $\mathbb{C}$ , then the union  $\bigcup_{U \in \mathcal{U}} U$  is also open.
- (b) Arbitrary intersection: if  $\mathcal{V}$  is any collection of closed subsets of  $\mathbb{C}$ , then intersection  $\bigcap_{V \in \mathcal{V}} V$  is also closed.

*Proof.* We take these one at a time.

- (a) Fix  $z \in \bigcup_{U \in \mathcal{U}} U$ . We need to show there is some  $r > 0$  such that

$$B(z, r) \stackrel{?}{\subseteq} \bigcup_{U \in \mathcal{U}} U.$$

Well, we know there must be some  $U_z \in \mathcal{U}$  such that  $z \in U_z$  by definition of the union. But now  $U_z$  is open, and therefore we are promised an  $r > 0$  such that

$$B(z, r) \subseteq U_z \subseteq \bigcup_{U \in \mathcal{U}} U,$$

so we are done.

- (b) Fix  $\mathcal{V}$  a collection of closed subsets of  $\mathbb{C}$ . We want to show that

$$\mathbb{C} \setminus \bigcap_{V \in \mathcal{V}} V$$

is open, which by de Morgan's law is equivalent to

$$\bigcup_{V \in \mathcal{V}} (\mathbb{C} \setminus V)$$

being open. However, each  $V \in \mathcal{V}$  is closed, so  $\mathbb{C} \setminus V$  will be open, so we are done by (a). ■

**Lemma 2.32.** The following are true.

- (a) Finite intersection: if  $\{U_k\}_{k=1}^n$  is a finite collection of open subsets of  $\mathbb{C}$ , then the intersection  $\bigcap_{k=1}^n U_k$  is also open.
- (b) Finite union: if  $\{V_k\}_{k=1}^n$  is a finite collection of closed subsets of  $\mathbb{C}$ , then  $\bigcup_{k=1}^n V_k$  is also closed.

*Proof.* We take these one at a time.

- (a) Fix  $z \in \bigcap_{k=1}^n U_k$  so that we need to find  $r > 0$  such that

$$B(z, r) \stackrel{\subseteq}{\subseteq} \bigcup_{k=1}^n U_k.$$

Well,  $z \in U_k$  for each  $k$ , and each  $U_k$  is open, so there is an  $r_k > 0$  such that  $B(z, r_k) \subseteq U_k$ . Thus, we set  $r := \min_k \{r_k\}$ ; because there are only finitely many  $r_k$ , we are assured that  $r > 0$ . Now, we observe that

$$B(z, r) \subseteq B(z, r_k) \subseteq U_k.$$

(Explicitly,  $|w - z| < r$  implies  $|w - z| < r_k$  because  $r \leq r_k$ .) Thus, it follows that

$$B(z, r) \subseteq \bigcap_{k=1}^n U_k,$$

as desired.

(b) We use de Morgan's laws. We want to show that

$$\mathbb{C} \setminus \bigcup_{k=1}^n V_k$$

is open, which by de Morgan's laws is the same thing as showing that

$$\bigcap_{k=1}^n (\mathbb{C} \setminus V_k)$$

is open. However, each  $\mathbb{C} \setminus V_k$  is open by hypothesis on the  $V_k$ , so the full intersection is open by (a). This finishes. ■

**Remark 2.33.** The finiteness is in fact necessary. For example,

$$\bigcap_{n \in \mathbb{N}} B(0, 1/n) = \{0\}.$$

Then one can check that each open ball is open while singletons in  $\mathbb{C}$  are not.

### 2.2.3 Interior, Closure

Let's see more definitions.

**Definition 2.34** (Interior). Given a subset  $X \subseteq \mathbb{C}$ , we define the *interior*  $X^\circ$  of  $X$  to be the union of all open sets contained in  $X$  (which will be open by Lemma 2.31).

**Remark 2.35.** In fact,  $X^\circ$  is the largest open subset of  $X$ , for any open subset  $U_0 \subseteq \mathbb{C}$  contained in  $X$  will have

$$U_0 \subseteq \bigcup_{\text{open } U \subseteq X} U = X^\circ.$$

It follows  $X$  is open if and only if  $X = X^\circ$ : if  $X = X^\circ$ , then  $X$  is open because  $X^\circ$  is open; if  $X$  is open, then  $X$  is the largest open subset of  $\mathbb{C}$  contained in  $X$ , so  $X = X^\circ$ .

**Definition 2.36** (Closure). Given a subset  $X \subseteq \mathbb{C}$ , we define the *closure*  $\overline{X}$  of  $X$  to be the intersection of all closed sets containing  $X$  (which will be closed by Lemma 2.31).

**Remark 2.37.** In fact,  $\overline{X}$  is the smallest closed set containing  $X$ , for any closed subset  $V_0 \subseteq \mathbb{C}$  containing  $X$  will have

$$V_0 \supseteq \bigcap_{\text{closed } V \supseteq X} V = \overline{X}.$$

It follows  $X$  is closed if and only if  $X = \overline{X}$ : if  $X = \overline{X}$ , then  $X$  is closed because  $\overline{X}$  is closed; if  $X$  is closed, then  $X$  is the smallest closed subset of  $\mathbb{C}$  containing  $X$ , so  $X = \overline{X}$ .

By the above definitions, it is not too hard to see that  $X^\circ \subseteq X \subseteq \overline{X}$ .

The interior and closure also let us define the boundary.

**Definition 2.38** (Frontier, boundary). Given a subset  $X \subseteq \mathbb{C}$ , we define the *frontier* or *boundary*  $\partial X$  of  $X$  to be  $\overline{X} \setminus X^\circ$ .

### 2.2.4 Connectivity

**Definition 2.39** (Disconnected). A subset  $X \subseteq \mathbb{C}$  is *disconnected* if and only if there exists nonempty disjoint open subsets  $U_1$  and  $U_2$  such that  $X \subseteq U_1 \cup U_2$  and  $X \cap U_1, X \cap U_2 \neq \emptyset$ . (In other words, the subspace of  $X \subseteq \mathbb{C}$  is (topologically) disconnected.) In this case, we say that  $U_1$  and  $U_2$  *disconnect*  $X$ .

Lastly, we say  $X$  is *connected* if and only if it is not disconnected.

**Example 2.40.** The set  $\emptyset$  is connected because it is impossible for  $U \cap \emptyset \neq \emptyset$  for any open set  $U$  of  $\mathbb{C}$ .

**Example 2.41.** Any singleton  $\{z\}$  is connected. In fact, one cannot decompose  $\{x\}$  into two disjoint sets at all, much less into disjoint sets of the form  $U \cap \{x\}$  with  $U$  open.

**Example 2.42.** Any open ball  $B(z, r)$  is connected. This is surprisingly annoying to check.

**Example 2.43.** The set  $\{1, 2\}$  is disconnected by  $U_1 = B(1, 1/2)$  and  $U_2 = B(2, 1/2)$ .

Connectivity plays nicely with the rest of our definitions as well.

**Lemma 2.44.** A given subset  $X \subseteq \mathbb{C}$  is connected if and only if the only subsets of  $X$  which are both open and closed (in the subspace topology) are  $\emptyset$  and  $X$ .

*Proof.* We take the directions independently. For the forwards direction, take  $X$  connected, and suppose that  $U \subseteq X$  is open and closed. In the subspace topology, we get that  $X \setminus U$  will also be open, and then the subsets  $U$  and  $X \setminus U$  are both open, disjoint and have

$$X = U \cup (X \setminus U).$$

Thus, we require  $U = \emptyset$  or  $X \setminus U = \emptyset$ , so  $U \in \{\emptyset, X\}$ .

We leave the reverse direction as an exercise. Suppose that  $X$  is disconnected, and we show that there is a nonempty proper closed and open subset of  $X$ . Well, because  $X$  is disconnected, we have disjoint open sets  $U_1$  and  $U_2$  of  $\mathbb{C}$  such that  $X \cap U_1, X \cap U_2 \neq \emptyset$  and  $X \subseteq U_1 \cup U_2$ . It follows that

$$X = (U_1 \cap X) \cup (U_2 \cap X). \quad (*)$$

However, now consider the open subset  $U := U_1 \cap X$  of  $X$ . We note that  $(U_1 \cap X) \cap (U_2 \cap X) = \emptyset$ , so by  $(*)$  we see that  $U_1 \cap X = X \setminus (U_2 \cap X)$ , so  $U_1 \cap X$  is closed as well.

To finish, we note that  $U \neq \emptyset$  is nonempty, and its complement is  $X \setminus U = U_2 \cap X$  is also nonempty, so  $U \neq X$  is proper. Thus,  $U$  is a proper nonempty closed and open subset of  $X$ . This finishes. ■

**Remark 2.45** (Nir). It is actually important that the open subsets in the above lemma are in the subspace topology and are not required to be  $\mathbb{C}$ -open. For example,  $X = \{1, 2\}$  is disconnected, but it has no nonempty  $\mathbb{C}$ -open subsets to witness this.

**Lemma 2.46.** Fix  $\mathcal{S}$  a collection of connected subsets of  $\mathbb{C}$ . If  $\bigcap_{S \in \mathcal{S}} S$  is nonempty, then  $\bigcup_{S \in \mathcal{S}} S$  will be connected.

*Proof.* Suppose  $\bigcup_{S \in \mathcal{S}} S$  is contained in the disjoint open subsets  $U_1$  and  $U_2$  of  $\mathbb{C}$ ; we claim  $U_1 \cap (\bigcup_{S \in \mathcal{S}} S) = \emptyset$  or  $U_2 \cap (\bigcup_{S \in \mathcal{S}} S) = \emptyset$ , which will finish.

Pick up some

$$z \in \bigcap_{S \in \mathcal{S}} S,$$

which exists because the intersection is nonempty. Without loss of generality, we may assume that  $z \in U_1$ .

Now,  $z \in S$  for each  $S \in \mathcal{S}$ , so we see  $U_1 \cap S \neq \emptyset$ , so because  $(U_1 \cap S) \cup (U_2 \cap S) = S$ , we see that  $U_2 \cap S = \emptyset$  by hypothesis on  $S$ 's connectivity. Thus, taking the union over the  $U_2 \cap S = \emptyset$ ,

$$U_2 \cap \left( \bigcup_{S \in \mathcal{S}} S \right) = \emptyset,$$

which finishes the proof. ■

**Remark 2.47.** The condition with nonempty intersection is necessary:  $\{0\}$  and  $\{1\}$  are connected, but  $\{0\} \cup \{1\}$  is not.

## 2.3 January 28

Hopefully we'll be in-person on Monday. Homework 2 will be released later today, due next Friday.

### 2.3.1 Sequences

Today we're talking about sequences, building towards a theory of sequences and series. Next week we will begin studying holomorphic functions and actually doing complex analysis.

Anyways, here is a series of definitions.

**Definition 2.48** (Sequence). A *sequence* of complex numbers is a function  $f : \mathbb{N} \rightarrow \mathbb{C}$ . Often we will notate this by  $\{z_n\}_{n \in \mathbb{N}}$  where  $z_n := f(n)$ .

By convention, all of our sequences will be sequences of complex numbers unless otherwise stated.

**Definition 2.49** (Subsequence). A sequence  $\{w_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  is a *subsequence* of a sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  if and only if there is some strictly increasing function  $g : \mathbb{N} \rightarrow \mathbb{N}$  such that  $w_n = z_{g(n)}$ .

**Definition 2.50** (Bounded). A sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  is *bounded* if and only if there exists a positive real number  $M > 0$  such that

$$\{z_n\}_{n \in \mathbb{N}} \subseteq B(0, M).$$

In other words,  $|z_n| < M$  for each  $n \in \mathbb{N}$ .

We are in particular interested in convergence in analysis.

**Definition 2.51** (Converges). A sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  *converges* to some  $z \in \mathbb{C}$  if and only if, for each  $\varepsilon > 0$ , there exists some  $N$  such that  $n > N$  implies

$$|z - z_n| < \varepsilon.$$

We will notate this by  $z_n \rightarrow z$  or  $\lim_{n \rightarrow \infty} z_n = z$ .

Note that the definition of the limit above says that

$$\lim_{n \rightarrow \infty} z_n = z \iff \lim_{n \rightarrow \infty} |z_n - z| = 0.$$

Intuitively, the distance between the  $z_n$  and the  $z$  has to “narrow in” on  $z$ .

We would like some real-analytic tools for our complex analysis. Here is a convergence lemma.

**Lemma 2.52.** Fix  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  a sequence. Then, letting  $z_n := x_n + y_n i$ , we have that  $z_n \rightarrow z$  where  $z = x + y i$  if and only if  $x_n \rightarrow x$  and  $y_n \rightarrow y$ .

*Proof.* This is essentially by definition of the metric on  $\mathbb{C}$ . We take the directions one at a time.

- Suppose that  $z_n \rightarrow z$  in  $\mathbb{C}$ . Then we claim that  $\operatorname{Re} z_n \rightarrow \operatorname{Re} z$  and  $\operatorname{Im} z_n \rightarrow \operatorname{Im} z$  in  $\mathbb{R}$ . Indeed, for any  $\varepsilon > 0$ , there is  $N$  such that

$$n > N \implies |z - z_n| < \varepsilon.$$

But now we see that  $|\operatorname{Re} z_n - \operatorname{Re} z|, |\operatorname{Im} z_n - \operatorname{Im} z| \leq \sqrt{(\operatorname{Re} z_n - \operatorname{Re} z)^2 + (\operatorname{Im} z_n - \operatorname{Im} z)^2}$ , so it follows

$$n > N \implies |\operatorname{Re} z_n - \operatorname{Re} z|, |\operatorname{Im} z_n - \operatorname{Im} z| < \varepsilon,$$

finishing.

- Suppose that  $\operatorname{Re} z_n \rightarrow x$  and  $\operatorname{Im} z_n \rightarrow y$ . We claim that  $z_n \rightarrow x + y i$ . Indeed, for any  $\varepsilon > 0$ , there exists  $N_x$  such that

$$n > N_x \implies |\operatorname{Re} z_n - x| < \varepsilon/2$$

and  $N_y$  such that

$$n > N_y \implies |\operatorname{Im} z_n - y| < \varepsilon/2.$$

It follows that

$$n > \max\{N_x, N_y\} \implies |z_n - (x + y i)| = \sqrt{|\operatorname{Re} z_n - x|^2 + |\operatorname{Im} z_n - y|^2} \leq \sqrt{\left(\frac{\varepsilon}{2}\right)^2 + \left(\frac{\varepsilon}{2}\right)^2} < \varepsilon.$$

This finishes. ■

Essentially, this means that checking convergence of complex numbers is the same as checking real and imaginary parts individually, so we can turn convergence questions into ones from real analysis.

We also have the following basic properties about convergence.

**Proposition 2.53.** Fix  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  a convergent sequence. The following are true.

- (a)  $\{z_n\}_{n \in \mathbb{N}}$  is bounded.
- (b) The limit of  $\{z_n\}_{n \in \mathbb{N}}$  is unique.
- (c) Every subsequence of  $\{z_n\}_{n \in \mathbb{N}}$  converges to  $z$ .

*Proof.* We take the claims one at a time. Let  $z \in \mathbb{C}$  be so that  $z_n \rightarrow z$ .

- (a) Fix  $\varepsilon = 1$  so that there exists  $N$  so that  $n > N$  implies  $|z_n - z| < 1$ . Now set

$$M := \max(\{|z_n| + 1 : n \leq N\} \cup \{|z| + 1\}).$$

We claim that  $|z_n| < M$  for each  $n \in \mathbb{N}$ . We have two cases.

- If  $n \leq N$ , then  $|z_n| < |z_n| + 1 \leq M$ .

- Otherwise,  $n > N$  so that

$$|z_n| \leq |z_n - z| + |z| < |z| + 1 \leq M,$$

so we are done.

- (b) Suppose that  $z_n \rightarrow z'$  for some  $z' \in \mathbb{C}$ , and we show  $z = z'$ . Indeed, if  $z = z'$ , then we are done, so suppose that  $z \neq z'$  so that  $|z - z'| \neq 0$ . Then we set  $\varepsilon := \frac{1}{2}|z - z'| > 0$ , and we are promised some  $N, N'$  such that

$$n > N \implies |z - z_n| < \frac{\varepsilon}{2} \quad \text{and} \quad n > N' \implies |z' - z_n| < \frac{\varepsilon}{2}.$$

In particular, we see that, for  $n > \max\{N, N'\}$ , we have

$$|z - z'| \leq |z - z_n| + |z_n - z'| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon = \frac{1}{2}|z - z'|.$$

But because  $0 \leq |z - z'|$ , we see that this forces  $|z - z'| = 0$ , so  $z = z'$  follows. (Technically we have hit contradiction, but we do not need to use this.)

- (c) Note that subsequences can be characterized by choosing a strictly increasing function  $f : \mathbb{N} \rightarrow \mathbb{N}$  so that we want to show  $z_{f(n)} \rightarrow z$ . Indeed, for any  $\varepsilon > 0$ , we are promised some  $N$  so that

$$n > N \implies |z - z_n| < \varepsilon.$$

Now, for each  $n \in \mathbb{N}$ , we have<sup>1</sup>  $f(n) \geq n$ , so we see that

$$n > N \implies f(n) > N \implies |z - z_{f(n)}| < \varepsilon,$$

which finishes. ■

Sequences themselves have an arithmetic.

**Proposition 2.54.** Fix  $\{z_n\}_{n \in \mathbb{N}}, \{w_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  sequences such that  $z_n \rightarrow z$  and  $w_n \rightarrow w$ . Then the following hold.

- (a)  $z_n + w_n \rightarrow z + w$ .
- (b)  $z_n w_n \rightarrow zw$ .
- (c) If  $w \neq 0$  and  $w_n \neq 0$  for each  $n \in \mathbb{N}$ , then  $\frac{1}{w_n} \rightarrow \frac{1}{w}$ .

*Proof.* We take these one at a time, essentially borrowing the proof from metric spaces.

- (a) Fix some  $\varepsilon > 0$ . We can find some  $N_z$  such that

$$n > N_z \implies |z - z_n| < \varepsilon/2$$

and some  $N_w$  such that

$$n > N_w \implies |w - w_n| < \varepsilon/2.$$

Now, taking  $N := \max\{N_z, N_w\}$  so that the triangle inequality gives

$$n > N \implies |(z + w) - (z_n + w_n)| \leq |z - z_n| + |w - w_n| < \varepsilon,$$

which finishes.

<sup>1</sup> We can show this by induction on  $n$ , for  $f(0) \geq 0$  and  $f(n+1) > f(n)$  forces  $f(n+1) \geq f(n) + 1$ .

- (b) We have to use the fact that the sequences are bounded here. Because  $w_n \rightarrow w$ , the sequence is bounded, so there is an  $M$  so that  $|w_n| < M$  for each  $n \in \mathbb{N}$ . Now, the key inequality is that

$$|z_n w_n - zw| \leq |z_n w_n - zw_n| + |zw_n - zw| \leq M|z_n - z| + |z| \cdot |w_n - w|. \quad (*)$$

So with this in mind, fix any  $\varepsilon > 0$ , and we see that we are promised  $N_z$  such that

$$n > N_z \implies |z_n - z| < \frac{\varepsilon}{2M}$$

and some  $N_w$  such that

$$n > N_w \implies |w_n - w| < \frac{\varepsilon}{2|z|}$$

so that  $(*)$  implies

$$n > \max\{N_z, N_w\} \implies |z_n w_n - zw| < \varepsilon,$$

finishing.

- (c) We begin with some motivating arithmetic. Observe that

$$\left| \frac{1}{w} - \frac{1}{w_n} \right| = \frac{|w_n - w|}{|w w_n|}.$$

We can upper-bound the numerator without tears, so we see the main difficulty is lower-bounding the denominator. Well, because  $w \neq 0$ , we can set  $\varepsilon = |w|/2$  so that there exists  $N_0$  such that

$$n > N_0 \implies |w_n - w| < |w|/2.$$

In particular, it follows that  $|w_n| \geq |w| - |w - w_n| = |w|/2$  for  $n > N_0$ .

With this in mind, fix any  $\varepsilon > 0$ . Then we are promised some  $N_1$  such that

$$n > N_1 \implies |w_n - w| < |w|^2 \varepsilon / 2$$

so that we see

$$n > \max\{N_0, N_1\} \implies \left| \frac{1}{w} - \frac{1}{w_n} \right| = \frac{|w_n - w|}{|w| \cdot |w_n|} \leq \frac{|w|^2 \varepsilon / 2}{|w| \cdot |w|/2} = \varepsilon,$$

finishing. ■

### 2.3.2 Limit Points

Here is our main character.

**Definition 2.55** (Limit point). Fix  $X \subseteq \mathbb{C}$  and some  $z \in \mathbb{C}$ . Then we say that  $z$  is a *limit point* if and only if there exists some sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  such that  $z_n \rightarrow z$ .

**Definition 2.56** (Accumulation point). Fix  $X \subseteq \mathbb{C}$  and some  $z \in \mathbb{C}$ . Then we say that  $z$  is an *accumulation point* if and only if there exists some sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq X \setminus \{z\}$  such that  $z_n \rightarrow z$ .

Essentially accumulation points do not allow isolated points while limit points do.

The above essentially gives a more directly topological definition of “closed set.” It also gives us a more directly topological definition of the closure.



**Lemma 2.57.** Fix  $X \subseteq \mathbb{C}$  and  $z \in \mathbb{C}$ . The following are equivalent.

- (a) We have that  $z \in \overline{X}$ .
- (b) For all  $\varepsilon > 0$ , we have  $B(z, \varepsilon) \cap X \neq \emptyset$ .
- (c) There exists  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  such that  $z_n \rightarrow z$ .

*Proof.* We show our directions one at a time.

- We show (a) implies (b). Suppose  $z \in \overline{X}$ , and for the sake of contradiction suppose we have  $\varepsilon > 0$  such that  $B(z, \varepsilon) \cap X = \emptyset$ . In particular,  $z \notin X$ .

Now,  $z \in \overline{X}$  implies that  $z$  is contained in every closed set containing  $X$  by definition of  $\overline{X}$ . But because  $B(z, \varepsilon)$  is open and is disjoint from  $X$ , we see

$$z \in \overline{X} \subseteq \mathbb{C} \setminus B(z, \varepsilon),$$

which is a contradiction.

- We show (b) implies (c). For each  $n \in \mathbb{N}$ , we know that  $B(z, 1/n) \cap X \neq \emptyset$ , so we find some  $z_n \in B(z, 1/n)$ . Now, for any  $\varepsilon > 0$ , choose  $N := 1/\varepsilon$  so that

$$n > N \implies |z_n - z| < \frac{1}{n} < \frac{1}{N} = \varepsilon,$$

so indeed  $z_n \rightarrow z$ .

- We show (b) implies (a). We proceed by contraposition. Suppose that  $z \notin \overline{X}$ . It follows that  $z \in \mathbb{C} \setminus \overline{X}$ , which is open, so there exists an  $r > 0$  such that

$$B(z, r) \subseteq \mathbb{C} \setminus \overline{X} \subseteq \mathbb{C} \setminus X.$$

It follows that  $B(z, r) \cap X = \emptyset$ .

- We show (c) implies (b). Suppose  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  has  $z_n \rightarrow z$  for some  $z \in \mathbb{C}$ . For any  $\varepsilon > 0$ , there exists  $N$  such that

$$n > N \implies |z_n - z| < \varepsilon,$$

so in particular, choosing any  $n := \lceil N \rceil + 1$  has  $z_n \in B(z, \varepsilon) \cap X$ , so  $B(z, \varepsilon) \cap X \neq \emptyset$ . ■

The above discussion can give us a more directly topological definition of "closed."

**Lemma 2.58.** A subset  $X \subseteq \mathbb{C}$  is closed in  $\mathbb{C}$  if and only if  $X$  contains all of its limit points.

*Proof.* By the previous lemma, we see that  $z \in \overline{X}$  if and only if  $z$  is a limit point of  $X$ , so  $\overline{X}$  is the set of limit points of  $X$ . Now,  $X$  is closed if and only if  $X = \overline{X}$ , so  $X$  is closed if and only if all limit points of  $X$  are in fact points of  $X$ . (Note that all points of  $X$  are automatically limit points essentially because  $X \subseteq \overline{X}$  for free.) ■

While we're here, we can pick up a nice topological result.

**Lemma 2.59.** Fix  $X \subseteq \mathbb{C}$  a connected subset. Then  $\overline{X}$  is also connected.

*Proof.* This argument is purely topological. We proceed by contraposition: suppose  $\overline{X}$  is disconnected by  $U_1, U_2 \subseteq \mathbb{C}$ . We claim that  $U_1, U_2$  disconnect  $X$ . Well, we already know that  $A \subseteq \overline{A} \subseteq U_1 \cup U_2$ , and we already know that  $U_1$  and  $U_2$  are disjoint.

We claim that, for  $U \subseteq \mathbb{C}$  an open subset, if  $U \cap \overline{X} \neq \emptyset$ , then  $U \cap X \neq \emptyset$  as well. Indeed, we proceed by contraposition: if  $U \cap X = \emptyset$ , then  $X \subseteq \mathbb{C} \setminus U$ , but  $\mathbb{C} \setminus U$  is closed, so

$$\overline{X} \subseteq \mathbb{C} \setminus U,$$

so  $\overline{X} \cap U = \emptyset$ .

Thus, it follows from  $U_1 \cap \overline{X}, U_2 \cap \overline{X} \neq \emptyset$  that  $U_1 \cap X, U_2 \cap X \neq \emptyset$ . This finishes the proof that  $U_1$  and  $U_2$  disconnect  $X$ . Indeed, ■

### 2.3.3 Cauchy Sequences

Just like in real analysis, we will be interested in Cauchy sequences.

**Definition 2.60** (Cauchy sequence). A sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  is a *Cauchy sequence* if and only if, for each  $\varepsilon > 0$ , there exists an  $N$  such that

$$n, m > N \implies |z_n - z_m| < \varepsilon.$$

We have the following results on Cauchy sequences.

**Proposition 2.61.** Fix  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  a sequence. If  $\{z_n\}_{n \in \mathbb{N}}$  converges, it is Cauchy.

*Proof.* This proof uses no special properties of  $\mathbb{C}$ . If  $z_n \rightarrow z$ , then for a given  $\varepsilon > 0$ , there exists  $N$  such that

$$n > N \implies |z_n - z| < \varepsilon/2.$$

It follows that

$$n, m > N \implies |z_n - z_m| < |z_n - z| + |z_m - z| < \varepsilon,$$

finishing. ■

**Proposition 2.62.** Every Cauchy sequence in  $\mathbb{C}$  converges.

*Proof.* If  $\{z_n\}_{n \in \mathbb{N}}$  is Cauchy, then we claim  $\{\operatorname{Re} z_n\}_{n \in \mathbb{N}}$  and  $\{\operatorname{Im} z_n\}_{n \in \mathbb{N}}$  are Cauchy sequences. Indeed, for any  $\varepsilon > 0$ , there exists  $N$  so that

$$n, m > N \implies |z_n - z_m| < \varepsilon,$$

but then  $|\operatorname{Re} z_n - \operatorname{Re} z_m| < |z_n - z_m|$  and  $|\operatorname{Im} z_n - \operatorname{Im} z_m| < |z_n - z_m|$ , so the same  $N$  witnesses that  $\{\operatorname{Re} z_n\}_{n \in \mathbb{N}}$  and  $\{\operatorname{Im} z_n\}_{n \in \mathbb{N}}$  are Cauchy in  $\mathbb{R}$ .

Now, Cauchy sequences in  $\mathbb{R}$  converge, so there are reals  $x, y \in \mathbb{R}$  such that  $\operatorname{Re} z_n \rightarrow x$  and  $\operatorname{Im} z_n \rightarrow y$ . It follows that  $z_n \rightarrow x + yi$ , finishing. ■

### 2.3.4 A Little More Topology

We close with one more topological definition.

**Definition 2.63** (Sequentially compact). A subset  $X \subseteq \mathbb{C}$  is *sequentially compact* if and only if every  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  has a convergent subsequence which converges in  $X$ .

**Remark 2.64.** This happens to be equivalent to  $X$  is compact because  $\mathbb{C} \cong \mathbb{R}^2$  satisfies the fact that all compact sets are closed and bounded.

**Example 2.65.** Every finite set is compact.

And here is a last definition.

**Definition 2.66** (Tends to infinity). A sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  *tends to infinity* (notated  $z_n \rightarrow \infty$ ) if and only if each  $M > 0$  has some  $N \in \mathbb{N}$  such that

$$n > N \implies |z_n| > M.$$

Essentially the points of  $\{z_n\}_{n \in \mathbb{N}}$  wander infinitely away.

## 2.4 January 31

So we are lecturing in-person today. Good morning everyone.

**Quote 2.67.** If I don't fall off the stage, I will consider it a major accomplishment.

Homework 2 is due Friday, the 4th of February. Office hours will occur at the usual times, but they will now occur in-person at Evans 749.

### 2.4.1 Series

Today we're mostly talking about series, and on Friday we'll talk about continuous functions.

**Definition 2.68** (Series). An *infinite series* over  $\mathbb{C}$  is an infinite sum

$$S := \sum_{n=1}^{\infty} z_n$$

where  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  is a sequence of complex numbers.

With respect to series, we really want to know when various series converge so that the series is well-defined.

**Definition 2.69** (Converge, diverge). Fix a sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  of complex numbers, we define the  $m$ th partial sum to be

$$S_m := \sum_{n=0}^m z_n.$$

Then we say that the infinite series *converges* if and only if the sequence  $\{S_m\}$  of partial sums converges. Otherwise, we say that  $S$  is *divergent*.

As usual, we start with some basic examples.

**Exercise 2.70.** Fix some  $z \in \mathbb{C}$  with  $|z| < 1$ , we define  $z_n := z^n$ . Then we have

$$S = \sum_{k=0}^{\infty} z^k = \frac{1}{1-z}.$$

*Proof.* Fix some partial sum

$$S_N := \sum_{k=0}^N z^k = 1 + z + z^2 + \cdots + z^N.$$

Multiplying by  $z$ , we see that

$$zS_N = z + z^2 + \cdots + z^N + z^{N+1}.$$

It follows that

$$S_N - zS_N = 1 - z^{N+1}.$$

Because  $|z| < 1$ , we have  $z \neq 1$ , so we may write

$$S_N = \frac{1}{1-z} - \frac{z^{N+1}}{1-z}.$$

However, we may note that as  $N \rightarrow \infty$ , the bad term  $z^{N+1}$  will have size

$$|z^{N+1}| = |z|^{N+1},$$

which goes to 0 (because  $|z| < 1$ ).<sup>2</sup> It follows that

$$\lim_{N \rightarrow \infty} S_N = \frac{1}{1-z},$$

which is what we wanted. ■

Anyways, here are some basic lemmas.

**Lemma 2.71** (Divergence test). Suppose that  $\{z_n\}_{n \in \mathbb{N}}$  is a sequence of complex numbers such that  $\sum z_n$  converges. Then  $z_n \rightarrow 0$  as  $n \rightarrow \infty$ .

*Proof.* Let  $S_n$  be the  $n$ th partial sum so that we are given  $S_n \rightarrow L$  for some  $L \in \mathbb{C}$ . But now we see that

$$z_{n+1} = \left( \sum_{k=0}^{N+1} z_k \right) - \left( \sum_{k=0}^N z_k \right) = S_{n+1} - S_n.$$

Using limit laws, we see that

$$\lim_{n \rightarrow \infty} z_{n+1} = \lim_{n \rightarrow \infty} S_{n+1} - \lim_{n \rightarrow \infty} S_n = L - L = 0.$$

Shifting the indices back gives  $z_n \rightarrow 0$  as  $N \rightarrow \infty$ . ■

Here is an important example of a divergent series.

**Exercise 2.72.** We claim that

$$S = \sum_{k=1}^{\infty} \frac{1}{k}$$

does not converge.

<sup>2</sup> This is surprisingly annoying to rigorize with an  $\varepsilon$ - $\delta$  proof, so we won't do so here. The interested can try to use induction to manually bound  $|z|^n$  by  $\frac{c}{n}$  for some  $c$ .

*Proof.* We will show that the sequence of partial sums  $\{S_n\}_{n=1}^{\infty}$  is not Cauchy, which will show that the series diverges. Well, observe that

$$S_{2^{n+1}} - S_{2^n} = \sum_{k=2^n+1}^{2^{n+1}} \frac{1}{k}$$

after cancelling out all of our terms. However, each term in the sum is at least  $\frac{1}{2^{n+1}}$ , so we bound

$$S_{2^{n+1}} - S_{2^n} \geq \frac{1}{2^{n+1}} (2^{n+1} - 2^n) = \frac{1}{2}.$$

We now show that the partial sums are not Cauchy. Fix  $\varepsilon$ . Supposing for the sake of contradiction that the sequence is Cauchy, there exists  $N$  so that  $n, m > N$  has

$$|S_n - S_m| < \frac{1}{2}.$$

However, we can find some power of 2 named  $2^r$  which exceeds  $N$ , in which case we find  $2^{r+1}, 2^r > N$  and

$$|S_{2^{r+1}} - S_{2^r}| \geq \frac{1}{2},$$

which is our contradiction. ■

**Remark 2.73.** Because a sequence will converge if and only if its real and imaginary parts do, we can turn a convergence test into a real-number test by taking the real and imaginary parts of the sum.

## 2.4.2 The Comparison Test

Recall the comparison test in  $\mathbb{R}$ .

**Theorem 2.74** (Comparison test). Fix  $\{x_n\}_{n \in \mathbb{N}}, \{y_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$  sequences of real numbers. Further, suppose that there exists a positive constant  $c > 0$  such that  $0 \leq x_n \leq cy_n$ . Then the following hold.

- If  $\sum y_n$  converges, then  $\sum x_n$  converges as well.
- If  $\sum x_n$  diverges, then  $\sum y_n$  diverges as well.

*Proof.* We appeal to real analysis. The interested can see Theorem 2.1.21 in Eterović. The main point is to use the Monotone sequence theorem. ■

Here is an example.

**Exercise 2.75.** Fix  $s > 1$  an integer. Then the series

$$S = \sum_{k=1}^{\infty} \frac{1}{k^s}$$

converges.

*Proof.* Because  $s$  is an integer, we have  $s \geq 2$ . Namely,  $\frac{1}{k^s} \leq \frac{1}{k^2}$ , so by the comparison test it suffices to just show the convergence of

$$S' := \sum_{k=1}^{\infty} \frac{1}{k^2}.$$

For this, we apply some trickery. In particular, for  $k > 1$ , we bound

$$\frac{1}{k^2} < \frac{1}{k(k-1)} = \frac{1}{k-1} - \frac{1}{k}.$$

In particular,

$$S' = 1 + \sum_{k=2}^{\infty} \frac{1}{k^2} < 1 + \sum_{k=2}^{\infty} \left( \frac{1}{k-1} - \frac{1}{k} \right).$$

Thus, by the comparison test, it suffices to show the convergence of

$$T := \sum_{k=2}^{\infty} \left( \frac{1}{k-1} - \frac{1}{k} \right).$$

But the  $n$ th partial sum will telescope, giving

$$T_n := \sum_{k=2}^n \left( \frac{1}{k-1} - \frac{1}{k} \right) = 1 - \frac{1}{n},$$

so  $T_n \rightarrow 1$  as  $n \rightarrow \infty$ , and  $T = 1$ . It follows that  $S'$  is upper-bounded by  $1 + T \leq 2$ . ■

### 2.4.3 Absolute Convergence

The following kind of convergence is nontrivially stronger, but that makes it better.

**Definition 2.76** (Absolute convergence). Fix a sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  of complex numbers. Then the sum  $S := \sum z_n$  converges *absolutely* if and only if the series

$$\sum_{n=0}^{\infty} |z_n|$$

also converges. In other words, the partial sums of the above series converges.

We have the following quick lemma to justify naming this “convergence.”

**Lemma 2.77.** If a series converges absolutely, then the series also converges.

*Proof.* The idea is to use the triangle inequality. Fix our series

$$S := \sum_{n=0}^{\infty} z_n$$

for which

$$T := \sum_{n=0}^{\infty} |z_n|$$

converges. Let  $S_n$  be the  $n$ th partial sum of  $S$  and  $T_n$  the  $n$ th partial sum of  $T$ .

Our goal is to show that  $\{S_n\}_{n \in \mathbb{N}}$  is Cauchy. Observe  $\{T_n\}_{n \in \mathbb{N}}$  is an increasing sequence of real numbers because  $|z| \geq 0$  always. To start off our arithmetic, we note that, for  $n, m \in \mathbb{N}$  with  $n > m$ , we have

$$|S_n - S_m| = \left| \sum_{k=m+1}^n z_k \right|,$$

which by the triangle inequality can be bounded by

$$|S_n - S_m| \leq \sum_{k=m+1}^n |z_k| = T_m - T_n.$$

But now we can use the fact that  $\{T_n\}_{n \in \mathbb{N}}$  must be Cauchy to finish: for any  $\varepsilon > 0$ , there exists some  $N$  such that  $n > m > N$  implies  $T_m - T_n < \varepsilon$ . But then this same  $N$  promises  $n > m > N$  implies

$$|S_n - S_m| < T_m - T_n < \varepsilon,$$

which is what we wanted. ■

Here is a surprise tool that will help us later.

**Lemma 2.78.** Fix a sequence  $\{a_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  of nonzero complex numbers. Further, suppose that the sequence  $\{a_n\}_{n \in \mathbb{N}}$  tends to infinity (i.e.,  $|a_n| \rightarrow \infty$  as  $n \rightarrow \infty$ ), then for any positive real number  $r \in \mathbb{R}^+$ , the series

$$\sum_{k=0}^{\infty} \left( \frac{r}{|a_k|} \right)^k$$

converges.

*Proof.* We need the  $a_n$  to be nonzero in order to allow division, so the real puzzle is to determine how to use the fact  $|a_n| \rightarrow \infty$ . Well, there exists some  $N$  such that  $n > N$  has

$$|a_n| > 2r.$$

But then  $\frac{r}{|a_n|} < \frac{1}{2}$  for each  $n > N$ , so we can use the comparison test as follows: observe that

$$\sum_{k=0}^{\infty} \frac{1}{2^k}$$

will converge, and there will exist some  $c > 1$  so that

$$\frac{r}{|a_k|} < \frac{c}{2^k}$$

for  $0 \leq k \leq N$ ; and then for  $n > N$ , we get the above inequality anyways as discussed earlier (observe we took  $c > 1$ ). ■

**Quote 2.79.** I can't break math on the first day of class. I can do it later on.

**Lemma 2.80.** Suppose that we have two series  $S := \sum_{k \in \mathbb{N}} z_k$  and  $T := \sum_{k \in \mathbb{N}} w_k$  are both absolutely convergent. Then the sum

$$P := \sum_{k=0}^{\infty} \left( \sum_{i+j=k} z_i w_j \right)$$

is absolutely convergent as well. In fact,  $P$  will converge to  $ST$ .

*Proof.* We sketch the result, and the remaining details are in Eterović. As usual, consider the partial sums

$$A_n := \sum_{k=0}^n |z_k| \quad \text{and} \quad B_n = \sum_{k=0}^n |w_k|,$$

both of which will converge as  $n \rightarrow \infty$ . Brazenly multiplying these together, we see that

$$A_n B_n = \sum_{i,j=0}^n |z_i w_j| = \sum_{k=0}^n \sum_{\substack{i+j=k \\ 0 \leq i,j \leq n}} |z_i w_j| + \sum_{k>n} \sum_{\substack{i+j=k \\ 0 \leq i,j \leq n}} |z_i w_j|.$$

In the first sum, observe that any time  $i + j = k$ , we will automatically have  $i, j \leq k \leq n$ . It follows that

$$A_n B_n = \sum_{k=0}^n \left( \sum_{i+j=k} z_i w_j \right) + \underbrace{\sum_{\substack{i+j>n \\ 0 \leq i,j \leq n}} |z_i w_j|}_{R_n}.$$

The key claim is that  $R_n \rightarrow 0$ . The main idea is that  $i + j > n$  implies that  $i \geq n/2$  or  $j \geq n/2$ , so we can write

$$|R_n| \leq \sum_{i=0}^n \sum_{j=n/2}^n |z_i w_j| + \sum_{i=n/2}^n \sum_{j=0}^n |z_i w_j| = \left( \sum_{i=0}^n |z_i| \right) \left( \sum_{j=n/2}^n |w_j| \right) + \left( \sum_{i=n/2}^n |z_i| \right) \left( \sum_{j=0}^n |w_j| \right).$$

Now, fix any  $\varepsilon > 0$ , and we show there exists  $X$  so that  $n > X$  has  $|R_n| < \varepsilon$ . Note  $A := \sum |z_k|$  and  $B := \sum |w_k|$  both converge and hence have Cauchy partial sums. Because the partial sums are increasing, we bound

$$|R_n| \leq A \left( \sum_{j=n/2}^n |w_j| \right) + B \left( \sum_{i=n/2}^n |z_i| \right)$$

So there exists  $N$  such that  $n > m > N$  has

$$\sum_{i=m+1}^n |z_i| < \frac{\varepsilon}{2B}$$

Similarly there exists  $M$  so that  $n > m > M$  has

$$\sum_{j=m+1}^n |w_j| < \frac{\varepsilon}{2A},$$

from which it follows that  $n > n/2 > \max\{N, M\}$  will have

$$|R_n| \leq A \cdot \frac{\varepsilon}{2A} + B \cdot \frac{\varepsilon}{2B} = \varepsilon,$$

which finishes.

Now, because  $R_n \rightarrow 0$ , we see

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \left( \sum_{i+j=k} |z_i| \cdot |w_j| \right) = \lim_{n \rightarrow \infty} A_n B_n - \lim_{n \rightarrow \infty} R_n,$$

which does indeed converge, so indeed the series

$$\sum_{k=0}^{\infty} \left( \sum_{i+j=k} |z_i| \cdot |w_j| \right)$$

will converge. By the comparison test (using the triangle inequality), it follows that

$$P = \sum_{k=0}^{\infty} \left( \sum_{i+j=k} z_i w_j \right)$$



will also absolutely converge.

To show that  $P$  converges to  $ST$ , we observe that the difference of the  $n$ th partial sum is

$$P_n - S_n T_n = \sum_{k=0}^n \left( \sum_{i+j=k} z_i w_j \right) - \sum_{i,j=0}^n z_i w_j = \sum_{k=0}^n \left( \sum_{i+j=k} z_i w_j \right) - \sum_{k=0}^n \left( \sum_{i+j=k}^n z_i w_j \right) + \sum_{\substack{0 \leq i,j \leq n \\ i+j < n}} z_i w_j,$$

so

$$P_n - S_n T_n = \sum_{\substack{0 \leq i,j \leq n \\ i+j < n}} z_i w_j.$$

But by the triangle inequality, we see  $|P_n - S_n T_n| \leq R_n$ , so  $P_n - S_n T_n \rightarrow 0$  as  $n \rightarrow \infty$ . It follows  $P_n$  and  $S_n T_n$  have the same limit as  $n \rightarrow \infty$  (which exists because  $S_n$  and  $T_n$  have a limit). So indeed,  $P = ST$ . ■

## 2.5 February 2

Good morning everyone. Here is some house-keeping.

- Homework #2 is due on Friday at 11:59, on GradeScope. The assignment has just been added.
- There are office hours to talk about the homework. Please come if you have questions.

### 2.5.1 Summation Review

Today we finish our discussion of series, for now. We quickly recall the definitions.

**Definition 2.81** (Series). An *infinite series* over  $\mathbb{C}$  is an infinite sum

$$S := \sum_{n=1}^{\infty} z_n$$

where  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  is a sequence of complex numbers.

**Definition 2.82** (Converge, diverge). Fix a sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  of complex numbers, we define the  $m$ th partial sum to be

$$S_m := \sum_{n=0}^m z_n.$$

Then we say that the infinite series *converges* if and only if the sequence  $\{S_m\}$  of partial sums converges. Otherwise, we say that  $S$  is *divergent*.

Today we are building towards proving Dirichlet's convergence theorem. We pick up the following lemmas.

**Lemma 2.83.** Fix sequences  $\{z_{k,\ell}\}_{k,\ell \in \mathbb{N}}$  a collection of complex numbers satisfying the following conditions.

- Fixing any  $k$ , the sum  $\sum_{\ell=0}^{\infty} |z_{k,\ell}|$  converges.
- The sum  $\sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}|$  converges.

Then the following are true.

- (a) Fix any  $\ell$ , the sum  $\sum_{k=0}^{\infty} |z_{k,\ell}|$  converges; i.e., the terms in the left sum below are well-defined.
- (b) We have that

$$\sum_{\ell=0}^{\infty} \sum_{k=0}^{\infty} z_{k,\ell} = \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} z_{k,\ell},$$

and both sums converge.

Intuitively, the first condition is requiring that the series horizontally does not grow too fast. The second condition is requiring an absolute convergence condition. Then the first conclusion says we can sum vertically instead, and the second conclusion says that we can move around the order of summation.

*Proof.* We will sketch this proof; we prove (a) and (b) more or less simultaneously. To turn the infinite double sum into something we can consider finite partial sums of, we set, for each natural  $N$ ,

$$S_n := \sum_{k=0}^n \sum_{\ell=0}^n |z_{k,\ell}|.$$

The main claim is that

$$\lim_{n \rightarrow \infty} S_n \stackrel{?}{=} \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}|.$$

Indeed, fix any  $\varepsilon > 0$ . Because the latter sum converges, there exists some natural  $A$  such that

$$\sum_{k>A} \sum_{\ell=0}^{\infty} |z_{k,\ell}| < \frac{\varepsilon}{2}.$$

Further, there exists some natural  $B_k$  such that

$$\sum_{\ell>B_k} |z_{k,\ell}| < \frac{\varepsilon}{2A}$$

for each  $k \in \mathbb{N}$ . Take  $B := \max_{0 \leq k < A} B_k$ . Now, we set  $N := \max\{A, B\}$ . To start off our inequalities, we note that

$$S_n = \sum_{k=0}^n \sum_{\ell=0}^n |z_{k,\ell}| \leq \sum_{k=0}^n \sum_{\ell=0}^{\infty} |z_{k,\ell}| \leq \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}|,$$

so we know the sign of our difference. In particular, for any  $n > N$ , we see that

$$S_n = \sum_{k=0}^N \sum_{\ell=0}^N |z_{k,\ell}| \geq \sum_{k=0}^K \sum_{\ell=0}^L |z_{k,\ell}|.$$

Thus,

$$0 \leq \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - S_n \leq \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - \sum_{k=0}^K \sum_{\ell=0}^L |z_{k,\ell}| = \sum_{k>K} \sum_{\ell=0}^{\infty} |z_{k,\ell}| + \sum_{k=0}^K \sum_{\ell>L} |z_{k,\ell}|$$

after some cancellation. But we can upper-bound the last quantity by  $\frac{\varepsilon}{2} + K \cdot \frac{\varepsilon}{2K} = \varepsilon$ , so we are done.

The main point of the above lemma is that we are told each  $\varepsilon > 0$  has some  $N$  so that  $n > N$  implies

$$\sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - S_n = \sum_{\substack{(k,\ell) \in \mathbb{Z}^2 \\ k > n \text{ or } \ell > n}} |z_{k,\ell}| < \varepsilon.$$

We now take the two parts in sequence.

- (a) Fix an index  $\ell'$ ; we show absolute convergence by showing that the partial sums of  $\sum_{k=0}^{\infty} |z_{k,\ell'}|$  are Cauchy. Indeed, fix some  $\varepsilon > 0$ , and we know there exists  $N$  so that each  $n > N$  has

$$\sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| - S_n < \varepsilon.$$

Now, we see that any  $n > m > N$  will have

$$\sum_{k=m+1}^n |z_{k,\ell'}| \leq \sum_{k=m+1}^N \sum_{\ell=0}^{\infty} |z_{k,\ell}| \leq \sum_{k=N+1}^{\infty} \sum_{\ell=0}^{\infty} |z_{k,\ell}| + \sum_{k=0}^N \sum_{\ell=N+1}^{\infty} |z_{k,\ell}| < \varepsilon,$$

so we are done.

- (b) As above, fix some  $\varepsilon > 0$ , and we are promised  $N$  so that

$$\sum_{\substack{(k,\ell) \in \mathbb{Z}^2 \\ k > N \text{ or } \ell > N}} |z_{k,\ell}| < \varepsilon/2.$$

Observe, for  $K, L > N$ , we have by the triangle inequality that

$$\left| \sum_{\ell=0}^L \sum_{k=0}^K z_{k,\ell} - S_N \right| < \varepsilon/2.$$

This bounds holds for any  $K$ , so we can send  $K$  arbitrarily large; that inner sum will converge, so in fact we can send  $K$  to  $\infty$  without ill effect. (Formally, the inner term is an increasing sequence bounded above, so it will converge as  $K \rightarrow \infty$ .) This gives

$$\left| \sum_{\ell=0}^L \sum_{k=0}^{\infty} z_{k,\ell} - S_N \right| \leq \varepsilon/2.$$

Again, the inner term is an increasing sequence as  $L \rightarrow \infty$  but still bounded above as  $\varepsilon/2$ , so the inner sum will converge as  $L \rightarrow \infty$  and still give the inequality

$$\left| \sum_{\ell=0}^{\infty} \sum_{k=0}^{\infty} z_{k,\ell} - S_N \right| < \varepsilon.$$

Now as we send  $\varepsilon \rightarrow 0$ , we see that  $\lim_{N \rightarrow \infty} S_N = \sum_{\ell=0}^{\infty} \sum_{k=0}^{\infty} z_{k,\ell}$ , which finishes. ■

### 2.5.2 Dirichlet Test

We now go directly for the Dirichlet test for convergence.

**Lemma 2.84** (Summation by parts). Fix sequences  $\{a_n\}_{n \in \mathbb{N}}$  and  $\{b_n\}_{n \in \mathbb{N}}$  sequences of complex numbers. Then we define

$$B_n := \sum_{k=0}^n b_k,$$

and  $B_{-1} = 0$  to be the empty sum. It follows that, for any  $n, m \in \mathbb{N}$  with  $n > m$ ,

$$\sum_{k=m}^n a_k b_k = a_n B_n - a_m B_{m-1} + \sum_{k=m}^{n-1} B_k (a_k - a_{k+1}).$$

*Proof.* This comes down to a direct computation. The main point is that  $b_k = B_k - B_{k-1}$ , which even works with  $k = 0$ . Indeed,

$$\begin{aligned}
 \sum_{k=m}^n a_k b_k &= \sum_{k=m}^n a_k (B_k - B_{k-1}) \\
 &= \sum_{k=m}^n a_k B_k - \sum_{k=m}^n a_k B_{k-1} \\
 &\stackrel{*}{=} a_n B_n + \sum_{k=m}^{n-1} a_k B_k - a_m B_{m-1} - \sum_{k=m}^n a_{k+1} B_k \\
 &= a_n B_n - a_m B_{m-1} + \sum_{k=m}^{n-1} B_k (a_k - a_{k+1}),
 \end{aligned}$$

which is what we wanted. The important step to pay attention to is the rearrangement we did in  $\stackrel{*}{=}$  in order to message the sums together. ■

And here is our theorem.

**Theorem 2.85** (Dirichlet's test). Fix  $\{a_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$  a sequence of real numbers and  $\{b_n\}_{n \in \mathbb{N}} \subseteq \mathbb{C}$  a sequence of complex numbers satisfying the following conditions.

- The sequence  $\{a_n\}_{n \in \mathbb{N}}$  is decreasing.
- We have  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ .
- Bounded partial sums: there exists a positive real number  $M$  such that

$$\left| \sum_{k=0}^n b_k \right| < M$$

for each  $n \in \mathbb{N}$ .

Then we claim that

$$\sum_{k=0}^{\infty} a_k b_k$$

converges.

*Proof.* As usual, fix our partial sums

$$S_n := \sum_{k=0}^n a_k b_k \quad \text{and} \quad B_n := \sum_{k=0}^n b_k.$$

We are given that the  $B_k$  are bounded, so we are going to want to use Lemma 2.84, which tells us that

$$S_n = a_n B_n + \sum_{k=0}^{n-1} B_k (a_k - a_{k+1}).$$

We examine the convergence of these terms individually.

- For the sum, we will show that it absolutely converges. We are given that the partial sums  $B_n$  are bounded by  $M$ , so we note  $|B_k (a_k - a_{k+1})| < M |a_k - a_{k+1}|$ , so it suffices to show that

$$M \sum_{k=0}^{n-1} |a_k - a_{k+1}|$$

converges as  $n \rightarrow \infty$ . It would be great if this would telescope, and indeed it does! Because the  $a_k$  are decreasing,

$$\sum_{k=0}^{\infty} |a_k - a_{k+1}| = \sum_{k=0}^{\infty} (a_k - a_{k+1}) = a_0 - a_{n+1}.$$

Because  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ , we see that this sum will converge to  $a_0$ . It follows that

$$\sum_{k=0}^{\infty} |B_k(a_k - a_{k+1})|$$

will converge by the Comparison test, so

$$\sum_{k=0}^{\infty} B_k(a_k - a_{k+1})$$

converges by absolute convergence.

- Note that the  $B_n$  are bounded in norm by  $M$ , so  $|a_n B_n| \leq M|a_n|$ , but  $|a_n| \rightarrow 0$  as  $n \rightarrow \infty$ , so  $|a_n B_n| \rightarrow 0$ . ■

Eterović has lots of different convergence tests in his notes, but we don't care about most of them. Here is one that we do care about.

**Theorem 2.86** (Integral test). Fix a decreasing function  $f : [1, \infty) \rightarrow \mathbb{R}^+$  and for which

$$\int_k^{k+1} f(x) dx$$

always exists. Then the sequence of integrals  $I_n := \int_1^n f(x) dx$  converges if and only if the summation

$$\sum_{k=1}^{\infty} f(k)$$

converges.

*Proof.* We omit this proof; it's a reasonably standard real-analytic test. ■

## 2.6 February 4

Today we are talking about continuity.



**Warning 2.87.** The first half of this lecture was transcribed from Professor Morrow's notes because I had to miss class for a job interview

### 2.6.1 Limits

Before defining continuity, we have the following definitions.

**Definition 2.88** (Limit). Fix  $f : X \rightarrow \mathbb{C}$  a function and  $z_0 \in \overline{X}$ . Then we say *the limit of  $f(z)$  as  $z$  approaches  $z_0$  equals  $w$* , denoted

$$\lim_{z \rightarrow z_0} f(z) = w,$$

if and only if, for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$|z - z_0| < \delta \implies |f(z) - w| < \varepsilon$$

for  $z \in X$ .

This is the standard  $\varepsilon$ - $\delta$  definition.

We also pick up the following convention as a surprise tool that may help us later.

**Definition 2.89** (Infinite limits). Fix  $f : X \rightarrow \mathbb{C}$  a function. Then we say *the limit of  $f(z)$  as  $z$  tends to infinity equals  $w$* , denoted

$$\lim_{z \rightarrow \infty} f(z) = w,$$

if and only if, for each  $\varepsilon > 0$ , there exists  $N > 0$  such that

$$|z| > N \implies |f(z) - w| < \varepsilon$$

for  $z \in X$ .

As in real analysis, the  $\varepsilon$ - $\delta$  definition of a limit can be translated to a statement about sequences.

**Proposition 2.90.** Fix  $\alpha \in \overline{X}$ . Then  $\lim_{z \rightarrow \alpha} f(z) = w$  if and only if, for each  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  such that  $z_n \rightarrow \alpha$  as  $n \rightarrow \infty$ , we have  $f(z_n) \rightarrow w$  as  $n \rightarrow \infty$ .

*Proof.* In the forwards direction, fix  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  such that  $z_n \rightarrow \alpha$ , and we show that  $f(z_n) \rightarrow w$ . Well, for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$|z - \alpha| < \delta \implies |f(z) - f(\alpha)| < \varepsilon,$$

where  $z \in X$ . But for this  $\delta > 0$ , there exists  $N$  such that

$$n > N \implies |z_n - \alpha| < \delta \implies |f(z_n) - f(\alpha)| < \varepsilon.$$

So indeed,  $f(z_n) \rightarrow f(\alpha)$ .

In the reverse direction, suppose that  $f(z)$  does not approach  $w$  as  $z \rightarrow \alpha$ . Then, there exists  $\varepsilon_0 > 0$  such that, for any  $\delta > 0$ , there is  $z \in X$  such that  $|z - \alpha| < \delta$  while  $|f(z) - w| > \varepsilon_0$ . Well, for any  $n \in \mathbb{N}$ , taking  $\delta = 1/(n+1)$ , we are promised  $z_n \in X$  such that

$$|z_n - \alpha| < \frac{1}{n+1} \text{ and } |f(z_n) - w| > \varepsilon_0.$$

So to finish, we claim that  $z_n \rightarrow \alpha$  as  $n \rightarrow \infty$ , but  $f(z_n)$  does not approach  $w$  as  $n \rightarrow \infty$ .

- For any  $\varepsilon > 0$ , we note that  $N := 1/\varepsilon$  has  $n > N$  implies

$$|z_n - \alpha| < \frac{1}{n+1} < \frac{1}{N+1} < \frac{1}{N} = \varepsilon,$$

so indeed  $z_n \rightarrow \alpha$  as  $N \rightarrow \infty$ .

- We note that  $\varepsilon_0 > 0$  satisfies that

$$|f(z_n) - w| > \varepsilon_0$$

for any  $n \in \mathbb{N}$ , so no  $N$  will have  $n > N$  implies  $|f(z_n) - w| < \varepsilon_0$ . Thus,  $f(z_n)$  does not approach  $w$  as  $n \rightarrow \infty$ .

The sequence  $\{z_n\}_{n \in \mathbb{N}}$  now completes the proof by showing the reverse direction by contraposition. ■

While we're here, we pick up the following definitions.

**Definition 2.91** (Bounded). A function  $f : X \rightarrow \mathbb{C}$  is *bounded* if there exists  $R > 0$  such that  $\text{im } f \subseteq B(0, R)$ .

**Definition 2.92** (Bounded near). Fix a nonempty open subset  $\Omega \subseteq \mathbb{C}$  and  $z_0 \in \Omega$ . Then  $f : \Omega \setminus \{z_0\} \rightarrow \mathbb{C}$  is *bounded near*  $z_0$  if and only if

$$\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0.$$

## 2.6.2 Continuity

And here is our central definition for today.

**Definition 2.93** (Continuous). A function  $f : X \rightarrow \mathbb{C}$  is *continuous at*  $z_0 \in X$  if and only if, for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$|z - z_0| < \delta \implies |f(z) - f(z_0)| < \varepsilon,$$

where  $z \in X$ . Further,  $f$  is *continuous on*  $X$  if and only if  $f$  is continuous at each  $z_0 \in X$ .

We have the following lemma of equivalent definitions.

**Lemma 2.94.** Suppose that  $f : X \rightarrow \mathbb{C}$ .

- (a) Then  $f$  is continuous at  $w$  if and only if every sequence  $\{z_n\} \subseteq X$  such that  $z_n \rightarrow z$  implies  $f(z_n) \rightarrow f(z)$ .
- (b) We have that  $f$  is continuous on  $X$  if and only if every open set  $U \subseteq \mathbb{C}$  has  $f^{-1}(U)$  open in  $X$ .
- (c) We have that  $f$  is continuous on  $X$  if and only if each closed set  $V \subseteq X$  has  $f^{-1}(V)$  closed in  $X$ .
- (d) Lastly, we have that  $f$  is continuous at if and only if, for each  $\varepsilon > 0$  and  $z \in \mathbb{C}$ , we have that  $f^{-1}(B(z, \varepsilon))$  is open in  $X$ .

*Proof.* We take the parts one at a time.

- (a) We could use Proposition 2.90, but we will just do this by hand. For the forwards direction, suppose that  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  converges to some  $w$ . Then let  $\varepsilon > 0$ . By assumption, there exists some  $\delta > 0$  such that

$$|z - w| < \delta \implies |f(z) - f(w)| < \varepsilon.$$

It follows from  $z_n \rightarrow w$  that there exists some  $N$  such that

$$n > N \implies |z_n - w| < \delta \implies |f(z_n) - f(w)| < \varepsilon,$$

so it follows that  $f(z_n) \rightarrow f(w)$ .

In the reverse direction, take  $f$  not continuous at  $w$ , so there exists  $\varepsilon > 0$  so that for all  $n \in \mathbb{N}$ , there exists some chosen  $z_n$  with

$$|z_n - w| < \delta \implies |f(z_n) - f(w)| > \varepsilon.$$

But as  $z_n \rightarrow w$ , we see that  $f(z_n)$  does not approach  $f(w)$ , so we are done.

- (b) In the forwards direction, suppose that  $U \subseteq \mathbb{C}$  is open, and we show that  $f^{-1}(U)$  is open in  $X$ . Well, suppose that  $z \in f^{-1}(U)$ , and we will find  $\delta > 0$  such that  $B(z, \delta) \subseteq f^{-1}(U)$ .

Well,  $f(z) \in U$ , so there exists  $\varepsilon > 0$  such that  $B(f(z), \varepsilon) \subseteq U$ . Thus, continuity of  $f$  requires some  $\delta > 0$  such that

$$|w - z| < \delta \implies |f(w) - f(z)| < \varepsilon,$$

which implies  $f(w) \in B(f(z), \varepsilon) \subseteq U$  implies  $w \in f^{-1}(U)$ . So indeed,  $B(z, \delta) \subseteq f^{-1}(U)$ .

In the reverse direction, suppose that each open  $U \subseteq \mathbb{C}$  has  $f^{-1}(U)$  is open. Now fix any  $z \in X$  and  $\varepsilon > 0$ . The set  $B(f(z), \varepsilon)$  is open, so

$$f^{-1}(B(f(z), \varepsilon))$$

is open. But  $z \in f^{-1}(B(f(z), \varepsilon))$ , so we can find  $\delta > 0$  such that  $B(z, \delta) \subseteq f^{-1}(B(f(z), \varepsilon))$ . Thus,  $|w - z| < \delta$  implies  $w \in f^{-1}(B(f(z), \varepsilon))$  implies  $f(w) \in B(f(z), \varepsilon)$  implies  $|f(w) - f(z)| < \varepsilon$ , finishing.

- (c) In the forwards direction, suppose  $f$  is continuous so that  $U \subseteq \mathbb{C}$  open implies  $f^{-1}(U) \subseteq X$  is open. But then, if  $V \subseteq \mathbb{C}$  is closed, then  $\mathbb{C} \setminus V$  is open, so<sup>3</sup>

$$f^{-1}(\mathbb{C} \setminus V) = f^{-1}(\mathbb{C}) \setminus f^{-1}(V) = X \setminus f^{-1}(V)$$

is open, so  $f^{-1}(V)$  is closed.

In the backwards direction, suppose  $f^{-1}$  preserves closed sets. Then, if  $U \subseteq \mathbb{C}$  is open,  $\mathbb{C} \setminus U$  is closed, so

$$f^{-1}(\mathbb{C} \setminus U) = f^{-1}(\mathbb{C}) \setminus f^{-1}(U) = X \setminus f^{-1}(U)$$

is closed, so  $f^{-1}(U)$  is open. Thus,  $f^{-1}$  preserves open sets, so  $f$  must be continuous.

- (d) In the forwards direction, fix  $\varepsilon > 0$  and  $z \in \mathbb{C}$ , so  $B(z, \varepsilon)$  is open, so  $f^{-1}(B(z, \varepsilon))$  is open in  $X$ , finishing. In the other direction fix  $\varepsilon > 0$  and  $z \in \mathbb{C}$  to consider  $B(f(z), \varepsilon) \subseteq U$ . Thus, continuity of  $f$  requires some  $\delta > 0$  such that

$$|w - z| < \delta \implies |f(w) - f(z)| < \varepsilon,$$

which implies  $f(w) \in B(f(z), \varepsilon) \subseteq U$  implies  $w \in f^{-1}(U)$ . So indeed,  $B(z, \delta) \subseteq f^{-1}(U)$ .

In the reverse direction, fix  $U$  open, and we show that  $f^{-1}(U)$  is open. Well, each  $z \in U$  has some  $\varepsilon_z$  such that  $B(z, \varepsilon_z) \subseteq U$ . But  $f^{-1}(B(z, \varepsilon_z))$  is open by hypothesis, so

$$f^{-1}(U) = f^{-1}\left(\bigcup_{z \in U} B(z, \varepsilon_z)\right) = \bigcup_{z \in U} f^{-1}(B(z, \varepsilon_z))$$

is an arbitrary union of open sets and hence open. ■

And here are some special examples.

**Example 2.95.** Fix some  $z_0 \in \mathbb{C}$ . Then  $f(z) := |z - z_0|$  is continuous on  $\mathbb{C}$ . Indeed, fix any  $w \in \mathbb{C}$ . Then for any  $\varepsilon > 0$ , we set  $\delta := \varepsilon$  so that  $|z - w| < \delta$  implies

$$|f(z) - f(w)| = \left| |z - z_0| - |w - z_0| \right| \leq |z - w| < \delta = \varepsilon.$$

**Example 2.96.** The functions  $\operatorname{Re}$  and  $\operatorname{Im}$  is continuous. Indeed, fix any  $w \in \mathbb{C}$ . Then, for any  $\varepsilon > 0$ , take  $\delta := \varepsilon$  so that  $|z - w| < \delta$  implies

$$|\operatorname{Re} z - \operatorname{Re} w| = |\operatorname{Re}(z - w)| \leq |z - w| < \delta = \varepsilon,$$

and similarly,

$$|\operatorname{Im} z - \operatorname{Im} w| = |\operatorname{Im}(z - w)| \leq |z - w| < \delta = \varepsilon,$$

Continuous functions also have some arithmetic.

<sup>3</sup> To see  $f^{-1}(A \setminus B) = f^{-1}(A) \setminus f^{-1}(B)$ , note that  $x \in f^{-1}(A \setminus B)$  if and only if  $f(x) \in A \setminus B$  if and only if  $f(x) \in A$  but  $f(x) \notin B$  if and only if  $x \in f^{-1}(A)$  but  $x \notin f^{-1}(B)$ .



**Proposition 2.97.** Fix  $f, g : X \rightarrow \mathbb{C}$  to functions continuous at  $z_0 \in X$ . Then  $f + g, f \cdot g$  are both continuous at  $z_0 \in X$ , and  $f/g$  is continuous at  $z_0$  provided  $g(z_0) \neq 0$ .

*Proof.* The point is to appeal to the corresponding results on convergence of sequences. In particular, we use the idea that  $f$  is continuous at  $z_0$  if and only if each sequence  $z_n \rightarrow z_0$  in  $X$  has  $f(z_n) \rightarrow f(z_0)$ . We omit the details because they are essentially the same as in a real analysis class. ■

**Corollary 2.98.** Every polynomial in one variable is a continuous function  $X \rightarrow \mathbb{C}$  for any  $X \subseteq \mathbb{C}$ .

*Proof.* Note that  $x \mapsto x$  is continuous, so by induction  $x \mapsto x^n$  is continuous for each  $n \in \mathbb{N}$ . Taking a  $\mathbb{C}$ -linear combination gives arbitrary polynomials. ■

Here is another sort of arithmetic.

**Lemma 2.99.** The composition of two continuous functions is continuous.

*Proof.* Omitted. ■

### 2.6.3 Connectedness

We want to build towards a particular type of continuous function.

**Proposition 2.100.** Fix  $X \subseteq \mathbb{C}$  a connected subset. Then a continuous function  $f : X \rightarrow \mathbb{C}$  has connected image  $f(X)$ .


*Proof.* The main point is to use the topological characterization of continuity. In particular, suppose that  $f(X)$  is disconnected, and we show that  $X$  is disconnected. In particular, suppose that  $U_1$  and  $U_2$  disconnect  $f(X)$ , and we have that  $f^{-1}(U_1)$  and  $f^{-1}(U_2)$  disconnect  $X$ . We will not run all the checks here; the main point is that  $f^{-1}(U_1)$  and  $f^{-1}(U_2)$  are open because  $f$  is continuous. ■

**Definition 2.101 (Path).** A *path* in  $\mathbb{C}$  is a continuous function  $\gamma : [a, b] \rightarrow \mathbb{C}$  where  $a < b$  are real numbers.

**Definition 2.102.** We say that a path  $\gamma$  is *closed* if and only if  $\gamma(a) = \gamma(b)$ . We say that  $\gamma$  is *simple* if and only if  $\gamma : (a, b) \rightarrow \mathbb{C}$  is injective.

**Remark 2.103.** The point of restricting  $\gamma$  to the open interval at the end so that closed, simple paths are allowed to exist.


**Example 2.104.** Here is a path.




**Example 2.105.** Here is a closed path.



**Example 2.106.** Here is a simple path.



**Example 2.107.** Here is a closed, simple path, also called a loop.




**Definition 2.108** (Concatenation). Fix  $\gamma_1 : [a, b] \rightarrow \mathbb{C}$  and  $\gamma_2 : [c, d] \rightarrow \mathbb{C}$  paths in  $\mathbb{C}$  such that  $\gamma_1(b) = \gamma_2(c)$ . Then we define the *concatenation* of  $\gamma_1$  and  $\gamma_2$  to be

$$(\gamma_1 * \gamma_2)(t) := \begin{cases} \gamma_1(t) & t \in [a, b], \\ \gamma_2(t) & t \in [b, d + c - b]. \end{cases}$$

The main point is that we are doing one path after the other.

**Example 2.109.** The following shows an example concatenation of  $\gamma_1 * \gamma_2$ , where  $\gamma_1, \gamma_2 : [0, 1] \rightarrow \mathbb{C}$ .



The entire path is  $\gamma_1 * \gamma_2$ .

Paths give us the following notion.

**Definition 2.110** (Path-connected). A subset  $X \subseteq \mathbb{C}$  is *path connected* if and only if, for any two  $x_0, x_1 \in X$ , there exists a path  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x_0$  and  $\gamma(1) = x_1$ .

**Lemma 2.111.** The open ball  $B(z, r)$  and closed ball  $\overline{B(z, r)}$  are both path-connected.

*Proof.* The point is that  $B(z, r)$  and  $\overline{B(z, r)}$  are both convex, so the path

$$\gamma(t) := z_0 + t(z_1 - z_0)$$

will work. ■

Here is the basic result.

**Proposition 2.112.** A space  $X$  is path-connected implies that  $X$  is connected. If  $X$  is open and connected, then  $X$  is path-connected.

*Proof.* We will show this next class. ■

## 2.7 February 7

Good morning everyone. A few announcements.

- Homework #3 is due on Friday.
- There will be no in-person class on Wednesday or Friday.
- Office hours this week are today (1:00PM–2:30PM) and tomorrow (2:00PM–3:30PM).

### 2.7.1 Connectedness

Today we're going to talk more about continuous functions.

Last time we ended with the following proposition.


**Proposition 2.113.** A space  $X$  is path-connected implies that  $X$  is connected. If  $X$  is open and connected, then  $X$  is path-connected.

*Proof.* We do these separately.

- Suppose that  $X = U_1 \sqcup U_2$  is disconnected, and we show that  $X$  is not path-connected. Namely, we have  $U_1, U_2 \subseteq X$  open subsets (in  $X$ ) which are disjoint and nonempty. Because  $U_1$  and  $U_2$  are nonempty, find  $x_1 \in U_1$  and  $x_2 \in U_2$ .

However, we claim there is no continuous path  $\gamma : [0, 1] \rightarrow X$  with  $\gamma(0) = x_1$  and  $\gamma(1) = x_2$ . Indeed, the image of  $\gamma([0, 1])$  must be connected, but then we can disconnect  $\gamma([0, 1])$  by  $U_1$  and  $U_2$ :  $\gamma([0, 1]) \subseteq U_1 \cup U_2$  and  $x_\bullet \in U_\bullet \cap \gamma([0, 1])$  and  $U_1 \cap U_2 = \emptyset$ .

At a high level, here is the image that a disconnected  $X$  cannot have a path between any two pair points: there is no possible red path below which stays in the gray region.



- Suppose we have a point  $z \in X$ , and we set

$$C(z) := \{w \in X : \text{there is a path from } z \text{ to } w\}.$$

We claim that  $C(z)$  is closed and open in  $X$ , which will force  $C(z) = X$  because  $X$  is connected and  $C(z)$  is nonempty ( $z \in C(z)$  by the trivial path  $\gamma : t \mapsto z$ ).

We start by showing  $C(z)$  is open: because  $X$  is open, there exists  $r > 0$  such that  $B(w, r) \subseteq X$ . But with  $w \in C(z)$ , there will be a path between any point in  $p \in B(z, w)$  and  $w$ , so there is a path from  $z$  to  $w$  to  $p$ . Here is the image.



Now we show that  $C(z)$  is closed. Suppose that  $w \in X \setminus C(z)$ , and we have to show that there is an open ball around  $w$  in  $X \setminus C(z)$ . To see this, fix an open ball  $B(w, r) \subseteq X$  for  $r > 0$ , but now there can be no path from  $z$  to anywhere in  $B(w, r)$ , for then we could just run the above argument again to show that  $w \in C(z)$ . ■

**Remark 2.114.** The proof for the second part merely needs  $X$  to be locally path-connected, not a metric space.

**Corollary 2.115.** We have that  $\mathbb{C}$  is path-connected and therefore connected.

*Proof.* Given any two points  $z, w \in \mathbb{C}$ , we choose the path  $\gamma : [0, 1] \rightarrow \mathbb{C}$  by

$$\gamma(t) = tz + (1 - t)w.$$

Indeed,  $\gamma(0) = w$  and  $\gamma(1) = z$ , and  $\gamma$  is somewhat clearly continuous by, say, checking component-wise. ■

## 2.7.2 Compactness

Let's do compactness better this time.

**Lemma 2.116.** Fix  $X \subseteq \mathbb{C}$  (sequentially) compact. Then  $X$  is both closed and bounded.

*Proof.* We start by showing  $X$  is closed. For this, we show that  $X$  contains all of its limit points.

Well, suppose that  $z \in X$  is a limit point so that we have a sequence  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  such that  $z_n \rightarrow z$ . But by the (sequential) compactness of  $X$ , this sequence has a convergent subsequence  $\{z_{\sigma n}\}_{n \in \mathbb{N}}$  which does converge in  $X$ . But any subsequence will converge to the same limit (!), so  $z_{\sigma n} \rightarrow z$  as well, so  $z \in X$  is forced.

We now show that  $X$  is bounded. We proceed by contraposition: if  $X$  is unbounded, then for any  $n \in \mathbb{N}$ , then we can find some  $z_n \in X \setminus B(0, n)$ . But then we can check that  $\{z_n\}_{n \in \mathbb{N}}$  has no convergent subsequence, essentially because it tends off to infinity. ■

Our goal for the rest of class is to prove the following two results.

**Proposition 2.117.** A subset  $X \subseteq \mathbb{C}$  is (sequentially) compact if and only if it is closed and bounded.

**Theorem 2.118.** [Heine–Borel] A subset  $X \subseteq \mathbb{C}$  is (sequentially) compact if and only if every open cover of  $X$  has a finite subcover.

On the homework, we showed the backward direction of Theorem 2.118.

**Remark 2.119.** Our hope is to have lots of equivalent characterizations of compactness so that we can have easier proofs of statements about compact sets.

To start off, here are some lemmas we will need.

**Lemma 2.120.** Fix  $X \subseteq \mathbb{C}$  (sequentially) compact. For any  $\varepsilon > 0$ , there exist only finitely many points  $z_1, \dots, z_n \in X$  such that

$$X \subseteq \bigcup_{k=1}^n B(z_k, \varepsilon).$$

*Proof.* The point is to build some inductive argument: one fixes an  $\varepsilon > 0$  and then continues choosing random points out of  $X$  until we cover  $X$ . Indeed, if the process does not terminate, then the sequence we generate has no convergent subsequence.

Rigorously, if  $X$  is empty, then just choose no points at all and be done. Otherwise, we can find some  $z_1 \in X$ . Inductively, suppose we have a sequence  $\{z_1, \dots, z_m\}$ . If

$$X \subseteq \bigcup_{k=1}^m B(z_k, \varepsilon),$$

then we are done. Otherwise, we can find  $z_{m+1} \in X \setminus \bigcup_{k=1}^m B(z_k, \varepsilon)$ .

If the above inductive process terminates, then we get the result. Otherwise, there is a sequence  $\{z_n\}_{n \in \mathbb{N}}$  such that

$$z_{n+1} \in X \setminus \bigcup_{k=1}^n B(z_k, \varepsilon).$$

We claim that  $\{z_n\}_{n \in \mathbb{N}}$  has subsequence converging in  $X$ . Indeed, suppose for the sake of contradiction that  $z_{\sigma n} \rightarrow z$  for some strictly increasing  $\sigma$  and  $z \in X$ . Then there exists  $N$  such that  $n > N$  implies

$$|z_{\sigma n} - z| < \varepsilon/2.$$

But then, finding some  $n+1, n > N$ , we have

$$|z_{\sigma(n+1)} - z_{\sigma n}| < |z_{\sigma(n+1)} - z| + |z_{\sigma n} - z| < \varepsilon,$$

so

$$z_{\sigma(n+1)} \in \bigcup_{k=1}^{\sigma(n+1)-1} B(z_k, \varepsilon),$$

which is our contradiction to the construction of  $z_n$ . ■

**Lemma 2.121.** Fix  $X \subseteq \mathbb{C}$  (sequentially) compact with some open cover  $\mathcal{U}$  of  $X$ . Then there is an  $\varepsilon > 0$  such that, for every  $z \in X$ , there is  $U \in \mathcal{U}$  such that  $B(z, \varepsilon) \subseteq U$ .

*Proof.* Suppose that, for all  $\varepsilon > 0$ , there exists some  $z \in X$  such that no  $U \in \mathcal{U}$  has  $B(z, \varepsilon) \subseteq U$ . We construct a sequence in  $X$  with no subsequence converging in  $X$ . Indeed, for any  $n \in \mathbb{N}$ , we find  $z_n \in X$  such that no  $U \in \mathcal{U}$  has  $B(z_n, 1/n) \subseteq U$ . We claim that  $\{z_n\}_{n \in \mathbb{N}}$  has no subsequence converging in  $X$ .

Indeed, suppose that we have  $z \in X$  and strictly increasing  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $z_{\sigma n} \rightarrow z$ . We will then be able to find some  $z_n$  such that  $B(z_n, 1/n) \subseteq U$  for some  $U \in \mathcal{U}$ , which will be a contradiction. Indeed,  $z \in X$ , and  $\mathcal{U}$  covers  $z$ , so there is some  $U \in \mathcal{U}$  with  $z \in U$ . In fact,  $U$  is open, so there is an  $\varepsilon > 0$  such that

$$B(z, \varepsilon) \subseteq U.$$

Now, there is  $N$  such that for  $n > N$ , we can guarantee that  $|z - z_n| < \varepsilon/2$ . Further, for  $n > 2/\varepsilon$ , we have  $1/n < \varepsilon/2$ . So  $n > \max\{N, 2/\varepsilon\}$  will have  $\sigma n > \max\{N, 2/\varepsilon\}$ , implying

$$|w - z_n| < 1/n < \varepsilon/2 \implies |w - z| < |w - z_n| + |z - z_n| = \varepsilon \implies w \in B(z, \varepsilon) \subseteq U,$$

so  $B(z_n, 1/n) \subseteq U$ . This contradiction finishes. ■

This is saying that there is a universal  $\varepsilon$  that we can find for our open cover.

**Lemma 2.122.** Fix  $X$  a bounded set. Then, for any  $\varepsilon > 0$ , there exist finitely many points  $z_1, \dots, z_n$  such that

$$X \subseteq \bigcup_{k=1}^n B(z_k, \varepsilon).$$

*Proof.* The point is to reduce this to the case of  $[-M, M]^2$  which can cover  $X$  because  $X$  is bounded, and then we can create the cover for  $X$  by hand. ■

Now let's attack one of our equivalent conditions for compactness.

**Proposition 2.117.** A subset  $X \subseteq \mathbb{C}$  is (sequentially) compact if and only if it is closed and bounded.

*Proof.* The forwards direction we have already done.

In the backwards direction, suppose that  $\{z_n\}_{n \in \mathbb{N}} \subseteq X$  is some sequence. Our main goal is to construct a convergent subsequence. Because  $X$  is bounded, we can choose  $w_{1,1}, \dots, w_{1,\ell_1}$  such that

$$X \subseteq \bigcup_{k=1}^{\ell_1} B(w_{1,k}, 1/2).$$

Now, because  $\{z_n\}_{n \in \mathbb{N}}$  is infinite, there must be some index  $w_1 := w_{1,k_1}$  such that

$$L_1 = \{n \in \mathbb{N} : z_n \in B(w_{1,k_1}, 1/2)\}$$

is infinite. The important point is that  $\{z_n\}_{n \in L_1}$  has formed a subsequence which lives inside a ball of radius  $1/2$ . We can continue this process: again using our bounded condition, we can find some  $w_{2,1}, \dots, w_{2,\ell_2} \in B(w_{1,k_1}, 1/2)$  such that

$$B(w_{1,k_1}, 1/2) \subseteq \bigcup_{k=1}^{\ell_2} B(w_{2,k}, 1/4).$$

Then we can choose  $L_2$  from here by choosing one of the  $w_{2,k}$  with infinitely many indices. Continuing this process forces our sequence to converge.

To more explicitly appeal to choice, we note that we can always find some sequence  $\{w_{k,i}\} \subseteq X$  such that

$$X \subseteq \bigcup_{k=1}^{\ell_n} B(w_{k,i}, 1/2^k),$$

but  $L_{i-1}$  is infinite, so there is a specific  $w_k := w_{k,i}$  such that

$$L_i := \{n \in L_{i-1} : |z_n - w_k| < 2^{-k}\}$$

is infinite. To actually construct our sequence from these infinite subsets, we define a choice function over our indices: define  $\varphi : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\varphi(n+1)$  is the smallest number exceeding  $\varphi(n)$  with  $\varphi(n+1) \in L_{n+1}$ . Then we know that

$$|z_{\varphi(n)} - w_k| < 2^{-n}$$

for each  $1 \leq k \leq n$ . Thus, for  $n \geq m > N$ , we have

$$|z_{\varphi(n)} - z_{\varphi(m)}| \leq |z_{\varphi(n)} - w_m| + |z_{\varphi(m)} - w_m| < 2 \cdot 2^{-m} < 2^{-N+1},$$

so for any  $\varepsilon > 0$ , we can choose  $N := 1 - \log_2 \varepsilon$  sufficiently large so that  $n, m > N$  implies

$$|z_{\varphi(n)} - z_{\varphi(m)}| < 2^{-N+1} = \varepsilon.$$

It follows that the subsequence defined by  $\varphi$  is Cauchy and hence converges. But because  $X$  is closed, any convergent sequence in  $X$  will be in  $X$ , so our sequence in  $X$  has a convergent subsequence. ■

## 2.8 February 9

### 2.8.1 More Compactness

To wrap up from last class, we show the following.

**Theorem 2.118.** [Heine–Borel] A subset  $X \subseteq \mathbb{C}$  is (sequentially) compact if and only if every open cover of  $X$  has a finite subcover.

*Proof.* The direction that sequentially compact implies closed and bounded was done on the homework.

We focus on the other direction. Fix  $\mathcal{U}$  an open cover of  $X$ . By Lemma 2.121, we know there exists  $\varepsilon > 0$  such that, for each  $z \in X$ , there is some  $U \in \mathcal{U}$  such that  $B(z, \varepsilon) \subseteq U$ . But in fact, with this  $\varepsilon > 0$ , Lemma 2.122 tells us that there exists finitely many points  $z_1, \dots, z_\ell$  such that

$$X \subseteq \bigcup_{k=1}^{\ell} B(z_k, \varepsilon).$$

But now, finding  $U_k$  such that  $B(z_k, \varepsilon) \subseteq U_k$  (possible by construction of  $\varepsilon$ ), we see that  $\{U_k\}_{k=1}^{\ell}$  will be our finite subcover. ■

**Remark 2.123.** The conclusion of the above theorem is the usual notion of compactness, so I will stop writing “(sequentially)” whenever I say “compact.”

Let’s see a use for our definitions of compactness.

**Corollary 2.124.** Let  $X \subseteq \mathbb{C}$  be a compact space and  $f : X \rightarrow \mathbb{C}$  continuous. Then  $f(X)$  is compact.

*Proof.* Give  $f(X)$  some open cover  $\mathcal{U}$ . Because  $f$  is continuous, we see that

$$\{f^{-1}(U)\}_{U \in \mathcal{U}}$$

is an open cover for  $X$ . But  $X$  is compact, so we can find some finite subcover  $\{U_k\}_{k=1}^n \subseteq \mathcal{U}$  so that  $\{f^{-1}(U_k)\}_{k=1}^n$  covers  $X$ . But then the  $\{U_k\}_{k=1}^n$  will cover  $f(X)$  by taking the union over our open subcover. ■

### 2.8.2 Uniform Continuity

The point of uniform convergence is to make fewer choices in our notion of continuity.

**Definition 2.125** (Uniform continuity). Fix  $X \subseteq \mathbb{C}$  a nonempty subset. Then a function  $f : X \rightarrow \mathbb{C}$  is *uniformly continuous* if and only if, for each  $\varepsilon > 0$ , there exists a single  $\delta > 0$  so that  $z, w \in X$  have

$$|z - w| < \delta \implies |f(z) - f(w)| < \varepsilon.$$

Importantly, this definition has  $\delta$  not depend on either  $z$  nor  $w$ , where continuity would allow  $\delta$  to depend on one of them.

Here is a nice result.

**Proposition 2.126.** Fix  $X$  a nonempty, compact subset. Then any continuous function  $f : X \rightarrow \mathbb{C}$  is uniformly continuous.

*Proof.* The point is to let  $\delta \rightarrow 0$  until we can fit some prescribed  $\varepsilon$  bound. Choose  $\delta = 1/n$  as  $n$  varies over positive integers, and we imagine fixing sequences  $\{z_n\}_{n=1}^\infty$  and  $\{w_n\}_{n=1}^\infty$  such that

$$|z_n - w_n| < 1/n.$$

Now we use the sequential compactness of  $X$ , which forces  $\{z_n\}_{n=1}^\infty$  to have a convergent subsequence, so we conjure  $\alpha \in X$  and a strictly increasing  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $z_{\sigma n} \rightarrow \alpha$  as  $n \rightarrow \infty$ .

We now claim that  $w_{\sigma n} \rightarrow \alpha$  as well. In particular, for any  $\delta > 0$ , there is some  $N_1$  so that  $n > N_1$  implies

$$|z_{\sigma n} - \alpha| < \delta/2.$$

Choosing  $N$  to be larger than  $N_1$  and  $2/\delta$ , we see that  $n > N$  will have

$$|w_{\sigma n} - \alpha| \leq |z_{\sigma n} - w_{\sigma n}| + |z_{\sigma n} - \alpha| < \frac{1}{\sigma n} + \frac{\delta}{2} \leq \frac{1}{n} + \frac{\delta}{2} < \frac{\delta}{2} + \frac{\delta}{2} = \delta,$$

so indeed  $w_{\sigma n} \rightarrow \alpha$  as  $n \rightarrow \infty$ .

Only now we suppose for the sake of contradiction we have some  $\varepsilon > 0$  such that any  $\delta > 0$  has some  $z$  and  $w$  such that  $|z - w| < \delta$  actually has  $|f(z) - f(w)| \geq \varepsilon$ . Taking  $\delta := 1/n$ , we are promised some sequences  $\{z_n\}_{n=1}^\infty$  and  $\{w_n\}_{n=1}^\infty$  so that

$$|z_n - w_n| < \delta \quad \text{and} \quad |f(z_n) - f(w_n)| \geq \varepsilon.$$

Using the above machinery, we see that  $z_{\sigma n} \rightarrow \alpha$  and  $w_{\sigma n} \rightarrow \alpha$  force  $f(z_{\sigma n}) \rightarrow f(\alpha)$  and  $f(w_{\sigma n}) \rightarrow f(\alpha)$  by continuity of  $f$ , but the sequences  $f(z_{\sigma n})$  and  $f(w_{\sigma n})$  are supposed to be  $\varepsilon$  far apart! Explicitly, we can find sufficiently large  $N_1$  and  $N_2$  such that

$$n > N_1 \implies |f(z_{\sigma n}) - \alpha| < \varepsilon/4,$$

$$n > N_2 \implies |f(w_{\sigma n}) - \alpha| < \varepsilon/4,$$

which by the triangle inequality means that any  $n > \max\{N_1, N_2\}$  will give

$$|f(z_{\sigma n}) - f(w_{\sigma n})| \leq |f(z_{\sigma n}) - \alpha| + |f(w_{\sigma n}) - \alpha| \leq \frac{\varepsilon}{4} + \frac{\varepsilon}{4} < \varepsilon,$$

which is a contradiction to the construction of  $z_{\sigma n}$  and  $w_{\sigma n}$ . ■

### 2.8.3 Uniform Convergence

We next talk about uniform convergence for functions. Here is our starter pack.

**Definition 2.127** (Sequence of functions). Fix  $X \subseteq \mathbb{C}$  a nonempty subset. Then a sequence of functions  $\{f_n\}_{n \in \mathbb{N}}$  is a function  $\varphi : \mathbb{N} \rightarrow (X \rightarrow \mathbb{C})$ . Namely, for each  $n \in \mathbb{N}$ , we are given a function  $\varphi(n) : X \rightarrow \mathbb{C}$ .



**Definition 2.128** (Pointwise convergence). Fix  $\{f_n\}_{n \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$ . Then  $\{f_n\}$  converges to some  $g : X \rightarrow \mathbb{C}$  pointwise if and only if, for each  $z \in X$ , we have  $f_n(z) \rightarrow g(z)$  as  $n \rightarrow \infty$ . We write this as  $f_n \rightarrow g$ .

This is called pointwise convergence because we are only checking convergence at each individual point  $z \in X$ , without caring about the larger structure of the function. This will cause problems later but not now.

**Definition 2.129** (Bounded). We say that a function  $f : X \rightarrow \mathbb{C}$  is *bounded* if and only if  $f(X) \subseteq \mathbb{C}$  is bounded. In other words, there is some  $M > 0$  so that  $f(X) \subseteq B(0, M)$ .

**Definition 2.130** (Uniform convergence). Fix  $\{f_n\}_{n \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$ . Then  $\{f_n\}$  converges to some  $g : X \rightarrow \mathbb{C}$  pointwise if and only if, for each  $\varepsilon > 0$ , there is some  $N$  so that

$$n > N \implies |f_n(z) - g(z)| < \varepsilon$$

for each  $z \in X$ .

The uniformity here is that the value of  $N$  is no longer allowed to depend on  $z$ . Here is an alternate definition.

**Proposition 2.131.** Fix  $\{f_n\}_{n \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$  and  $g : X \rightarrow \mathbb{C}$  some function. Then  $\{f_n\}_{n \in \mathbb{N}}$  converges uniformly to  $g$  if and only if

$$\lim_{n \rightarrow \infty} \sup_{z \in X} \{|f_n(z) - g(z)|\} = 0.$$

*Proof.* We take the directions independently.

- In the forward direction, we know that there is an  $N_1$  so that  $n > N_1$  implies each  $z \in X$  has

$$|f_n(z) - g(z)| < 1.$$

In particular, for  $n > N_1$ , the set  $\{|f_n(z) - g(z)| : z \in X\}$  is bounded above by 1, so the supremum will exist; set  $y_n := \sup\{|f_n(z) - g(z)| : z \in X\}$  so that we want to show  $y_n \rightarrow 0$  as  $n \rightarrow \infty$ .

More generally, for any  $\varepsilon > 0$ , there exists some  $N$  so that  $n > N_0$  implies

$$|f_n(z) - g(z)| < \varepsilon/2.$$

So  $n > \max\{N_0, N_1\}$ , we will have that  $y_n = \sup\{|f_n(z) - g(z)| : z \in X\}$  both exists and has  $y_n \leq \varepsilon/2 < \varepsilon$ . So we do get  $y_n \rightarrow 0$  as  $n \rightarrow \infty$ .

- In the reverse direction, set  $y_n := \sup\{|f_n(x) - g(x)| : x \in X\}$  so that  $y_n \rightarrow 0$  as  $n \rightarrow \infty$ . Namely, for each  $\varepsilon > 0$ , there exists some  $N$  so that  $n > N$  has  $y_n < \varepsilon$ . In particular, we see  $n > N$  has

$$|f_n(x_0) - g(x_0)| \leq \sup\{|f_n(x) - g(x)| : x \in X\} = y_n < \varepsilon$$

for each  $x_0 \in X$ . So indeed,  $f_n$  converges to  $g$  uniformly. ■

## 2.8.4 Distances Between Functions

Later in life it will be nice to view functions as forming a metric under  $d(f, g) := \sup\{|f(x) - g(x)|\}$ . However, this supremum need not only exist; here is one condition in which it does.

**Lemma 2.132.** Fix  $f, g : X \rightarrow \mathbb{C}$  bounded functions. Then  $\sup\{|f(x) - g(x)| : x \in X\}$  exists.

*Proof.* Because  $f$  is bounded, there exists  $M_f$  so that each  $x \in X$  has  $|f(x)| < M_f$ . Similarly, because  $g$  is bounded, there exists  $M_g$  so that each  $x \in X$  has  $|g(x)| < M_g$ . It follows that, for each  $x \in X$ ,

$$|f(x) - g(x)| \leq |f(x)| + |g(x)| \leq M_f + M_g,$$

so the set  $\{|f(x) - g(x)| : x \in X\}$  is bounded above and in particular has a supremum. ■

**Proposition 2.133.** Fix  $f, g, h : X \rightarrow \mathbb{C}$  all bounded functions. Then

$$\sup_{x \in X} \{|f(x) - h(x)|\} \leq \sup_{x \in X} \{|f(x) - g(x)|\} + \sup_{x \in X} \{|g(x) - h(x)|\}.$$

Note that all the suprema above exist by Lemma 2.132

*Proof.* The point is to reduce to the typical triangle inequality. Indeed, for any  $x \in X$ , we see that

$$|f(x) - h(x)| \leq |f(x) - g(x)| + |g(x) - h(x)|.$$

Thus,

$$\begin{aligned} \sup_{x \in X} \{|f(x) - h(x)|\} &\leq \sup_{x \in X} \{|f(x) - g(x)| + |g(x) - h(x)|\} \\ &\leq \sup_{x \in X} \{|f(x) - g(x)|\} + \sup_{x \in X} \{|g(x) - h(x)|\}, \end{aligned}$$

which is what we wanted. We have used the fact that  $\sup(A + B) \leq \sup A + \sup B$  for  $A, B \subseteq \mathbb{R}$ , which is not hard to show: if  $a + b \in A + B$ , then  $a \leq \sup A$  and  $b \leq \sup B$ , so  $a + b \leq \sup A + \sup B$ ; thus,  $\sup(A + B) \leq \sup A + \sup B$ .<sup>4</sup> ■

**Remark 2.134** (Nir). Viewing Lemma 2.132 as providing a distance metric, the above proposition proves the triangle inequality for this metric.

We can also build a Cauchy criterion for uniform convergence.

**Proposition 2.135.** A sequence of functions  $\{f_n\}_{n \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$ . Then  $\{f_n\}_{n \in \mathbb{N}}$  converges to some function uniformly if and only if the quantity  $\sup_{x \in X} \{f_n(x) - f_m(x)\}$  exists and, for any  $\varepsilon > 0$ , there exists some  $N$  so that  $n, m > N$  implies

$$\sup_{x \in X} \{|f_n(x) - f_m(x)|\} < \varepsilon$$

for any  $x \in X$ .

We note that the hypothesis that the supremum exists can be removed if the functions are presupposed to be bounded.

*Proof.* We again take the directions independently.

<sup>4</sup> In fact,  $\sup A + \sup B \leq \sup(A + B)$  as well. We show  $\sup A \leq \sup(A + B) - \sup B$ . Fixing  $a \in A$ , we need  $a \leq \sup(A + B) - \sup B$ , so we show  $\sup B \leq \sup(A + B) - a$ . Fixing  $b \in B$ , we need  $b \leq \sup(A + B) - a$ , which is clear.

- Suppose that the sequence of functions  $\{f_n\}_{n \in \mathbb{N}}$  converges to a function  $g$  uniformly. Then, for any  $\varepsilon > 0$ , we are promised some  $N$  so that  $n > N$  will have

$$|f_n(x) - g(x)| < \varepsilon/4$$

for any  $x \in X$ . In particular, for any  $n, m > N$ , we see

$$|f_n(x) - f_m(x)| < |f_n(x) - g(x)| + |f_m(x) - g(x)| < \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \frac{\varepsilon}{2},$$

so

$$\sup_{x \in X} \{|f_n(x) - f_m(x)|\} \leq \frac{\varepsilon}{2} < \varepsilon,$$

which is what we wanted.

- There are two steps.
  - We begin by constructing  $g$ . Well, for each  $x \in X$ , we note that any  $\varepsilon > 0$  will have some  $N$  so that  $n, m > N$  implies

$$|f_n(x) - f_m(x)| \leq \sup_{x \in X} \{|f_n(x) - f_m(x)|\} < \varepsilon,$$

so the sequence  $\{f_n(x)\}_{n \in \mathbb{N}}$  is a Cauchy sequence and hence converges in  $\mathbb{C}$ . We define  $g(x)$  to be the limit of  $f_n(x)$  as  $n \rightarrow \infty$ .

- Next we show the uniform convergence. Fix some  $\varepsilon > 0$ . Then we are promised some  $N$  so that  $n, m > N$  has

$$\sup_{x \in X} \{|f_n(x) - f_m(x)|\} < \varepsilon.$$

In particular, for any  $x \in X$

$$f_n(x) - g(x) = \lim_{m \rightarrow \infty} (f_n(x) - f_m(x)),$$

so because  $z \mapsto |z|$  is continuous, any  $n > N$  will have

$$|f_n(x) - g(x)| = \lim_{m \rightarrow \infty} |f_n(x) - f_m(x)| \leq \lim_{m \rightarrow \infty} \sup_{z \in X} \{|f_n(z) - f_m(z)|\} < \lim_{m \rightarrow \infty} \varepsilon = \varepsilon,$$

where the last inequality holds by taking  $m$  sufficiently large (that is,  $m > N$ ). So we have been provided uniform convergence. ■

**Remark 2.136** (Nir). In the language of metric topology, the above proposition asserts that the space of (bounded) functions is metrically complete. For this, one must technically show that  $\{f_n\}_{n \in \mathbb{N}}$  being bounded implies that the convergent  $g$  is bounded, but this is not hard: there is  $N$  so that  $n > N$  has  $|f_n(x) - g(x)| < 1$  for each  $x \in X$ .

**Remark 2.137.** In lecture, Professor Morrow asserted that we require these functions to be bounded. I do not think this is the case; indeed, the above proof never uses this hypothesis.

We close with one result which shows that uniform continuity is nice.

**Proposition 2.138.** Fix  $\{f_n\}_{n \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$  all continuous at some  $x \in X$ . If  $\{f_n\}_{n \in \mathbb{N}}$  converges uniformly to some function  $g : X \rightarrow \mathbb{C}$ , then  $g$  is also continuous.

*Proof.* The idea is to well-approximate  $g$  by  $f_n$ s. Fix any  $\varepsilon > 0$ . By the uniform convergence, there will be some  $N$  so that

$$|f_n(z) - g(z)| < \varepsilon/3$$

for any  $n > N$  and  $z \in X$ ; fix some  $n > N$ . Because  $f_n$  is continuous, we are promised some  $\delta > 0$  (allowed to vary with our chosen  $x \in X$ ) so that

$$|z - x| < \delta \implies |f_n(z) - f_n(x)| < \varepsilon/3$$

for any  $z \in X$ . Well, if  $|z - x| < \delta$ , then the triangle inequality gives

$$|g(z) - g(x)| \leq |g(z) - f_n(z)| + |f_n(z) - f_n(x)| + |f_n(x) - g(x)| < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon,$$

which is what we needed. ■

**Remark 2.139** (Nir). In fact, if the  $\{f_n\}_{n \in \mathbb{N}}$  are uniformly continuous, then  $g$  will also be uniformly continuous. The argument is similar.

---

## THEME 3: HOLOMORPHIC FUNCTIONS

---

*I turn with terror and horror from this lamentable scourge of continuous functions with no derivatives.*

—Charles Hermite

### 3.1 February 11

The wheel of time marches on. Today, we actually start talking about complex analysis.

#### 3.1.1 Differentiability

We are going to talk about holomorphic functions.

**Convention 3.1.** We set  $\Omega$  to be some open subset of  $\mathbb{C}$ .

This gives the following definition.

**Definition 3.2** (Differentiable). Fix an open subset  $\Omega \subseteq \mathbb{C}$  and  $f : \Omega \rightarrow \mathbb{C}$  a function. Then  $f$  is *complex differentiable* at  $z_0 \in \Omega$  with derivative  $\alpha \in \mathbb{C}$  if and only if

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \alpha.$$

We write this as  $f'(z_0) = \alpha$ .

If  $f'$  is itself a differentiable function, then  $f$  would be “twice” differentiable, and we denote this function by  $f''$ . In general, if  $f$  can be differentiated  $n$  times, we denote the corresponding function by  $f^{(n)}$ .



**Warning 3.3.** In the definition of complex differentiability, we are taking the limit with  $h \in \mathbb{C}$ , not  $h \in \mathbb{R}$ . This will make complex differentiability significantly more structured.

Differentiability gives rise to the following definition.

**Definition 3.4** (Holomorphic, entire). Fix an open subset  $\Omega \subseteq \mathbb{C}$  and  $f : \Omega \rightarrow \mathbb{C}$  a function. Then  $f$  is *holomorphic* on  $\Omega$  if and only if  $f$  is complex differentiable at each  $z_0 \in \Omega$ . If  $\Omega = \mathbb{C}$ , then we say  $f$  is *entire*.

Here is a small usual lemma.

**Lemma 3.5.** Fix an open subset  $\Omega \subseteq \mathbb{C}$  and  $f : \Omega \rightarrow \mathbb{C}$  a function. Then if  $f$  is differentiable at  $z_0 \in \Omega$ , then  $f$  is continuous at  $z_0 \in \mathbb{C}$ .

*Proof.* We compute that

$$\begin{aligned}
 \lim_{z \rightarrow z_0} (f(z) - f(z_0)) &= \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \cdot \lim_{z \rightarrow z_0} (z - z_0) \\
 &\stackrel{*}{=} \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \cdot \lim_{z \rightarrow z_0} (z - z_0) \\
 &= f'(z_0) \cdot 0 \\
 &= 0.
 \end{aligned}$$

It follows by rearrangement that  $\lim_{z \rightarrow z_0} f(z) = f(z_0)$ , which is what we wanted. Notably,  $\stackrel{*}{=}$  sets  $h := z - z_0$ . ■

### 3.1.2 Basic Properties

As usual, differentiable functions have an arithmetic.

**Proposition 3.6.** Fix an open subset  $\Omega \subseteq \mathbb{C}$  and  $f, g : \Omega \rightarrow \mathbb{C}$  functions differentiable at  $z_0 \in \mathbb{C}$ .

- (a) We have that  $(af + bg)'(z_0) = af'(z_0) + bg'(z_0)$ , where  $a, b \in \mathbb{C}$ .
- (b) We have that  $(fg)'(z_0) = f'(z_0)g(z_0) + f(z_0)g'(z_0)$ .
- (c) If  $g'(z_0) \neq 0$ , then

$$(f/g)'(z_0) = \frac{f'(z_0)g(z_0) - f(z_0)g'(z_0)}{g(z_0)^2}.$$

*Proof.* We copy the proofs from real analysis.

- (a) We check that

$$\begin{aligned}
 \lim_{h \rightarrow 0} \frac{(af + bg)(z_0 + h) - (af + bg)(z_0)}{h} &= a \cdot \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} + b \cdot \lim_{h \rightarrow 0} \frac{g(z_0 + h) - g(z_0)}{h} \\
 &= a \cdot f'(z_0) + b \cdot g'(z_0),
 \end{aligned}$$

which is what we wanted.

- (b) The key idea is to add and subtract  $f(z_0)g(z_0 + h)$ . Indeed, we see

$$\begin{aligned}
 \lim_{h \rightarrow 0} \frac{(fg)(z_0 + h) - (fg)(z_0)}{h} &= \lim_{h \rightarrow 0} \frac{f(z_0 + h)g(z_0 + h) - f(z_0)g(z_0 + h)}{h} \\
 &\quad + \lim_{h \rightarrow 0} \frac{f(z_0)g(z_0 + h) - f(z_0)g(z_0)}{h} \\
 &= \left( \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \right) \left( \lim_{h \rightarrow 0} g(z_0 + h) \right) \\
 &\quad + f(z_0) \left( \lim_{h \rightarrow 0} \frac{g(z_0 + h) - g(z_0)}{h} \right) \\
 &= f'(z_0)g(z_0) + f(z_0)g'(z_0),
 \end{aligned}$$

which is what we wanted.

- (c) This will follow from applying the product rule to  $f \cdot \frac{1}{g}$ , where we can compute the derivative of  $\frac{1}{g}$  by the chain rule. We refer to Eterović's notes for the details. ■

**Remark 3.7** (Nir). Technically part (c) will require us to compute the derivative of  $f(z) := \frac{1}{z}$  for  $z \neq 0$  to finish the proof. Well, for any  $z \in \mathbb{C} \setminus \{0\}$ , we see that

$$\frac{f(z+h) - f(z)}{h} = \frac{\frac{1}{z+h} - \frac{1}{z}}{h} = \frac{z - (z+h)}{hz(z+h)} = -\frac{1}{z(z+h)}.$$

Taking  $h \rightarrow 0$  reveals that the derivative is in fact  $f'(z) = -\frac{1}{z^2}$ .

Let's give some examples of entire functions.

**Exercise 3.8.** Fix  $n$  some positive integer. We show that the function  $f(z) := z^n$  is entire with derivative  $f'(z) := nz^{n-1}$ .

*Proof.* We employ the usual proof involving the binomial theorem. Note that

$$f(z+h) = (z+h)^n = \sum_{k=0}^n \binom{n}{k} z^{n-k} h^k,$$

so

$$\frac{f(z+h) - f(z)}{h} = \sum_{k=1}^n \binom{n}{k} z^{n-k} h^{k-1},$$

where notably the  $k=0$  term was killed by the  $-f(z)$ . Thus,

$$\lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} = \sum_{k=1}^n \binom{n}{k} z^{n-k} \left( \lim_{h \rightarrow 0} h^{k-1} \right),$$

but all terms except  $k=1$  will now vanish as  $h \rightarrow 0$ , so we are left with  $nz^{n-1}$  as our limit. ■

**Remark 3.9** (Nir). One could also show this by induction, using the product rule.

**Corollary 3.10.** Any polynomial function is entire.

*Proof.* Polynomials are (finite) linear combinations of the monomials  $f_n(z) := z^n$ , so this follows from combining the above two results. ■

### 3.1.3 Advanced Properties

We also have a notion of L'Hôpital's rule.

**Proposition 3.11** (L'Hôpital's rule). Fix  $\Omega \subseteq \mathbb{C}$  an open subset with  $f, g : \Omega \rightarrow \mathbb{C}$  holomorphic functions. Then, given  $z_0 \in \Omega$  with  $f(z_0) = g(z_0) = 0$  while  $g'(z_0) \neq 0$ , we have that

$$\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{f'(z_0)}{g'(z_0)}.$$

*Proof.* Note that, because  $f(z_0) = g(z_0) = 0$ , we see that

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z)}{z - z_0} \quad \text{and} \quad g'(z_0) = \lim_{z \rightarrow z_0} \frac{g(z)}{z - z_0}.$$

Dividing, we see that

$$\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \lim_{z \rightarrow z_0} \frac{f(z)/(z - z_0)}{g(z)/(z - z_0)} = \lim_{z \rightarrow z_0} \frac{f'(z_0)}{g'(z_0)} = \frac{f'(z_0)}{g'(z_0)},$$

which is what we wanted. ■

**Remark 3.12** (Nir). The above proof technically does not work because we have not ruled out the possibility that  $g$  might vanish arbitrarily close to  $z_0$ , thus making the limits not actually make sense. We will not fix this problem, but we will remark that a holomorphic function will only have finitely many zeroes on a compact set, which we could use to create a neighborhood for  $z_0$  on which  $g$  doesn't vanish.

And here is our chain rule.

**Proposition 3.13** (Chain rule). Fix  $\Omega$  and  $U$  open subsets of  $\mathbb{C}$  with functions  $f : \Omega \rightarrow U$  and  $g : U \rightarrow \mathbb{C}$ . Further, suppose that  $f$  is differentiable at  $z_0 \in \Omega$  and that  $g$  is differentiable at  $f(z_0) \in U$ . Then  $(g \circ f)$  is differentiable at  $z_0$  with derivative

$$(g \circ f)'(z_0) = g'(f(z_0))f'(z_0).$$

*Proof.* This proof is long, so we will try to be brief. The main idea is to consider the auxiliary function  $r : U \setminus \{f(z_0)\} \rightarrow \mathbb{C}$  defined by

$$r(w) := \frac{g(w) - g(f(z_0))}{w - f(z_0)} - g'(f(z_0)).$$

We extend  $r$  to  $f(z_0)$  by setting  $r(f(z_0)) := 0$ . Now, the differentiability of  $g$  at  $f(z_0)$  implies that

$$\lim_{z \rightarrow z_0} \frac{g(z) - g(f(z_0))}{z - z_0} = g'(f(z_0)),$$

so in particular rearranging implies that  $r$  is continuous on at  $f(z_0) \in U$ .

The reason we used the letter  $r$  is that we should think of  $r$  as a remainder term. Indeed, we see

$$g(w) - g(f(z_0)) = g'(f(z_0))(w - f(z_0)) + r(w)(w - f(z_0)).$$

Plugging in  $w = f(z)$ , we get

$$g(f(z)) - g(f(z_0)) = g'(f(z_0))(f(z) - f(z_0)) + r(f(z))(f(z) - f(z_0)),$$

so

$$\frac{g(f(z)) - g(f(z_0))}{z - z_0} = g'(f(z_0)) \cdot \frac{f(z) - f(z_0)}{z - z_0} + r(f(z)) \cdot \frac{f(z) - f(z_0)}{z - z_0}.$$

Sending  $z \rightarrow z_0$  makes the rightmost term vanish by continuity because  $r(f(z_0)) = 0$  and the limit is  $f'(z_0)$ , so we are left with

$$(g \circ f)'(z_0) = g'(f(z_0))f'(z_0),$$

which is what we wanted. ■

**Remark 3.14** (Nir). Let's complete the proof of quotient rule. Note that the derivative of  $\frac{1}{g(z)}$  will be, by the chain rule,  $-\frac{1}{g(z)^2} \cdot g'(z)$ . Thus, the derivative of  $\frac{f(z)}{g(z)} = f(z) \cdot \frac{1}{g(z)}$  will be

$$f'(z) \cdot \frac{1}{g(z)} - f(z) \cdot \frac{g'(z)}{g(z)^2} = \frac{f'(z)g(z) - f(z)g'(z)}{g(z)^2}.$$

And we finish with a result which is less common in real analysis, essentially saying that differentiable functions are "approximately" linear.



**Proposition 3.15** (Carathéodory). Fix  $\Omega \subseteq \mathbb{C}$  an open subset with a function  $f : \Omega \rightarrow \mathbb{C}$  and point  $z_0 \in \Omega$ . Then  $f$  is differentiable at  $z_0$  if and only if there exists a function  $h : \Omega \rightarrow \mathbb{C}$  which is continuous at  $z_0$  such that

$$f(z) - f(z_0) = h(z)(z - z_0).$$

In particular,  $h(z_0) = f'(z_0)$ .

*Proof.* We show the directions independently.

- Suppose  $f$  is differentiable at  $z_0$ . We construct the function  $h$  manually. We define

$$h(z) := \begin{cases} (f(z) - f(z_0))/(z - z_0) & z \in \Omega \setminus \{z_0\}, \\ f'(z_0) & z = z_0. \end{cases}$$

In particular, we note that  $h$  is continuous at  $z_0$  because  $h(z) \rightarrow f'(z_0)$  as  $z \rightarrow z_0$  by differentiability of  $f$ .

- Suppose  $h$  is such a function. Then

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \lim_{z \rightarrow z_0} h(z) = h(z_0)$$

by continuity. Formally, the first equality is holding for the limit in  $\Omega \setminus \{z_0\}$ , and the second equality is continuity for  $h|_{\Omega \setminus \{z_0\}}$ .

To finish, we note that the second part shows that  $h(z_0) = f'(z_0)$ . ■

## 3.2 February 14

Happy Valentine's Day, I suppose. Homework #4 is due on Sunday. Homework #5 will be released on Friday.

### 3.2.1 Motivating Cauchy–Riemann Equations

Today we're talking about the Cauchy–Riemann equations.



**Idea 3.16.** The Cauchy–Riemann equations are necessary conditions for a function to be holomorphic.

In fact, they will be sufficient as well, but we will only see this next class.

Throughout today's class, we will fix  $\Omega \subseteq \mathbb{C}$  a nonempty open subset. We recall that a function  $f : \Omega \rightarrow \mathbb{C}$  is “differentiable” at some  $z_0 \in \Omega$  if and only if the limit

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists. If it exists, we denoted it by  $f'(z_0)$ , though we will not assume it exists yet. If we fix  $\Delta z := z - z_0$ , then we can write the above as

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{\Delta z}.$$

Now, to motivate our discussion, we recall that under the isomorphism  $\mathbb{C} \cong \mathbb{R}^2$  with basis  $\{1, i\}$ , we can define  $u(x, y) := \operatorname{Re} f(x + yi)$  and  $v(x, y) := \operatorname{Im} f(x + yi)$  where  $u, v : \mathbb{R}^2 \rightarrow \mathbb{R}$  so that

$$f(x + yi) = u(x, y) + iv(x, y).$$

The point of this is to encode some geometry directly into our set-up.

**Example 3.17.** Given  $f(z) = z^2$ , we can plug in

$$f(x + yi) = (x + yi)^2 = \underbrace{x^2 - y^2}_u + i \cdot \underbrace{2xy}_v.$$

Now that we're moving things to  $\mathbb{R}^2$ , we will fix  $z_0 := x_0 + y_0i$  for  $x_0, y_0 \in \mathbb{R}$  with  $z = x + yi$  so that  $\Delta z = (x - x_0) + (y - y_0)i = \Delta x + i\Delta y$ . And for a little more convenience, we fix  $\Delta w := f(z_0 + \Delta z) - f(z_0)$  so that

$$f'(z_0) \stackrel{?}{=} \lim_{z \rightarrow z_0} \frac{\Delta w}{\Delta z},$$


if the limit exists. Expanding out  $f$  into real and imaginary parts, we find

$$\Delta w := (u(x_0 + \Delta x, y_0 + \Delta y) - u(x_0, y_0)) + i(v(x_0 + \Delta x, y_0 + \Delta y) - v(x_0, y_0)).$$

Now assume that  $f$  is in fact differentiable at  $z_0$  so that  $f'(z_0)$  will actually exist. Our key idea to continue is to split up the limit into real and imaginary parts because it will exist if and only if the limits of the real and imaginary parts exist. So we note

$$\begin{aligned} f'(z_0) &= \lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z} \\ &= \lim_{(\Delta x, \Delta y) \rightarrow 0} \operatorname{Re} \left( \frac{\Delta w}{\Delta z} \right) + i \lim_{(\Delta x, \Delta y) \rightarrow 0} \operatorname{Im} \left( \frac{\Delta w}{\Delta z} \right) \end{aligned} \quad (*)$$

We will now compute this limit in two ways to get the Cauchy–Riemann equations, as follows.



These are probably the easiest two limits that we could think of, so it's nice that they will be so useful. Anyways, here is our working out.

- We set  $\Delta y = 0$  so that  $\Delta z = \Delta x$ . This gives

$$\frac{\Delta w}{\Delta x} = \frac{u(x_0 + \Delta x, y_0) - u(x_0, y_0)}{\Delta x} + i \cdot \frac{v(x_0 + \Delta x, y_0) - v(x_0, y_0)}{\Delta x}.$$

On one hand, we can use  $(*)$  to show the real part comes out to

$$\operatorname{Re} f'(z_0) = \lim_{(\Delta x, \Delta y) \rightarrow 0} \operatorname{Re} \left( \frac{\Delta w}{\Delta x} \right) = \lim_{\Delta x \rightarrow 0} \frac{u(x_0 + \Delta x, y_0) - u(x_0, y_0)}{\Delta x}.$$

This limit must exist because  $f$  is differentiable at  $z_0$ , and when this limit exists, the rightmost limit is called the partial derivative  $u_x(x_0, y_0)$ .

On the other hand, the imaginary part comes out to

$$\operatorname{Im} f'(z_0) = \lim_{(\Delta x, \Delta y) \rightarrow 0} \operatorname{Im} \left( \frac{\Delta w}{\Delta x} \right) = \lim_{\Delta x \rightarrow 0} \frac{v(x_0 + \Delta x, y_0) - v(x_0, y_0)}{\Delta x},$$

which comes out to  $v_x(x_0, y_0)$  because we know that the limit exists.

So in total, we see  $f'(z_0) = u_x(x_0, y_0) + i \cdot v_x(x_0, y_0)$ .

- We set  $\Delta x = 0$  so that  $\Delta z = i\Delta y$ . Be warned that an unexpected sign is about to appear from this  $i$ . This time we get

$$\frac{\Delta w}{\Delta z} = \frac{u(x_0, y_0 + \Delta y) - u(x_0, y_0)}{i\Delta y} + i \cdot \frac{v(x_0, y_0 + \Delta y) - v(x_0, y_0)}{i\Delta y}.$$

To “rationalize” the denominators, we write

$$\frac{\Delta w}{\Delta z} = \frac{v(x_0, y_0 + \Delta y) - v(x_0, y_0)}{\Delta y} - i \cdot \frac{u(x_0, y_0 + \Delta y) - u(x_0, y_0)}{\Delta y},$$

where we are using  $1/i = -i$ . Note that the  $u$ s and  $v$ s have swapped from the last computation!

We now compute our limits. On one hand,

$$\operatorname{Re} f'(z_0) = \lim_{(\Delta x, \Delta y) \rightarrow 0} \operatorname{Re} \left( \frac{\Delta w}{\Delta z} \right) = \lim_{\Delta y \rightarrow 0} \frac{v(x_0, y_0 + \Delta y) - v(x_0, y_0)}{\Delta y},$$

which is  $v_y(x_0, y_0)$  because the limit exists. On the other hand,

$$\operatorname{Im} f'(z_0) = \lim_{(\Delta x, \Delta y) \rightarrow 0} \operatorname{Im} \left( \frac{\Delta w}{\Delta z} \right) = \lim_{\Delta y \rightarrow 0} -\frac{u(x_0, y_0 + \Delta y) - u(x_0, y_0)}{\Delta y},$$

which is  $-u_y(x_0, y_0)$  because the limit exists.

So in total, we see  $f'(z_0) = v_y(x_0, y_0) - iu_y(x_0, y_0)$ .

**Remark 3.18.** Either equation itself is pretty useful to actually compute formulae for the derivatives.

Synthesizing, we see

$$f'(z_0) = u_x(x_0, y_0) + i \cdot v_x(x_0, y_0) = v_y(x_0, y_0) - iu_y(x_0, y_0).$$

Comparing real and imaginary parts, we get the following.

**Theorem 3.19.** [Cauchy–Riemann] Fix  $\Omega \subseteq \mathbb{C}$  a nonempty open subset and  $f : \Omega \rightarrow \mathbb{C}$  a function differentiable at some  $z_0 = x_0 + y_0i \in \mathbb{C}$ . If we write  $f(x + yi) = u(x, y) + i(v(x, y))$ , then

$$\begin{cases} u_x(x_0, y_0) = v_y(x_0, y_0), \\ v_x(x_0, y_0) = -u_y(x_0, y_0). \end{cases}$$

In fact,  $f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0) = v_y(x_0, y_0) - iu_y(x_0, y_0)$ .

*Proof.* This follows from the above discussion. ■

### 3.2.2 Examples

Let’s see some examples to be convinced of the utility of Theorem 3.19. Let’s start by checking that something is differentiable.

**Example 3.20.** Take  $f(z) = z^2$  so that

$$(x + yi)^2 = (x + yi)(x + yi) = (x^2 - y^2) + i(2xy)$$

so that  $u(x, y) = x^2 - y^2$  and  $v(x, y) = 2xy$  has  $f(x + yi) = u(x, y) + iv(x, y)$ . We know that  $f$  is entire (it's impossible), so picking up any  $z = x + yi \in \mathbb{C}$ , we compute

$$u_x(x, y) = 2x = v_y(x, y) \quad \text{and} \quad v_x(x, y) = 2y = -(-2y) = -u_y(x, y),$$

verifying Theorem 3.19. In fact, we can see that  $f'(z) = u_x(x, y) + v_x(x, y)i = 2x + 2yi = 2z$ .

And now let's see something which isn't differentiable.

**Example 3.21.** Take  $f(z) = |z|^2$  so that

$$f(x + yi) = |x + yi|^2 = (x + yi)(x - yi) = x^2 + y^2,$$

which only has a real part! Namely, we have  $u(x, y) = x^2 + y^2$  and  $v(x, y) = 0$  to make  $f(x + yi) = u(x, y) + iv(x, y)$ . Now suppose for the sake of contradiction that  $f$  were differentiable at some  $z = x + yi \in \mathbb{C}$ . Then we are forced into

$$2x = u_x(x, y) = v_x(x, y) = 0 \quad \text{and} \quad 0 = v_y(x, y) = -u_y(x, y) = -2y,$$

which means  $x = y = 0$ . So  $f$  is differentiable at nowhere outside  $\mathbb{C} \setminus \{0\}$ .

Observe that the above example does not show that  $f$  is differentiable at  $0 \in \mathbb{C}$ , though this is true. To be explicit, Theorem 3.19 does not tell us that satisfying the Cauchy–Riemann equations implies differentiability.

**Remark 3.22.** Extending Example 3.21, we can show that the only entire real-valued function is constant.

Let's also close with an application of Theorem 3.19.

**Corollary 3.23.** Fix  $\Omega \subseteq \mathbb{C}$  a connected nonempty open subset and  $f : \Omega \rightarrow \mathbb{C}$  a function differentiable on all of  $\Omega$  so that  $f'(z) = 0$  for all  $z \in \Omega$ . Then  $f$  is constant.

*Proof.* By Theorem 3.19, we see that, for any  $z = x + yi$ , we see

$$u_x(x, y) = v_y(x, y) = \operatorname{Re} f'(z) = 0 \quad \text{and} \quad v_x(x, y) = -u_y(x, y) = \operatorname{Im} f'(z) = 0.$$

In particular, for some function  $g : C \rightarrow \mathbb{R}$  for some  $C \subseteq \mathbb{R}^2$  connected and open, having  $g_x = 0$  forces  $g$  to be constant as a function of  $x$  on any connected horizontal line, and  $g_y = 0$  forces  $g$  to be constant as a function of  $y$ .

Now, because any path between two points in an open subset can be approximated by vertical and horizontal line segments contained in neighborhoods of points, we see that the endpoints of any path in  $C$  must have the same value.<sup>1</sup> But  $C$  is open and connected and hence path-connected, so  $C$  any two points can be connected by path, so  $g$  must be constant on all of  $C$ .

Returning to  $f$ , we see that  $u$  and  $v$  will be constant on the embedding of  $\Omega$  into  $\mathbb{R}^2$  (recall that  $\mathbb{C} \cong \mathbb{R}^2$  topologically, so  $\Omega \subseteq \mathbb{R}^2$  remains open and connected), so  $f$  is constant on  $\Omega$ . This is what we wanted. ■

**Remark 3.24.** We do need the connected hypothesis: we could take  $\Omega = \mathbb{C} \setminus \mathbb{R}$  and with  $f(z) = 1_{\operatorname{Re} z > 0}$ .

<sup>1</sup> Please don't ask me to rigorize this.

### 3.3 February 16

We talk more about the Cauchy–Riemann equations today. For our announcements, Homework #4 is due on Sunday. There is a midterm next Friday; we will get a review sheet and some practice problems in the next few days. There will be no homework, and there will be extra office hours.

#### 3.3.1 Introducing Necessary Conditions

The slogan for today as follows.



**Idea 3.25.** The Cauchy–Riemann equations provide a sufficient condition for differentiability.

Recall our theorem.

**Theorem 3.19.** [Cauchy–Riemann] Fix  $\Omega \subseteq \mathbb{C}$  a nonempty open subset and  $f : \Omega \rightarrow \mathbb{C}$  a function differentiable at some  $z_0 = x_0 + y_0i \in \mathbb{C}$ . If we write  $f(x + yi) = u(x, y) + i(v(x, y))$ , then

$$\begin{cases} u_x(x_0, y_0) = v_y(x_0, y_0), \\ v_x(x_0, y_0) = -u_y(x_0, y_0). \end{cases}$$

In fact,  $f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0) = v_y(x_0, y_0) - iu_y(x_0, y_0)$ .

These are sufficient conditions for differentiability. Today we are discussing necessary conditions for differentiability.

**Theorem 3.26.** Fix  $\Omega \subseteq \mathbb{C}$  a nonempty open subset and  $f : \Omega \rightarrow \mathbb{C}$  a function. Writing  $f(x + yi) = u(x, y) + iv(x, y)$  and fixing some  $z_0 := x_0 + y_0i$ , then suppose we have the following.

- We have  $u_x, u_y, v_x, v_y$  all exist and are continuous (!).
- We have

$$\begin{cases} u_x(x_0, y_0) = v_y(x_0, y_0), \\ v_x(x_0, y_0) = -u_y(x_0, y_0). \end{cases}$$

Then  $f$  is differentiable at  $z_0$ .

**Remark 3.27.** It is possible to construct functions which are differentiable at  $z_0$  but do not have continuous first partial derivatives.

Let's do some examples of Theorem 3.26 to see its utility.

**Example 3.28.** Fix  $f(x + yi) = x^2 + y + i(y^2 - x)$ . Here,  $u(x, y) = x^2 + y$  and  $v(x, y) = y^2 - x$ , so we see

$$u_x(x, y) = 2x, \quad u_y(x, y) = 1, \quad v_x(x, y) = -1, \quad \text{and} \quad v_y(x, y) = 2y.$$

So all first partial derivatives are continuous. To satisfy the Cauchy–Riemann equations, we see that we need  $u_x = v_y$  and  $u_y = -v_x$ , which is equivalent to  $2x = 2y$  and  $1 = -(-1)$ . It follows from Theorem 3.26 that  $f$  is differentiable on the line  $y = x$ , and  $f$  is not differentiable anywhere else by Theorem 3.19.

**Remark 3.29.** Another type of question is to be given  $u(x, y)$  and be asked for what  $v(x, y)$  might be.

### 3.3.2 Proving Necessary Conditions

Let's go ahead and prove Theorem 3.26.

*Proof of Theorem 3.26.* As with last time, we fix  $\Delta z := z - z_0$  and  $\Delta x = x - x_0$  and  $\Delta y = y - y_0$  so that our difference quotient is

$$\frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} = \underbrace{\frac{u(x_0 + \Delta x, y_0 + \Delta y) - u(x_0, y_0)}{\Delta z}}_{\Delta u / \Delta z :=} + i \cdot \underbrace{\frac{v(x_0 + \Delta x, y_0 + \Delta y) - v(x_0, y_0)}{\Delta z}}_{\Delta v / \Delta z :=}.$$

So our goal is to show that

$$\lim_{\Delta z \rightarrow 0} \left( \frac{\Delta u}{\Delta z} + i \cdot \frac{\Delta v}{\Delta z} \right)$$

exists and is equal to  $u_x(x_0, y_0) + i v_x(x_0, y_0)$ . So we need to force our first partial derivatives into the limit.

We start with  $\Delta u / \Delta z$ . To make our partial derivatives appear, we write

$$\begin{aligned} \frac{\Delta u}{\Delta z} &= \frac{u(x_0 + \Delta x, y_0 + \Delta y) - u(x_0, y_0)}{\Delta z} \\ &= \frac{u(x_0 + \Delta x, y_0 + \Delta y) - u(x_0, y_0 + \Delta y)}{\Delta z} + \frac{u(x_0, y_0 + \Delta y) - u(x_0, y_0)}{\Delta z}. \end{aligned}$$

To get our partial derivatives, we apply the Mean value theorem (!): define

$$F(x) := u(x, y_0 + \Delta y) \quad \text{and} \quad F(y) := u(x_0, y).$$

We do our applications one at a time.

- Note that  $F(x)$  is differentiable everywhere from  $x_0$  to  $x_0 + \Delta x$ , so the Mean value theorem provides some  $x_0^*$  between  $x_0$  and  $x_0 + \Delta x$  such that

$$F(x_0 + \Delta x) - F(x_0) = F'(x_0^*) \Delta x.$$

- Similarly,  $F(y)$  is differentiable everywhere from  $y_0$  to  $y_0 + \Delta y$ , so the Mean value theorem provides some  $y_0^*$  between  $y_0$  and  $y_0 + \Delta y$  such that

$$F(y_0 + \Delta y) - F(y_0) = F'(y_0^*) \Delta y.$$

Synthesizing and plugging in, we get

$$\frac{\Delta u}{\Delta z} = \frac{u_x(x_0^*, y_0) \Delta x}{\Delta z} + \frac{u_y(x_0, y_0^*) \Delta y}{\Delta z}.$$

We now use continuity of our first partial derivative. Our hope is that sending  $\Delta z \rightarrow 0$  will send  $u_x(x_0^*, y_0) \rightarrow u_x(x_0, y_0)$  and  $u_y(x_0, y_0^*) \rightarrow u_y(x_0, y_0)$ . To show this, we show the difference will be small. We write

$$\frac{\Delta u}{\Delta z} = \frac{u_x(x_0, y_0) \Delta x}{\Delta z} + \frac{u_y(x_0, y_0) \Delta y}{\Delta z} + E_{ux} + E_{uy},$$

where

$$E_{ux} = (u_x(x_0^*, y_0) - u_x(x_0, y_0)) \frac{\Delta x}{\Delta z} \quad \text{and} \quad E_{uy} = (u_y(x_0, y_0^*) - u_y(x_0, y_0)) \frac{\Delta y}{\Delta z}.$$

We now remark that we can repeat the entire above argument for  $\frac{\Delta v}{\Delta z}$ . Namely, running the above machinery lets us write

$$\frac{\Delta v}{\Delta z} = \frac{v_x(x_0, y_0) \Delta x}{\Delta z} + \frac{v_y(x_0, y_0) \Delta y}{\Delta z} + E_{vx} + E_{vy},$$

where

$$E_{vx} = (v_x(x_0^{**}, y_0) - v_x(x_0, y_0)) \frac{\Delta x}{\Delta z} \quad \text{and} \quad E_{vy} = (v_y(x_0, y_0^{**}) - v_y(x_0, y_0)) \frac{\Delta y}{\Delta z}.$$

We now show that the various  $E_\bullet$  terms vanish as  $\Delta z \rightarrow 0$ . Note that, as  $\Delta z \rightarrow 0$ , the following happen.

- Because  $x_0^*$  and  $x_0^{**}$  are bounded between  $x_0$  and  $x_0 + \Delta x$ , they will approach  $x_0$ .
- Because  $y_0^*$  and  $y_0^{**}$  are bounded between  $y_0$  and  $y_0 + \Delta y$ , they will approach  $y_0$ .
- We will have  $\left| \frac{\Delta x}{\Delta z} \right| \leq 1$  and  $\left| \frac{\Delta y}{\Delta z} \right| \leq 1$  by direct expansion of the norm because  $\operatorname{Re} \Delta z = \Delta x$  and  $\operatorname{Im} \Delta z = \Delta y$ .

It follows that each of the  $E_\bullet$  do indeed vanish as  $\Delta z \rightarrow 0$ . For example,

$$\left| (u_x(x_0^*, y_0) - u_x(x_0, y_0)) \frac{\Delta x}{\Delta z} \right| \leq |u_x(x_0^*, y_0) - u_x(x_0, y_0)|$$

will go to 0 as  $\Delta z \rightarrow 0$  by the continuity of  $u_x$  at  $(x_0, y_0)$ .

Now we return to our difference quotient. We see

$$\begin{aligned} \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} &= \lim_{\Delta z \rightarrow 0} \left( \frac{\Delta u}{\Delta z} + i \cdot \frac{\Delta v}{\Delta z} \right) \\ &= \lim_{\Delta z \rightarrow 0} \left( \frac{u_x(x_0, y_0)\Delta x}{\Delta z} + \frac{u_y(x_0, y_0)\Delta y}{\Delta z} + i \cdot \frac{v_x(x_0, y_0)\Delta x}{\Delta z} + i \cdot \frac{v_y(x_0, y_0)\Delta y}{\Delta z} \right) \\ &\quad + \lim_{\Delta z \rightarrow 0} E_{ux} + \lim_{\Delta z \rightarrow 0} E_{uy} + \lim_{\Delta z \rightarrow 0} E_{vx} + \lim_{\Delta z \rightarrow 0} E_{vy} \\ &= \lim_{\Delta z \rightarrow 0} \left( \frac{u_x(x_0, y_0)\Delta x}{\Delta z} + \frac{u_y(x_0, y_0)\Delta y}{\Delta z} + i \cdot \frac{v_x(x_0, y_0)\Delta x}{\Delta z} + i \cdot \frac{v_y(x_0, y_0)\Delta y}{\Delta z} \right), \end{aligned}$$

using the fact that our error terms all vanish. At this point we use the Cauchy–Riemann equations. We see

$$\begin{aligned} \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} &= \lim_{\Delta z \rightarrow 0} \left( \frac{u_x(x_0, y_0)\Delta x}{\Delta z} - \frac{v_x(x_0, y_0)\Delta y}{\Delta z} + i \cdot \frac{v_x(x_0, y_0)\Delta x}{\Delta z} + i \cdot \frac{u_x(x_0, y_0)\Delta y}{\Delta z} \right) \\ &= \lim_{\Delta z \rightarrow 0} \left( u_x(x_0, y_0) \cdot \frac{\Delta x + i\Delta y}{\Delta z} \right) + i \cdot \lim_{\Delta z \rightarrow 0} \left( v_x(x_0, y_0) \cdot \frac{\Delta x + i\Delta y}{\Delta z} \right), \end{aligned}$$

which finishes after evaluating our first partial derivatives. ■

## 3.4 February 18

Good morning everyone. Here are some announcements.

- Homework #4 is due on Sunday.
- Next Friday is our midterm. A review sheet has been posted. Some practice problems and a practice midterm will be released today or tomorrow.
- Next week will have office hours every day.
- Next Wednesday will be a review class.

### 3.4.1 Power Series

Today we are building towards a discussion of analytic functions. We won't actually define what "analytic" means, but it will be important, so we will spend today setting up the definitions and results.

**Definition 3.30** (Complex power series). A complex power series is a formal expression of the form

$$S(z) := \sum_{k=0}^{\infty} a_k x^k$$

where  $\{a_k\}_{k \in \mathbb{N}} \subseteq \mathbb{C}$  and  $z$  is a (formal) variable taking complex values.

Our main goal for today is to be able to answer the following question.

**Question 3.31.** For which  $z$  will  $S(z)$  converge?

The answer to this question is essentially the same as for real analysis: it's the radius of convergence.

**Definition 3.32** (Radius of convergence). The *radius of convergence* of a complex power series  $S(z) = \sum_{k=0}^{\infty} a_k z^k$  is defined to be equal to the radius of convergence of the real power series

$$T(x) = \sum_{k=0}^n |a_k| x^k.$$

Concretely, we define

$$R := \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}}.$$

We should probably check convergence in the radius of convergence.

**Proposition 3.33.** Fix a complex power series  $S(z) = \sum_{k=0}^{\infty} a_k z^k$  with radius of convergence  $R$ . Then the following hold.

- (a) The sequence of partial sums  $\sum_{k=0}^n |a_k z^k|$  converge for any  $z$  with  $|z| < R$ . In other words,  $S(z)$  converges absolutely.
- (b) The series  $S(z)$  will diverge for  $z$  with  $|z| > R$ .

*Proof.* We take these one at a time. The point is to imitate the proofs from real analysis.

- (a) We note that, if  $R = 0$ , there is nothing to prove here. Otherwise, fix  $z$  with  $|z| < R$  so that there exists some  $\rho \in \mathbb{R}$  with  $|z| < \rho < R$ . For example,  $\rho := \frac{|z|+R}{2}$  will do.

Now, because  $\rho < R$ , we see that  $\frac{1}{\rho} > \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$  (this is legal because  $R \neq 0$ ), so there exists some  $N$  for which any fixed  $n > N$  has

$$\sup_{k \geq n} \sqrt[k]{|a_k|} < \frac{1}{\rho}.$$

In particular, each  $k > N$  will have  $\sqrt[k]{|a_k|} < 1/\rho$ , so  $|a_k| \rho^k < 1$ . So, setting

$$M := \max(\{1\} \cup \{|a_k| : k \leq N\}),$$

we see that  $|a_k| \rho^k \leq M$  for each  $k \in \mathbb{N}$ .

But because  $|z| < \rho$ , we note that  $|z|/\rho < 1$ , so we bound

$$|a_k z^k| = |a_k \rho^k| \cdot \left| \frac{z^n}{\rho^n} \right| \leq M \left| \frac{z}{\rho} \right|^n.$$

However,  $|z/\rho| < 1$ , so the series  $\sum_{k=0}^{\infty} |z/\rho|^k$  will converge as a geometric series, so we are done by the comparison test.

- (b) We proceed by contraposition. Suppose that  $S(z)$  converges, so by Lemma 2.71,  $a_k z^k \rightarrow 0$  as  $k \rightarrow \infty$ . We show that  $|z| \leq R$ . If  $z = 0$ , there is nothing to say; otherwise, it will suffice to show that

$$\frac{1}{|z|} \stackrel{?}{\geq} \limsup_{k \rightarrow \infty} \sqrt[k]{|a_k|}.$$

For this, fix  $\varepsilon = 1$ , so we are granted some  $N$  for which  $k > N$  has

$$|a_k z^k| < 1.$$



In particular, this rearranges into  $1/|z| > \sqrt[k]{|a_k|}$ . So for each  $n > N$ , we see  $1/|z| > \sqrt[k]{|a_k|}$  for  $k > n$ , so  $1/|z| \geq \sup\{\sqrt[k]{|a_k|} : k > n\}$ , so

$$\frac{1}{|z|} \geq \lim_{n \rightarrow \infty} \sup\{\sqrt[k]{|a_k|} : k > n\} = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|},$$

which is what we wanted. ■

**Remark 3.34** (Nir). The proof of (b) might feel weird because we are not using the full power of  $S(z)$  converging, just that its terms go to 0. However, a power series will “essentially” converge whenever its terms go to 0 (aside from boundary cases), so it is not too surprising that this is all that we need.

We have the following warning.



**Warning 3.35.** Proposition 3.33 is agnostic to the case of  $|z| = R$ .

In general, the behavior need not be uniform, as with  $\sum_{k=0}^{\infty} z^k = \frac{1}{1-z}$ .

### 3.4.2 Series of Functions

We will be interested in series of functions, which generalize power series.

**Definition 3.36** (Series of functions). Fix  $X \subseteq \mathbb{C}$  a nonempty set and  $\{f_k\}_{k \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$ . Then we define the *series of functions*

$$S(z) = \sum_{k=0}^{\infty} f_k(z)$$

for each  $z \in X$ .

Observe that the partial sums of some  $S(z) = \sum_{k=0}^{\infty} f_k(z)$  will look like some finite sum

$$S_m(z) = \sum_{k=0}^m f_k(z),$$

which defines a sequence of functions  $\{S_m\}_{m \in \mathbb{N}}$  where  $S_m : X \rightarrow \mathbb{C}$ . We are interested in the convergence of  $S$  as a function.

**Definition 3.37** (Uniform convergence). Fix  $X \subseteq \mathbb{C}$  a nonempty subset and  $\{f_k\}_{k \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$  defining a series of functions  $S(z) = \sum_{k=0}^{\infty} f_k(z)$ . Then  $S(z)$  *converges uniformly* if and only if the sequence of partial sums converges uniformly to a function on  $X$ .

Uniform convergence will be nice because (say) it will preserve continuity, but before talking about utility, we discuss a way to check uniform convergence.

**Theorem 3.38** (Weierstrass  $M$ -test). Fix  $X \subseteq \mathbb{C}$  a nonempty subset and  $\{f_k\}_{k \in \mathbb{N}}$  a sequence of functions  $X \rightarrow \mathbb{C}$  defining a series of functions  $S(z) = \sum_{k=0}^{\infty} f_k(z)$ . Further, suppose that, for each  $k \in \mathbb{N}$ , there exists some  $M_k$  such that

$$|f_k(z)| < M_k$$

for each  $z \in X$ . If  $\sum_{k=0}^{\infty} M_k$  converges, then  $S(z)$  converges uniformly.

In other words, we can determine uniform convergence of a series of functions by bounding the functions individually.

*Proof of Theorem 3.38.* This is not as hard as it looks. Let  $S_m$  denote the  $m$ th partial sum. By Proposition 2.135, it suffices to show that, for each  $\varepsilon > 0$ , there exists some  $N$  such that  $n \geq m > N$  implies

$$\sup_{x \in X} \{|S_n(z) - S_m(z)|\} < \varepsilon.$$

Well, we know that the series  $\sum_{k=0}^{\infty} M_k$  converges, so its partial sums are Cauchy, so there exists some  $N$  such that  $n \geq m > N$  implies

$$\sum_{k=m+1}^n M_k < \varepsilon,$$

where the left-hand side is the difference between the  $n$ th and  $m$ th partial sums. So now we bound

$$|S_n(z) - S_m(z)| = \left| \sum_{k=m+1}^n f_k(z) \right| \leq \sum_{k=m+1}^n |f_k(z)| \leq \sum_{k=m+1}^n M_k,$$

for any  $z \in X$ . Thus,

$$\sup_{z \in X} \{|S_n(z) - S_m(z)|\} \leq \sum_{k=m+1}^n M_k < \varepsilon.$$

This finishes the proof. ■

And now let's apply the Weierstrass  $M$ -test to power series.

**Corollary 3.39.** Fix  $S(z) = \sum_{k=0}^{\infty} a_k z^k$  a power series with positive radius of convergence  $R > 0$ . We have the following.

- (a) For any  $r$  such that  $0 < r < R$ , the power series  $S(z)$  converges uniformly in  $\overline{B(0, r)}$ .
- (b) The power series  $S(z)$  is continuous on  $B(0, r)$ .

*Proof.* Most of our work will be done in (a), which comes from the Weierstrass  $M$ -test.

- (a) Fix some  $r$  with  $0 < r < R$ . Note that  $S(r)$  converges absolutely by Proposition 3.33. To use the Weierstrass  $M$ -test, we set  $f_k(z) := a_k z^k$  with  $M_k := |a_k| r^k$  so that  $|z| \leq r$  implies

$$|f_k(z)| = |a_k z^k| = |a_k| \cdot |z|^k \leq |a_k| r^k.$$

But we know that  $S(r)$  converges absolutely, so

$$\sum_{k=0}^{\infty} |a_k r^k| = \sum_{k=0}^{\infty} M_k$$

converges, so now Theorem 3.38 promises that  $S(z)$  will converge uniformly for each  $z \in \overline{B(0, r)}$ .

- (b) Note that, for every  $k$ , the function  $f_k(z) = a_k z^k$  is a polynomial and hence entire and hence continuous on  $B(0, R)$ .

The trick is to apply (a) by starting with a fixed  $z_0 \in B(0, R)$  with  $r$  such that  $|z_0| < r < R$ . In particular, by restriction, it suffices to show that  $S|_{B(0, r)}$  is continuous at  $z_0$ . (For example,  $r = \frac{|z_0| + R}{2}$  will work.) So now we note that the continuous partial sums of  $S(z)$  converge uniformly to  $S(z)$  on  $B(0, r)$  by (a), so Proposition 2.138 forces  $S(z)$  itself to be continuous on  $B(0, r)$ . This finishes. ■

We remark that the restriction to  $S|_{B(0, r)}$  only works because  $B(0, r)$  is an open set. Here is the exact lemma we just used.

**Lemma 3.40.** Fix  $f : X \rightarrow \mathbb{C}$  a function and  $U \subseteq \mathbb{C}$  an open subset  $X$  with  $x \in U \cap X$ . Then  $f$  is continuous at  $x$  if and only if the restriction  $f|_{U \cap X} : U \cap X \rightarrow \mathbb{C}$  is continuous at  $x$ .

An alternate way to give the hypothesis on  $U$  is that  $U \cap X$  is an open subset of  $X$ .

*Proof.* We show the directions independently.

- Suppose that  $f$  is continuous at  $x$ . We show that  $f|_{U \cap X}$  is continuous at  $x$ . Well, for any  $\varepsilon > 0$ , we are promised some  $\delta > 0$  so that any  $z \in X$  has

$$|z - x| < \delta \implies |f(z) - f(x)| < \varepsilon.$$

In particular, any  $z \in X \cap U$  has

$$|z - x| < \delta \implies |f|_{U \cap X}(z) - f|_{U \cap X}(x)| = |f(z) - f(x)| < \varepsilon.$$

- Suppose that  $f|_{U \cap X}$  is continuous at  $x$ . Fix any  $\varepsilon > 0$ . Because  $x \in U$  and  $U$  is open, there exists  $r > 0$  such that  $B(x, r) \subseteq U$ . Because  $f|_{U \cap X}$  is continuous at  $x$ , there exists some  $\delta_0 > 0$  such that

$$|z - x| < \delta_0 \implies |f(z) - f(x)| = |f|_{U \cap X}(z) - f|_{U \cap X}(x)| < \varepsilon$$

for  $z \in U \cap X$ . However, taking  $\delta := \min\{r, \delta_0\}$ , we see that any  $z \in X$  with  $|x - z| < \delta$  will have  $z \in B(x, \delta) \subseteq U$ , so  $z \in U \cap X$  automatically. So  $|z - x| < \delta$  will still imply

$$|f(z) - f(x)| < \varepsilon,$$

and we are done. ■