18.755: Lie Groups and Lie Algebras II

Nir Elber

Spring 2026

Contents

How strange to actually have to see the path of your journey in order to make it.

—Neal Shusterman, [Shu16]

Contents 2

1 How to Use Highest Weights 5
1.1 February 2 . . . 5
1.1.1 Review of Lie Groups . . . 5
1.1.2 Review of Lie Algebras . . . 7
1.1.3 Fundamental Theorems . . . 8
1.1.4 Representations of Lie Algebras . . . 9
1.1.5 Structure Theory of Lie Algebras . . . 11
1.1.6 Root Systems . . . 12
1.2 February 4 . . . 13
1.2.1 Weights . . . 13
1.2.2 The Dynkin Diagram . . . 14
1.2.3 Back to Representations of Lie Algebras . . . 16
1.2.4 Weyl Formulae . . . 17
1.2.5 Representation Theory of $\mathfrak{sl}_n(\mathbb{C})$ . . . 19
1.3 February 9 . . . 22
1.3.1 Representations of $\mathrm{GL}_n(\mathbb{C})$ . . . 22
1.3.2 Schur–Weyl Duality . . . 23
1.3.3 Schur Functors . . . 25
1.4 February 11 . . . 27
1.4.1 Finding Invariant Polynomials . . . 27
1.4.2 Schur Polynomials . . . 29
1.4.3 Howe Duality . . . 31
1.5 February 17 . . . 32
1.5.1 Miniscule Weights . . . 32
1.5.2 Miniscule Representations . . . 34
1.5.3 Applications of Miniscule Representations . . . 35
1.6 February 18 . . . 37
1.6.1 The Conjugate Partition . . . 37

1.6.2 More on Miniscule Weights in Classical Types . . . 38
1.6.3 Miniscule Representations . . . 39
1.6.4 Fundamental Representations . . . 40
1.7 February 23 . . . 42
1.7.1 Spin Groups . . . 42
1.7.2 Clifford Algebras . . . 43
1.7.3 Spin Representations . . . 45
1.7.4 Dual Representations . . . 46
1.7.5 Maximal Root . . . 46
1.8 February 25 . . . 47
1.8.1 The Principal $\mathfrak{sl}_2$-triple . . . 47
1.8.2 Real, Complex, and Quaternionic Representations . . . 49

2 Integration **52**
2.1 March 2 . . . 52
2.1.1 Review of Topology . . . 52
2.1.2 Partition of Unity . . . 53
2.1.3 Integration on Manifolds . . . 55
2.1.4 Integration on Lie Groups . . . 56
2.1.5 Applications to Compact Lie Groups . . . 57
2.2 March 2 . . . 59
2.2.1 Stating the Peter–Weyl Theorem . . . 59
2.2.2 Compact Operators . . . 62
2.3 March 9 . . . 64
2.3.1 Proof of the Peter–Weyl Theorem . . . 64
2.3.2 Faithful Representations . . . 65
2.3.3 Haar Measures for Compact Groups . . . 67
2.4 March 11 . . . 70
2.4.1 Examples of Compact Groups . . . 70
2.4.2 Representation Theory of Compact Groups . . . 71
2.4.3 The Hydrogen Atom . . . 72
2.5 March 16 . . . 73
2.5.1 The Eigenfunctions . . . 74
2.5.2 Coulomb Waves . . . 76
2.6 March 18 . . . 77
2.6.1 The Pauli Exclusion Principle . . . 77
2.6.2 The Automorphism Group . . . 78

3 Structure Theory **81**
3.1 March 30 . . . 81
3.1.1 Forms of Semisimple Lie Algebras . . . 81
3.1.2 Real Forms . . . 83
3.1.3 Inner Forms . . . 83
3.1.4 The Compact Form . . . 84
3.2 April 1 . . . 86
3.2.1 Real Forms of Classical Types . . . 86
3.2.2 Counting from the Compact Form . . . 88
3.2.3 Vogan Diagrams . . . 89
3.2.4 Real Forms of Type A . . . 90
3.3 April 6 . . . 90
3.3.1 Linear Algebra over the Quaternions . . . 91
3.3.2 Real Forms of Classical Types . . . 92
3.4 April 8 . . . 94
3.4.1 Real Forms of Types F_4 and G_2 . . . 94

3.4.2 Real Forms of Type E . . . 95
3.5 April 13 . . . 98
3.5.1 Classification of Compact Connected Lie Groups . . . 98
3.5.2 Polar Decomposition . . . 101
3.6 April 15 . . . 102
3.6.1 Reductive and Linear Groups . . . 103
3.6.2 Maximal Tori . . . 104
3.6.3 Maximal Abelian Subspaces . . . 106
3.7 April 22 . . . 107
3.7.1 The Cartan Decomposition . . . 107
3.7.2 Maximal Compact Subgroups . . . 107
3.7.3 Cartan Subalgebras . . . 110

4 Cohomological Methods **112**
4.1 April 27 . . . 112
4.1.1 The Weyl Integration Formula . . . 112
4.1.2 Review of de Rham Cohomology . . . 113
4.1.3 Group Actions on Cohomology . . . 114
4.1.4 The Cohomology of a Lie Group . . . 116
4.1.5 Using Hopf Algebras . . . 117
4.2 April 29 . . . 119
4.2.1 The Exponents . . . 119
4.2.2 The Cohomology of Homogeneous Spaces . . . 121

Bibliography **125**

List of Definitions **126**

Theme 1

How to Use Highest Weights

1.1 February 2

Here we go.

1.1.1 Review of Lie Groups

We start with some quick review. Here are our groups.

Definition 1.1 (Lie group)**.** A *Lie group* is a group object G in the category of manifolds. One may specify a "real" or "complex" Lie group, which means that we are taking the category of real or complex manifolds. Explicitly, we are asking for G to be equipped with regular maps $m\colon G \times G \to G$, $i\colon G \to G$, and an identity. A *homomorphism of Lie groups* is a morphism of the group objects.

Example 1.2. One has the usual examples: $\mathbb{R}^n$, $\mathrm{U}(n)$, $\mathrm{Sp}_{2n}(\mathbb{R})$, $\mathrm{O}(p,q)$, and $\mathrm{SU}(n)$ are all real Lie groups.

Example 1.3. There are classical groups over $\mathbb{C}$, such as $\mathrm{SL}_n(\mathbb{C})$, which are all Lie groups.

Definition 1.4. If G is a Lie group, then its connected component G° is a normal Lie subgroup.

Remark 1.5. The quotient $\pi_0 G := G/G^\circ$ is a discrete topological group.

Remark 1.6. Given a Lie group G, the universal cover $\widetilde{G} \to G$ can be checked to a Lie group via some universal properties, so we receive a homomorphism $\pi\colon \widetilde{G} \to G$. It turns out that the kernel is a central discrete subgroup $Z \subseteq \widetilde{G}$. It notably follows that $\pi_1(G)$ is abelian.

Remark 1.7. One can check that G° is generated by any open neighborhood of the identity. Indeed, the generated subgroup can be seen to be both open and closed.

Example 1.8. With $G = S^1$, we have the universal cover $\widetilde{G} = \mathbb{R}$, and the kernel is $\mathbb{Z} \subseteq \mathbb{R}$.

We also have subgroups.

Definition 1.9 (Lie subgroup)**.** A *Lie subgroup* is an immersed submanifold $H \subseteq G$ which is also a subgroup, meaning that $H \hookrightarrow G$ admits injective differentials. A *closed Lie subgroup* is an embedded submanifold $H \subseteq G$ which is also a subgroup.

Remark 1.10. It turns out that closed Lie subgroups are in fact closed subsets, which can be checked locally.

Example 1.11. The subgroup $\mathbb{Q}^n \subseteq \mathbb{R}^n$ is a Lie subgroup, but it is not a closed Lie subgroup. The only closed Li subgroups are vector spaces.

Example 1.12. The subgroup $\mathrm{O}_n(\mathbb{R}) \subseteq \mathrm{GL}_n(\mathbb{R})$ is a closed real Lie subgroup.

Remark 1.13. It turns out that a closed subgroup of G is in fact a closed Lie subgroup. We will prove this later in the semester.

Definition 1.14 (quotient)**.** Fix a closed Lie subgroup $H \subseteq G$. Then G/H is a manifold with transitive G-action. If H is normal, then G/H is further a Lie group.

Remark 1.15. In general, if G acts transitively on a manifold X, then for any $x \in X$, $\mathrm{Stab}_G(x) \subseteq G$ is a closed Lie subgroup, and the quotient is isomorphic to X.

Remark 1.16. If G acts on a space X which is not transitive, then for any $x \in X$, the subset $Gx \subseteq X$ is at least an immersed submanifold.

Example 1.17. The group $\mathbb{R}$ has an action on $\mathbb{R}^2/\mathbb{Z}^2$ by $t\colon x \mapsto tx$. The orbit of (say), $(1/2, \sqrt{2}/2)$ is an immersed but not closed submanifold.

Definition 1.18 (representation)**.** Fix a Lie group G. A *representation* of a Lie group is a homomorphism $G \to \mathrm{GL}_n(\mathbb{C})$.

Example 1.19. Let G act on itself by conjugation. Then each $g \in G$ acts on $T_1G \to T_1G$, so we receive an adjoint representation $\mathrm{Ad}_\bullet\colon G \to \mathrm{GL}(T_1G)$.

As usual, one can define morphisms of representations, subrepresentations, direct sums, duals, tensor products, irreducible representations, and so on. We also have a Schur's lemma.

Lemma 1.20. Fix irreducible representations V and W of G.

(a) Then a G-equivariant map $\varphi\colon V \to W$ is either zero or an isomorphism.

(b) Any G-equivariant map $A \to A$ is a scalar.

Proof. Omitted. ■

Definition 1.21 (unitary)**.** A *unitary representation* is one admitting a G-invariant positive-definite Hermitian form.

Remark 1.22. Any unitary representation admits a decomposition into irreducible representations by taking orthogonal complements.

Non-Example 1.23. Let $B \subseteq \mathrm{GL}_2(\mathbb{C})$ be the subgroup of upper-triangular matrices. Then the standard representation of B does not admit a decomposition into irreducibles, so it cannot be made unitary.

Example 1.24. If G is finite, then any representation V admits a unitary structure: given any unitary structure $\langle -, - \rangle_0$, one can define an invariant unitary structure

$$\langle v, w \rangle := \frac{1}{|G|} \sum_{g \in G} \langle gv, gw \rangle_0,$$

where dg is a choice of Haar measure.

Theorem 1.25 (Maschke)**.** Fix a finite group G. Then all representations admit decomposition into irreducible representations.

Proof. This follows from Example 1.24. ■

1.1.2 Review of Lie Algebras

We now linearize our story.

Remark 1.26. Note that G acts on itself by left translations ℓ_g, so the tangent bundle TG can be given a global frame by the induced isomorphisms $d\ell_g \colon T_1 G \to T_g G$.

Notation 1.27. For each $a \in T_1 G$, we define the vector field L_a by

$$L_a := ga \in T_a G.$$

Remark 1.28. One can check that all left-invariant vector fields take the form L_a.

Definition 1.29 (commutator)**.** Fix a Lie group G. For each $a, b \in T_1 G$, we may take the commutator $[L_a, L_b]$ to produce another left-invariant vector field, which we label $L_{[a,b]}$.

Remark 1.30. The formalism of the commutator tells us that $[-,-]$ is antisymmetric and satisfies the Jacobi identity

$$[a, [b, c]] + [b, [c, a]] + [c, [a, b]] = 0.$$

Definition 1.31 (Lie algebra)**.** Fix a vector space $\mathfrak{g}$ over a field k. Then a *Lie algebra* is such a vector space $\mathfrak{g}$ equipped with an antisymmetric pairing $[-,-] \colon \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ satisfying the Jacobi identity

$$[a, [b, c]] + [b, [c, a]] + [c, [a, b]] = 0.$$

Example 1.32. For any Lie group G, we have seen that we may equip $\operatorname{Lie} G := T_1 G$ with the structure of a Lie group.

Example 1.33. If $G = \mathrm{GL}_n(\mathbb{C})$, then $\mathfrak{g} = M_n(\mathbb{C})$, and one can check that $[X, Y] = XY - YX$.

We now define Lie subalgebras and morphisms of Lie algebras in the expected way.

Definition 1.34 (Lie ideal)**.** Fix a Lie algebra $\mathfrak{g}$. Then a *Lie ideal* $\mathfrak{h} \subseteq \mathfrak{g}$ is a subspace for which $[\mathfrak{h}, \mathfrak{g}] \subseteq \mathfrak{h}$.

Example 1.35. For any closed Lie subgroup $H \subseteq G$, we see that $\operatorname{Lie} H \subseteq \operatorname{Lie} G$ is a Lie subalgebra. If H is normal, then $\operatorname{Lie} H$ is a Lie ideal.

As expected, there is some representation theory.

Definition 1.36. Fix a Lie algebra $\mathfrak{g}$ over a field k. Then a *representation* of $\mathfrak{g}$ is a morphism $\mathfrak{g} \to \mathfrak{gl}_n(k)$.

One can relate $\operatorname{Lie} G$ to G more directly via exponentiation.

Definition 1.37 (exponential)**.** Fix a Lie group G with Lie algebra $\mathfrak{g}$. We define a map $\exp\colon \mathfrak{g} \to G$ as follows. For each $a \in \mathfrak{g}$, one can check that the differential equation

$$\begin{cases} e'(t) = e(t) \cdot a, \\ e(0) = 1, \end{cases}$$

admits a unique solution; we then define $\exp(ta) := e(t)$. (This is independent of the choice of t.) It turns out that $t \mapsto \exp(ta)$ is a group homomorphism.

Example 1.38. If $G = \mathrm{GL}_n(\mathbb{C})$, then $\exp\colon M_n(\mathbb{C}) \to \mathrm{GL}_n(\mathbb{C})$ is the usual matrix exponential.

Remark 1.39. It turns out that $\exp$ is a local diffeomorphism (though not necessarily injective), so there is a local inverse $\log\colon U \to \mathfrak{g}$, where U is some open neighborhood of the identity.

Remark 1.40. For small a and b, it turns out that

$$\log(\exp(a)\exp(b)) = a + b + \frac{1}{2}[a, b] + \cdots,$$

where $\cdots$ denotes cubic terms. For example, if G is commutative, then we see that the Lie bracket $[-,-]$ vanishes; conversely, if $[-,-]$ vanishes, then G can be checked to commute in an open neighborhood of the identity, so G commutes.

1.1.3 Fundamental Theorems

In a first course, one checks the following two fundamental theorems.

Theorem 1.41. Fix a Lie group G. Then there is a bijection between connected closed Lie subgroups $H \subseteq G$ and Lie subalgebras $\mathfrak{h} \subseteq \operatorname{Lie} G$.

Theorem 1.42. Fix Lie groups G and K, with G simply connected. Then taking the differential

$$\mathrm{Hom}(G, K) \to \mathrm{Hom}(\mathrm{Lie}\, G, \mathrm{Lie}\, K)$$

is an isomorphism.

There is a third fundamental theorem, which we will prove later.

Theorem 1.43. For any finite-dimensional Lie algebra $\mathfrak{g}$ (over $\mathbb{R}$ or $\mathbb{C}$), then there is a Lie group G with $\mathrm{Lie}\, G \cong \mathfrak{g}$.

The three theorems provide an equivalence between the category of simply connected Lie groups and the category of Lie algebras, thereby classifying the former.

Remark 1.44. It follows that one may classify connected Lie groups as quotients of simply connected Lie groups by discrete central subgroups.

1.1.4 Representations of Lie Algebras

Let's start with the representation theory of $\mathfrak{sl}_2(\mathbb{C})$.

Theorem 1.45. Fix the usual basis $e := \left[\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right]$, $f := \left[\begin{smallmatrix} 0 & 0 \\ 1 & 0 \end{smallmatrix}\right]$, and $h := [e, f]$ of $\mathfrak{sl}_2(\mathbb{C})$.

(a) Then all irreducible representations of $\mathfrak{sl}_2(\mathbb{C})$ can be parameterized as $\{V_n\}_{n \geq 0}$, where V_n is the representation of homogeneous polynomials in x and y of degree n.

(b) Every representation is a direct sum of irreducible representations.

(c) Clebsch–Gordon rule: for any n and m, we have

$$V_n \otimes V_m = \bigoplus_{i=0}^{\min\{m,n\}} V_{|m-n|+2i}.$$

It will be helpful to turn representation theory of Lie algebras into a module category.

Definition 1.46 (universal enveloping algebra)**.** Fix a Lie algebra $\mathfrak{g}$. Then we define $U\mathfrak{g}$ as the quotient of the tensor algebra by the relation

$$[x, y] = x \otimes y - y \otimes x.$$

Remark 1.47. It turns out that $\mathrm{Rep}\, \mathfrak{g}$ is the same category as $\mathrm{Mod}\, U\mathfrak{g}$.

Even though we have taken a quotient by an inhomogeneous relation, $U\mathfrak{g}$ still receives a natural filtration by degree.

Theorem 1.48 (Poincaré–Birkhoff–Witt)**.** Fix a Lie algebra $\mathfrak{g}$, and equip $U\mathfrak{g}$ with the natural filtration. For any basis $\{x_1, \ldots, x_n\}$ of $\mathfrak{g}$, the ordered monomials in the basis form a basis of $U\mathfrak{g}$.

To continue our story, we need some adjectives for Lie algebras.

Definition 1.49 (solvable)**.** A Lie algebra $\mathfrak{g}$ is *solvable* if and only if the derived series eventually vanishes. Here, the derived series is defined inductively by $D^0(\mathfrak{g}) := \mathfrak{g}$ and $D^{n+1}(\mathfrak{g}) := [D^n(\mathfrak{g}), D^n(\mathfrak{g})]$ for each $n \geq 0$.

Definition 1.50 (nilpotent)**.** A Lie algebra $\mathfrak{g}$ is *nilpotent* if and only if the lower central series eventually vanishes. Here, the derived series is defined inductively by $L_0(\mathfrak{g}) := \mathfrak{g}$ and $L_{n+1}(\mathfrak{g}) := [L_n(\mathfrak{g}), \mathfrak{g}]$ for each $n \geq 0$.

Remark 1.51. One can see that nilpotent implies solvable.

The representation theory of solvable Lie algebras is quite easy.

Theorem 1.52 (Lie)**.** Fix a finite-dimensional solvable Lie algebra $\mathfrak{g}$ over an algebraically closed field of characteristic zero.

(a) Then every irreducible representation of $\mathfrak{g}$ is one-dimensional.

(b) Every representation admits a basis on which $\mathfrak{g}$ acts by upper-triangular matrices.

Theorem 1.53 (Engel)**.** Fix a finite-dimensional Lie algebra $\mathfrak{g}$. Then $\mathfrak{g}$ is nilpotent if and only if $\mathrm{ad}_X : \mathfrak{g} \to \mathfrak{g}$ is nilpotent for all $X \in \mathfrak{g}$.

Thus, we see that we will want to ignore solvable and nilpotent pieces.

Definition 1.54 (radical)**.** Fix a Lie algebra $\mathfrak{g}$. Then the *radical* $\operatorname{rad} \mathfrak{g}$ is the sum of all solvable ideals of $\mathfrak{g}$.

Remark 1.55. One can check that $\operatorname{rad} \mathfrak{g}$ is a solvable ideal, so it is automatically the largest solvable ideal.

Definition 1.56 (semisimple)**.** Fix a Lie algebra $\mathfrak{g}$. Then $\mathfrak{g}$ is *semisimple* if and only if $\operatorname{rad} \mathfrak{g} = 0$.

Remark 1.57. It turns out that

$$\mathfrak{g}_{\mathrm{ss}} := \frac{\mathfrak{g}}{\operatorname{rad} \mathfrak{g}}$$

is always semisimple. It turns out that the induced exact sequence splits, so there is a decomposition $\mathfrak{g} = \mathfrak{g}_{\mathrm{ss}} \ltimes \operatorname{rad} \mathfrak{g}$, which is known as the Levi decomposition; we will prove this later.

Having defined semisimple, we should define "simple."

Definition 1.58 (simple)**.** A Lie algebra $\mathfrak{g}$ is *simple* if and only if its only ideals are 0 and $\mathfrak{g}$.

Remark 1.59. One can check that semisimple Lie algebras are precisely the sums of simple Lie algebras.

It turns out to be convenient to allow a little radical.

Definition 1.60 (reductive)**.** A Lie algebra $\mathfrak{g}$ is *reductive* if and only if its radical is its center.

Example 1.61. One can check that $\mathfrak{sl}_n(\mathbb{C})$ is simple, and $\mathfrak{gl}_n(\mathbb{C})$ is reductive.

To test for a Lie algebra being semisimple (and other adjectives), we introduce the Killing form.

Definition 1.62 (Killing form)**.** Fix a Lie algebra $\mathfrak{g}$. Then we define the *Killing form* by

$$K(x, y) := \operatorname{tr}(\mathrm{ad}_x \circ \mathrm{ad}_y).$$

Remark 1.63. One can check that K is $\mathfrak{g}$-invariant.

Theorem 1.64 (Cartan criteria)**.** Fix a Lie algebra $\mathfrak{g}$.

(a) $\mathfrak{g}$ is solvable if and only if $[\mathfrak{g}, \mathfrak{g}] \subseteq K$.

(b) $\mathfrak{g}$ is semisimple if and only if K is non-degenerate.

Proposition 1.65. A Lie algebra $\mathfrak{g}$ is reductive if and only if it admits a representation $\rho\colon \mathfrak{g} \to \mathfrak{gl}(V)$ for which the bilinear form

$$B_V(X, Y) := \operatorname{tr}(\rho_X \circ \rho_Y)$$

is non-degenerate.

We may as well state one of the main theorems of our representation theory.

Theorem 1.66. Every finite-dimensional representation of a semisimple Lie algebra is completely reducible.

1.1.5 Structure Theory of Lie Algebras

Here is another piece of structure theory.

Definition 1.67 (adjoint)**.** Fix a semisimple Lie algebra $\mathfrak{g}$. Then we define the *adjoint* Lie group G^{ad} by $G^{\mathrm{ad}} := \operatorname{Aut}(\mathfrak{g})^\circ \subseteq \mathrm{GL}(\mathfrak{g})$.

Remark 1.68. It turns out that $\operatorname{Lie} G^{\mathrm{ad}} = \mathfrak{g}$.

In our setting, one can generalize the Jordan decomposition.

Definition 1.69 (semisimple, nilpotent)**.** An element $X \in \mathfrak{g}$ is *semisimple* or *nilpotent* if and only if the operator $\operatorname{ad}_X X$ is.

Theorem 1.70. Fix a Lie algebra $\mathfrak{g}$. Then any $X \in \mathfrak{g}$ can be written uniquely as a sum of a semisimple and nilpotent element.

Remark 1.71. It turns out that semisimple elements always act semisimply on representations, and nilpotent elements always act nilpotently on representations.

The notion of semisimple elements is important to define Cartan subalgebras.

Definition 1.72 (Cartan)**.** Fix a semisimple Lie algebra $\mathfrak{g}$. Then a *Cartan subalgebra* is a maximal commutative subalgebra of

Proposition 1.73. Fix a semisimple Lie algebra $\mathfrak{g}$. All Cartan subalgebras are conjugate by G^{ad}.

Definition 1.74. Fix a semisimple Lie algebra $\mathfrak{g}$. Then the *rank* of $\mathfrak{g}$ is the dimension of the Cartan subalgebras.

A choice of Cartan subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$ produces a root decomposition, which we write as

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \mathfrak{h}^* \setminus \{0\}} \mathfrak{g}_\alpha.$$

Definition 1.75 (root system)**.** Fix a semisimple Lie algebra $\mathfrak{g}$ and a Cartan subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$. Then the *root system* of $\mathfrak{g}$ consists of those nonzero eigenvalues $\alpha \in \mathfrak{h}^*$ for the adjoint action of $\mathfrak{h}$ on $\mathfrak{g}$. We write $\mathfrak{g}_\alpha$ for this eigenspace, and we write $\Phi(G)$ for the root system.

Remark 1.76. One can check that $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] \subseteq \mathfrak{g}_{[\alpha,\beta]}$. In fact, $\mathfrak{g}_\alpha$ and $\mathfrak{g}_\beta$ are orthogonal for the Killing form except when $\alpha = -\beta$, where it is a perfect pairing.

Remark 1.77. It turns out that $\dim \mathfrak{g}_\alpha = 1$ for each α. It follows that

$$\#\Phi(\mathfrak{g}) = \dim \mathfrak{g} - \operatorname{rank} \mathfrak{g}.$$

Remark 1.78. There are the usual pictures of root systems of various types.

1.1.6 Root Systems

It is useful to write down what properties are satisfied by these root systems.

Definition 1.79 (root system)**.** Fix a Euclidean space E. Then a finite subset $\Phi \subseteq E$ is a *root system* if and only if

(a) Φ spans E,

(b) for each $\alpha, \beta \in \Phi$, the number

$$n_{\alpha\beta} := \frac{2(\alpha, \beta)}{(\alpha, \alpha)}$$

is an integer,

(c) for each $\alpha, \beta \in \Phi$, the reflection

$$s_\alpha(\beta) := \beta - n_{\alpha\beta}\beta$$

is in Φ.

We say that Φ is *reduced* if and only if $\alpha \in \Phi$ implies that $2\alpha \notin \Phi$.

Definition 1.80 (reducible)**.** A root system Φ is *reducible* if and only if it can be written as a disjoint union of root systems coming from a decomposition of the Euclidean space into a product of Euclidean spaces.

The reflections are important enough to be placed into a group.

Definition 1.81 (Weyl group)**.** Fix a root system $\Phi \subseteq E$. Then the *Weyl group* W is the subgroup of $\mathrm{GL}(E)$ generated by the reflections.

Example 1.82. The Weyl group associated to the root system of $\mathfrak{sl}_{n+1}$ consists of the permutation matrices in $\mathrm{GL}_n(\mathbb{R})$. Indeed, each reflection corresponds to a transposition. This root system is said to be of type A_n, where n refers to the rank.

Remark 1.83. There is a determinant $\det$ on the Weyl group, which we define to be the determinant of the natural representation of W in $\mathrm{GL}(E)$.

Example 1.84. The root system associated to $\mathfrak{so}_{2n+1}$ is B_n. The root system associated to $\mathfrak{sp}_{2n}$ is C_n. Lastly, the root system associated to $\mathfrak{so}_{2n}$ is D_n.

Remark 1.85. There are also various exceptional reduced root systems, which we may say something about later.

We can even break down irreducible root systems into more controlled pieces.

Definition 1.86 (positive)**.** Fix a root system $\Phi \subseteq E$. For a choice of $t \in E$ for which $(t, \alpha) \neq 0$ for all $\alpha \in E$, we say that a root in Φ is *positive* if and only if $(t, \alpha) > 0$. Similarly, α is *negative* if and only if $(t, \alpha) < 0$. We let Φ^+ and Φ^- denote the sets of positive and negative roots, respectively.

Definition 1.87. Fix a root system $\Phi \subseteq E$. A positive root is *simple* if and only if it is not a sum of other positive roots (with positive integer coefficients). We let Π denote the set of simple roots.

Proposition 1.88. Fix a root system $\Phi \subseteq E$. Then Π is a basis, and every positive root $\alpha \in \Phi^+$ can be written as a unique sum of elements of Π with positive integer coefficients.

Each root system also admits a dual.

Definition 1.89 (dual root system)**.** Fix a root system $\Phi \subseteq E$. Then we define the *dual root system* $\Phi^\vee \subseteq E^\vee$ to be given by the points

$$\alpha^\vee = \frac{2(\alpha, -)}{(\alpha, \alpha)}$$

for each $\alpha \in \Phi$.

Remark 1.90. The reduced root system B_n is dual to C_n.

1.2 February 4

We continue our review today.

1.2.1 Weights

It will be helpful to have some lattices from our root systems.

Definition 1.91. Fix a root system $\Phi \subseteq E$.

- The *root lattice* Q is spanned by Φ.
- The *coroot lattice* $Q^\vee$ is spanned by the $\alpha^\vee$.
- The *weight* lattice $P \subseteq E$ is $(Q^\vee)^*$.
- The *coweight lattice* $P^\vee \subseteq E^*$ is Q^*.

In general, $Q \subseteq P$, but equality does not have to hold.

Example 1.92. For $\mathfrak{sl}_n$, the quotient P/Q is isomorphic to $\mathbb{Z}/n\mathbb{Z}$.

It is useful to have a basis for the weight lattice

Definition 1.93 (fundamental weight)**.** Fix a root system $\Phi \subseteq E$. A *fundamental weight* is an element of the dual basis (in the weight lattice P) of the $\alpha_\bullet^\vee \in Q^\vee$ where α is a simple root. Explicitly, if $\{\alpha_1, \ldots, \alpha_r\}$ are the simple roots, then the fundamental weights $\{\omega_1, \ldots, \omega_r\}$ satisfy

$$(\omega_i, \alpha_j^\vee) = 1_{i=j}.$$

Definition 1.94 (dominant)**.** Fix a root system $\Phi \subseteq E$. A weight λ is *dominant* if and only if $\langle \lambda, \alpha_i^\vee \rangle$ is a nonnegative integer for all i.

Remark 1.95. One can check that the dominant weights are exactly the $\mathbb{Z}_{\geq 0}$-span of the fundamental weights.

1.2.2 The Dynkin Diagram

Let's start trying to classify our Lie algebras.

Definition 1.96 (Cartan matrix)**.** Fix a root system $\Phi \subseteq E$, and order the simple roots as $\{\alpha_1, \ldots, \alpha_r\}$. The matrix A with entries

$$a_{ij} := \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)}$$

is the *Cartan matrix*.

Remark 1.97. The Cartan matrix satisfies the following properties, all essentially by construction.

- We have $a_{ii} = 2$ for each i.
- For distinct i and j, we have $a_{ij} \leq 0$, and a_{ij} vanishes if and only if a_{ji} vanishes.
- Because the inner product $(-,-)$ on the $\alpha_\bullet$s is positive definite, the pairing $(v, w) := v^\intercal A w$ is positive definite.

Remark 1.98. For any i and j, a piece of the Cartan matrix

$$\begin{bmatrix} 2 & a_{ij} \\ a_{ji} & 2 \end{bmatrix}$$

must have positive determinant by the positive-definiteness. Thus, $4 - a_{ij}a_{ji} > 0$, so $a_{ij}a_{ji} \in [0, 1, 2, 3]$ follows.

Example 1.99. The Cartan matrix for $\mathfrak{sl}_4$ is

$$\begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}.$$

The data of a Cartan matrix can be encoded combinatorially into a Dynkin diagram.

Definition 1.100 (Dynkin diagram)**.** Fix a root system $\Phi \subseteq E$, and order the simple roots as $\{\alpha_1, \ldots, \alpha_r\}$. Let A be the Cartan matrix. Then we define the *Dynkin diagram* to have r vertices and the following connections.

- If $a_{ij} = 0$, then vertices i and j have no connection.
- If $a_{ij} = -1$ and $a_{ji} = -1$, we draw a single line between i and j.
- If $a_{ij} \leq -2$, then we draw an arrow from vertex i to j with $|a_{ij}|$ heads.

There are the usual pictures of all Dynkin diagrams.

Example 1.101. For example, A_{n-1} is a path with $n - 1$ vertices.

$$\bullet \text{———} \bullet \text{———} \bullet \text{———} \cdots \text{———} \bullet$$

Remark 1.102. It turns out that the root system determines the Lie algebra, the Cartan matrix determines the root system, and the Dynkin diagram determines the Cartan matrix.

Remark 1.103 (Coxeter group)**.** One can read the Weyl group off of the Cartan matrix: namely, the reflection $s_i s_j$ has order

$$m_{ij} := \begin{cases} 2 & \text{if } a_{ij}a_{ji} = 0, \\ 3 & \text{if } a_{ij}a_{ji} = 1, \\ 4 & \text{if } a_{ij}a_{ji} = 2, \\ 6 & \text{if } a_{ij}a_{ji} = 3. \end{cases}$$

Indeed, this claim amounts to checking that $s_i s_j$ is a rotation with specified degree, which can be seen on the inner products. It turns out that the Weyl group is generated by the relations $s_i^2 = 1$ and $(s_i s_j)^{m_{ij}} = 1$ for each pair of distinct i and j.

To prove Remark 1.102, a difficult step is to show that every Dynkin diagram does in fact give rise to a Lie algebra. This is the content of the following theorem.

Theorem 1.104 (Serre presentation)**.** Fix a finite-dimensional simple Lie algebra $\mathfrak{g}$ over $\mathbb{C}$ with root system Φ, and choose an ordered set of simple roots $\Pi = \{\alpha_1, \ldots, \alpha_r\}$. Select basis vectors e_i and f_i in each space $\mathfrak{g}_{\alpha_i}$ and $\mathfrak{g}_{-\alpha_i}$, and define $h_i := [e_i, f_i]$. Then we have the following relations.

(a) We have $[h_i, h_j] = 0$, $[h_i, e_j] = a_{ij} e_j$, and $[h_i, f_j] = -a_{ij} f_j$.

(b) We have $[e_i, f_j] = 1_{i=j} h_i$.

(c) Serre relations: $\mathrm{ad}_{e_i}^{1-a_{ij}} e_j = 0$ and $\mathrm{ad}_{f_i}^{1-a_{ij}} f_j = 0$.

In fact, the free Lie algebra defined by these generators and relations (for any root system Φ) produces a finite-dimensional Lie algebra.

1.2.3 Back to Representations of Lie Algebras

Throughout, we fix a finite-dimensional Lie algebra $\mathfrak{g}$, choose a Cartan subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$. Then we choose a collection of positive roots for the induced root system, which lets us split $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{n}_+ \oplus \mathfrak{n}_-$, where $\mathfrak{n}_+$ is the sum of the positive root spaces, and $\mathfrak{n}_-$ is the sum of the negative root spaces.

Definition 1.105 (Verma module)**.** Fix a weight $\lambda \in \mathfrak{h}^*$. Then we define the *Verma module* M_λ to be generated as a $U\mathfrak{g}$-module by the vector v_λ, given the relations $ev_\lambda = 0$ for $e \in \mathfrak{n}_+$ and $hv_\lambda = \lambda(h)v_\lambda$ for $h \in \mathfrak{h}$.

Remark 1.106. In other words, $M_\lambda = U\mathfrak{g} \otimes_{U(\mathfrak{h} \oplus \mathfrak{n}_+)} \mathbb{C}_\lambda$, where $\mathbb{C}_\lambda$ is the representation of $\mathfrak{h} \oplus \mathfrak{n}_+$ on which $\mathfrak{h}$ acts by λ and $\mathfrak{n}_+$ acts by zero. For example, the PBW theorem gives us an isomorphism

$$U\mathfrak{n}_- \otimes U(\mathfrak{h} \oplus \mathfrak{n}_+) \to U\mathfrak{g}$$

of vector spaces, so M_λ is a free module over $U\mathfrak{n}_-$ of rank 1.

Remark 1.107. For generic λ, it turns out that M_λ is irreducible.

Definition 1.108 (highest weight)**.** Fix a representation V of $\mathfrak{g}$. Then a weight λ is a *highest weight* if and only if V is a nonzero quotient of M_λ. In other words, there is a vector $v \in V$ on which $\mathfrak{h}$ acts by λ and $\mathfrak{n}_+$ acts by zero.

Our classification result is as follows.

Theorem 1.109. Fix everything as above.

(a) The Verma modules M_λ admit a unique irreducible quotient.

(b) The representation L_λ is finite-dimensional if and only if λ is a dominant weight.

(c) Every irreducible finite-dimensional representation of $\mathfrak{g}$ takes the form L_λ for a dominant weight λ.

Proof. The proof of (a) is not too hard, and it is also not hard to show that L_λ being finite-dimensional, then λ is dominant. The hard part is to show that λ being dominant implies that L_λ is finite-dimensional!

Lastly, by iteratively acting by $\mathfrak{n}_+$, one can show that every irreducible finite-dimensional representation V of $\mathfrak{g}$ admits some highest weight λ. Indeed, given a finite-dimensional irreducible representation V of $\mathfrak{g}$,

the representation theory of $\mathfrak{sl}_2$ implies that the action of $\mathfrak{h}$ on V diagonalizes and admits integer eigenvalues. It follows that there is a surjection $M_\lambda \twoheadrightarrow V$, so there is a surjection $L_\lambda \twoheadrightarrow V$, which must then be an isomorphism. ■

Remark 1.110. Here is another way to select the highest weight: one can give the weight lattice a partial ordering, and then the highest one is the maximal element among the weights λ for which $V[\lambda] \neq 0$. In particular, the maximality implies that $\mathfrak{n}_+$ acts by zero on $V[\lambda]$.

There is also a construction of the quotient L_λ by taking relations.

Theorem 1.111. Fix everything as above, and choose a dominant weight λ. Then L_λ is the quotient of M_λ by the relations

$$f_i^{(\lambda,\alpha_i^\vee)+1} v_\lambda = 0.$$

Remark 1.112. These relations are forced by passing to the representation theory of $\mathfrak{sl}_2$.

1.2.4 Weyl Formulae

Our construction of L_λ is not so explicit; for example, what is the dimension of L_λ? To answer this question, we will need the Weyl dimension formula. We start with the Weyl character formula.

Definition 1.113 (character)**.** Fix a representation V of a semisimple Lie algebra $\mathfrak{g}$ admitting a weight decomposition

$$V = \bigoplus_{\mu \in P} V[\mu].$$

Then we define the *character*

$$\operatorname{ch} V := \sum_{\mu \in P} \dim V[\mu] e^\mu$$

living in the free power series vector space with basis given by the e^μs, with a relation $e^{\mu+\lambda}$.

Remark 1.114. If V is finite-dimensional, then it admits a weight decomposition (by the representation theory of $\mathfrak{sl}_2$), and the character is a finite sum.

Remark 1.115. Let's explain why this is a character. Suppose that V is a representation of a Lie group G with Lie algebra $\mathfrak{g}$. For $h \in \mathfrak{h}$, the exponential $\exp(h) \in G$ acts on V diagonally, with trace

$$\operatorname{tr}_V \exp(h) = \sum_{\mu \in P} \dim V[\mu] e^{\mu(h)},$$

which is exactly $\operatorname{ch} V(h)$. In fact, these formulae determine the entire character because semisimple elements can all be found in Cartan subalgebras and are dense in G.

We are now ready to state the Weyl character formula.

Theorem 1.116 (Weyl character formula)**.** Fix a semisimple Lie algebra $\mathfrak{g}$ with root system Φ, and let Φ^+ be a set of positive roots. Further, set $\rho := \frac{1}{2}\sum_{\alpha\in\Phi^+}\alpha$. Then for each dominant weight λ, we have

$$\operatorname{ch} L_\lambda = \frac{\displaystyle\sum_{w\in W}\det(w)e^{w(\lambda+\rho)-\rho}}{\displaystyle\prod_{\alpha\in\Phi_+}(1-e^{-\alpha})}.$$

Proof. Omitted. The idea is that the character of M_λ is easy to compute, and one can build a resolution of L_λ in terms of Verma modules. ■

Remark 1.117. One can write the right-hand side more symmetrically as

$$\frac{\displaystyle\sum_{w\in W}\det(w)e^{w(\lambda+\rho)}}{\displaystyle\prod_{\alpha\in\Phi_+}\left(e^{\alpha/2}-e^{-\alpha/2}\right)}.$$

Remark 1.118. A priori, the right-hand side is a rational function, but one can use the anti-symmetry of the Weyl group action in the numerator to show that the denominator does in fact divide the numerator.

Example 1.119 (Weyl denominator)**.** If $\lambda = 0$, then L_λ is the trivial representation, so it follows that

$$\sum_{w\in W}\det(w)e^{w\rho} = \prod_{\alpha\in\Phi_+}\left(e^{\alpha/2}-e^{-\alpha/2}\right).$$

Example 1.120 (Vandermonde determinant)**.** Suppose that $\mathfrak{g} = \mathfrak{sl}_n$ so that $\alpha_{ij} = x_i - x_j$. Then we set $X_i := e^{x_i}$, so we find that

$$\sum_{w\in S_n}\left(\operatorname{sgn}(w)\prod_{i=1}^n X_{w(i)}^{i-1}\right) = \prod_{i<j}(X_i - X_j).$$

Note that the right-hand side is precisely the determinant of the matrix

$$\begin{bmatrix} 1 & X_1 & \cdots & X_1^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & X_n & \cdots & X_n^{n-1} \end{bmatrix}.$$

By Theorem 1.116, we could attempt to compute $\dim L_\lambda$ by plugging in $h = 0$ to the polynomial.

Theorem 1.121 (Weyl dimension formula)**.** Fix a semisimple Lie algebra $\mathfrak{g}$ with root system Φ, and let Φ^+ be a set of positive roots. Then for any dominant weight λ,

$$\dim L_\lambda = \prod_{\alpha\in\Phi_+}\frac{(\alpha^\vee, \lambda+\rho)}{(\alpha^\vee, \rho)}.$$

Proof. This is slightly complicated because the given denominator vanishes at $h = 0$, so we are left to compute the limit

$$\dim L_\lambda = \lim_{h \to 0} \frac{\sum_{w \in W} \det(w) e^{(w(\lambda+\rho),h)}}{\prod_{\alpha \in \Phi_+} \left(e^{(\alpha,h)/2} - e^{-(\alpha,h)/2}\right)}.$$

This limit looks difficult to compute, but we may merely compute it on the line $h := 2th_\rho$, where h_ρ is the dual element for $\rho \in h^*$. In particular, it turns out that the numerator will factor in this case! To see this, note $\alpha(h_\rho) = (\alpha, \rho)$, so

$$\sum_{w \in W} \det(w) e^{(w(\lambda+\rho),2th_\rho)} = \sum_{w \in W} \det(w) e^{2t(w(\lambda+\rho),\rho)},$$

which upon moving the w around and plugging into Example 1.119 gives

$$\prod_{\alpha \in \Phi_+} \left(e^{t(\alpha,\lambda+\rho)} - e^{-t(\alpha,\lambda+\rho)}\right).$$

The denominator is now

$$\prod_{\alpha \in \Phi_+} \left(e^{(\alpha,h)/2} - e^{-(\alpha,h)/2}\right) = \prod_{\alpha \in \Phi_+} \left(e^{t(\alpha,\rho)} - e^{-t(\alpha,\rho)}\right).$$

Sending $t \to 0$ (and cancelling out a factor of two) proves the result. ■

1.2.5 Representation Theory of $\mathfrak{sl}_n(\mathbb{C})$

The dominant weights ω_i are the "simplest," so the simplest interesting representations will be the L_{ω_i}s. For example, we may hope to find all other representations inside them.

Proposition 1.122. Fix a semisimple Lie algebra $\mathfrak{g}$ with root system Φ, and let $\Pi = \{\alpha_1, \ldots, \alpha_r\}$ be an ordered set of simple roots. Further, let $\{\omega_1, \ldots, \omega_r\}$ be the set of dominant weights. Then the category finite-dimensional representations of $\mathfrak{g}$ is $\otimes$-generated by the representations $\{L_{\omega_i}\}_{i=1}^r$.

Proof. Because any finite-dimensional representation is a direct sum of the irreducible representations of the form L_λ, where λ is a dominant weight, it is enough to find these representations inside some tensor product of L_{ω_i}s. Well, λ is dominant, so we may write $\lambda = \sum_{i=1}^r k_i \omega_i$ for some nonnegative integers $\{k_1, \ldots, k_r\}$. We then consider the representation

$$V := \bigotimes_{i=1}^r L_{\omega_i}^{k_i}.$$

For any dominant weight μ, recall that μ is the highest weight of L_μ, meaning that the weights of L_μ are concentrated in $\mu - P^+$, where P^+ consists of the dominant weights. Additionally, the nature of the $\mathfrak{g}$-action on the tensor product implies that $W[\mu] \otimes W'[\mu'] = (W \otimes W')[\mu + \mu']$ for any W and W' and μ and μ'. Thus, we can see that the highest weight of V is $\lambda = \sum_{i=1}^r k_i \omega_i$, and it has multiplicity 1. It follows that there is a map $M_\lambda \to V$, which must descend to an embedding $L_\lambda \hookrightarrow V$, as desired. ■

Remark 1.123. The proof also shows that the other L_μs appearing in the representation V have $\mu < \lambda$. Indeed, one can simply calculate the weights appearing in V/L_λ and note that they are all strictly smaller than λ. The multiplicities of the various L_μ appearing in V can be interesting to calculate.

Remark 1.124. In fact, we can see that L_λ is the irreducible subrepresentation generated by the highest weight vector

$$v_\lambda = v_{\omega_1}^{\otimes k_1} \otimes \cdots \otimes v_{\omega_r}^{\otimes k_r}$$

in V. Indeed, the image of L_λ in V certainly contains this vector because this is the unique line with weight λ, so it follows that the generated subrepresentation must be exactly the irreducible representation L_λ.

Let's work out our general theory in an interesting example.

Example 1.125. Let V_{std} be the standard representation of $\mathfrak{sl}_n(\mathbb{C})$. There is a choice of ordered simple roots of $\mathfrak{sl}_n(\mathbb{C})$ for which

$$L_{\omega_i} \cong \wedge^i V_{\mathrm{std}}.$$

Proof. We proceed in steps.

1. We set up notation. Here, $\mathfrak{h}$ is the set of diagonal matrices in $\mathfrak{sl}_n(\mathbb{C})$, which we embed into $\mathbb{C}^n$ as the subspace with sum zero. Let $\{e_1, \ldots, e_n\}$ be the natural basis of $\mathbb{C}^n$, so we find that

   $$\Phi = \{e_i - e_j : i \neq j\}.$$

   Indeed, the elementary matrix E_{ij} has weight $e_i - e_j$. By choosing the vector $t := (n, n-1, n-2, \ldots, 1, 0)$ in $\mathbb{C}^n$, we see that Φ^+ consists of those $e_i - e_j$ with $i < j$. Then the simple roots are given by $\alpha_i := e_i - e_{i+1}$, which we can check is in fact a basis.

   For example, we see that this implies that the positive spaces in $\mathfrak{g}$ are given by

   $$\mathfrak{n}_+ := \bigoplus_{p<q} \mathbb{C}E_{pq}.$$

2. We calculate the fundamental weights. Now, the dual space $\mathfrak{h}^*$ is then the quotient of $\mathbb{C}^n$ by the diagonal subspace $\mathbb{C}(1, \ldots, 1)$. To compute our fundamental weights ω_i, we see that we are on the hunt for some vectors ω_i for which

   $$(\omega_i, e_j - e_{j+1}) = 1_{i=j}.$$

   Well, $\omega_i = (1, \ldots, 1, 0, \ldots, 0)$ works, where there are i total 1s. Accordingly, our dominant weights are the $\mathbb{Z}_{\geq 0}$-linear combinations of the ω_i, so they are the decreasing sequences of nonnegative integers.

3. We claim that the representation $\wedge^i V_{\mathrm{std}}$ is irreducible. Well, let $W \subseteq \wedge^i V_{\mathrm{std}}$ be some nonzero subrepresentation, and we want to show that $W = \wedge^i V_{\mathrm{std}}$. Give V_{std} the standard basis $\{v_1, \ldots, v_n\}$, and we note that this basis is an eigenbasis for the $\mathfrak{h}$-action on V_{std}. Thus,

   $$\{v_{n_1} \wedge \cdots \wedge v_{n_i} : n_1 < \cdots < n_i\}$$

   is an eigenbasis for the $\mathfrak{h}$-action on $\wedge^i V_{\mathrm{std}}$. Because $W \subseteq V$ is a $\mathfrak{g}$-subrepresentation, we conclude that it is an $\mathfrak{h}$-subrepresentation, so W must contain some eigenvector $v_{n_1} \wedge \cdots \wedge v_{n_i}$.

   Now, for any $p \neq q$, we see that $E_{pq} \in \mathfrak{sl}_n(\mathbb{C})$ acts by zero on all basis vectors $v_\bullet$ except it sends $v_q \mapsto v_p$. Thus, we see that we may apply such elementary matrices to any given eigenvector $v_{n_1} \wedge \cdots \wedge v_{n_i}$ to move it to $v_1 \wedge \cdots \wedge v_i$ (namely, apply E_{1n_1}, then E_{2n_2}, and so on), and one can apply the process in reverse to move any $v_1 \wedge \cdots \wedge v_i$ to any other eigenvector. We conclude that W being stable under $\mathfrak{g}$ forces $W = \wedge^i V$.

4. We are now ready to show that $L_{\omega_i} \cong \wedge^i V_{\mathrm{std}}$. Because $\wedge^i V_{\mathrm{std}}$ is already irreducible, it is enough to check that its highest weight is ω_i. Well, we claim that the highest weight vector is

   $$w := v_1 \wedge \cdots \wedge v_i.$$

Well, for some $p < q$, we can see that $E_{pq}w = 0$ because E_{pq} can only ever send a basis vector v_q to the "earlier" basis vector v_p, which either causes w to vanish outright (if $> i$) or introduces a multiplicity into w, still causing $E_{pq}w = 0$. Thus, we do indeed see that $\mathfrak{n}_+ w = 0$. As for the weight, one merely needs to calculate

$$\operatorname{diag}(x_1, \ldots, x_n)w = (x_1 + \cdots + x_i)w,$$

which is precisely the action by ω_i! ■

Remark 1.126. The moral is the story is that the category $\operatorname{Rep}\mathfrak{sl}_n(\mathbb{C})$ is $\otimes$-generated by the representations $\wedge^i V_{\text{std}}$. In fact, because $\wedge^i V_{\text{std}}$ is a quotient of $V_{\text{std}}^{\otimes i}$, it follows that the category is $\otimes$-generated by V_{std}.

Remark 1.127. There is a perfect pairing

$$\wedge^i V_{\text{std}} \otimes \wedge^{n-i} V_{\text{std}} \to \wedge^n V_{\text{std}} = \mathbb{C}$$

given simply by $w \otimes w' \mapsto w \wedge w'$; indeed, we can see this is a perfect pairing where a basis vector $v_{n_1} \wedge \cdots \wedge v_{n_i}$ has dual basis vector $v_{m_1} \wedge \cdots \wedge v_{m_{n-i}}$, where $\{n_1, \ldots, n_i\} \cup \{m_1, \ldots, m_{n-i}\} = \{1, \ldots, n\}$. Thus, we see that $L_{\omega_i}^* = L_{\omega_{n-i}}$ for each i.

Here is another interesting family of representations.

Example 1.128. Let V_{std} be the standard representation of $\mathfrak{sl}_n(\mathbb{C})$. There is a choice of ordered simple roots of $\mathfrak{sl}_n(\mathbb{C})$ for which

$$L_{m\omega_1} \cong \operatorname{Sym}^m V_{\text{std}}.$$

Proof. We continue from Example 1.125.

1. We claim that the representation $\operatorname{Sym}^m V_{\text{std}}$ is irreducible. As in Example 1.125, we see that

$$\{v_{i_1} \cdots v_{i_m} : i_1 \leq \cdots \leq i_m\}$$

is an eigenbasis for the action of $\mathfrak{h}$ on V_{std}. Thus, any nonzero subrepresentation W of V_{std} contains such an eigenvector. However, given one such eigenvector $v_{i_1} \cdots v_{i_m}$, one can iteratively apply E_{1i_1}, then E_{1i_2}, and so on, eventually proving that $v_1^m \in W$. Moving this basis vector around, we see that $v_i^m \in W$ for each i.

We now claim that each basis vector $v_{i_1} \cdots v_{i_m}$ lives in W. Indeed, by hitting v_1^m with $E_{11} - E_{22} + \sum_{i=2}^n c_i E_{1i}$, we see that

$$\left(v_1 + \sum_{i=2}^n c_i v_2\right)^m \in W.$$

Each coefficient varies with a different multivariate polynomial in the $c_\bullet$s, so by choosing specializations carefully (and generically), we see that each monomial lives in W. Formally, one can inductively allow more and more of the c_is to be nonzero, one at a time.

2. We complete the proof. In light of the previous step, it is enough to show that v_1^m is the highest weight vector of $\operatorname{Sym}^m V_{\text{std}}$ and has weight $m\omega_1$. Certainly E_{pq} kills v_1^m for each $p < q$ because $q > 1$. As for the weight calculation, we note that

$$\operatorname{diag}(x_1, \ldots, x_n) \cdot v_1^m = m x_1 \cdot v_1^m,$$

as desired. ■

Remark 1.129. Another way to check that $\operatorname{Sym}^m V_{\text{std}}$ is irreducible would be to use the Weyl dimension formula (Theorem 1.121) to compute that $\dim L_{m\omega_1} = \dim \operatorname{Sym}^m V_{\text{std}}$. Then the second step provides a canonical map $L_{m\omega_1} \to V_{\text{std}}$, which must be injective and hence an isomorphism.

1.3 February 9

Today we discuss Schur–Weyl duality.

1.3.1 Representations of $\mathrm{GL}_n(\mathbb{C})$

Here is our result.

Proposition 1.130. Irreducible representations of $\mathrm{GL}_n(\mathbb{C})$ are parameterized by pairs (m, λ), where $m \in \mathbb{Z}$ and $\lambda = (\lambda_1, \ldots, \lambda_n)$ is a decreasing sequence of nonnegative integers with $\lambda_n = 0$, and

$$m = \sum_i \lambda_i + nr$$

for some integer r. The highest weight of the corresponding irreducible representation $L_{m,\lambda}$ is $(\lambda_1 + r, \ldots, \lambda_n + r)$.

Proof. It is not quite the case that $\mathrm{GL}_n(\mathbb{C})$ is $\mathbb{G}_m \times \mathrm{SL}_n(\mathbb{C})$, but the natural covering map

$$\mathbb{G}_m(\mathbb{C}) \times \mathrm{SL}_n(\mathbb{C}) \to \mathrm{GL}_n(\mathbb{C})$$

has finite kernel $\mu_n \subseteq \mathbb{C}^\times$. Thus, the irreducible representations of $\mathrm{GL}_n(\mathbb{C})$ consist of the irreducible representations of $\mu_n \times \mathrm{SL}_n(\mathbb{C})$ for which μ_n acts by the identity. Notably, such an irreducible representation can be described as a tensor product of a representation of $\mathbb{G}_m(\mathbb{C})$ and a representation of $\mathrm{SL}_n(\mathbb{C})$, with some condition on the diagonal.

However, irreducible representations of $\mathbb{C}^\times$ are one-dimensional. Passing to the Lie algebra, we see that we are basically given the data of an operator h for which $e^{2\pi i h} = 1$, which means that the operator admits integer eigenvalues, so we see that the corresponding representations of $\mathbb{C}^\times$ are given by the characters $\chi_m(z) := z^m$.

Thus, irreducible representations of $\mathbb{C}^\times \mathrm{SL}_n(\mathbb{C})$ can be described as tensor products

$$L_{m,\lambda} := \chi_m \otimes L_\lambda,$$

where λ is some weight of $\mathfrak{sl}_n(\mathbb{C})$. Recall that such weights can be described as decreasing sequences of n integers with $\lambda_n = 0$. Notably, μ_n acts on $L_{m,\lambda}$ by sending the scalar z to $z^{-m+\sum_i \lambda_i}$, which needs to be divisible by n for our representation to descend to n. We can compute the highest weight of $L_{m,\lambda}$ as $(\lambda_1 + r, \ldots, \lambda_n + r)$. ■

Remark 1.131. Equivalently, we may say that representations of $\mathrm{GL}_n(\mathbb{C})$ are parameterized by decreasing sequences of integers $\lambda = (\lambda_1, \ldots, \lambda_n)$, which is the highest weight.

Example 1.132. We can compute that there is now a fundamental weight $(1, \ldots, 1)$, which is the determinant representation.

Here is an important class of our representations.

Definition 1.133 (polynomial)**.** An irreducible representation L_λ of $\mathrm{GL}_n(\mathbb{C})$ is *polynomial* if and only if $\lambda_n \geq 0$.

Remark 1.134. We can think about the sequence λ as instead being a partition

$$\mu := (\lambda_1 - \lambda_2, \ldots, \lambda_{n-1} - \lambda_n)$$

of λ_1. We may then write $L_\mu := L_\lambda$.

Remark 1.135. Partitions are labeled by Young diagrams, which we notably have no use for.

Remark 1.136. One can check that L_λ is polynomial if and only if L_λ embeds into $V^{\otimes N}$ for some N, where V is the standard representation. In particular, we see that L_λ is contained in $V^{\otimes|\mu|}$, which we can see by the same argument as in Proposition 1.122. The point is that the determinant representation can still be found in $V^{\otimes n}$, but the inverse of the determinant cannot!

Remark 1.137. One can check that polynomial representations are precisely those which extend continuously to $M_n(\mathbb{C})$; equivalently, we may say that these representations are those for which the matrix coefficients are polynomial (not just rational). Again, the point is that the determinant representation has polynomial matrix coefficients, but the inverse of the determinant does not.

1.3.2 Schur–Weyl Duality

Having said something about the representations of $\mathrm{GL}_n(\mathbb{C})$, we are ready to relate them to the symmetric group. Set $V := \mathbb{C}^n$. Then we note that both $\mathrm{GL}_n(\mathbb{C})$ and S_N act on $V^{\otimes N}$, where notably S_N acts by permuting the coordinates.

Theorem 1.138 (Schur–Weyl duality)**.** Fix positive integers n and N, and set $V := \mathbb{C}^n$. Let A be the image of the natural map $U\mathfrak{gl}(V) \to \mathrm{End}_{\mathbb{C}} V^{\otimes N}$, and let B be the image of the natural map $\mathbb{C}[S_N] \to \mathrm{End}_{\mathbb{C}} V^{\otimes N}$.

(a) The algebras A and B are centralizers of each other.

(b) There is a decomposition
$$V^{\otimes N} = \bigotimes_{\lambda \vdash N} L_\lambda \otimes \pi_\lambda$$
of $V^{\otimes N}$ into irreducible representations of $A \times B$; the indexing consists of partitions of N with at most n parts.

(c) As λ varies over partitions of at most n parts, then the π_λs are pairwise non-isomorphic irreducible representations of S_N.

(d) If $n \geq N$, then the π_λs exhaust all irreducible representations of S_N.

Remark 1.139. Note that $n = N$ implies that all permutations of N have at most n parts, so both (c) and (d) apply.

Remark 1.140. The existence of the decomposition in (b) should not be a surprise: for any group G and completely reducible representation Y of G, the natural map
$$\bigoplus_{W \in \mathrm{IrRep}(G)} W \otimes \mathrm{Hom}_G(W, Y) \to Y$$
is an isomorphism; indeed, to prove this, one simply decomposes Y and checks that the statement is true on irreducibles, where it follows from Schur's lemma. For (b), we see that we can simply let π_λ be the space $\mathrm{Hom}_{\mathrm{GL}_n(\mathbb{C})}(L_\lambda, V^{\otimes N})$.

Definition 1.141 (Schur algebra)**.** The algebra A in Theorem 1.138 is the *Schur algebra*, denoted $\mathcal{S}_{n,N}$. The algebra B is then justly called the *centralizer algebra*, denoted $C_{n,N}$.

We are going to need a few lemmas. Our first goal is to show that A is the centralizer of B.

Lemma 1.142. Let U be a complex vector space. Then $S^N U$ is spanned by the pure tensors

$$\{x \otimes \cdots \otimes x : x \in U\}.$$

Proof. Any $u \in S^N U$ has finitely many terms, so by taking the finite number of vectors contained therein, we may reduce to the finite-dimensional case. Now, the span of our pure tensors forms a nonzero subrepresentation of $S^N U$ (for the group $\mathrm{GL}(U)$), but this representation is already known to be irreducible by Example 1.128, so the span must cover everything. ■

Lemma 1.143. Let R be an algebra over $\mathbb{C}$, and let $S^N R$ be the Nth symmetric power, which we note is also an algebra. Now, define $\Delta\colon R \to S^N R$ by

$$\Delta(x) := (x \otimes 1 \otimes \cdots \otimes 1) + (1 \otimes x \otimes \cdots \otimes 1) + \cdots + (1 \otimes 1 \otimes \cdots \otimes x).$$

Then R is generated as an algebra by elements of the form $\Delta(x)$.

Proof. By Newton's identities, there is a polynomial P_N (with rational coefficients) for which

$$z_1 \cdots z_N = P_N\left(\sum_{i=1}^N z_i^1, \ldots, \sum_{i=1}^N z_i^N\right).$$

(This is an identity that takes place in $\mathbb{Q}[z_1, \ldots, z_N]$.) It follows that

$$x \otimes \cdots \otimes x = P_N\left(\Delta(x), \ldots, \Delta(x^N)\right)$$

for any $x \in R$, so we may conclude by Lemma 1.142. ■

Lemma 1.144. In the context of Theorem 1.138, the algebra A is the centralizer of B.

Proof. Note that certainly A and B commute with each other with no effort because B only permutes the factors of V. On the other hand, note that the centralizer Z_B of B is

$$\begin{aligned} Z_B &= \operatorname{End}\left(V^{\otimes N}\right)^{S_N} \\ &\overset{*}{=} \left(\operatorname{End}_{\mathbb{C}}(V)^{\otimes N}\right)^{S_N} \\ &= S^N \operatorname{End}_{\mathbb{C}}(V). \end{aligned}$$

Here, $\overset{*}{=}$ holds because there is a natural map going upwards, and it can be seen to be an isomorphism on a basis. Thus, by Lemma 1.143, it is enough to show that the elements $\Delta(\varphi)$ are in A for each $\varphi \in \operatorname{End}_{\mathbb{C}}(V)$, which is true by definition of A. ■

The rest of Schur–Weyl duality follows from the following piece of algebra.

Lemma 1.145 (Double centralizer)**.** Fix a finite-dimensional vector space V over any field k, and choose two algebras $A, B \subseteq \operatorname{End}_k V$. Suppose that B is a sum of matrix algebras, and A is the centralizer of B.

(a) The algebra A is isomorphic to a sum of matrix algebras.

(b) The algebra B is the centralizer of A.

(c) There is a decomposition
$$V = \bigoplus_{i \in I} (X_i \otimes Y_i),$$
where the lists $\{X_i\}_{i \in I}$ and $\{Y_i\}_{i \in I}$ are exactly lists of irreducible representations of A and B, respectively.

Proof. Start by letting $\{Y_i\}_{i \in I}$ be the list of irreducible representations of B so that we have a decomposition
$$V = \bigoplus_{i \in I} (X_i \otimes Y_i),$$
where $X_i := \operatorname{Hom}_B(Y_i, V)$. Because B embeds in $\operatorname{End}_k V$, we see that Y_i does admit an embedding into V: note that B is a finite-dimensional semisimple algebra, so one can see from the structure theory that any faithful representation contains every irreducible representation. Now, A is the centralizer of B, which is $\operatorname{End}_B(V)$, but this is
$$\operatorname{End}_B(V) = \bigoplus_{i \in I} \operatorname{End}_k(W_i)$$
because the V_is cannot map to distinct other representations. Then (a) is immediate, and (b) and (c) follow by a computation reversing everything in sight to compute decompositions over A. Namely, one will find that $B = \bigoplus_{i \in I} \operatorname{End}_k Y_i$. ■

Proof of Theorem 1.138. Note that B is a quotient of $\mathbb{C}[S_N]$ and is therefore a sum of matrix algebras by the representation theory of finite groups. We now apply Lemma 1.145, which we may do by Lemma 1.144. The decomposition in (b) follows immediately because the irreducible representations of A are labeled by partitions λ of at most n parts, and we see that (c) also immediately follows.

It remains to show (d). It is enough to show that the projection $\mathbb{C}[S_N] \to B$ is injective so that irreducible representations of B are irreducible representations of S_N. Well, let $\{e_1, \ldots, e_n\}$ be a basis of V, and then we see that $\mathbb{C}[S_N]$ acts faithfully on $e_1 \otimes \cdots \otimes e_N$ because the S_N-action produces linearly independent vectors! Thus, the map $\mathbb{C}[S_N] \to \operatorname{End}_{\mathbb{C}} V^{\otimes N}$ is injective, and we are done. ■

Remark 1.146. Theorem 1.138 provides a parameterization of irreducible representations of S_N by partitions of N for every $n \geq N$. This parameterization does not depend on the choice of n, which one can see via the natural diagonal embedding $\mathrm{GL}_n(\mathbb{C}) \to \mathrm{GL}_{n+1}(\mathbb{C})$. This is not too interesting, so we will not write it out.

1.3.3 Schur Functors

We would now like to describe the representations L_λ of $\mathrm{GL}_n(\mathbb{C})$ more explicitly.

Definition 1.147 (Schur functor)**.** Fix a partition λ of a positive integer N. Then we define the *Schur functor* by
$$S^\lambda := \operatorname{Hom}_{S_N}\left(\pi_\lambda, (-)^{\otimes N}\right).$$

Example 1.148. If $\lambda = (N)$, then $L_\lambda = S^N V$, so π_λ is the trivial representation, so $S^\lambda V = S^N V$.

Example 1.149. If $\lambda = (1, \ldots, 1)$, then $L_\lambda = \wedge^N V$ (which is the determinant representation when $\dim V = N$), so π_λ is the sign representation, so $S^\lambda V = \wedge^N V$.

Remark 1.150. In general, Theorem 1.138 has given us a decomposition

$$V^{\otimes N} = \bigoplus_{\lambda \vdash N} S^\lambda V \otimes \pi_\lambda.$$

Example 1.151. We have a decomposition

$$V^{\otimes 2} = \left(S^2 V \otimes \mathbb{C}_{\text{triv}}\right) \oplus \left(\wedge^2 V \otimes \mathbb{C}_{\text{sgn}}\right).$$

Note that we have silently applied Examples 1.148 and 1.149.

Example 1.152. We have a decomposition

$$V^{\otimes 3} = \left(S^3 V \otimes \mathbb{C}_{\text{triv}}\right) \oplus \left(\wedge^3 V \otimes \mathbb{C}_{\text{sgn}}\right) \oplus \left(S^{(2,1)} V \otimes \pi_{(2,1)}\right).$$

By referencing a character table of S_3, we see that $\pi_{(2,1)}$ must be the standard representation of S_3, which is its action on the trace-zero hyperplane in $\mathbb{C}^3$. For example, if we restrict to the group $\mathrm{GL}_3(\mathbb{C})$, then we have a decomposition

$$V^{\otimes 3} = S^3 V \oplus \wedge^3 V \oplus S^{(2,1)} V^{\oplus 2}.$$

On the other hand, the left-hand side is $V^{\otimes 2} \otimes V = \left(S^2 V \otimes V\right) \oplus \left(\wedge^2 V \otimes V\right)$. The former factor has a copy of $S^3 V$, and the latter factor has a copy of $\wedge^3 V$; none of these inclusions are equalities, so it follows that each has one copy of $S^{(2,1)} V$. This allows us to describe $S^{(2,1)} V \subseteq S^2 V \otimes V$ as the subset whose total symmetrization is zero. (There is a similar description of $S^{(2,1)} V$ as embedded in $\wedge^2 V \otimes V$.)

While it is a little annoying to describe the $S^\lambda V$s explicitly (though Example 1.152 gives a method), there is a formula for their dimension.

Proposition 1.153. Fix $V := \mathbb{C}^n$. Then

$$\dim S^\lambda V = \prod_{1 \le i < j \le n} \frac{(\lambda_i - \lambda_j) + (j - i)}{j - i}.$$

Proof. Recall that $\rho = (n-1, n-2, \ldots, 0)$ because the positive roots are $e_i - e_j$ where $i < j$. Thus, we see that $(\rho, e_i - e_j) = (j - i)$, so $(\lambda + \rho, e_i - e_j) = (\lambda_i - \lambda_j) + (j - i)$, and Theorem 1.121 gives the result. ■

We take a moment to give some simplifications. If λ has only k parts, then we could split the product into

$$\prod_{1 \le i < j \le k} \frac{(\lambda_i - \lambda_j) + (j - i)}{j - i} \cdot \prod_{1 \le i \le k < j \le n} \frac{\lambda_i + (j - i)}{j - i}.$$

The latter product now telescopes after λ_i terms, allowing us to rewrite this as

$$\prod_{1 \le i < j \le k} \frac{(\lambda_i - \lambda_j) + (j - i)}{j - i} \cdot \prod_{i=1}^{k} \frac{(n + 1 - i) \cdots (n + \lambda_i - i)}{(k + 1 - i) \cdots (k + \lambda_i - i}.$$

For example, it follows that the dimension is a polynomial P_λ with rational coefficients, whose roots are integers contained in $[1 - \lambda, \ldots, k - 1]$.

1.4 February 11

Today we discuss the fundamental theorem of invariant theory.

1.4.1 Finding Invariant Polynomials

Here is the basic problem: suppose that we have a finite-dimensional complex vector space V, and we are given various tensors $T_i \in V^{\otimes m_i} \otimes (V^*)^{\otimes n_i}$. Then we would like to find the invariant polynomial functions of $T_1, \ldots, T_k$, where "invariant" means that they are invariant under the ambient action of $\mathrm{GL}(V)$.

Suppose for concreteness that we are hunting for homogeneous polynomial functions $F(T_1, \ldots, T_k)$ which is homogeneous of degree d_i in the variable T_i. Here is one method of construction.

1. Set down some vertices i, which we equip with m_i inward arrows and n_i outward arrows, which we number.

2. We then draw d_i such vertices of color i, and we connect only those vertices which can preserve the numbering. This then gives a graph Γ, which exists if and only if the number of incoming and outgoing arrows is the same, meaning that
$$\sum_i d_i m_i = \sum_i d_i n_i.$$
(Certainly this condition is necessary by a degree argument. The converse is some graph theory.) If Γ does not exist, then notably there is no such polynomial because hitting $F(T_1, \ldots, T_k)$ with the scalar λ will not be invariant.

3. Now, once we have found the graph Γ, we construct the desired polynomial by contracting the tensors along the edges.

Example 1.154. Suppose we are given $T_1 \in V^{\otimes 2} \otimes V^*$ and $T_2 \in V^{*\otimes 2}$. This means that we start with the following.

$$\begin{tikzcd}
? \arrow[dr, "1"] & & ? \arrow[dr, "1"] & & ? & \\
& 1 & & 1 & & 2 \arrow[ul, "1"'] \arrow[dl, "2"] \\
? \arrow[ur, "2"] & & ? \arrow[ur, "2"] & & ? &
\end{tikzcd}$$

One possible way to connect these arrows is as follows.

$$\begin{tikzcd}
1 \arrow[d] & \\
1 \arrow[loop below] & 2 \arrow[ul, bend right] \arrow[ul, bend left]
\end{tikzcd}$$

Accordingly, if we write $T_1 = (u_k^{ij})$ and $T_2 = (v_{pq})$, then our polynomial is
$$u_{i_1}^{i_1 j_1} u_{j_2}^{i_2 j_2} u_{j_1 j_2}.$$

Theorem 1.155. Fix everything as above. Then the constructed polynomials F_Γ span the algebra of invariant polynomials.

Proof. Let A be the full algebra of invariant polynomials, and let A_d be the subset which are homogeneous of degree $d = (d_1, \dots, d_k)$. By taking a dual, a polynomial F can be viewed as an element of

$$\bigotimes_i \left(V^{*\otimes m_i} \otimes V^{\otimes n_i}\right) = V^{\otimes \sum d_i n_i} \otimes V^{*\otimes \sum_i d_i m_i}.$$

We are on the hunt for $\mathrm{GL}(V)$-invariants, so the space becomes

$$\mathrm{Hom}_{\mathrm{GL}(V)}\left(V^{\otimes \sum d_i m_i}, V^{\otimes \sum d_i n_i}\right).$$

Note that this vanishes for weight reasons if $\sum_i d_i m_i \neq \sum_i d_i n_i$. On the other hand, if the two sums equal some N, then Theorem 1.138 tells us that this space is

$$\mathrm{Hom}_{\mathrm{GL}(N)}\left(V^{\otimes N}, V^{\otimes N}\right) = \mathbb{C}[S_N]/I$$

for some ideal I. Accordingly, $A_{d_1,\dots,d_k}$ is spanned by some polynomials which correspond to permutations $\sigma \in S_N$. These permutations σ amount to the data of a graph Γ because we are basically matching where certain edges go. Explicitly, we can number all the ingoing and outgoing edges "at the vertex" and then use the permutation σ to tell us which ingoing edge should match with which outgoing edge. ■

Remark 1.156. Given N, if $\dim V$ is large enough (namely, $\dim V \geq N$), then it turns out that the polynomials F_Γ are equal if and only if the graphs Γ are isomorphic. (Indeed, the ideal I vanishes when $\dim V \geq N$ by Theorem 1.138, so different permutations σ will give linearly independent polynomials. One then needs to pass to graphs.) Here, a graph isomorphism is required to preserve the bi-numbering of the edges (introduced in the proof) and the labels of the vertices.

Example 1.157. Suppose $T_1, \dots, T_k$ are all in $V \otimes V^*$. Then all vertices look like $\to 1 \to$, so all the labels do not matter. By tracking the possible graphs, one can see that the algebra of invariant polynomials is generated by traces T_w of cyclic words w in the $T_\bullet$s. For example, the two-cycle $1 \to 2$ produces a factor of $\mathrm{tr}(T_1T_2)$, and the three-cycle $1 \to 2 \to 3$ produces a factor of $\mathrm{tr}(T_1T_2T_3)$. As a further specialization, note with just $k = 1$, we recover the fact that the algebra of functions on $V \otimes V^* \cong M_{\dim V}(\mathbb{C})$ which are invariant for the $\mathrm{GL}(V)$-action is generated by the trace.

Remark 1.158. Continuing the above example, it further turns out that these generators are "asymptotically algebraically independent," in the sense that any polynomial P which vanishes on all the T_ws for all possible dimensions of V must have $P = 0$. Indeed, P has only finitely many terms in the spanning set of Theorem 1.155, and these terms become linearly independent as $\dim V$ goes to infinity (namely, $\dim V$ larger than the total degree of $P(T_{w_1}, \dots, T_{w_k})$ will suffice). Thus, $P = 0$ is forced!

Corollary 1.159. There is no polynomial identity P that holds for all matrices of all size. In other words, choose a polynomial $P \in \mathbb{C}\langle X_1, \dots, X_r\rangle$ in the free non-commutative polynomial ring. If we have $P(A_1, \dots, A_r) = 0$ for all choices of square matrices $A_1, \dots, A_r$ of given dimension, then $P = 0$.

Proof. If $P(A_1, \dots, A_r) = 0$, then note that

$$\mathrm{tr}\, P(A_1, \dots, A_r) A_{r+1} = 0$$

for any other choice of matrix A_{r+1}. Remark 1.158 now implies that P must vanish! ■

Remark 1.160. If one fixes the size of matrices, then there are in fact many polynomial relations. For example, all 1×1 matrices satisfy the relation $AB - BA = 0$. As another example, for 2×2 matrices, $(AB - BA)^2$ is a scalar matrix, so $(AB - BA)^2 C - C(AB - BA)^2 = 0$. In general, for any n and commutative ring R, then any $X_1, \ldots, X_{2n} \in M_n(R)$ satisfies

$$\sum_{\sigma \in S_{2n}} \operatorname{sgn}(\sigma) X_{\sigma(1)} \cdots X_{\sigma(2n)} = 0.$$

1.4.2 Schur Polynomials

We now introduce some Schur polynomials.

Definition 1.161 (Schur polynomial)**.** Let λ be a partition of n, and let s_λ be the trace of $\operatorname{diag}(x_1, \ldots, x_n) \in \mathrm{GL}_n(\mathbb{C})$ acting on the irreducible representation L_λ. Then s_λ is a *Schur polynomial*.

Remark 1.162. By Theorem 1.116, we find that

$$s_\lambda(x_1, \ldots, x_n) = \frac{\displaystyle\sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) x_{\sigma(1)}^{\lambda_1 + n - 1} \cdots x_{\sigma(n)}^{\lambda_n}}{\displaystyle\prod_{i<j} (x_i - x_j)}.$$

(To make the formula look correct, one should multiply the numerator and denominator of Theorem 1.116 by e^ρ, which makes the denominator into $\prod_{\alpha \in \Phi_+} \left(e^{\alpha/2} - e^{-\alpha/2}\right)$.) Note that the numerator may be written as the Vandermonde determinant $\det \left[x_i^{\lambda_i + n - j}\right]_{ij}$.

Remark 1.163. One can consider the Schur polynomials as polynomials in infinitely many variables by imagining that $s_\lambda(x_1, \ldots, x_n)$ is a polynomial $s_\lambda(x_1, \ldots, x_n, 0, \ldots, 0)$.

Example 1.164. The Schur polynomial s_λ for $\lambda = (N)$ corresponds to the representation $L_\lambda = S^N V$. Thus, by expanding $\operatorname{diag}(x_1, \ldots, x_n)$ on a basis $e_{i_1} \cdots e_{i_N}$ of V, we see that

$$s_\lambda(x_1, \ldots, x_n) = \sum_{i_1 + \cdots + i_N = n} x_1^{i_1} \cdots x_N^{i_N}.$$

This is a complete symmetric function. By rewriting a partition, we can also write this as

$$\sum_{1 \le j_1 \le \cdots \le j_N \le n} x_{j_1} \cdots x_{j_N}.$$

Example 1.165. The Schur polynomial s_λ for $\lambda = (1, \ldots, 1)$ corresponds to the representation $L_\lambda = \wedge^N V$. Computing the trace on a basis $e_{i_1} \wedge \cdots \wedge e_{i_N}$ of V recovers the elementary symmetric polynomial

$$s_\lambda(x_1, \ldots, x_n) = \sum_{1 \le j_1 < j_2 < \cdots < j_N \le n} x_{j_1} \cdots x_{j_N}.$$

We now use these Schur polynomials to compute the characters of S_N.

Proposition 1.166. Fix a representation π_λ of S_N, where λ is a partition of N with at most n parts. Let $\sigma \in S_N$ be a permutation with m_i cycles of length i, and set $\chi_\lambda(\sigma) := \operatorname{tr}(\sigma; \pi_\lambda)$. Then $\chi_\lambda(\sigma)$ is the coefficient of the monomial $x_1^{\lambda_1+n-1} \cdots x_n^{\lambda_n}$ in the polynomial

$$\prod_{i<j}(x_i - x_j) \prod_k \left(x_1^i + \cdots + x_n^i\right)^{m_i}.$$

Proof. The idea is to compute the trace of the operator $g \otimes \sigma := g^{\otimes N} \circ \sigma$ on $V^{\otimes N}$, where $V := \mathbb{C}^n$. We also may as well suppose that $g = \operatorname{diag}(x_1, \ldots, x_n)$. Thus, σ permutes the Vs, and g then acts diagonally on a corresponding basis.

Example 1.167. The permutation $\sigma = (123)(45)$ will move around the Vs as in the following diagram.



We now compute our trace two ways.

- One way to compute the trace of the action of $g \otimes \sigma$ is to decompose σ into cycles via its action on $V^{\otimes N}$. As such, we may assume that σ is a single cycle of length m acting on $V^{\otimes m}$. Then σ permutes the standard basis of pure tensors $e_{i_1} \otimes \cdots \otimes e_{i_m}$ along cycles (while $g^{\otimes m}$ acts diagonally), so the only way for such a basis vector to produce a nontrivial contribution to the trace would be if σ fixes $e_{i_1} \otimes \cdots \otimes e_{i_m}$, which means $i_1 = \cdots = i_m$. It follows that the trace of $g \otimes \sigma$ is $x_1^m + \cdots + x_n^m$.

  Now returning to the case of general σ, we multiply through the cycle contributions to find

$$\operatorname{tr}\left(g \otimes \sigma; V^{\otimes N}\right) = \prod_i \left(x_1^i + \cdots + x_n^i\right)^{m_i}.$$

- Alternatively, we use Schur–Weyl duality. Theorem 1.138 tells us that we have a decomposition

$$V^{\otimes N} = \bigoplus_{\lambda \vdash N} L_\lambda \otimes \pi_\lambda,$$

  where g only acts on L_λ, and σ only acts on π_λ. Thus,

$$\operatorname{tr}\left(g \otimes \sigma; V^{\otimes N}\right) = \sum_{\lambda \vdash N} s_\lambda(x_1, \ldots, x_n)\chi_\lambda(\sigma).$$

Comparing the two formulae gives

$$\prod_i \left(x_1^i + \cdots + x_n^i\right)^{m_i} = \sum_{\lambda \vdash N} s_\lambda(x_1, \ldots, x_n)\chi_\lambda(\sigma).$$

We now multiply through by the Weyl denominator, which gives

$$\prod_{i<j}(x_i - x_j) \prod_i \left(x_1^i + \cdots + x_n^i\right)^{m_i} = \sum_{\lambda \vdash N} \left(\sum_{\tau \in S_n} \operatorname{sgn}(\tau) x_{\tau(1)}^{\lambda_1+n-1} \cdots x_{\tau(n)}^{\lambda_n} \chi_\lambda(\sigma) \right).$$

Taking coefficients completes the proof. In particular, we note that the monomial $x_1^{\lambda_1+n-1} \cdots x_n^{\lambda_n}$ is featured only once on the right-hand side: if one has $x_1^{\lambda_1+n-1} \cdots x_n^{\lambda_n} = x_{\tau(1)}^{\mu_1+n-1} \cdots x_{\tau(n)}^{\mu_n}$, then comparing the highest degrees forces $\lambda_1 + n - 1 = \mu_1 + n - 1$ and $\tau(1) = 1$, so one can remove x_1 from both sides and induct. ■

1.4.3 Howe Duality

Here is another application of Schur–Weyl duality.

Proposition 1.168 (Howe duality)**.** Fix complex vector spaces V and W. Then

$$S^n(V \otimes W) = \bigoplus_{\lambda \vdash n} S^\lambda V \otimes S^\lambda W.$$

Proof. Recall that $(V \otimes W)^{\otimes n} = V^{\otimes n} \otimes W^{\otimes n}$. But then Theorem 1.138 tells us that

$$(V \otimes W)^{\otimes n} = \left(\bigoplus_{\lambda \vdash n} S^\lambda V \otimes \pi_\lambda \right) \otimes \left(\bigoplus_{\mu \vdash n} S^\mu W \otimes \pi_\mu \right).$$

Now, we may take S_n-invariants on both sides to see that

$$S^n(V \otimes W) = \bigoplus_{\lambda, \mu \vdash n} S^\lambda V \otimes S^\mu W \otimes (\pi_\lambda \otimes \pi_\mu)^{S_n}.$$

It remains to get rid of the factor of $(\pi_\lambda \otimes \pi_\mu)^{S_n}$. Well, χ_λ has integral values by Proposition 1.166, so it is self-dual, so $(\pi_\lambda \otimes \pi_\mu)^{S_n} = \mathrm{Hom}_{S_n}(\pi_\lambda, \pi_\mu)$, which we see vanishes if $\lambda \neq \mu$ by Schur's lemma (and is $\mathbb{C}$ otherwise). The result follows. ■

Corollary 1.169. Let A_V and A_W be the subalgebras of $\mathrm{End}_{\mathbb{C}}(S^n(V \otimes W))$ generated by $\mathrm{End}(V)$ and $\mathrm{End}(W)$. Then A_V and A_W are centralizers of each other.

Proof. By Proposition 1.168, we see that $S^n(V \otimes W)$ decomposes into a tensor product of irreducibles $S^\lambda V \otimes S^\lambda W$ for $A_V \times A_W$, so it is enough to check the result for these irreducibles. For example, any endomorphism of $S^\lambda V \otimes S^\lambda W$ which preserves A_W must then be a scalar on $S^\lambda W$ and therefore acts only on $S^\lambda V$, so it comes from A_V. ■

We can give a purely combinatorial application of Howe duality.

Proposition 1.170 (Cauchy identity)**.** One has

$$\sum_\lambda s_\lambda(x_1, \ldots, x_n) s_\lambda(y_1, \ldots, y_m) z^{|\lambda|} = \prod_{i=1}^n \prod_{j=1}^m \frac{1}{1 - z x_i y_j}.$$

We will want the following lemma.

Lemma 1.171 (Molien)**.** Let $A \colon V \to V$ be an operator on a finite-dimensional vector space. Then

$$\sum_{r \geq 0} \mathrm{tr}(S^r A; S^r V) T^r = \frac{1}{\det(1 - zA)}.$$

Proof. It is enough to show this over $\mathbb{C}$ because we can reduce this to an equality of some polynomials, which can be checked on complex points. Then we are trying to check an equality of two continuous functions on the complex space $\mathrm{GL}(V)$, so it is enough to check it on the dense subset of diagonalizable matrices $A = \mathrm{diag}(x_1, \ldots, x_n)$. Then the trace of $S^r A$ is simply

$$\sum_{i_1 + \cdots + i_r = n} x_1^{i_1} \cdots x_n^{i_n},$$

so our left-hand side is

$$\sum_{i_1,\ldots,i_n\geq 0} x_1^{i_1}\cdots x_n^{i_n} z^{i_1+\cdots+i_n}.$$

However, we can rearrange this to

$$\left(\sum_{i_1\geq 0}(x_1z)^{i_1}\right)\cdots\left(\sum_{i_n\geq 0}(x_nz)^{i_n}\right)=\frac{1}{1-x_1z}\cdots\frac{1}{1-x_nz},$$

which is exactly $1/\det(1-zA)$. ■

Proof of Proposition 1.170. We will compute the $\sum_{N\geq 0}\operatorname{tr} S^N(x\otimes y)z^N$, where $x:=\operatorname{diag}(x_1,\ldots,x_n)$ and $y:=\operatorname{diag}(y_1,\ldots,y_m)$ are in $\mathrm{GL}(V)$ and $\mathrm{GL}(W)$, respectively.

- By Lemma 1.171, we see that

$$\sum_{N\geq 0}\operatorname{tr} S^N(x\otimes y)z^N=\frac{1}{\det(1-z(x\otimes y)},$$

  which is

$$\prod_{i=1}^{n}\prod_{j=1}^{m}\frac{1}{1-zx_iy_j}.$$

- On the other hand, by Proposition 1.168, we find that

$$\operatorname{tr} S^N(x\otimes y)=\sum_{\lambda\vdash N}s_\lambda(x)s_\lambda(y)z^{|\lambda|}.$$

Setting these equal completes the proof. ■

1.5 February 17

Today we discuss miniscule weights.

1.5.1 Miniscule Weights

Here is our definition.

Definition 1.172 (miniscule)**.** Fix a semisimple Lie algebra $\mathfrak{g}$. A dominant integral weight ω of $\mathfrak{g}$ is *miniscule* if and only if $(\omega,\beta)\leq 1$ for all coroots β.

Remark 1.173. It is enough to check the condition $(\omega,\beta)\leq 1$ for positive coroots.

Example 1.174. The weight 0 is always miniscule.

Example 1.175. For $\mathfrak{g}=\mathfrak{gl}_n(\mathbb{C})$, all the fundamental weights ω_i are miniscule. Indeed, the fundamental weights are

$$\omega_i=(\underbrace{1,\ldots,1}_{i},0,\ldots,0).$$

But now $(\omega_i,e_j-e_k)\in\{0,+1\}$ for each $j<k$, depending on the relative placement of i between j and k. Indeed, it is zero if j and k are both below or both above i, and it is one if i is between them.

Remark 1.176. Any nonzero miniscule weight ω is fundamental; we may assume that $\mathfrak{g}$ is simple. Let $\{\alpha_1, \ldots, \alpha_r\}$ be a collection of simple roots, and let θ^* be the maximal coroot, which takes the form $\sum_i m_i \alpha_i^\vee$ where each m_i is positive. (One way to find this (co)weight is as the highest weight of the (co)adjoint representation. It then turns out that this θ^* is a coroot—the weights of the adjoint representation are all roots—for which $\theta^* - \alpha_i^\vee$ is a positive coroot always.) Then

$$(\omega, \theta^*) = \sum_i m_i (\omega, \alpha_i^\vee).$$

Thus, there is at most one index i with $(\omega, \alpha_i^\vee) = 1$, which means that $\omega = 0$ or ω equals one of the ω_is (because the fundamental weights form a dual basis).

It is helpful to characterize our miniscule weights.

Lemma 1.177. Fix a semisimple Lie algebra $\mathfrak{g}$, and choose a fundamental weight ω_i. Then ω_i is miniscule if and only if $m_i = 1$, where m_i comes from the expansion $\theta^* = \sum_j m_j \alpha_j^\vee$ of the maximal coroot.

Proof. Certainly if ω_i is miniscule, then $(\omega, \theta^*) \leq 1$ implies that $m_i \leq 1$ and so $m_i = 1$. Conversely, if $m_i = 1$, then for any positive coroot

$$\beta = \sum_j n_j \alpha_j^\vee,$$

we see that we must have $n_j \leq m_j$ for each j, so $(\omega_i, \beta) \leq 1$ follows. ■

Example 1.178. A direct calculation of the maximal coroot shows that the Lie algebras with exceptional types E_2, F_4, and E_8 admit no miniscule weights.

Lemma 1.179. Fix a semisimple Lie algebra $\mathfrak{g}$. Suppose that we have $\omega \in Q$ for which $(\omega, \beta) \leq 1$ for all coroots β. Then $\omega = 0$.

Proof. Because ω is in the root lattice, we may write

$$\omega = \sum_i k_i \alpha_i$$

for some integers $k_\bullet$. For the sake of contradiction, assume that ω is nonzero; we may assume that ω is a minimal counterexample, in the sense that it minimizes $\sum_i |k_i|$. Because the inner product is positive-definite, we know that (ω, ω) is positive, so

$$\sum_i k_i (\omega, \alpha_i) > 0.$$

Thus, one of the summands is positive, so we may find an index j such that k_j and (ω, α_j) have the same sign; by possibly replacing ω with $-\omega$, we may assume that $k_j > 0$ and $(\omega, \alpha_j) > 0$. It follows that $(\omega, \alpha_j^\vee) > 0$, so $(\omega, \alpha_j^\vee) = 1$ by hypothesis.

Thus, $s_i \omega = \omega - \alpha_i$, so $s_i \omega$ is another vector in the root lattice, and we can see that $s_i \omega$ also satisfies the hypothesis. Indeed, the hypothesis $(\omega, \beta) \leq 1$ and being in the root lattice is invariant under the Weyl group because the Weyl group preserves the set of coroots and the inner product. As such, we have arrived at our contradiction because the sum of the coordinates of $s_i \omega$ is strictly smaller than the sum of the coordinates of ω. ■

1.5.2 Miniscule Representations

Miniscule weights are interesting to us because their corresponding representations are easy to understand.

Proposition 1.180. Fix a semisimple Lie algebra $\mathfrak{g}$, and choose a dominant integral weight ω. Then the following are equivalent.

(a) The weight ω is miniscule.

(b) All weights of L_ω are contained in the orbit $W\omega$.

(c) If λ is a dominant integral weight such that $\omega - \lambda \in Q_+$, then $\lambda = \omega$.

Proof. We show our implications in sequence.

- We show (a) implies (c), which we will do by induction on the rank of $\mathfrak{g}$. (There is nothing to do if the rank is zero.) We proceed in steps.

  1. If $\omega = 0$, then $\lambda \in -Q_+$, so $(\lambda, \rho) \leq 0$; but because λ is dominant, if it is nonzero, then $(\lambda, \rho) > 0$. We may now suppose that ω is nonzero.
  2. Then ω is a fundamental weight ω_i (by Remark 1.176). Now, write $\omega - \lambda = \sum_j n_j \alpha_j$, where the n_js are nonnegative. If n_j vanishes for some j with $j \neq i$, then we may pass to a smaller Lie algebra (by deleting the jth vertex of the Dynkin diagram), so we are done by the induction.

     Thus, we may assume that $n_j > 0$ for each $j \neq i$.
  3. Now, for any positive coroot β, we know that
     $$(\omega_i - \lambda, \beta) = (\omega_i, \beta) - (\lambda, \beta) \leq (\omega_i, \beta) \leq 1.$$
     Additionally, if $\alpha_i^\vee$ is not found in the expansion of β, then we must actually have $(\omega_i, \beta) \leq 0$; for example, $(\omega_i - \lambda, \alpha_j^\vee) \leq 0$ for all $j \neq i$.
  4. Now, observe that
     $$(\omega_i - \lambda, \omega_i - \lambda) = \sum_j n_j (\omega_i - \lambda, \alpha_i^\vee).$$
     If this is nonpositive, then $\omega_i - \lambda$ vanishes, and we are done. But each of the n_js are nonnegative for each $j \neq i$, and we just showed $(\omega_j - \lambda, \alpha_j^\vee) \leq 0$ for $j \neq i$, so the only way to avoid nonpositivity is for $n_i > 0$ and $(\omega_i - \lambda_i, \alpha_i^\vee) > 0$.

     We are thus reduced to the case that $n_j > 0$ for all j, including $j = i$. We are now ready to conclude. Recall θ^* is a dominant coweight, so $(\omega_i - \lambda, \theta^*) \geq 1$ because we just showed that $(\omega_i - \lambda, \alpha_i^\vee) > 0$ for all i. (In a few more words, one can expand $\omega_i - \lambda = \sum_j n_j \alpha_j$ and $\theta^* = \sum_j p_j \omega_j^\vee$, where the n_js are positive and the p_js are nonnegative, with at least one nonzero.) On the other hand, $(\omega_i, \theta^*) \leq 1$, so $(\lambda, \theta^*) \leq 0$, so $\lambda = 0$, so $\omega_i \in Q_+$, which contradicts Lemma 1.179.

- We show that (c) implies (b). Let μ be a weight of L_ω. Then we can find $w \in W$ to push μ into the dominant chamber, meaning that $\lambda := w\mu$ is a dominant weight. But the construction of L_ω implies that $\omega - \lambda \in Q_+$, so $\omega = \lambda$ follows by (c).

- We show that (b) implies (a) by contraposition. Indeed, suppose that ω is not miniscule, which implies that there is a positive root α such that $(\omega, \alpha^\vee) > 1$. Unravelling $\alpha^\vee$, we see that $2(\omega, \alpha) > (\alpha, \alpha)$. Thus, $\omega - \alpha$ is a weight of L_ω, witnessed by the weight vector $f_\alpha v_\omega$, which we note is nonzero by the representation theory of $\mathfrak{sl}_{2,\alpha}$ (because $2(\omega, \alpha) > (\alpha, \alpha)$).

  We will be done as soon as we can check that $\omega - \alpha$ is not Weyl-conjugate to ω, but this is not hard: indeed,
  $$(\omega - \alpha, \omega - \alpha) = (\omega, \omega) - 2(\omega, \alpha) + (\alpha, \alpha) < (\omega, \omega).$$

The above implications complete the proof. ■

Corollary 1.181. Fix a miniscule weight ω of a semisimple Lie algebra $\mathfrak{g}$. Then

$$\operatorname{ch} L_\omega = \sum_{\gamma \in W\omega} e^\gamma.$$

Proof. This follows from part (b) of Proposition 1.180. Note that the multiplicities of these weight spaces must be 1 because the multiplicity of the highest weight space needs to be 1, and the Weyl group permutes the weight spaces $L[\gamma]$ as γ varies over $W\omega$ (by the definition of the Weyl group action). ■

Proposition 1.182. Fix a dominant integral weight ω of a semisimple Lie algebra $\mathfrak{g}$. Then ω is miniscule if and only if the restriction of L_ω to any simple $\mathfrak{sl}_{2,\alpha} \subseteq \mathfrak{g}$ is a direct sum of one-dimensional and two-dimensional representations for each α.

Proof. In the forward direction, we may suppose that ω is miniscule. Then any weight vector $v \in L_\omega$ which is a highest weight vector for some $\mathfrak{sl}_{2,\alpha}$ has $h_\alpha v = (\omega, \alpha^\vee)v$. But then $|(\omega, \alpha^\vee)| \leq 1$, so the result follows.

In the backward direction, if ω is not miniscule, then we may find a positive root α such that $(\omega, \alpha^\vee) \geq 2$. Then the highest weight vector $v \in L_\omega$ admits $h_\alpha v = (\omega, \alpha^\vee)v$, so v generates a $\mathfrak{sl}_{2,\alpha}$-representation of dimension $(\omega, \alpha^\vee) + 1$. ■

1.5.3 Applications of Miniscule Representations

We will get significant mileage out of the following application of the Weyl character formula.

Corollary 1.183. Fix a miniscule weight ω of a semisimple Lie algebra $\mathfrak{g}$. Then for all dominant integral weights λ,

$$L_\omega \otimes L_\lambda = \bigoplus_{\gamma \in W\omega} L_{\lambda+\gamma},$$

where $L_{\lambda+\gamma}$ vanishes (by convention) if and only if $\lambda + \gamma$ is not dominant.

Proof. By iteratively removing highest weight vectors, it is enough to check that the two representations have the same character. Well, by the Weyl character formula, we see that

$$\begin{aligned}
\operatorname{ch}(L_\omega \otimes L_\lambda) &= \operatorname{ch}(L_\omega)\operatorname{ch}(L_\lambda) \\
&= \sum_{\gamma \in W\omega} e^\gamma \cdot \frac{\sum_{w \in W} \operatorname{sgn}(w) e^{w(\lambda+\rho)}}{\prod_{\alpha \in \Phi_+}(e^{\alpha/2} - e^{-\alpha/2})} \\
&= \frac{1}{\prod_{\alpha \in \Phi_+}(e^{\alpha/2} - e^{-\alpha/2})} \sum_{\substack{w \in W \\ \gamma \in W\omega}} \operatorname{sgn}(w) e^{w(\lambda+\rho)+\gamma}.
\end{aligned}$$

We now replace γ with $w\gamma$, merely moving around some orbit, which gives

$$\operatorname{ch}(L_\omega \otimes L_\lambda) = \sum_{\gamma \in W\omega} \frac{1}{\prod_{\alpha \in \Phi_+}(e^{\alpha/2} - e^{-\alpha/2})} \sum_{w \in W} \operatorname{sgn}(w) e^{w(\lambda+\rho+\gamma)}.$$

Each internal sum does look like $\operatorname{ch} L_{\lambda+\gamma}$, so we will be done as soon as we check that it vanishes if $\lambda+\gamma$ fails to be dominant. Well, suppose that $\lambda+\gamma$ is not dominant; then we can find $\alpha_i^\vee$ such that $(\lambda+\gamma, \alpha_i^\vee) < 0$. On the other hand, $(\gamma, \alpha_i^\vee) \geq -1$ because $\gamma \in W\omega$, and ω is miniscule, so $(\lambda+\gamma, \alpha_i^\vee) = -1$, so $(\lambda+\gamma+\rho, \alpha_i^\vee) = 0$. Thus, s_i fixes $\lambda+\gamma+\rho$, so the terms $\operatorname{sgn}(w)e^{w(\lambda+\rho+\gamma)}$ and $\operatorname{sgn}(s_i w)e^{w(\lambda+\rho+\gamma)}$ in our sum will cancel each other out. ■

Let's apply this to $\mathfrak{sl}_n(\mathbb{C})$.

Example 1.184. We work with $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$. Let V be the standard representation of $\mathrm{GL}_n(\mathbb{C})$, which is a miniscule representation by Example 1.175. The weights of V are the standard basis of the diagonal Cartan $\mathfrak{h} \subseteq \mathfrak{sl}_n(\mathbb{C})$. Then for any partition λ, we see that

$$V \otimes L_\lambda = \bigoplus_{i=1}^{n} L_{\lambda+e_i},$$

where $L_{\lambda+e_i}$ vanishes by convention if $\lambda + e_i$ fails to be a partition. In other words, $V \otimes L_\lambda$ consists of those L_μs, where the Young diagram for μ is obtained by adding a box (suitably) to the Young diagram for λ. For example,

$$V \otimes L_{(6,4,4,2,1)} = L_{(7,4,4,2,1)} \oplus L_{(6,5,4,2,1)} \oplus L_{(6,4,4,3,1)} \oplus L_{(6,4,4,2,2)} \oplus L_{(6,4,4,2,1,1)}.$$

Example 1.185. Continue in the setting of Example 1.184. Then $\wedge^i V$ is the miniscule representation attached to the fundamental weight ω_i (see Example 1.175). Thus, for any partition λ, we see that $\wedge^i V \otimes L_\lambda$ contains those permutations μ which are obtained by adding one to i different coordinates in the partition λ, in a way that ensures that μ is still a partition. Combinatorially, we are trying to add k different boxes to different rows to the Young diagram of λ in a way that ensures that we still have a Young diagram afterward.

We can now combine this discussion with Schur–Weyl duality.

Corollary 1.186. Fix some positive integer N.

(a) For any partition $\lambda \vdash N$, we have

$$\mathbb{C}[S_{N+1}] \otimes_{\mathbb{C}[S_N]} \pi_\lambda = \bigoplus_{\mu \in \lambda + \square} \pi_\mu,$$

where μ varies over those Young diagrams obtained by suitably adding a box to the Young diagram of λ.

(b) For any partition $\mu \vdash N+1$, we have

$$\pi_\mu|_{S_N} = \bigoplus_{\lambda \in \mu - \square} \pi_\lambda,$$

where λ varies over those Young diagrams obtained by suitably removing a box to the Young diagram of λ.

Proof. We quickly explain that (a) implies (b) by Frobenius reciprocity. (A similar argument also shows that (b) implies (a).) Indeed, by Frobenius reciprocity, some π_λ has multiplicity m in the restriction $\pi_\mu|_{S_N}$ if and only if π_μ has multiplicity m in the induction $\mathbb{C}[S_{N+1}] \otimes_{\mathbb{C}[S_N]} \pi_\lambda$. But (a) tells us exactly that this multiplicity vanishes except when μ is obtained by adding a single box to λ (in which case the multiplicity is one). Unwinding this produces the required decomposition of π_μ.

Thus, we will content ourselves with showing (a). Well, define $V := \mathbb{C}^n$, where n is large (e.g., larger than $N+1$). Frobenius reciprocity implies that

$$\mathrm{Hom}_{S_{N+1}}\left(\mathbb{C}[S_{N+1}] \otimes_{\mathbb{C}[S_N]} \pi_\lambda, V^{\otimes(N+1)}\right) = \mathrm{Hom}_{S_N}\left(\pi_\lambda, V \otimes V^{\otimes N}\right),$$

and this latter representation is simply $V \otimes S^\lambda V$ by definition of the Schur functor S^λ (and the fact that S_N

is acting trivially on V). On the other hand,

$$\mathrm{Hom}_{S_{N+1}}\left(\bigoplus_{\mu\in\lambda+\square}\pi_\mu, V^{\otimes(N+1)}\right) = \bigoplus_{\mu\in\lambda+\square} S^\mu V$$

by definition of the Schur functors. Now, Example 1.184 has shown that our two calculations have produced the same representation of $\mathrm{GL}(V)$, so Theorem 1.138 tells us that the corresponding representations of S_{N+1} must also be the same. ■

1.6 February 18

We continue providing some applications of miniscule weights.

1.6.1 The Conjugate Partition

Partitions come with a duality.

Definition 1.187 (conjugate)**.** Given a partition λ, we define the *conjugate partition* $\lambda^\dagger$ to have Young diagram which is the transpose of λ. Explicitly, the ith component of $\lambda^\dagger$ is the number of components of λ which are greater than or equal to i.

Example 1.188. We have that $(3,3,2,1)^\dagger = (4,3,2)$, which we can see as follows.

$$\begin{array}{cccc} \bullet & \bullet & \bullet & \\ \bullet & \bullet & \bullet & \\ \bullet & \bullet & & \\ \bullet & & & \end{array} \Rightarrow \begin{array}{cccc} \bullet & \bullet & \bullet & \bullet \\ \bullet & \bullet & \bullet & \\ \bullet & \bullet & & \end{array}$$

Proposition 1.189. Fix a partition $\lambda \vdash N$. Then

$$\pi_\lambda \otimes \mathbb{C}_{\mathrm{sgn}} = \pi_{\lambda^\dagger}.$$

Proof. This is fairly tricky due to our construction of π_λ. We induct on N. With $N = 1$, this has no content because all representations are trivial.

For the induction, we may suppose that $\nu \vdash (N+1)$, and remove a box of the Young diagram of ν to obtain some $\lambda \in \nu - \square$. Now, note that

$$(\mathbb{C}[S_{N+1}] \otimes_{\mathbb{C}[S_N]} \pi_\lambda) \otimes \mathbb{C}_{\mathrm{sgn}} = \mathbb{C}[S_{N+1}] \otimes_{\mathbb{C}[S_N]} (\pi_\lambda \otimes \mathbb{C}_{\mathrm{sgn}}),$$

which by the induction is $\mathbb{C}[S_{N+1}] \otimes_{\mathbb{C}[S_N]} \pi_\lambda^\dagger$. But Corollary 1.186 then explains that this representation is

$$\bigoplus_{\eta\in\lambda^\dagger+\square} \pi_\eta.$$

It follows that $\pi_\nu \otimes \mathbb{C}_{\mathrm{sgn}} = \pi_{\overline{\nu}}$ for some $\overline{\nu} \in \lambda^\dagger + \square$.

It remains to identify which partition in $\lambda^\dagger + \square$. For this, we recall from the homework that there is a Jucys–Murphy element $c := \sum_{i<j}(ij) \in \mathbb{C}[S_{N+1}]$ which is central and acts on π_ν by the scalar

$$c(\nu) := \sum_{(i,j)\in\nu} (j-i).$$

Accordingly, it acts on $c(\overline{\nu})$ by $-c(\nu)$ (because $-\otimes\mathbb{C}_{\text{sgn}}$ sends (ij) to $-(ij)$). Similarly, c acts on $\pi_{\nu^\dagger}$ by $c(\nu^\dagger) = -c(\nu)$, which we can see by a direct calculation (namely, flipping the Young diagram and thus flipping (i,j) to (j,i)). Thus, $c(\overline{\nu}) = c(\nu^\dagger)$.

We are thus done because the contents $c(\mu)$ are all different for all $\mu \in \lambda^\dagger + \square$: indeed, each place to add a square to the Young diagram of $\lambda^\dagger$ adds a different amount to the content. To see this, note that the places to add a box go from the top-right to the bottom-left, and any such move downwards or leftwards strictly decreases the content in the box. ■

Corollary 1.190 (Skew Howe duality)**.** Fix finite-dimensional complex representations V and W. Then there is a decomposition into $\mathrm{GL}(V)\times\mathrm{GL}(W)$-representations

$$\lambda^N(V\otimes W) = \bigoplus_{\lambda\vdash N} S^\lambda V\otimes S^{\lambda^\dagger}W.$$

Proof. This is on the homework. The idea is to argue as in Proposition 1.168. ■

1.6.2 More on Miniscule Weights in Classical Types

We continue with a semisimple Lie algebra $\mathfrak{g}$ with root lattice Q and weight lattice P. We also fix a collection (α_i) of simple roots so that we have our Cartan matrix $A = ((\alpha_i,\alpha_j^\vee))_{ij}$. Notably,

$$\alpha_i = \sum_j A_{ij}\omega_j,$$

so $P = AQ$, so $|P/Q| = \det A$.

Proposition 1.191. Fix a semisimple Lie algebra $\mathfrak{g}$. Then every coset in P/Q admits a unique miniscule weight.

Proof. Let's start with existence. Fix a coset $C \subseteq P/Q$, and choose some $\omega \in C\cap P_+$ (which exists by adding enough fundamental weights to any representative of C). We may even minimize ω so that $(\omega,\rho^\vee)$ is minimal, where we recall $\rho^\vee := \frac{1}{2}\sum_{\alpha\in\Phi_+}\alpha^\vee$.

We claim that ω is miniscule. Well, let λ be any dominant weight present in the weight decomposition of L_ω. We want to show that $\omega=\lambda$, which completes the proof by Proposition 1.180. Because λ is a weight in L_ω, it follows that $\lambda-\omega\in Q_+$, so $\lambda\in C\cap P_+$. Thus, $(\lambda,\rho^\vee)\ge(\omega,\rho^\vee)$ by minimality, so

$$(\lambda-\omega,\rho^\vee)\ge 0.$$

On the other hand, $\omega-\lambda\in Q_+$, so we may expand $\omega-\lambda=\sum_i m_i\alpha_i$ for nonnegative m_is, so $(\omega-\lambda,\rho^\vee)\le 0$. Thus, $(\omega-\lambda,\rho^\vee)$ actually vanishes, and all the m_is are required to vanish, so $\omega=\lambda$.

We now turn to uniqueness. Suppose we have two distinct miniscule weights ω_1 and ω_2 in our coset C. Then $\omega_1-\omega_2\in Q$ is nonzero, so Lemma 1.179 grants us a coroot β with $(\omega_1-\omega_2,\beta)\ge 2$. Thus, $(\omega_1,\beta)=1$ and $(\omega_2,\beta)=-1$ because these weights are miniscule, which imply that β is positive and negative, respectively (by dominance). This is a contradiction, so we are done. ■

Remark 1.192. For example, it follows that the number of miniscule weights is $|P/Q| = \det A$.

1.6.3 Miniscule Representations

Let's write down some fundamental weights for our classical types. We will not write out all the calculations.

Example 1.193. We write down the fundamental weights for type C_n.

Proof. Here, $\mathfrak{g} = \mathfrak{sp}_{2n}$. The positive roots are given by

$$\{e_i \pm e_j : i < j\} \sqcup \{2e_i\}_i.$$

As such, our simple positive roots are given by $\alpha_1 = e_1 - e_2, \ldots, \alpha_{n-1} = e_{n-1} - e_n$ and $\alpha_n = 2e_{2n}$. The coroots are the same, except $\alpha_n^\vee = e_n$. We can thus see that the fundamental weights are

$$\omega_i = (\underbrace{1, \ldots, 1}_{i}, 0, \ldots, 0)$$

because $(\omega_i, \alpha_j^\vee) = 1_{i=j}$.

It follows that P/Q has two classes,[1] so there is a unique nonzero miniscule weight. We can see that L_{ω_1} is the standard representation of $\mathfrak{sp}_{2n}$, whose weights are given by $\{\pm e_i\}_i$ (indeed, choose the diagonal torus of $\mathfrak{sp}_{2n}$), which is exactly the orbit of the dominant weight e_1 under the Weyl group $S_n \ltimes \{\pm 1\}^n$. Thus, ω_1 is the unique nonzero miniscule weight by Proposition 1.180. ■

Example 1.194. We write down the fundamental weights and miniscule representations for type B_n.

Proof. Here, $\mathfrak{g} = \mathfrak{so}_{2n+1}$. The positive roots are given by

$$\{e_i \pm e_j : i < j\} \sqcup \{e_i\}_i.$$

As such, our simple positive roots are given by $\alpha_1 = e_1 - e_2, \ldots, \alpha_{n-1} = e_{n-1} - e_n$ and $\alpha_n = e_{2n}$. The coroots are the same, except $\alpha_n^\vee = 2e_n$. We can thus see that the fundamental weights are

$$\omega_i = \begin{cases} (\underbrace{1, \ldots, 1}_{i}, 0, \ldots, 0) & \text{if } i < n, \\ (1/2, \ldots, 1/2) & \text{if } i = n, \end{cases}$$

because $(\omega_i, \alpha_j^\vee) = 1_{i=j}$.

It follows that P/Q has two classes (the present situation is dual to type C_n), so there is a unique nonzero miniscule weight. This time, the miniscule weight is ω_n, which can be checked directly by computing the inner product with all the coroots (which have already been described). Thus, by Proposition 1.180 shows finds that L_{ω_n} has character with weights which are just signs applied to ω_n.

This is the "spin" representation S. We claim that S does not lift to a representation of $\mathrm{SO}_{2n+1}(\mathbb{C})$ but instead lifts to a representation of its universal cover $\mathrm{Spin}_{2n+1}(\mathbb{C})$. Indeed, weights of all representations of $\mathrm{SO}_{2n+1}(\mathbb{C})$ should be integral: using the bilinear form $\sum z_i z_{n+i} + z_{n+1}^2$, we see that any

$$g \coloneqq \operatorname{diag}\left(a_1, \ldots, a_n, a_1^{-1}, \ldots, a_n^{-1}, 1\right)$$

acts on a weight space $L[\lambda]$ by $a_1^{\lambda_1} \cdots a_n^{\lambda_n}$. Thus, if L comes from a representation of $\mathrm{SO}_{2n+1}(\mathbb{C})$, then the $\lambda_\bullet$s are required to be integers for the representation to be holomorphic. ■

[1] The root lattice consists of the tuples in $\mathbb{Z}^n$ with even sum, but the weight lattice has everything in $\mathbb{Z}^n$.

Example 1.195. We write down the fundamental weights for type D_n.

Proof. Here, $\mathfrak{g} = \mathfrak{so}_{2n}$. The positive roots are given by

$$\{e_i \pm e_j : i < j\}.$$

As such, our simple positive roots are given by $\alpha_1 = e_1 - e_2, \ldots, \alpha_{n-1} = e_{n-1} - e_n$ and $\alpha_n = e_{n-1} + e_n$. The coroots are the same. We can thus see that the fundamental weights are

$$\omega_i = \begin{cases} (\underbrace{1, \ldots, 1}_{i}, 0, \ldots, 0) & \text{if } i \leq n-2, \\ (1/2, \ldots, 1/2, -1/2) & \text{if } i = n, \\ (1/2, \ldots, 1/2, 1/2) & \text{if } i = n, \end{cases}$$

because $(\omega_i, \alpha_j^\vee) = 1_{i=j}$.

It turns out that P/Q has up to four elements,[2] which can be computed from $\det A$. Thus, there are three nonzero miniscule weights, which are L_{ω_1} (which is the standard representation) and the two "spin representations" $S_+ := L_{\omega_{n-1}}$ and $S_- := L_{\omega_n}$. The fact that these are miniscule can be computed in the usual way by taking inner products with coroots.

Now, by taking a Weyl action, one can compute that the weights of $L_{\omega_{n-1}}$ (respectively, L_{ω_n}) are simple tuples of $\pm 1/2$ with an odd number (respectively, even number) of negative signs. Indeed, the Weyl group consists is the semidirect product of S_n acting on the kernel of $\det\colon \{\pm 1\}^n \to \{\pm 1\}$. As before, S_+ and S_- do not lift to $\mathrm{SO}_{2n}(\mathbb{C})$. ■

While we're here, let's say something about our exceptional types.

Example 1.196. For G_2, F_4, and E_8, one directly calculates that $\det A = 1$, so $P = Q$, so there are no nonzero miniscule weights.

Example 1.197. For E_6, one finds that $\det A = 3$, so there are two nonzero miniscule weights. On the Dynkin diagram, they are the weights ω_1 and ω_6 as follows.

$$\begin{array}{ccccccccc} \omega_1 & - & \bullet & - & \bullet & - & \bullet & - & \omega_6 \\ & & & & | & & & & \\ & & & & \bullet & & & & \end{array}$$

It turns out that L_{ω_1} has dimension 27, which is related to the lines on a cubic surface; it further turns out that L_{ω_6} is the dual of L_{ω_1}. There are many ways to construct L_{ω_1}.

Example 1.198. For E_7, one finds that $\det A = 2$, so there is only one nonzero miniscule weight. It turns out to be the weight ω_1 as follows.

$$\begin{array}{ccccccccccc} \omega_1 & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\ & & & & & & | & & & & \\ & & & & & & \bullet & & & & \end{array}$$

One has $\dim L_{\omega_1} = 56$.

1.6.4 Fundamental Representations

Let's work out our fundamental representations.

[2] It may actually be easier to calculate some candidate miniscule weights first, which we do in the following sentence, and then one can check that they represent all cosets.

Example 1.199. We work out the fundamental representations for type C_n.

Proof. Here, $\mathfrak{g} = \mathfrak{sp}_{2n}$. We already know that $L_{\omega_1} = V$, where V is the standard representation. We also know that $\wedge^2 V$ contains a copy of L_{ω_2}, but $\wedge^2 V$ is not irreducible: note that the bilinear form is in $\wedge^2 V^*$, so there is a copy of B^{-1} in $\wedge^2 V$. It turns out that the embedding

$$L_{\omega_2} \to \wedge^2 V \twoheadrightarrow \wedge_0^2 V,$$

for example by comparing with the dimension formula.

The general discussion is on the homework. Namely, for any $i \geq 2$, there is a contraction $\iota_B \colon \wedge^i V \to \wedge^{i-2} V$ given by contracting along B. Its kernel is a subrepresentation $\wedge_0^i V \subseteq \wedge^i B$, and finds that $\wedge_0^i B \cong L_{\omega_i}$. In fact, there is more structure: there is another operator $m_B \colon \wedge^i V \to \wedge^{i+2} V$ given by taking a wedge product with B^*. Suitably normalized, $f := \iota_B$ and $e := m_B$ produce an $\mathfrak{sl}_2$-triple on the exterior algebra $\wedge^\bullet V$, so $\wedge^\bullet V$ receives an action by $\mathfrak{sl}_2 \times \mathfrak{sp}_{2n}$. Thus, we may decompose

$$\wedge^\bullet V = \bigoplus_{i=1}^{n} \wedge_0^i V \otimes L_i,$$

where L_i is an irreducible representation of $\mathfrak{sl}_2$ with highest weight $m - i$. (Accordingly, we have an instance of Lemma 1.145.) ■

Remark 1.200. For abelian varieties A of dimension n, we see that $\mathfrak{sp}_{2n}$ admits a standard action on $V = \mathrm{H}^1(A; \mathbb{C})$. The decomposition $\wedge^\bullet V$ turns out to agree with the Hodge decomposition.

Remark 1.201. It follows from this discussion and Proposition 1.122 that all finite-dimensional representations of $\mathfrak{sp}_{2n}$ are found in tensor powers of the standard representation.

Example 1.202. We work out the fundamental representations for type B_n.

Proof. Let V be the standard representation of $\mathfrak{so}_{2n+1}$. On the homework, we will show that the representations $\wedge^i V$ are all irreducible for $i \leq n-1$, so $L_{\omega_i} = \wedge^i V$. ■

Example 1.203. We work out the fundamental representations for type D_n.

Proof. Let V be the standard representation of $\mathfrak{so}_{2n+1}$. Again, on the homework, one shows that $\wedge^i V$ all irreducible for $i \leq n-2$, so $L_{\omega_i} = \wedge^i V$. ■

Remark 1.204. In type D_n, it also turns out that $\wedge^{n-1} V$ is irreducible, but it is not fundamental because $(1, \ldots, 1, 0)$ is not a fundamental weight: it equals $\omega_{n-1} + \omega_n$. It follows that

$$\wedge^{n-1} V \subseteq S_+ \otimes S_-.$$

(Indeed, recall the construction of Proposition 1.122.)

Remark 1.205. Note that the spin representations are not found in tensor powers of the standard representation!

1.7 February 23

Here we go.

1.7.1 Spin Groups

Here is the easiest example of the spin representation.

Example 1.206. By the "belt trick," one has that $\pi_1(\mathrm{SO}_3(\mathbb{C})) \cong \mathbb{Z}/2\mathbb{Z}$, and it turns out that its universal cover is $\mathrm{SL}_2(\mathbb{C})$. (Indeed, $\mathfrak{sl}_2 = \mathfrak{so}_3$.) In particular, the adjoint representation defines a covering $\mathrm{SL}_2(\mathbb{C}) \to \mathrm{SO}_3(\mathbb{C})$, and it turns out that the kernel is $\{\pm 1\}$. (We will also show that $\mathrm{SL}_2(\mathbb{C})$ is simply connected in Example 1.209.) Thus, $\mathrm{Spin}_3(\mathbb{C}) = \mathrm{SL}_2(\mathbb{C})$, and the spin representation is just the standard representation.

It remains to handle $n \geq 3$.

Proposition 1.207. For $n \geq 3$, we have $\pi_1(\mathrm{SO}_n(\mathbb{C})) \cong \mathbb{Z}/2\mathbb{Z}$.

We will want the following lemma.

Lemma 1.208. Let $X_n \subseteq \mathbb{C}^n$ be the hypersurface cut out by the equation

$$z_1^2 + \cdots + z_n^2 = 1.$$

For any $k \in \{1, \ldots, n-2\}$, one has $\pi_k(X_n) = 0$.

Proof. One can retract $X_n(\mathbb{C})$ to $X_n(\mathbb{R})$, which is S^{n-1}, and then the result follows from calculations of lower-dimensional homotopy groups. To do the contraction, define $f_\bullet \colon X_n \times [0,1] \to X_n$ by

$$f_t(x+iy) = \frac{x+tiy}{\sqrt{x^2 - t^2y^2}}.$$

Indeed, note that $x + iy \in X_n$ (where $x, y \in \mathbb{R}^n$) is equivalent to having $|x|^2 - |y|^2 = 1$ and $xy = 0$, and we can see that the output continues to have this property. But now f_1 is the identity and f_0 has real output, so we are done. ■

Example 1.209. Adjusting basis, one sees that

$$\mathrm{SL}_2(\mathbb{C}) = \{(a,b,c,d) : ad - bc = 1\}$$

is simply connected (by taking $n = 4$). In fact, we also see that $\pi_2(\mathrm{SL}_2(\mathbb{C})) = 0$.

Example 1.210. At $n = 3$, there is a short exact sequence

$$1 \to \mu_2 \to \mathrm{SL}_2(\mathbb{C}) \to \mathrm{SO}_3(\mathbb{C}) \to 1.$$

This is a homotopy fiber sequence because the latter action is free, so taking the long exact sequence in homotopy and using Example 1.209 shows that $\pi_1(\mathrm{SO}_3(\mathbb{C})) = \mathbb{Z}/2\mathbb{Z}$.

Proof of Proposition 1.207. We proceed by induction; Example 1.210 allows us to assume $n \geq 4$.

We now show the general case. Note that $\mathrm{SO}_n(\mathbb{C})$ acts transitively on X_n, and the stabilizer of the vector $(1, 0, \ldots, 0)$ is exactly $\mathrm{SO}_{n-1}(\mathbb{C})$ embedded diagonally. Thus, we have a homotopy fiber sequence

$$\mathrm{SO}_{n-1}(\mathbb{C}) \to \mathrm{SO}_n(\mathbb{C}) \to X_n.$$

Taking the long exact sequence in homotopy yields the exact sequence

$$\pi_2(X_n) \to \pi_1(\mathrm{SO}_{n-1}(\mathbb{C})) \to \pi_1(\mathrm{SO}_n(\mathbb{C})) \to \pi_1(X_n).$$

The end terms vanish for $n \geq 4$ by Lemma 1.208, so we are done by induction. ■

We may thus make the following definition.

Definition 1.211 (spin)**.** For a positive integer $n \geq 3$, let $\mathrm{Spin}_n(\mathbb{C})$ be the universal cover of $\mathrm{SO}_n(\mathbb{C})$ for each n.

Remark 1.212. By Proposition 1.207, we see that the canonical projection $\mathrm{Spin}_n(\mathbb{C}) \to \mathrm{SO}_n(\mathbb{C})$ is a double cover.

Example 1.213. The short exact sequence in Example 1.210 shows that $\mathrm{Spin}_3(\mathbb{C}) = \mathrm{SL}_2(\mathbb{C})$.

Example 1.214. Note that $\mathfrak{so}_4 = \mathfrak{sl}_2 \oplus \mathfrak{sl}_2$ (e.g., by looking at Dynkin diagrams), so $\mathrm{Spin}_4(\mathbb{C}) \cong \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C})$ because there is a unique simply connected Lie group attached to a given finite-dimensional Lie algebra. One can calculate that the spin representations are given by $S_+ = \mathbb{C}^2 \boxtimes \mathbb{C}$ and $S_- = \mathbb{C} \boxtimes \mathbb{C}^2$. By tracking through the isomorphism $\mathfrak{so}_4 = \mathfrak{sl}_2 \oplus \mathfrak{sl}_2$, one finds that the projection $\mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C}) \to \mathrm{SO}_4(\mathbb{C})$ has kernel $(-1, -1)$. Additionally, the standard representation of $\mathrm{SO}_4(\mathbb{C})$ is $\mathbb{C}^2 \boxtimes \mathbb{C}^2$.

Example 1.215. One has that $\mathfrak{so}_5 \cong \mathfrak{sp}_4$, which one can see by looking at Dynkin diagrams. One can show as before that Sp_{2n} is simply connected, so $\mathrm{Spin}_5(\mathbb{C}) = \mathrm{Sp}_4(\mathbb{C})$, and the double cover $\mathrm{Sp}_4(\mathbb{C}) \to \mathrm{SO}_5(\mathbb{C})$ has kernel $\{\pm 1\}$. (This is the kernel because the kernel needs to be central, and the nontrivial element needs to act by -1 on the spin representations in order to avoid descending to SO.) The spin representation of $\mathrm{SO}_5(\mathbb{C})$ turns out to be the standard representation of $\mathrm{Sp}_4(\mathbb{C})$.

Example 1.216. One has $\mathfrak{so}_6 \cong \mathfrak{sl}_4$, so $\mathrm{Spin}_6(\mathbb{C}) = \mathrm{SL}_4(\mathbb{C})$ because $\mathrm{SL}_4(\mathbb{C})$ is simply connected. The double cover $\mathrm{SL}_4(\mathbb{C}) \to \mathrm{SO}_6(\mathbb{C})$ has kernel $\{\pm 1\}$. The spin representations S_+ and S_- are the standard representation and its dual. (Note that these are the only possible irreducible representations of $\mathrm{SL}_4(\mathbb{C})$ with the correct dimension.) On the other hand, the standard representation of $\mathrm{SO}_6(\mathbb{C})$ is the second exterior power of the standard representation.

For $n \geq 7$, the group $\mathrm{Spin}_n(\mathbb{C})$ is harder to describe, and it is our next goal.

1.7.2 Clifford Algebras

Throughout, V is an inner product space over a field k which has characteristic not equal to 2.

Definition 1.217. Fix an inner product space V over a field k with $\operatorname{char} k \neq 2$. Then the *Clifford algebra* $\operatorname{Cl} V$ is the quotient of the universal tensor algebra by the ideal generated by the elements

$$v \otimes v - \frac{1}{2}(v, v).$$

Remark 1.218. For any $a, b \in V$, it follows that $2(ab+ba) = (a+b)^2 - a^2 - b^2 = (a+b, a+b) - (a,a) - (b,b) = 2(a,b)$, so $ab + ba = (a,b)$. Setting $a = b$ recovers the relation $2a^2 = (a,a)$.

Remark 1.219. The Clifford algebra admits a natural filtration $\{F_i\}$ induced by the standard filtration on the tensor algebra. In particular, $F_0 = k$, $F_1 = k + V$, $F_2 = (k+V)\cdot(k+V)$, and so on.

Remark 1.220. Because the relations we took a quotient by for $\operatorname{Cl} V$ live in even degree, there is a $\mathbb{Z}/2\mathbb{Z}$-grading on $\operatorname{Cl} V$, which we label $\operatorname{Cl}_+ V \oplus \operatorname{Cl}_- V$.

Remark 1.221. Recall that one can define the exterior algebra $\wedge V$ as the quotient of the tensor algebra TV by the ideal generated by the elements v^2. Thus, the natural map $TV \to \operatorname{Cl} V$ descends to a map $\wedge V \to \operatorname{gr} \operatorname{Cl} V$. This map is further a surjection: fixing an orthonormal basis $\{e_i\}$ of V, we see that $\operatorname{Cl} V$ is spanned by the vectors of the form

$$e_{i_1} \cdots e_{i_r}.$$

Using the relations $e_i e_j + e_j e_i = 1_{i=j}$, we see that we may even assume $i_1 < \cdots < i_r$. But then these vectors are found from $\wedge V$, so we are done.

Here is a basis, which is a variant of the Poincaré–Birkhoff–Witt theorem.

Theorem 1.222. Fix an inner product space V.

(a) If $\dim V = 2n$, then $\operatorname{Cl} V \cong M_{2^n}(\mathbb{C})$.

(b) If $\dim V = 2n+1$, then $\operatorname{Cl} V \cong M_{2^n}(\mathbb{C}) \oplus M_{2^n}(\mathbb{C})$.

Proof. Let's start with the case $\dim V = 2n$, and choose a basis $\{a_1, \ldots, a_n, b_1, \ldots, b_n\}$ so that $(a_i, a_j) = 0$, $(b_i, b_j) = 0$ and $(a_i, b_j) = 1_{i=j}$. We now define operators on $M := \wedge(a_1, \ldots, a_n)$ as follows: for each i, we define $A_i v := a_i v$ and $B_i v := \frac{\partial}{\partial a_i} v$, where

$$\frac{\partial}{\partial a_i}(a_{i_1} \wedge \cdots \wedge a_{i_n}) := \begin{cases} 0 & \text{if } i \notin \{i_1, \ldots, i_n\}, \\ (-1)^{k-1}(a_{i_1} \wedge \cdots \wedge \widehat{a}_{i_k} \wedge \cdots \wedge a_{i_n}) & \text{if } i = i_k. \end{cases}$$

One can check that $A_i B_j + B_j A_i = 1_{i-j}$ and $A_i A_j + A_j A_i = 0$ and $B_j B_i + B_i B_j = 0$, so we get a natural map $\rho\colon \operatorname{Cl} V \to \operatorname{End} M$ given by $\rho(a_i) := A_i$ and $\rho(b_i) := B_i$.

Now, on the homework, we will show that increasing sequences $\{i_\bullet\}$ and $\{j_\bullet\}$ produce linearly independent elements

$$A_{i_1} \cdots A_{i_r} B_{j_1} \cdots B_{j_s}.$$

It follows that ρ is an embedding because the elements $a_{i_1} \cdots a_{i_r} b_{j_1} \cdots b_{j_s}$ are already known to be a spanning set, and we can see that they are sent to a linearly independent set. But then note that $\dim \operatorname{Cl} V = 2^{2n}$ while $\dim M = 2^{2n}$, so it actually follows that ρ is an isomorphism.

We now turn to the case where $\dim V = 2n+1$. Then we can pick a basis $\{a_1, \ldots, a_n, b_1, \ldots, b_n, z\}$, where the $a_\bullet$s and $b_\bullet$s have relations as above, and $(a_i, z) = (b_i, z) = 0$ for all i and $(z, z) = 2$. In particular, we see that $z a_i = -a_i z$ and $z b_i = -b_i z$ and $z^2 = 1$. Now, set $M := \wedge(a_1, \ldots, a_n)$. There are two ways to extend the action of $\operatorname{Cl} V_0$ (where $V_0 := \operatorname{span}\{a_1, \ldots, a_n, b_1, \ldots, b_n\}$) to $\operatorname{Cl} V$. Indeed, either one can take

$$z(a_{i_1} \cdots a_{i_r}) = \pm(-1)^r (a_1 \wedge \cdots \wedge a_{i_r}).$$

Notably, the choice of sign $\pm$ produces non-isomorphic modules $M_\pm$ for $\operatorname{Cl} V$ because there is a unique vector v with $b_i v = 0$ for all i, but then $zv = \pm v$ in $M_\pm$.

Thus, one receives a natural map

$$\rho\colon \operatorname{Cl} V \to \operatorname{End} M_+ \oplus \operatorname{End} M_-,$$

and one checks as before that it is an isomorphism. For example, one can check that M_+ and M_- are irreducible, which makes ρ a surjection, and then it becomes an isomorphism via some dimension calculations. ■

Remark 1.223. The above proof has actually shown that the natural map $\wedge V \to \operatorname{gr} \operatorname{Cl} V$ is an isomorphism.

Remark 1.224. Here is another approach: one can try to check that M is an irreducible module for $\operatorname{Cl} V$ instead of trying to show that the elements $A_{i_1} \cdots A_{i_r} B_{j_1} \cdots B_{j_s}$ are linearly independent. In particular, irreducibility implies that ρ is surjective by some density theorem.

1.7.3 Spin Representations

We are now ready to construct some spin representations.

Proposition 1.225. Fix an inner product space V over a field of characteristic not 2. View $\mathfrak{so}(V)$ as $\wedge^2 V$ via the identification $V \cong V^*$. Then the natural map $\xi\colon \wedge^2 V \to \operatorname{Cl} V$ defined by

$$\xi(a \wedge b) = \frac{1}{2}(ab - ba)$$

is a Lie algebra homomorphism. Here, $\operatorname{Cl} V$ has been given the natural Lie bracket from being an associative algebra.

Proof. This is a direct calculation. Note

$$\begin{aligned}
[\xi(a \wedge b), \xi(c \wedge d)] &= \left[ab - \frac{1}{2}(a,b), cd - \frac{1}{2}(c,d)\right] \\
&\overset{*}{=} [ab, cd] \\
&= abcd - cdab,
\end{aligned}$$

where $\overset{*}{=}$ has used the Clifford algebra relations. Now, one can successively switch vectors around with some error terms. For example, $abcd = abcd + acbd - acbd = (b,c)ad - acbd$. Continuing this process until we get to the permutation $cdab$, we see that this equals

$$(b,c)ad - (b,d)ac + (a,c)db - (a,d)cb,$$

which can be seen to be

$$(b,c)\xi(a \wedge b) - (b,d)\xi(a \wedge c) + (a,c)\xi(d \wedge b) - (a,d)\xi(c \wedge b),$$

which is $\xi([a \wedge b, c \wedge d])$ in $\mathfrak{so}(V)$. ■

Notably, one now sees that the map $U\mathfrak{so}(V) \to \operatorname{Cl}_+ V$ is surjective, so we can use the modules M constructed in Theorem 1.222 to find our spin representations.

- When $\dim V = 2n$, it turns out that the $\mathbb{Z}/2\mathbb{Z}$-grading on $\operatorname{Cl} V$ produces a $\mathbb{Z}/2\mathbb{Z}$-grading $M = M_0 \oplus M_1$ on M. Thus, we may view $M_0 \oplus M_1$ as a representation of $\mathfrak{so}_{2n}$, and these representations have dimension 2^{n-1}. By finding a suitable highest weight vector and comparing dimensions, one finds that these representations are S_+ and S_-, respectively.
- When $\dim V = 2n+1$, it turns out that both M_+ and M_- are both the same spin representation S, which one again checks by find a suitable highest weight vector and comparing dimensions. Notably, it turns out that M_+ and M_- remain irreducible when restricted $\mathfrak{so}(V)$.

1.7.4 Dual Representations

We return to the setting where $\mathfrak{g}$ is a finite-dimensional simple Lie algebra over $\mathbb{C}$.

Remark 1.226. Fix an irreducible representation L_λ of $\mathfrak{g}$, where (as usual) λ is some dominant integral weight. Then $L_\lambda^* \cong L_{\overline{\lambda}}$ for some other dominant weight $\overline{\lambda}$. Notably, the highest weight of L_λ^* is just the negative of the lowest weight of L_λ. To compute this, recall that there is a maximal element $w_0 \in W$ of length $|\Phi_+|$, meaning that $w_0(\Phi_+) = \Phi_-$. Thus, w_0 maps the dominant chamber P_+ to $-P_+$. It follows that $w_0\lambda$ is exactly the lowest weight of L_λ, so $\overline{\lambda} = -w_0\lambda$. (Indeed, writing $w_0\mu = \lambda - \sum_i k_i\alpha_i$ for nonnegative k_i, then $\mu = w_0\lambda - \sum_i k_i w_0\alpha_i = w_0\lambda + \sum_i k_i\alpha_{\sigma(i)}$, where σ is some permutation of the simple roots induced by w_0. The minimality of $w_0\lambda$ follows.)

It is potentially interesting to calculate w_0 in some special cases. Note that w_0 maps the positive roots to the negative roots, so $-w_0$ permutes the simple roots, so it is a permutation of the vertices of the Dynkin diagram. Further, because Weyl group elements preserve the inner product, we see that $-w_0$ will further preserve the edges of the Dynkin diagram. Thus, $-w_0$ is a graph automorphism!

Example 1.227. For the types A_1, B_n, C_n, G_2, F_4, E_7, and E_8, the Dynkin diagram admits no automorphism, so $-w_0$ is the identity.

The remaining cases are slightly trickier. Let's remove a few more. Note that W acts trivially on the quotient P/Q: indeed, we see that

$$s_i\lambda = \lambda - (\lambda, \alpha_i^\vee)\alpha_i,$$

and the latter term is in Q. Thus, $-w_0$ acts by an inversion on P/Q, so we conclude that $-w_0 \neq 1$ whenever P/Q is not a product of $(\mathbb{Z}/2\mathbb{Z})$s.

Example 1.228. In type A_{n-1} for $n \geq 3$, we see that $P/Q \cong \mathbb{Z}/n\mathbb{Z}$, so $-w_0$ must act by flipping the entire Dynkin diagram. One can also see explicitly that this longest Weyl element is the full flip in S_n.

The remaining cases in D_n are harder. If n is odd, then $P/Q \cong (\mathbb{Z}/4\mathbb{Z})$, and so $-w_0$ acts by -1 already, so it is nontrivial. If n is even, then $P/Q \cong (\mathbb{Z}/2\mathbb{Z})^2$, but one can calculate that $-w_0$ acts by 1. This last calculation must be done explicitly. The Weyl group in this case is $S_n \ltimes \{\pm 1\}_0^n$, where $\{\pm 1\}_0^n$ is the kernel of the multiplication map $\{\pm 1\}^n \to \{\pm 1\}$. Then w_0 is the element sending positive roots to negative roots, which when n is even is simply $-\operatorname{id}$. (When n is odd, this element is not in the Weyl group!) Thus, $-w_0$ is trivial!

Example 1.229. For $\mathfrak{so}(4k+2)$, one can calculate $S_+^* = S_-$. But for $\mathfrak{so}(4k)$, one finds that $S_+^* = S_+$ and $S_-^* = S_-$.

Example 1.230. For type E_6, one finds that $P/Q \cong \mathbb{Z}/3\mathbb{Z}$, so $-w_0$ is nontrivial and flips the whole Dynkin diagram. In particular, it switches the two miniscule weights.

1.7.5 Maximal Root

We have already recalled a little work with the maximal root when we discussed miniscule representations, but we will now study the maximal root in more detail.

Definition 1.231 (maximal root)**.** Fix a simple Lie algebra $\mathfrak{g}$. Then the *maximal root* θ of $\mathfrak{g}$ is the highest weight of the adjoint representation of $\mathfrak{g}$.

Remark 1.232. Because the adjoint representation has weights equal to the root system, we see that θ is in fact a root. Its maximality follows because it is the highest weight among the roots.

Example 1.233. In type A_{n-1}, we can see that $\theta = (1, 0, \ldots, 0, -1)$, which is not fundamental.

Example 1.234. In type C_n, one sees that the adjoint representation $\mathfrak{g}$ is $S^2 V$, which is $L_{2\omega_1}$. Thus, θ is not fundamental.

Proposition 1.235. For all simple Lie algebras $\mathfrak{g}$ which are not in types A or C, the maximal root θ is fundamental.

Proof. In the case $\mathfrak{g} = \mathfrak{so}_N$, we need to have $N \geq 7$ to avoid the exceptional isomorphisms. In this case, one finds that $\mathfrak{g} = \wedge^2 V$, which is L_{ω_2}. We will not say much about the exceptional types, but they can be computed explicitly. ■

Remark 1.236. There is a notion of extended Dynkin diagram which allows one to provide a more algebraic proof of this fact.

1.8 February 25

We continue.

1.8.1 The Principal $\mathfrak{sl}_2$-triple

Let $\mathfrak{g}$ be a simple finite-dimensional Lie algebra.

Definition 1.237. Let $\mathfrak{g}$ be a semisimple finite-dimensional Lie algebra. Then we define $e := \sum_i e_i$ and $h := 2\rho^\vee$. There are coefficients

$$f := \sum_i c_i f_i$$

such that (e, f, h) is an $\mathfrak{sl}_2$-triple. This is the *principal $\mathfrak{sl}_2$-triple*.

Remark 1.238. The coefficients are given by $c_i := (2\rho^\vee, \omega_i)$, which one can check because this is what is needed for $[e, f] = \sum_i c_i h_i$ to equal $h = \sum_i (2\rho^\vee, \omega_i) h_i$.

Example 1.239. For $\mathfrak{gl}_n$, one finds that

$$e = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 0 \end{bmatrix}$$

and $h = \operatorname{diag}(n-1, n-3, \ldots, 3-n, 1-n)$. On the homework, we will calculate f.

It is somewhat important to decompose $\mathfrak{g}$ as a representation of this principal $\mathfrak{sl}_2$.

Definition 1.240 (height)**.** Fix a semisimple finite-dimensional Lie algebra $\mathfrak{g}$. Then the *height* $|\alpha|$ of a root α is the sum of the coefficients when α is expressed a sum of simple roots. We will let r_m be the number of positive roots with height m. By convention, we let r_0 be the rank of $\mathfrak{g}$.

Remark 1.241. View $\mathfrak{g}$ as a representation of the principal $\mathfrak{sl}_2$-triple. One can check that $(2\rho^\vee, \alpha) = |\alpha|$, so $[h, e_\alpha] = 2\,|\alpha|\, e_\alpha$. Thus, the weights of $\mathfrak{g}$ are all even, and

$$r_m = \dim \mathfrak{g}[2m].$$

Thus, the representation theory of $\mathfrak{sl}_2$ assures us that $\{r_m\}$ is a decreasing sequence.

Tracking through the representation of $\mathfrak{sl}_2$, we are thus motivated to study the differences $r_m - r_{m+1}$.

Definition 1.242 (exponent)**.** Fix a semisimple finite-dimensional Lie algebra $\mathfrak{g}$. Then an *exponent* of $\mathfrak{g}$ is a positive integer m for which $r_m > r_{m+1}$. Its *multiplicity* is $r_m - r_{m+1}$. We may enumerate the exponents (with multiplicity) by $m_1, m_2, \ldots$.

Example 1.243. Note that r_1 is the rank of $\mathfrak{g}$ because roots of height one are (by definition) simple.

Example 1.244. Suppose $\mathfrak{g}$ is simple. Note that r_2 is the number of roots of the form $\alpha_i + \alpha_j$, which happens exactly when i and j are connected. (Indeed, if there is no connection, then the roots are perpendicular.) But the Dynkin diagram is a tree, so $r_2 = r_1 - 1$, so 1 is an exponent with multiplicity 1.

Example 1.245. For n large enough, we see that $r_n = 0$, so the number of exponents (enumerated with multiplicity) is exactly r.

Example 1.246. The maximal root θ corresponds to the exponent $m_r = (\theta, \rho^\vee)$. By definition, $(\theta, \rho^\vee) + 1$ is the "Coxeter number."

Example 1.247. For $\mathfrak{g} = \mathfrak{sl}_n$, we see that roots of height ℓ are those of the form $e_i - e_{i+\ell}$. Thus, the exponents are $\{1, 2, \ldots, n-1\}$, and the Coxeter number is n.

Remark 1.248. By the representation theory of $\mathfrak{sl}_2$, we see that

$$\mathfrak{g} = \bigoplus_{i=1}^{r} L_{2m_i}.$$

For example, this implies that $\sum_{i=1}^r (2m_i + 1) = \dim \mathfrak{g}$, so $\sum_{i=1}^r m_i$ is the number of positive roots.

Example 1.249. For example, $\mathfrak{sl}_n = L_2 \oplus L_4 \oplus \cdots \oplus L_{2n-2}$. Another way to see this decomposition is to apply the Clebsch–Gordan rule to compute $L_{n-1} \otimes L_{n-1}^* - \mathbb{C}$, where "$-\mathbb{C}$" refers to the fact that we are removing the trivial representation.

Here are the exponents for the other Lie algebras.

Remark 1.250. It turns out that the exponents depends only on the undirected graph isomorphism of the Dynkin diagram, which one can see by Chevalley's theorem.

Example 1.251. Here are the exponents for the classical types.

- The exponents of $\mathfrak{sp}_{2n}$ are $\{1, 3, \ldots, 2n-1\}$.
- The exponents of $\mathfrak{so}_{2n+1}$ are $\{1, 3, \ldots, 2n-1\}$.
- The exponents of $\mathfrak{so}_{2n+2}$ are $\{1, 3, \ldots, 2n-1\} \sqcup \{n\}$, where the extra n at the end is added with multiplicity when n is odd.

Example 1.252. Here are the exponents for the exceptional types.

- The exponents of F_4 are $\{1, 5, 7, 11\}$.
- The exponents of E_6 are $\{1, 4, 5, 7, , 11\}$.
- The exponents of E_7 are $\{1, 5, 7, 9, 11, 13, 17\}$.
- The exponents of E_8 are $\{1, 7, 11, 13, 17, 19, \}$. These are the

Remark 1.253. Here is a curious property of these exponents: they are symmetric, in the sense that $m_i + m_{r-i}$ is constant. We will prove this later.

We close our discussion by saying a little more about the aforementioned Coxeter number.

Definition 1.254 (Coxeter number)**.** Fix a simple Lie algebra $\mathfrak{g}$. Then the *Coxeter number* $h_{\mathfrak{g}}$ is $(\theta, \rho^\vee) + 1$, which is one more than the height of the maximal root θ. The *dual Coxeter number* $h_{\mathfrak{g}}^\vee$ is $(\theta^\vee, \rho) + 1$.

Remark 1.255. If $(-,-)$ is normalized so that $(\theta, \theta) = 2$, then we see that the dual Coxeter number is also $(\theta, \rho) + 1$. This normalization is natural because the quadratic Casimir element C for this normalization acts on the adjoint representation by

$$(\theta, \theta + 2\rho) = 2h_{\mathfrak{g}}^\vee.$$

Remark 1.256. If $\mathfrak{g}$ is simply laced (i.e., among the types A, D, or E), then $h_{\mathfrak{g}} = h_{\mathfrak{g}}^\vee$ because θ and $\theta^\vee$ are the same in these simply laced cases. However, they are not the same in general. Indeed, $h_{B_n}^\vee = 2n-1$, $h_{C_n}^\vee = n+1$, $h_{G_2}^\vee = 4$, and $h_{F_4}^\vee = 9$. Notably, this dual Coxeter number is also not the dual Coxeter number of the dual!

1.8.2 Real, Complex, and Quaternionic Representations

As an application of our principal $\mathfrak{sl}_2$, we say something about dividing our representations into various types.

Definition 1.257 (real, complex, quaternionic)**.** Fix a finite-dimensional irreducible representation of a group G. Then V has *complex type* if and only if $V \not\cong V^*$. Continuing, given an isomorphism $A\colon V \to V^*$, this induces a dual map $A^*\colon V^* \to V$, so there is a scalar λ such that $A^* = \lambda A$ after these identifications. If $A^* = A$, then V has *real type*, and if $A^* = -A$, then V has *quaternionic type*.

Remark 1.258. If we forget about the complex structure of V, merely viewing it in $\mathrm{Rep}_{\mathbb{R}}(V)$, then it turns out that $\mathrm{End}_{\mathbb{R}[G]}(V)$ is $\mathbb{C}$ for complex type, $M_2(\mathbb{R})$ for the real type, and $\mathbb{H}$ for quaternionic type. We will prove this on the homework.

Remark 1.259. On the homework, we will show that when G is finite, V has real type if and only if V admits a basis in which all elements of G act by matrices with real coefficients.

Example 1.260. Consider $G = \mathrm{SL}_2(\mathbb{C})$, and consider the irreducible representation $L_n = \mathrm{Sym}^n \mathbb{C}^2$. Then L_1 is of quaternionic type, so one can check that L_n has quaternionic type for n odd and real type for n even.

Example 1.261. For $G = \mathrm{SL}_n(\mathbb{C})$ for $n \geq 3$, one has that V and V^* are non-isomorphic, so V has complex type.

It is thus interesting to know which representations of a Lie group admit which types.

Proposition 1.262. Fix a Lie simply connected Lie group G with Lie algebra $\mathfrak{g}$, and choose a dominant weight λ.

(a) If $\lambda \neq -w_0\lambda$, then L_λ is of complex type.

(b) If $\lambda = -w_0\lambda$, then L_λ is of real type if $(\lambda, 2\rho^\vee)$ is even and is of quaternionic type if $(\lambda, 2\rho^\vee)$ is odd.

Proof. Part (a) follows because $L_\lambda^* = L_{-w_0\lambda}$ by Remark 1.226.

For (b), we are granted an isomorphism $A\colon L_\lambda \to L_\lambda^*$. We now restrict L_λ to the principal $\mathfrak{sl}_2$-triple (e, f, h) of $\mathfrak{g}$. Then the largest eigenvalue of h on L_λ comes from its maximal weight, so this eigenvalue is $m := (2\rho^\vee, \lambda)$; furthermore, we see that $L_\lambda[m]$ is one-dimensional. The other weight vectors in L_λ have weights

$$(2\rho^\vee, \lambda - \beta) = m - 2\sum_i (\omega_i, \beta),$$

which we note is strictly less than m. Thus, A must carry L_m to L_m^*, so we are done by Example 1.260! ■

Let's use Proposition 1.262 to determine the types of the spin representations.

Example 1.263. Consider $\mathfrak{g} = \mathfrak{so}_{2n}$ so that $\rho^\vee = \rho = (n-1, n-2, \ldots, 0)$. Then one sees that $S_+ = L_{\omega_n}$ and $S_- = L_{\omega_{n-1}}$, where $\omega_n = (1/2, \ldots, 1/2)$ and $\omega_{n-1} = (1/2, \ldots, 1/2, -1/2)$. We know that S_+ and S_- are self-dual only when n is even (by looking at the action of $-w_0$). For example, we can calculate

$$(\omega_n, 2\rho^\vee) = \frac{n(n-1)}{2},$$

so S_+ is of real type for $n \equiv 0 \pmod 4$ and of quaternionic type for $n \equiv 2 \pmod 4$.

Example 1.264. Consider $\mathfrak{g} = \mathfrak{so}_{2n+1}$ so that $\rho^\vee = (n, n-1, \ldots, 1)$ and $S = L_{\omega_n}$ has $\omega_n = (1/2, \ldots, 1/2)$. Then S is always self-dual because $-\omega_0$ is trivial, and

$$(\omega_n, 2\rho^\vee) = \frac{n(n+1)}{2}.$$

Thus, S is of quaternionic type when $n \equiv 1, 2 \pmod 4$ and is of real type when $n \equiv 0, 3 \pmod 4$.

The above two examples combine into the following.

Theorem 1.265 (Real Bott periodicity)**.** The type of the spin representations of $\mathfrak{so}_n$ is uniquely determined by $n \pmod 8$.

Proof. Combine Examples 1.263 and 1.264. ■

Remark 1.266. This is related to Bott periodicity in homotopy theory, but I do not know how.

Theme 2
Integration

2.1 March 2

We'll need to review a little topology.

2.1.1 Review of Topology

We end our class by saying something about integration.

Definition 2.1 (locally compact)**.** Fix a Hausdorff topological space X. Then X is *locally compact* if and only if any $p \in X$ admits an open neighborhood U for which the closure $\overline{U}$ is compact.

Example 2.2. The base of balls in $\mathbb{R}^n$ shows that $\mathbb{R}^n$ is locally compact. Because manifolds are locally homeomorphic to $\mathbb{R}^n$, we see that manifolds are locally compact.

Lemma 2.3. Fix a second countable locally compact space X. Then X admits a filtration $\{K_i\}$ of compact sets for which $X = \bigcup_i K_i$, and each $x \in X$ admits an open neighborhood contained in some K_i.

Proof. The idea is that X being second countable means that any open cover can be refined (and shrunken) into a countable one. Indeed, for an open cover $\mathcal{U}$, define a function from the base $\mathcal{B}$ to $\mathcal{U}$ by sending a basic open subset B to one in $\mathcal{U}$ which contains it, if one exists. The image of this map is the required subcover.

Thus, for each $x \in X$, we find an open neighborhood U_x such that $\overline{U}_x$ is compact. Then $\{U_x\}_{x \in X}$ can be refined to a countable subcover $\{U_i\}_{i=1}^n$, and defining

$$K_j := \bigcup_{i \leq j} \overline{U_i}$$

will do the trick. ■

Definition 2.4 (locally finite)**.** An open cover $\mathcal{U}$ of a topological space X is *locally finite* if and only if each $x \in X$ admits an open neighborhood V for which

$$\{U \in \mathcal{U} : U \cap V \neq \varnothing\}$$

is finite for each $x \in U$.

Lemma 2.5. Fix a second countable locally compact topological space X. Then every base of X has a countable, locally finite subcover.

Proof. By Lemma 2.3, we may filter X as $X = \bigcup_i K_i$ by compact sets as described.

Let our base be $\mathcal{B}$, and we will build our subcover $\mathcal{U}$ by removing elements from $\mathcal{B}$, starting with $\mathcal{U} = \mathcal{B}$. We may immediately assume that it is countable because X is second countable, so we may enumerate $\mathcal{U}$ as $\{U_i\}_{i\in\mathbb{N}}$. We now apply the following inductive process for each n, starting with $n = 1$.

1. Because K_n is compact, we are granted a positive integer N for which $\{U_i\}_{i\leq N}$ covers K_n, which we keep in $\mathcal{U}$.

2. We then remove from $\mathcal{U}$ all other open subsets which intersect with K_n, and go back to the previous step now with K_{n+1}. Note that because $\mathcal{B}$ was a base, any element $x \notin K_n$ will admit an open neighborhood avoiding K_n, so $\mathcal{U}$ continues to be a base on $X \setminus K_n$.

At the end of this process, we see that $\mathcal{U}$ is certainly a countable cover of X, and it is locally finite because any point $x \in X$ admits an open neighborhood U contained in one of the K_ns, and each K_n only has intersection with finitely many open subsets in $\mathcal{U}$. ■

Let's also say something about differential forms.

Definition 2.6 (differential form)**.** Fix a manifold M of dimension n. For a positive integer $k \leq n$, we define $\Omega^k(M)$ to be the space of *differential forms* of degree k. Explicitly, $\Omega^k(M) := \wedge^k T^*M$.

Remark 2.7. In local coordinates $(x_1, \ldots, x_n)$ of M, a local frame of T^*M is given by $\{dx_1, \ldots, dx_n\}$. Thus, some $\omega \in \Omega^k(M)$ takes the form

$$\omega = \sum_{i_1<\cdots<i_k} f_{i_1\ldots i_k}(x)\, dx_{i_1} \wedge \cdots \wedge dx_{i_k}.$$

If we wanted to change coordinates to some $(y_1, \ldots, y_n)$, where we now think of x_i as a function of the ys, then

$$\omega = \sum_{\substack{i_1<\cdots<i_k \\ j_1<\cdots<j_k}} f_{i_1\ldots i_k} \det\left[\frac{\partial(x_{i_1}, \ldots, x_{i_k}}{\partial(y_{j_1}, \ldots, y_{j_k})}\right] dy_{j_1} \wedge \cdots \wedge dy_{j_k}.$$

Definition 2.8 (de Rham cohomology)**.** Fix a manifold M. There is a differential map $d\colon \mathcal{O}(M) \to \Omega(M)$ given by sending a function f to the covector df; this extends to a map $d\colon \Omega^k(M) \to \Omega^{k+1}(M)$. Then the *$k$th de Rham cohomology* of M is given by

$$\mathrm{H}^k_{\mathrm{dR}}(M) = \frac{\ker\left(d\colon \Omega^k(M) \to \Omega^{k+1}(M)\right)}{\operatorname{im}\left(d\colon \Omega^{k-1}(M) \to \Omega^k(M)\right)}.$$

2.1.2 Partition of Unity

It is frequently desirable to pass some result to an open cover, for which one uses partitions of unity.

Definition 2.9 (partition of unity)**.** Fix an open cover $\mathcal{U}$ of a manifold M. Then a *partition of unity* subordinate to $\mathcal{U}$ is a collection of smooth nonnegative functions $\{f_i\}_{i\in I}$ satisfying the following.

- For each i, we have $\operatorname{supp} f_i \subseteq U_i$ for some $U_i \in \mathcal{U}$.
- Each $y \in M$ admits an open neighborhood V such that all but finitely many of the f_is are nonzero on V.
- We have $\sum_i f_i = 1$.

Remark 2.10. For each $U \in \mathcal{U}$, we may then define a function

$$F_U := \sum_{\substack{i\in I \\ U_i = U}} f_i$$

is also a partition of unity, where $\operatorname{supp} F_U \subseteq U$ for each U.

Remark 2.11. Let $\{f_i\}$ be a partition of unity. Then for each $x \in M$, we see that the sum $\sum_i f_i(x)$ is a finite sum by the second point. Furthermore, only countably many of the f_is are allowed to be nonzero because any open cover of M admits a countable subcover, which we apply to the open cover coming from the second point.

Proposition 2.12. Every open cover $\mathcal{U}$ of a manifold M admits a smooth partition of unity subordinate to $\mathcal{U}$.

Proof. We may as well assume that M is connected. Then set $n := \dim M$. The function

$$h(t) := \begin{cases} e^{1/(t-1)} & \text{if } t \in [0,1), \\ 0 & textift \geq 1, \end{cases}$$

is a smooth function $[0,\infty) \to [0,1]$, supported on $[0,1]$, and has $h(0) = 1$. We can then define $H(x) := h\big(|x|^2\big)$ to be a smooth function $H\colon \mathbb{R}^n \to [0,1]$ supported on the unit ball satisfying $H(0) = 1$.

Briefly, for any closed embedding $\overline{B(0,1)} \to M$, we let the image be called a closed ball. The point is that any closed ball $\overline{B} \subseteq M$ admits a smooth function $H_B\colon M \to [0,1]$ defined by extending the composite

$$\overline{B} \cong \overline{B(0,1)} \xrightarrow{H} [0,1]$$

to all of M by extending by zero. In particular, this continues to be smooth because H and all its derivatives vanish at the boundary.

Now, for each $x \in M$, one can find an open neighborhood of x contained in some $U_x \in \mathcal{U}$ which is homeomorphic to $\mathbb{R}^n$. Thus, we can find a closed ball $\overline{B}_x$ around x contained in U_x. Then $\{B_x\}_{x\in M}$ is a cover for M, which by Lemma 2.5 can be reduced to a locally finite countable subcover $\{\overline{B}_i\}_{i\in I}$. Then note that the function

$$F := \sum_{i\in I} H_{B_i}$$

is locally a finite sum and thus well-defined. (Namely, by construction, each $x \in M$ admits an open neighborhood U admitting only finitely many of the B_is.) Further, it is smooth by construction (which is checked locally where this is a finite sum), and it is positive everywhere because H_{B_i} is positive on B_i. We can now define $f_i := H_{B_i}/F$ for each i, and we find that $\{f_i\}_{i\in I}$ is our desired partition of unity. For example, note that H_{B_i} is supported inside some $U_i \in \mathcal{U}$ by construction of B_i! ■

Remark 2.13. Even if M is merely a locally compact topological space, one can find continuous partitions of unity. Of course, a different argument is required.

2.1.3 Integration on Manifolds

We are now ready to define some notion of integration. Recall that one can integrate any continuous nonnegative function f on an open subset $U \subseteq \mathbb{R}^n$, notated

$$\int_U f(x_1, \ldots, x_n)\, dx_1 \wedge \cdots \wedge dx_n.$$

(This is explained in a typical calculus class.) Note that changing variables in the integral from some $x_\bullet$s to some $y_\bullet$s introduces a factor of

$$|\det [\partial x_i / \partial y_j]|,$$

but $dx_1 \wedge \cdots \wedge dx_n$ will introduce the same factor without the absolute values. Thus, our integral is only well-defined up to changes of variables with positive Jacobians. This amounts to a choice of orientation on the manifold.

Definition 2.14 (integral)**.** Let M be an oriented manifold of dimension n, and choose a continuous n-form ω which is nonnegative on oriented bases of the tangent spaces. Further, fix an orientation-preserving atlas $\{(U_i, \varphi_i)\}$ on M, and let $\{f_i\}$ be a partition of unity subordinate to the cover $\{U_i\}$. Then we define

$$\int_M \omega := \sum_{i \in I} \int_{\varphi(U_i)} \varphi_i^*(f_i \omega).$$

Remark 2.15. One can check that this integral does not depend on the choice of atlas and also on the choice of partition of unity. Indeed, if we want to compare the integral to one coming from an atlas $\{(V_j, \psi_j)\}_j$ with partition of unity $\{g_j\}_j$, one has an atlas $\{(U_i \cap V_j)\}_{ij}$ refining both with partition of unity $\{f_i g_j\}_{ij}$. But one can then calculate that refining an atlas and partition of unity will not change the answer of our integral simply by splitting up integrals in $\mathbb{R}^n$ into various open covers.

Remark 2.16. One can even make a definition without the hypothesis on ω. Indeed, for any ω, we can define

$$|\omega|(v_1, \ldots, v_n) :- |\omega(v_1, \ldots, v_n)|$$

for any oriented basis $\{v_1, \ldots, v_n\}$ of a tangent space. Then we may integrate $|\omega|$.

Remark 2.17. If ω is a complex-valued differential form, then we can still integrate ω by integrating the real and complex parts separately.

Remark 2.18. There is a bundle of "densities" on M, denoted $|\wedge^n TM|$, which is a real line bundle on M. The point is that "things we can integrate" live in this bundle. Formally, the real line bundle has transition functions given by $|g_{ij}|$, where the g_{ij}s are the transition functions for $\wedge^n TM$. It turns out that this bundle is always trivial, which one sees by trivializing it locally and then gluing the pieces together via a partition of unity; its positive sections are (smooth) measures on M.

It is worthwhile to explain the source of orientations for our purposes.

Remark 2.19. Any choice of continuous differential top form $\omega \in \Omega^n(M)$ which is non-vanishing everywhere, then we can use ω to define an orientation simply by saying that a basis $\{v_1, \ldots, v_n\}$ of a tangent space is oriented if and only if $\omega(v_1, \ldots, v_n) > 0$. (For example, this implies that nonorientable manifolds have no non-vanishing top forms.) Notably, this choice of ω then induces a measure μ on M given by sending U to the integral $\int_U \omega$; note that this measure is finite on small enough open subsets, which we can see by passing to Euclidean space.

Notation 2.20. Fix a non-vanishing top differential form ω of a manifold M. Then we let μ_ω denote the measure

$$\mu_\omega(U) := \int_U \omega.$$

The moral is that we may integrate general nonnegative continuous functions on M against the measure μ_ω, using measure theory.

Remark 2.21. If M is compact, then one can check that $\int_M \omega$ is finite. Indeed, compactness allows us to give M a finite cover by closed balls, and integrals over closed balls are known to be compact from calculus (for example, they have maximums and so on).

The last thing we should say about integration is Stokes's theorem.

Theorem 2.22 (Stokes)**.** Let M be an oriented manifold (with boundary) of dimension n. If ω is a smooth $(n-1)$-differential form with compact support, then

$$\int_M d\omega = \int_{\partial M} \omega.$$

Example 2.23. If $\partial M = \varnothing$, then $\int_M d\omega = 0$.

Sketch. Using a partition of unity, we may reduce to the case where M is a subset of Euclidean space. Namely, for a given partition of unity $\{f_i\}$, one finds that

$$\int_{\partial M} \omega = \sum_{i \in I} \int_{\partial M} f_i \omega,$$

and

$$\int_{\partial M} f_i \omega = \int_M (df_i \wedge \omega + f_i \, d\omega)$$

by a local calculation. (This reduces to the Fundamental theorem of calculus, once one finds a way to pass from arbitrary subsets of Euclidean space to boxes.) Now, summing over each i finds that $\sum_i df_i = d(1) = 0$, so the left summand cancels out. Summing over the right summand completes the proof. ■

2.1.4 Integration on Lie Groups

Here is our construction.

Lemma 2.24. Real lie groups are orientable.

Proof. Fix a real Lie group G with Lie algebra $\mathfrak{g}$ of dimension n. We can then choose any nonzero $\xi_1 \in \wedge^n \mathfrak{g}^*$, and then ξ_1 extends to a full left-invariant top differential form by

$$\xi_g := L_g^* \xi_1.$$

This continues to be a smooth top differential form, so we receive an orientation by Remark 2.19. ■

Remark 2.25. Note that this choice of left-invariant top differential form ξ is unique up to rescaling. Indeed, $\wedge^n \mathfrak{g}^*$ is a one-dimensional vector space, so any two ξ and ξ' will differ by a scalar at the identity, which then spreads out.

Definition 2.26 (Haar measure)**.** Fix a real Lie group G. Then we let μ_L be the left-invariant Haar measure induced by the left-invariant top differential form, which is unique up to scalar. We define μ_R similarly.

One may hope that μ_L and μ_R agree, but this is not always the case.

Proposition 2.27. Fix a real Lie group G. Then $\mu_L = \mu_R$ (up to scalar) if and only if $|\wedge^n \mathfrak{g}^*|$ s a trivial representation of G.

Proof. Note that $\mu_L = \mu_R$ is equivalent to μ_R being a left Haar measure, which is equivalent to μ_R being conjugation-invariant. Thus, we would like for the corresponding density $|\xi| \in |\wedge^n \mathfrak{g}^*|$ to be conjugation-invariant. But G is acting by the adjoint anyway, so having a density be conjugation-invariant implies that the full representation is trivial. ■

Remark 2.28. This representation is simply given by acting by $\det \mathrm{Ad}_g$, so it is equivalent to ask for $\det \mathrm{Ad}_g \in \{\pm 1\}$ for all g.

Example 2.29. Let G be the group of real matrices of the form $\left[\begin{smallmatrix} a & b \\ 0 & 1 \end{smallmatrix}\right]$, where $a > 0$. Then $\mathfrak{g}$ consists of matrices of the form $\left[\begin{smallmatrix} a & b \\ 0 & 0 \end{smallmatrix}\right]$, and one can check that toral elements $\mathrm{diag}(t, 1) \in G$ have nontrivial determinant adjoint action on $\mathfrak{g}$: indeed, $\mathrm{diag}(t, 1)$ sends $\left[\begin{smallmatrix} a & b \\ 0 & 0 \end{smallmatrix}\right] \mapsto \left[\begin{smallmatrix} a & tb \\ 0 & 0 \end{smallmatrix}\right]$.

This test is fairly easy to execute in some cases.

Definition 2.30 (unimodular)**.** A real Lie group G is *unimodular* if and only if $\mu_L = \mu_R$ (up to scalar).

Example 2.31. If G is compact, then we claim that G is unimodular. Indeed, the representation $g \mapsto |\det \mathrm{Ad}_g|$ is some character $G \to \mathbb{R}^+$. But the image is a compact subgroup of $\mathbb{R}^+$ is $\{1\}$, so we conclude that this character must be trivial.

Notation 2.32. If G is a compact real Lie group, we let μ be the Haar measure (which is both left- and right-invariant), normalized so that $\int_G \mu = 1$.

Indeed, the Haar measures agree by Example 2.31, and we may normalize $\int_G \mu$ because this value is finite by Remark 2.21.

2.1.5 Applications to Compact Lie Groups

Here is the usual application of our integration theory: the unitary trick from finite groups upgrades to compact groups.

Proposition 2.33 (Weyl's unitary trick)**.** Let G be a compact real Lie group, and let V be a continuous finite-dimensional complex representation. Then V is unitary.

Proof. Start with any positive-definite Hermitian form $\langle -, - \rangle_0$. Then we define the Hermitian form $\langle -, - \rangle$ on V by

$$\langle v, w \rangle := \int_G \langle gv, gw \rangle \, dg.$$

One can see directly that this is in fact Hermitian and positive-definite, and it is G-invariant by construction. We remark that this integral converges by measure theory (not differential topology): we are computing the integral of some continuous function on the compact space G. ■

Corollary 2.34. Let G be a compact real Lie group G. Then all continuous finite-dimensional complex representations are semisimple.

Proof. Choose any finite-dimensional complex representation V. For any choice of subrepresentation $W \subseteq V$, we need to find a complement. Well, Proposition 2.33 grants an invariant positive-definite Hermitian form $\langle -, - \rangle$ on V, so we may use the orthogonal complement $W^\perp \subseteq V$. The invariance of the form implies that $W^\perp$ is a subrepresentation, and linear algebra can verify that the natural map $W \oplus W^\perp \to V$ is an isomorphism. ■

Example 2.35. Note that the group SU_n is simply connected: indeed, an argument similar to Proposition 1.207 can show that the quotient $\mathrm{SU}_n/\mathrm{SU}_{n-1}$ is the sphere S^{n-1}. Thus, $\mathrm{Rep}_{\mathbb{C}} \mathrm{SU}_n$ is equivalent to the category $\mathrm{Rep}_{\mathbb{C}} \mathfrak{su}_n$, which is equivalent to the category $\mathrm{Rep}_{\mathbb{C}}(\mathfrak{su}_n)_{\mathbb{C}}$. But $(\mathfrak{su}_n)_{\mathbb{C}} = \mathfrak{sl}_n(\mathbb{C})$. We thus get a new proof that the category $\mathrm{Rep}_{\mathbb{C}} \mathfrak{sl}_n(\mathbb{C})$ is semisimple by Corollary 2.34!

Remark 2.36. It turns out that the method of the previous example can work for arbitrary reductive Lie algebras.

We can even provide a notion of matrix coefficients.

Definition 2.37. Fix a continuous finite-dimensional representation V of a Lie group G. For any $v \in V$ and $\ell \in V^*$, we define the *matrix coefficient* $m_{\ell,v} \colon G \to \mathbb{C}$ by

$$m(g) := \ell(gv).$$

Here is a remarkable upgrading result.

Proposition 2.38. Fix a continuous finite-dimensional representation V of a Lie group G. Then the matrix coefficients are smooth functions on G.

Proof. Say that a vector $v \in V$ is a smooth vector if and only if the matrix coefficients $m_{\ell,v}$ are smooth for all $\ell \in V^*$. Letting V^{sm} be the locus of smooth vectors, it is not too hard to check that V^{sm} is some subrepresentation of V. We would like to show that $V^{\mathrm{sm}} = V$. Because V is finite-dimensional it is enough to show that V^{sm} is merely dense in V.

The difficulty lies in constructing any interesting vectors in V^{sm}. Fix a smooth compactly supported function φ on G, which can for example be obtained from a partition of unity. Then for each $v \in V$, we define

$$v_\varphi := \int_G \varphi(g)(gv) \, dg.$$

It is not too hard to check that w is smooth: for any $h \in G$ and $\ell \in V^*$, we calculate that

$$\ell(hw) = \int_G \varphi(g)\ell(hgw)\,dg.$$

But this is $\int_G \varphi\left(h^{-1}g\right) gv\,dg$, which is a smooth function in $h \in G$ because φ is smooth.

Let's show that the vectors we have constructed are in fact dense. We now fix $v \in V$ and let φ vary. In particular, choose a sequence $\{\varphi_n\}$ of smooth compactly supported functions on G with the following properties.

- For any open neighborhood U of 1, one has $\operatorname{supp}\varphi_n \subseteq U$ for all but finitely many n.
- One has $\int_U \varphi_n = 1$ for all n.

We then set $w_n := v_{\varphi_n}$, and one can check that $w_n \to v$ as $n \to \infty$. (This is a fairly technical point: by the continuity of the representation, one knows that gv is close to v for g in some small open neighborhood U of the identity. Choosing n large enough so that $\operatorname{supp}\varphi_n \subseteq U$, one can then bound the difference $|v - w_n|$.) The density follows! ■

Remark 2.39. Even if V is infinite-dimensional, this argument successfully shows that V^{sm} is dense in V.

2.2 March 2

We now move towards the Peter–Weyl theorem.

2.2.1 Stating the Peter–Weyl Theorem

Last time, we showed that matrix coefficients of continuous representations are smooth.

Notation 2.40. Fix a continuous finite-dimensional irreducible representation V of a compact Lie group G, and let $\{v_i\}$ be an orthonormal basis for some choice of invariant unitary form on V. Then we define $\psi_{V,ij}$ by

$$\psi_{V,ij}(g) := v_j^*(gv_i).$$

Remark 2.41. The $\psi_{V,ij}$s are independent of the choice of invariant unitary form (which we recall exists from Proposition 2.33): indeed, the unitary form is unique up to scalar, and scaling the form merely adjusts the orthonormal basis.

Proposition 2.42. Choose irreducible finite-dimensional continuous representations V and W of a compact Lie group G. Then

$$\int_G \psi_{V,ij}(g)\overline{\psi_{W,k\ell}(g)}\,dg = \begin{cases} 0 & \text{if } V \not\cong W, \\ 1_{i=k}1_{j=\ell}/\dim V & \text{if } V = W. \end{cases}$$

Proof. We imitate the proof for finite groups. Let $\rho_V : G \to \mathrm{GL}(V)$ and $\rho_W : G \to \mathrm{GL}(W)$ be the associated continuous maps. Fix our appropriate orthonormal bases $\{v_i\}$ and $\{w_k\}$, so we see

$$\begin{aligned}\int_G \psi_{V,ij}(g)\overline{\psi_{W,k\ell}(g)}\,dg &= \int_G ((\rho_V(g) \otimes \rho_{\overline{W}}(g))(v_i \otimes w_k), v_j \otimes w_\ell)\,dg \\ &= \left(\int_G (\rho_V(g) \otimes \rho_{\overline{W}}(g))\,dg(v_i \otimes w_k), (v_j \otimes w_\ell)\right).\end{aligned}$$

Here, the large integral in the last expression is interpreted as taking place in the large finite-dimensional vector space of endomorphisms of $V \otimes \overline{W}$. Now, because W is unitary, we see that $\overline{W} \cong W^*$. Thus, P is some endomorphism of $V \otimes W^*$, and we can see by construction that it outputs to $(V \otimes W^*)^G$. For example, if $V \ncong W$, then $V \otimes W^* = \mathrm{Hom}(W, V)$ admits no G-invariants, so $P = 0$ follows.

It remains to handle the case where $V = W$. Then $\mathrm{Hom}_G(V, V) = \mathbb{C}$ by Schur's lemma, which we can trace back to $(V \otimes W^*)^G$ being one-dimensional and spanned by $u := \sum_i (v_i \otimes v_i)$. (Recall that $\{v_i\}$ is an orthonormal basis.) Now, $(u, u) = \dim V$ by a direct calculation. Thus, we see that P is a projector onto $\mathbb{C}u$, so it must be

$$P(x) = \frac{(x, u)}{(u, u)} u = \frac{(x, u)u}{\dim V}.$$

We conclude that our original integral is

$$\begin{aligned} \int_G \psi_{V,ij}(g)\overline{\psi_{W,k\ell}(g)}\, dg &= \sum_t \left(\frac{(v_i \otimes v_j, u)}{\dim V} (v_t \otimes v_t, v_j \otimes v_\ell) \right) \\ &= \frac{(v_i \otimes v_j, u)}{\dim V} 1_{j=\ell} \\ &= \frac{1}{\dim V} 1_{i=k} 1_{j=\ell}, \end{aligned}$$

as required. ■

We are now ready to state the Peter–Weyl theorem, which is an analog of the fact that matrix coefficients form a basis of $\mathrm{End}\, \mathbb{C}[G]$ for finite groups G.

Theorem 2.43 (Peter–Weyl)**.** Fix a compact Lie group G. As V varies over continuous irreducible representations, the matrix coefficients $\psi_{V,ij}$ form an orthogonal basis of $L^2(G)$ (as a Hilbert space).

Proposition 2.42 has already shown that our matrix coefficients are orthogonal, so it only remains to show that we have enough matrix coefficients to span. This will require an analytic input, which is not totally surprising.

Example 2.44. Let G be the circle group $\mathrm{U}(1)$. Because G is abelian, its continuous representations are all one-dimensional, and one can check that they all look like $\chi_n \colon G \to \mathbb{C}^\times$ given by $\chi_n(z) := z^n$. Then Theorem 2.43 asserts that the functions $\theta \mapsto e^{in\theta}$ form an orthonormal basis of $L^2(\mathbb{R}/\mathbb{Z})$ (where we have implicitly identified $\mathbb{R}/\mathbb{Z}$ with S^1). This is the main theorem on Fourier series!

Remark 2.45. The previous example explains why one may call Theorem 2.43 the genesis of "non-abelian" Fourier analysis. In particular, we see that Fourier analysis is therefore fairly well-understood for compact groups, but the corresponding story for locally compact groups is a subject of ongoing research.

Let's start by reformulating Theorem 2.43.

Notation 2.46. Let V be an irreducible finite-dimensional continuous representation of a compact Lie group G. Then there is a map $\psi_V \colon V \otimes V^* \to L^2(G)$ given by

$$\psi_V(v \otimes \ell)(g) := \ell(gv).$$

Remark 2.47. By Proposition 2.42, we see that a choice of orthonormal basis for all the irreducible representations V will exhibit an embedding

$$\bigoplus_{V \in \operatorname{IrRep} G} (V \otimes V^*) \to L^2(G).$$

Explicitly, Proposition 2.42 has explained that an orthonormal basis of the source goes to orthogonal (and nonzero) matrix coefficients in the target.

Definition 2.48. Let G be a compact Lie group. Then we define $L^2_{\text{alg}}(G) \subseteq L^2(G)$ to be the image of the map

$$\bigoplus_{V \in \operatorname{IrRep} G} (V \otimes V^*) \to L^2(G).$$

Remark 2.49. On the homework, we will show that $\varphi \in L^2(G)$ lives in $L^2_{\text{alg}}(G)$ if and only if φ generates a finite-dimensional representation under left translation. In fact, this is equivalent to generating a finite-dimensional representation under just right translation or under both left and right translation.

Remark 2.50. In light of Proposition 2.42, Theorem 2.43 is equivalent to asserting that any function in $L^2(G)$ which is orthogonal to all matrix coefficients must vanish. This is equivalent to being orthogonal to the span of all the matrix coefficients, which is $L^2_{\text{alg}}(G)$ by the above discussion.

Remark 2.51. Because $L^2(G)$ is some Hilbert space admitting a countable orthonormal basis, we see that the presence of this embedding implies that there can only be countably many irreducible representations of G!

Before we go further, here is an application.

Corollary 2.52. For each continuous finite-dimensional representation V of a compact Lie group G, define the character $\chi_V \colon G \to \mathbb{C}$ by $\chi_V(g) := \operatorname{tr}(g; V)$. As V varies over irreducible representations, the characters χ_V form an orthonormal basis of $L^2(G)^G$.

Proof. To begin, note that

$$\chi_V(g) = \sum_i \psi_{V,ii}(g).$$

In particular, by summing with Proposition 2.42, we see that $\langle \chi_V, \chi_W \rangle$ is

$$\int_G \chi_V(g)\overline{\chi_W(g)}\, dg = \begin{cases} 0 & \text{if } V \not\cong W, \\ 1 & \text{if } V = W. \end{cases}$$

Thus, we do have an orthonormal subset of $L^2(G)$. Additionally, we can see that χ_V lives in $L^2(G)^G$ because traces are independent of conjugation.

Now, because χ_V is exactly the image of $(V \otimes V^*)^G$ in $L^2(G)$ (as calculated above from an orthonormal basis), we see that $L^2_{\text{alg}}(G)^G$ is exactly the span of the χ_Vs. Thus, we want to show that $L^2_{\text{alg}}(G)^G$ is dense in $L^2(G)^G$. This will require a trick: choose some $\psi \in L^2(G)^G$, and then Theorem 2.43 grants us a sequence of functions $\{\psi_n\}_n$ in $L^2_{\text{alg}}(G)$ such that $\psi_n \to \psi$ as $n \to \infty$. We would like to push the ψ_ns to $L^2_{\text{alg}}(G)^G$, so we define

$$\widetilde{\psi}_n(g) := \int_G \psi_n\left(gxg^{-1}\right)\, dg.$$

This function now lives in $L^2(G)^G$, and a short calculation with Remark 2.49 can verify that it lives in $L^2_{\text{alg}}(G)$ as well. Furthermore, one can calculate that $\left\|\widetilde{\psi}_n - \psi\right\| \leq \|\psi_n - \psi\|$ (because ψ started out as an invariant), so $\widetilde{\psi}_n \to \psi$ as $n \to \infty$. ■

2.2.2 Compact Operators

Let's review a little functional analysis before we get going in earnest. For us, all Hilbert spaces are separable.

Definition 2.53 (bounded)**.** Let H be a Hilbert space. A linear operator $A\colon H \to H$ is *bounded* if and only if there is a constant $C \geq 0$ for which

$$\|Av\| \leq C\,\|v\|$$

for all $v \in H$. We define the *norm* $\|A\|$ to be the infimum of C satisfying the above inequality.

Remark 2.54. An operator A is bounded if and only if it is continuous.

Remark 2.55. By definition, $\|Av\| \leq (\|A\| + \varepsilon)\,\|v\|$ for any $\varepsilon > 0$ and any vector v. But we may take $\varepsilon \to 0^+$ to see that $\|Av\| \leq \|A\| \cdot \|v\|$.

Remark 2.56. The norm on bounded linear operators gives the algebra of bounded operators on H the structure of a Banach space.

Definition 2.57 (compact)**.** Let H be a Hilbert space. A linear operator $A\colon H \to H$ is *compact* if and only if there is a sequence of operators $\{A_n\}_n$ on H of finite rank such that $\|A_n - A\| \to 0$ as $n \to \infty$.

Remark 2.58. The subspace of finite-rank operators in $\operatorname{End} H$ is some open subset, and by definition, we see that the compact operators are the closure of this open subspace.

Here is what we will need about compact operators.

Definition 2.59 (precompact)**.** A subset U of a Hilbert space H is *precompact* if and only if any sequence admits a convergent subsequence.

Lemma 2.60. Let A be a compact operator on a Hilbert space H. Then A maps bounded sets to precompact sets.

Proof. Let $\{v_n\}_n$ be a bounded sequence, and we want to show that $\{Av_n\}_n$ admits a convergent subsequence. By scaling, we may assume that $\|v_n\| \leq 1$ for all n. Using the compactness of A (and by passing to a subsequence), we may find operators $\{A_n\}_n$ of finite rank so that $\|A_n - A\| < 1/n$ for all n.

Now, let $\{v_n^1\}$ be a subsequence of $\{v_n\}$ for which the sequence $\{A_1 v_n^1\}$ converges, which exists because this sequence is bounded and belongs to a finite-dimensional space. Then let $\{v_n^2\}$ be a subsequence of $\{v_n^1\}$ for which $\{Av_n^2\}$ also converges, and we may continue this process indefinitely to produce subsequences $\{v_n^m\}$.

This infinite process looks like it may fail to produce a sequence in the end, but we remedy this by considering the subsequence

$$w_n \coloneqq v_n^n.$$

We would like to show that the sequence $\{Aw_n\}$ converges. Well, note that

$$\begin{aligned}\left\|Av_i^k - Av_j^k\right\| &\le \left\|A_k v_i^k - A_k v_j^k\right\| + \|A - A_k\| \cdot \left\|v_i^k - v_j^k\right\| \\ &\le \left\|A_k v_i^k - A_k v_j^k\right\| + \frac{2}{k} - \varepsilon_k,\end{aligned}$$

for some $\varepsilon_k > 0$ chosen so that $\|A - A_j\| = \frac{2}{k} - \varepsilon_k/2$. Now, $\left\{A_k v_i^k\right\}$ converges, so there is a large M_k for which $i, j > M_k$ will have $\left\|A_k v_i^k - A_k v_j^k\right\| < \varepsilon_k$. Plugging this in, we see that $i, j > M_k$ has

$$\left\|Av_i^k - Av_j^k\right\| \le \frac{2}{k}.$$

Now, note that w_n is a subsequence of $\{v_n^k\}$ starting from the kth term, so there is a large N_k (matching up with M_k) for which $\|Aw_i - Aw_j\| \le 2/k$ for $i, j > N_k$. We conclude that $\{Aw_i\}$ is Cauchy, so it converges because we are in a Hilbert space. ■

Non-Example 2.61. If H is infinite-dimensional, then the identity operator is not compact.

Remark 2.62. The converse of Lemma 2.60 is also true, but we will not need it.

Our next task is to find some compact operators.

Lemma 2.63. Let M be a compact manifold, and let dx be a continuous measure. For a continuous function $K\colon M \times M \to \mathbb{R}$, define the kernel operator A on $L^2(M; dx)$ by

$$A\psi(y) := \int_M K(x, y)\psi(x)\, dx.$$

Then A is a compact operator.

Proof. Embed M into a tubular neighborhood of $[0, 1]^n$ for some larger n, so it suffices to prove the result for $[0, 1]^n$; alternatively, one can use a partition of unity argument by covering M with such boxes.[1] Up to rescaling the measure, we may assume that M has volume 1.

Then for each large N, tile $[0, 1]$ with intervals of length $1/N$, and approximate K by the corresponding Riemann sum: define

$$K_N(x, y) := \max_{(x', y')} K(x', y'),$$

where (x', y') varies over a chosen box around (x, y) with side length $1/N$. Let A_N be the corresponding kernel operator, which we see has finite rank because each K_N is given by its finitely many points. It remains to show that $A_N \to A$. Well,

$$\|A - A_N\| \le \sup_{(x,y)} |K(x, y) - K_N(x, y)|.$$

However, the source $M \times M$ is compact, so K's continuity upgrades to uniform continuity, so the right-hand side goes to 0 as N becomes large (because the distance between (x, y) and (x', y') in $K_N(x, y)$ is controlled). ■

Here is our main application, which is a spectral theorem.

Definition 2.64 (adjoint)**.** A bounded operator A on a Hilbert space H admits an *adjoint* $A^\dagger\colon H \to H$ for which

$$(A^\dagger v, w) = (v, Aw).$$

We say that A is *self-adjoint* if and only if $A^\dagger = A$.

[1] This reduction is not actually necessary: we just need to have some Riemannian metric on M.

Theorem 2.65 (Hibert–Schmidt)**.** Let A be a compact self-adjoint operator on a Hilbert space H. Then H admits an orthogonal decomposition

$$H = \ker A \oplus \widehat{\bigoplus_{\lambda}} H_\lambda,$$

where H_λ is the eigenspace for the eigenvalues of λ. In fact, H_λ is finite-dimensional, the eigenvalues are real, and the collection of eigenvalues is either finite or admits 0 as its only limit point.

Proof. Let's start by proving the result for A^2. Let β be the norm of A^2; if $\beta = 0$, there is nothing to do. We will hunt for an eigenvector for A^2 with eigenvalue β. Because A is compact, we have a sequence $\{A_n\}$ of finite-rank operators converging to A; by replacing A_n with $\frac{1}{2}(A_n + A_n^\dagger)$, we may even assume that the A_ns are self-adjoint. Now, the norm β_n of $\|A_n\|^2$ is the maximal eigenvalue of A_n (by linear algebra of finite-dimensional spaces), and because taking the norm is continuous, we see that $\beta_n \to \beta$.

Let's begin our hunt for an eigenvector. We know that each A_n^2 admits an eigenvector v_n with eigenvalue β_n. We may rescale each v_n to have norm 1, so the sequence $\{A^2 v_n\}$ admits a limit point, which we call $w \in H$. Notably, $A_n^2 v_n$ should also approach w, so because $A_n^2 v_n = \beta_n v_n$, we see that $\beta_n v_n \to w$, so $v_n \to \beta^{-1} w$. We thus see that $Aw = \beta w$, so A has an eigenvector w.

We can now replace H with $\{w\}^\perp$ and redo this process (inductively). If A has finite rank, then this process will terminate, and there is nothing more to do after noting that whatever we are left with is $\ker A$. Otherwise, we have a sequence of eigenvalues $\{\beta_1, \beta_2, \ldots\}$ for A^2, which is decreasing. Note that the compactness of A implies that no eigenvalue can be seen infinitely many times because then the restriction of A to this eigenspace is a scalar, which is not a compact operator! A similar argument shows that $\beta_n \to 0$ as $n \to \infty$ (otherwise, we have infinitely many eigenvectors with a lower-bounded eigenvalue, and the restriction to this sum of eigenspaces is not compact). We conclude that the eigenspaces H_{β_k} are all finite-dimensional. Taking an orthogonal complement, we see that

$$H = H' \oplus \widehat{\bigoplus_{k}} H_{\beta_k}.$$

We would like to check that H' is $\ker A$, which is not hard: for any $u \in H'$, we see that u is orthogonal to the eigenvectors with eigenvalues $\{\beta_1, \ldots, \beta_k\}$ by construction, so by definition of β_{k+1}, we see that $\left\|A^2 u\right\| \le \beta_{k+1} \|u\|$; sending $k \to \infty$ shows that $A^2 u = 0$, so $A^2 u = 0$ follows. Because A is self-adjoint, we see that A cannot have any vector u with $Au \ne 0$ while $A^2 u = 0$, for then the restriction of A to $\operatorname{span}\{u, Au\}$ is not self-adjoint. We conclude that $Au = 0$.

It remains to further refine this decomposition to work for A, which is not hard: each A preserves the given decomposition into H_{β_k}s, and then A restricts as a self-adjoint operator on this finite-dimensional space where A^2 is a scalar. Thus, A diagonalizes into eigenspaces with eigenvalues $\sqrt{\beta_k}$ and $-\sqrt{\beta_k}$. ■

2.3 March 9

Today, we will prove the Peter–Weyl theorem.

2.3.1 Proof of the Peter–Weyl Theorem

We will prove the following equivalent version of Theorem 2.43 (see Remark 2.50).

Theorem 2.66. Let G be a compact Lie group. Then $L^2_{\text{alg}}(G)$ is dense in $L^2(G)$.

Proof. Choose a "δ-like" sequence $\{h_N(x)\}$ of real-valued functions $h_N \colon G \to \mathbb{R}_{\ge 0}$ such that $\int_G h_N(x)\,dx = 1$ for all N and such that any open neighborhood U of 1 has $\operatorname{supp} h_N \subseteq U$ for all N large enough. (Such a sequence can be seen to exist by passing to Euclidean space in a small open neighborhood of $1 \in g$.)

Quickly, we note that we may assume that $h_N(x)$ satisfies $h_N(x) = h_N\left(x^{-1}\right)$ by replacing each h_N with $h'_N(x) := \frac{1}{2}\left(h_N(x) + h_N\left(x^{-1}\right)\right)$. We now let B_N be the corresponding integral operator, which is compact by Lemma 2.63. By staring at the definition

$$B_N\psi(y) := \int_G h_N(x)\psi\left(x^{-1}y\right)\,dx = \int_G h_N\left(yz^{-1}\right)\psi(z)\,dz,$$

and because h_N satisfies $h_N(x) = h_N\left(x^{-1}\right)$, one finds that B_N is self-adjoint. Additionally, B_N commutes with the G-action by right translation.

Now, Theorem 2.65 allows us to decompose $L^2(G)$ into eigenspaces for B_N as

$$L^2(G) = \ker B_N \oplus \widehat{\bigoplus_\lambda} H_{N,\lambda}.$$

Notably, $H_{N,\lambda} \subseteq L^2(G)$ is finite-dimensional for each λ, and it is a G-invariant subspace (because B_N commutes with the G-action), so we conclude that $H_{N,\lambda} \subseteq L^2_{\text{alg}}(G)$. It follows that $\operatorname{im} B_N$ is contained in $\overline{L^2_{\text{alg}}(G)}$.

We now do a careful limiting process to $N \to \infty$. Well, we for all $\psi \in L^2(G)$, we may find N for which $B_N\psi \in \overline{L^2_{\text{alg}}(G)}$, meaning that there is ψ_N for which $\|B_N\psi - \psi_N\|_2 < 1/N$. For example, if ψ is continuous, then one sees that $B_N\psi \to \psi$ uniformly by some arguments with the small supports of the h_Ns. Thus, for N large enough, we see that $\|B_N\psi - \psi\| < 1/N$, so $\|\psi - \psi_N\|_2 < 2/N$, so the ψ_Ns converge to ψ. It follows that ψ is also in the closure of $L^2_{\text{alg}}(G)$!

The previous paragraph has shown that $C(M)$ is in the closure of $L^2_{\text{alg}}(G)$, which is enough because $C(M)$ is dense in $L^2(M)$. Indeed, one can see this with a partition of unity argument, reducing to the case of a box. ■

2.3.2 Faithful Representations

Our next goal is to show that any compact group G can be realized in some unitary group; by Proposition 2.33, it is enough to show that G merely admits a faithful representation. Notable, one already knows that any finite group G admits a faithful representation V (namely, the regular representation), which can then be made into a unitary representation.

Lemma 2.67. Fix a compact Lie group G, and let $\{G_n\}_{n\geq 0}$ be a descending chain of closed Lie subgroups. Then this chain stabilizes.

Proof. Note that the sequence $\{\dim G_n\}_n$ is decreasing and hence stabilizes, so we may as well assume that it is constant. Because we have a chain of closed Lie subgroups, we also have a chain of closed Lie subalgebras $\{\operatorname{Lie} G_n\}_n$, and they now all have the same dimension, so they are all equal! Set $\mathfrak{g} := \operatorname{Lie} G_0$ to be this Lie algebra.

Now, let H be the identity component of any G_n, which all agree because all the $G_n s$ have the same Lie algebra. Then the sequence of quotients $\{G_n/H\}_n$ is a descending chain of finite sets (these sets are finite because our groups are compact), so it must stabilize. Thus, for n large enough, we see that the canonical inclusion $G_{n+1} \subseteq G_n$ is an isomorphism on the identity component and also on the irreducible components, so this inclusion must be surjective! Thus, our chain has stabilized. ■

Remark 2.68. The analogous statement for algebraic groups follows from the descending chain condition for ideals.

Proposition 2.69. Any compact Lie group G admits a faithful representation.

Proof. By Theorem 2.43, we know that G admits countably many irreducible representations, which we enumerate by $\{\rho_n\}_n$; say V_0 is the trivial representation. Then we let G_1 be the kernel of ρ_1, we let G_2 be $G_1 \cap \ker \rho_2$, and so on. In particular, G_n is the kernel of $\rho_1 \oplus \cdots \oplus \rho_n$, so it is enough to show that G_n is trivial for n large enough.

Well, we have a descending chain $\{G_n\}_n$ of subgroups, which must stabilize to some subgroup $H \subseteq G$ by Lemma 2.67. By construction, we know that H acts trivially on all irreducible representations of G, so H acts trivially on $L^2_{\text{alg}}(G)$ and thus trivially on $L^2(G)$ by Theorem 2.43. It follows that H is trivial: the only element $g \in G$ which acts trivially on $L^2(G)$ must be trivial, which one can see by choosing inside $L^2(G)$ a sufficiently small bump function. Indeed, any compactly supported function which is 1 at the identity and 0 at g will suffice. ■

Example 2.70. Proposition 2.69 is false for non-compact connected Lie groups. We will consider the group $\widetilde{G}$ which is the universal cover of $\mathrm{PSL}_2(\mathbb{R})$. Note the action of $\mathrm{PSL}_2(\mathbb{R})$ on the upper-half plane realizes $\mathrm{PSL}_2(\mathbb{R})/\mathrm{PSO}_2(\mathbb{R})$ as the upper-half plane, so $\pi_1(\mathrm{PSL}_2(\mathbb{R})) = \mathbb{Z}$ by using the long exact sequence in homotopy; thus, we see that $\widetilde{G}$ fits into a central extension

$$0 \to \mathbb{Z} \to \widetilde{G} \to \mathrm{PSL}_2(\mathbb{R}) \to 1.$$

Choose $z \in \widetilde{G}$ to generate the central copy of $\mathbb{Z}$, and we claim that z^2 acts by the identity on any complex finite-dimensional continuous representation $\rho\colon G \to \mathrm{GL}(V)$; this will complete the proof. Indeed, V naturally admits an action by the Lie algebra $\mathrm{Lie}\,\widetilde{G}_{\mathbb{C}} = \mathfrak{sl}_2(\mathbb{C})$; but then V admits a weight decomposition for this action, so ρ factors through $\mathrm{SL}_2(\mathbb{C})$. But now z^2 maps to the identity in $\mathrm{SL}_2(\mathbb{C})$ because $\mathrm{SL}_2(\mathbb{R}) \to \mathrm{PSL}_2(\mathbb{R})$ is a double cover.

Here is an application of having a faithful representation: we may sharpen our Peter–Weyl theorem.

Corollary 2.71. Fix a compact Lie group G. Then $L^2_{\text{alg}}(G)$ is dense in $C(G)$, where $C(G)$ has been given the supremum norm.

Proof. Let $\mathcal{A}$ be the closure of $L^2_{\text{alg}}(G)$ in $C(G)$, where the topology on $C(G)$ is induced by the supremum norm.

We would like to show that $\mathcal{A} = C(G)$. Well, $\mathcal{A}$ is a closed subalgebra, and $\mathcal{A}$ is closed under complex conjugation. Thus, by the Stone–Weierstrass theorem tells us that it is enough to check that $\mathcal{A}$ separates points. We will actually show that $L^2_{\text{alg}}(G)$ separates points. Note that $L^2_{\text{alg}}(G)$ separates points in G by Proposition 2.69: indeed, matrix coefficients of a faithful representation separate points from the identity (because of the trivial kernel). ■

Example 2.72. With $G = S^1$, we have shown that we can uniformly approximate periodic continuous functions $\mathbb{R}/\mathbb{Z} \to \mathbb{C}$ by polynomials in the exponentials $t \mapsto e^{2\pi nt}$. Restricting further to the even functions recovers an approximation of continuous functions by polynomials. Indeed, taking a quotient of $\mathbb{R}/\mathbb{Z}$ by the action of -1 produces a closed unit interval, and the corresponding characters $t \mapsto e^{2\pi int}$.

Here is another result of this flavor.

Corollary 2.73. Fix a compact Lie group G. Let $A \subseteq L^2_{\text{alg}}(G)$ be a subalgebra. If A is left-invariant, invariant under complex conjugation, and separates points, then $A =^2_{\text{alg}} (G)$.

Proof. By the Stone–Weierstrass theorem, we know that A is dense in $C(G)$ and hence in $L^2(G)$. Thus, for any irreducible representation V of G, we know that $\mathrm{Hom}_G(V, A)$ is dense in

$$\mathrm{Hom}_G\left(V, L^2(G)\right),$$

but the latter is isomorphic to V^* by Theorem 2.43. Thus, using semisimplicity of representations of G, we see that

$$A = \bigoplus_{V \in \mathrm{IrRep}(G)} (V \otimes V^*),$$

which is $L^2_{\mathrm{alg}}(G)$. ■

One can use the above corollary to show that any faithful representation (along with its dual) $\otimes$-generates the category of finite-dimensional representations of G.

Definition 2.74 (unimodular)**.** Fix a Lie group G. A finite-dimensional representation V of G is *unimodular* if and only if $\wedge^{\dim V} V$ is isomorphic to the trivial representation.

Remark 2.75. It is equivalent to ask for the action of G on V to factor through $\mathrm{SL}(V)$.

Proposition 2.76. Let V be a faithful finite-dimensional representation of a compact Lie group G.

(a) Let A be the subalgebra of $C(G)$ generated by the matrix coefficients. If V is unimodular, then $A = L^2_{\mathrm{alg}}(G)$.

(b) If Y is an irreducible representation of G, then there exist m and n such that Y is contained in $V^{\otimes n} \otimes V^{*\otimes m}$ for some n and m.

(c) If Y is an irreducible representation of G, and V is unimodular, then there exist n such that Y is contained in $V^{\otimes n}$.

Proof. We show the parts separately. Set $d := \dim V$ for brevity.

(a) We use Corollary 2.73. Because V is faithful, we automatically know that A separates points, and A is automatically left-invariant (because our matrix coefficients are arbitrary). Thus, it remains to show that A is invariant under complex conjugation. Well, note that $\overline{A}$ is generated by matrix coefficients of $\overline{V}$, which by Proposition 2.33 is isomorphic to V^*. But V is unimodular, so $V^* = \wedge^{d-1} V$ (using the canonical $\wedge$-pairing). But the matrix coefficients of $\wedge^{d-1} V$ are polynomials in the matrix coefficients of V, so we conclude that $\overline{A} \subseteq A$.

(b) By replacing V with $V \oplus V^*$, we may as well assume that V is unimodular, so we reduce to proving (c).

(c) Because V is unimodular, our argument in (a) has shown that $V^* = \wedge^{d-1} V$ and that $\mathrm{Sym}(V \otimes V^*)$ surjects onto A, which we know is $L^2_{\mathrm{alg}}(G)$. Thus, we know that Y is a subrepresentation of $\mathrm{Sym}(V \otimes V^*)$, so there is N large enough with $Y \subseteq \mathrm{Sym}^N(V \otimes V^*)$. The result follows after replacing V^* with $\wedge^{d-1} V$. ■

Example 2.77. We work with the group $G = S^1$, and we let V be the tautological representation $S^1 \subseteq \mathbb{C}^\times = \mathrm{GL}_1(\mathbb{C})$. Even though V is faithful, it is not unimodular, so its tensor powers do not generate Then the matrix coefficient of V is just $e^{i\theta}$, which generates the subalgebra $\mathbb{C}\left[e^{i\theta}\right]$ of $L^2_{\mathrm{alg}}(G) = \mathbb{C}\left[e^{i\theta}, e^{-i\theta}\right]$.

2.3.3 Haar Measures for Compact Groups

Let's explain how one could generalize our theory to arbitrary compact topological groups, such as $\mathrm{GL}_n(\mathbb{Z}_p)$. Here are the groups of interest.

Definition 2.78 (compact)**.** A *compact group* G is a separable compact Hausdorff topological group.

The main input was the presence of an integration theory. For compact Lie groups, we appealed to differential topology (using the Lie algebra). We will find a Haar measure with our means.

This will require some heavier measure theory.

Theorem 2.79 (Riesz–Markov–Kakutani)**.** Fix a compact Hausdorff separable topological space X. The collection of finite Borel measures μ on X is in bijection with the collection of positive continuous linear functionals $I_\mu \colon C(X;\mathbb{R}) \to \mathbb{R}$. In particular, for such a finite Borel measure μ, this map is given by

$$I_\mu(f) := \int_X f \, d\mu.$$

Here, a continuous linear functional $I \colon C(X;\mathbb{R}) \to \mathbb{R}$ is positive if I is nonnegative when the input function is nonnegative.

Before going further, we remark that we can normalize our measures.

Definition 2.80 (probability)**.** Fix a compact Hausdorff separable topological space X. Then a *probability measure* μ is a finite Borel measure for which $\mu(X) = 1$.

Remark 2.81. Because X is compact, any finite Borel measure μ has $\mu(X) < \infty$, so one can always normalize our finite Borel measure to be a probability measure.

Here is our main theorem.

Theorem 2.82 (Haar–von Neumann)**.** Fix a compact topological group G. Then there is a unique left-invariant probability measure μ on G. In fact, μ is right-invariant.

Remark 2.83. There are also left-invariant measures for locally compact groups. They do not have to be right-invariant.

Proof. Quickly, note that right-invariance follows from the uniqueness: if μ is a left-invariant probability measure, then for any $g \in G$, the measure $S \mapsto \mu(Sg)$ continues to be left-invariant and probability, so it must equal μ! Right-invariance follows.

Let's start with the existence of this measure. By Theorem 2.79, it is enough to find a positive left-invariant functional $I \colon C(X;\mathbb{R}) \to \mathbb{R}$. The idea is to compute the integral of f as an "average value," which we will compute via some iterated Markov chain.

1. Because G is separable, it admits a countable dense sequence $\{g_i\}_i$, and we also choose positive real numbers $\{p_i\}$ with $\sum_i p_i = 1$. Now, define the averaging operator A on $C(G;\mathbb{R})$ by

$$Af(x) := \sum_i p_i f(xg_i).$$

(This series converges because $\sum_i p_i = 1$, and f is bounded because G is compact.) By construction, $\|Af\| \le \|f\|$, so $\|A\| \le 1$; further, by plugging in a constant function, we see that $\|A\| = 1$.

Remark 2.84. Here is how to motivate the operator A: one should think about this as coming from a Markov chain on G given by sending $x \in G$. Then $Af(x)$ is the expectation of the function f, starting at x, of proceeding one step along this Markov chain. In particular, one may hope for stabilization, as one knows for Markov chains with finitely many states.

2. Let's explain a weak stabilization. Set $\nu(f) := \frac{1}{2}(\max f - \min f)$, which provides some seminorm, and we let $L \subseteq C(G;\mathbb{R})$ be the subspace of constant functions. In particular, $\nu(f)$ is the distance between a function f and L; for example, $\nu(f) = 0$ if and only if $f \in L$. Our stabilization claim is that any $f \notin L$ has
$$\nu(Af) < \nu(f).$$
To show this, we will show that $\max Af < \max f$; by symmetry (for example, using $-f$), we find that $\min Af > \min f$, so $\nu(Af) < \nu(f)$ follows. It remains now to show that $\max Af < \max f$. Well, for $x \in G$, choose an index j so that $f(xg_i) < \max f$; this exists because f is non-constant. (If f is constant on an orbit of the g_is, then the density of the g_is forces f to be full constant.) We now find
$$Af(x) = p_j f(xg_j) + \sum_{i \neq j} f(xg_i)$$
is bounded by
$$Af(x) \leq p_j f(xp_j) + (1 - p_j) \max f < \max f,$$
so $\max Af < \max f$.

3. We now iterate A to produce our stabilization. For $f \in C(G;\mathbb{R})$, we claim that $\{A^n f\}_n$ converges to some constant function, whose value we define to be $I(f)$. This map $f \mapsto I(f)$ will produce our Haar measure by Theorem 2.79. Set $f_n := A^n f$ for brevity.

   To start, note that $\|f_n\| \leq \|f\|$ for all n because $\|A\| = 1$, and we claim that this sequence is equicontinuous; then the Arzelá–Ascoli theorem grants us a convergent subsequence. In other words, we must check that any $\varepsilon > 0$ admits an open neighborhood U of the identity for which
$$|f_n(ux) - f_n(x)| \overset{?}{<} \varepsilon$$
for all $u \in U$ and $x \in G$. Because each f_n is just some finite linear combination of shifts of f, it is enough to check that f is uniformly continuous. This follows from the compactness of G. Indeed, we must show that there is an open neighborhood U of 1 for which $|f(ux) - f(x)| < \varepsilon$ for all $x \in G$ and $u \in U$. To see this, suppose not, and then there is $\varepsilon > 0$ so that any open neighborhood U_n of 1 admits $x_n \in G$ and $u_n \in U$ for which $|f(u_n x_n) - f(x_n)| \geq \varepsilon$. But then the $x_\bullet$s live in a compact space, so we may replace them with a convergent subsequence. By requiring $\bigcap U_n = \{1\}$, we see that $f(x_n) \to f(x)$ and so $f(u_n x_n) \to f(x)$, which is a contradiction.

   The Arzelá–Ascoli theorem now grants us a convergent subsequence $\{f_{n(m)}\}_m$, which will converge to some continuous function h in the supremum norm. But then note that
$$\nu(f_{n(m)}) \geq \nu(Af_{n(m)}) \geq \nu(f_{n(m+1)})$$
by the previous step, so we conclude that $\nu(h) = \nu(Ah)$ by sending $m \to \infty$, so $h \in L$ by the previous step. We now define $I(f) := h$, where we are identifying the constant function h with its value in $\mathbb{R}$.

   Before going on to the next step, we check that I is well-defined, for which we should check that $I(f)$ does not depend on the choice of convergent subsequence. Well, we actually claim the sequence of functions $\{f_n\}$ genuinely converges to h. Indeed, note that ν is continuous for the supremum norm, and we know that the sequence $\{\nu(f_n)\}_n$ is decreasing with a subsequence going to 0. It follows that $\{\nu(f_n)\}_n$ actually goes to zero, which means that $f_n - \frac{1}{2}(\max f_n + \min f_n) \to 0$, so the functions $\{f_n\}$ converge to the same limit as the constant functions $\frac{1}{2}(\max f_n + \min f_n)$. These constant functions have a convergent subsequence which goes to h, so we conclude that $f_n \to h$.

4. We start running some checks on I. For continuity, suppose that we have a sequence of functions $\{f_n\}$ converging to a function f. Linearity follows because A and taking limits are both linear: given functions f_1 and f_2 and scalars c_1 and c_2, we see that the sequence of functions $\{A^n(c_1 f_1 + c_2 f_2)\}$ will have the same limit as $\{c_1 A^n f_1 + c_2 A^n f_2\}$ (provided these limits exist, which holds by the previous step), so it follows that
$$I(c_1 f_1 + c_2 f_2) = c_1 \underbrace{\lim_{n\to\infty} A^n f_1}_{I(f_1)} + c_2 \underbrace{\lim_{n\to\infty} A^n f_2}_{I(f_2)}.$$

Continuing, note that $I(1) = 1$ because $A(1) = 1$, and I is left-invariant because A is left-invariant. Lastly, I is positive: if f is nonnegative, then Af is nonnegative, so the entire sequence of functions $\{A^n f\}$ is nonnegative, so $I(f) \geq 0$.

5. We complete the proof. A small adjustment to the definition of A reveals that the same argument produces a positive right-invariant continuous linear functional I_* with $I_*(1) = 1$, so at least such a functional exists. It remains to show uniqueness. Well, if J is any right-invariant positive continuous linear functional on $C(G; \mathbb{R})$ with $J(1) = 1$, then

$$J(f) = J(R_g f)$$

for any $g \in G$. Taking a linear combination of this equation, we see that $J(f) = J(Af)$, so $J(f)$ agrees with J applied to the limit of $\{A^n f\}$, so $J(f) = J(I(f))$, which is just $I(f)$. The uniqueness of the integral follows. ■

Remark 2.85. In fact, the proof at the end has shown that our left-invariant integral I is automatically right-invariant.

2.4 March 11

We began class by completing the proof of Theorem 2.82.

2.4.1 Examples of Compact Groups

To justify our study of compact groups, we should probably give some interesting examples which are not Lie groups.

Definition 2.86 (profinite)**.** Let $\{G_i\}$ be a countable collection of finite groups equipped with surjective homomorphisms $G_{i+1} \twoheadrightarrow G_i$ for each i. Then we define the *profinite* group G to consist of sequences $(g_i) \in \prod_i G_i$ for which $g_i \in G_i$ is the image of g_{i+1}. Thus, by the Axiom of choice, there are canonical surjections $\pi_n \colon G \twoheadrightarrow G_n$, and we may give G a topology by declaring the kernels of each π_n to be a neighborhood basis of the identity.

Remark 2.87. One may equivalently define the topology to be the subspace topology from the canonical injection $G \to \prod_i G_i$. It is not hard to check, then, that G is closed in $\prod_i G_i$, so it follows that G is compact. In general, any inverse limit of compact groups will produce a compact group.

Remark 2.88. One can check that $G^\circ = \{1\}$.

Remark 2.89. The group G is infinite as long as $\lim_n |G_n| = \infty$. In fact, as soon as $\lim_n |G_n| = \infty$, one can check that G is uncountable and has no isolated points.

Example 2.90. For any prime p, we can define the group $\mathbb{Z}_p$ to be given by the countable collection

$$\cdots \to \mathbb{Z}/p^3\mathbb{Z} \to \mathbb{Z}/p^2\mathbb{Z} \to \mathbb{Z}/p\mathbb{Z}.$$

In fact, $\mathbb{Z}_p$ is a ring, so for any algebraic group G over $\mathbb{Z}$, one also has a profinite group $G(\mathbb{Z}_p)$.

Example 2.91. The group $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ can be constructed as the limit of the groups $\mathrm{Gal}(K/\mathbb{Q})$, where $K/\mathbb{Q}$ is a Galois extension. Thus, $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ admits a natural structure of a profinite group.

Here are some more exotic examples.

Example 2.92. The inverse limit of the groups $\{\mathbb{R}/\mathbb{Z} \times \mathbb{Z}/p^\bullet\mathbb{Z}\}$ is $\mathbb{R}/\mathbb{Z} \times \mathbb{Z}_p$, which is a compact group.

Example 2.93. Let G be the inverse limit $\lim \mathbb{R}/p^\bullet\mathbb{Z}$, where the projections $\mathbb{R}/p^{n+1}\mathbb{Z} \to \mathbb{R}/p^n\mathbb{Z}$ are the canonical ones (induced by $\mathrm{id}\colon \mathbb{R} \to \mathbb{R}$). Then G is an inverse limit of compact groups, so it is a compact group. We claim that G lives in a short exact sequence

$$0 \to \mathbb{Z}_p \to G \to \mathbb{R}/\mathbb{Z} \to 0.$$

Indeed, the map $G \to \mathbb{R}/\mathbb{Z}$ is the canonical one, so it only remains to compute the kernel. Well, for each n, we see that the kernel of $\mathbb{R}/p^n\mathbb{Z} \to \mathbb{R}/\mathbb{Z}$ is represented by $\mathbb{Z}/p^n\mathbb{Z}$, so taking the limit gives $\mathbb{Z}_p$.

2.4.2 Representation Theory of Compact Groups

Our integration theory now has the following applications.

Corollary 2.94. Fix a separable compact topological group G. Then the finite-dimensional representations of G are unitary and completely reducible.

Proof. Repeat the proof of Corollary 2.34. ■

Theorem 2.95 (Peter–Weyl)**.** Fix a separable compact topological group G.

(a) The collection of finite-dimensional irreducible representations $\mathrm{IrRep}\, G$ is countable, and

$$L^2(G) = \widehat{\bigoplus_\rho}(V \otimes V^*).$$

(b) Set $L^2_{\mathrm{alg}}(G) := \bigoplus_\rho (V \otimes V^*)$. Then $L^2_{\mathrm{alg}}(G)$ is dense in $C(G)$, where $C(G)$ has been given the supremum norm.

Proof. The same arguments for Theorem 2.43 and Corollary 2.71 apply, as soon as we have a continuous approximation of the identity. Well, let $\{U_i\}$ be a neighborhood basis of the identity; we may assume that $\overline{U}_{i+1} \subseteq U_i$ for each i because G is locally compact. Then Urysohn's lemma grants us functions $f_i\colon G \to [0,1]$ such that $\operatorname{supp} f_i \subseteq U_i$ but $f_i|_{U_{i+1}} = 1$. Thus, $\int_G f(g)\,dg \geq 0$, and we can rescale these functions to be our continuous approximation of the identity. ■

Remark 2.96. Suppose G is profinite, then any continuous representation of G maps to $\mathrm{GL}(V)$ for some complex vector space V. A "no small subgroups" argument is able to show that this representation thus factors through a finite quotient of G.

As an application, we may classify our compact groups!

Corollary 2.97. Any compact group G is an inverse limit of compact Lie groups.

Proof. By Theorem 2.95, we may enumerate the irreducible representations of G as $\{\rho_i\}_{i\in\mathbb{N}}$. Let K_m be the kernel of the representation $\bigoplus_{i\le m}\rho_i$, and define $G_m := G/K_m$. We now combine two observations.

- Note that G/K_m embeds as a closed subgroup of $\mathrm{U}(\rho_1\oplus\cdots\oplus\rho_m)$, which turns out to be a Lie subgroup. Thus, G_m is a compact Lie group.
- On the other hand, matrix coefficients of the ρ_is separate points by Theorem 2.95. Indeed, it is enough to show that $L^2_{\mathrm{alg}}(G)$ separates points, but this is dense in $C(G)$, which certainly separates points (for example, by our construction of approximation to identity).

Combining the above two points shows us that the canonical map

$$G \to \lim_{m\to\infty} G_m$$

embeds G into an inverse limit of compact Lie groups. We claim that this map is actually surjective, which will complete the proof. Indeed, choose a sequence (g_m) in the target. For each m, we see that the pre-image of $g_m \in G/K_m$ is the closed subset $g_mK_m \subseteq G$. By construction, this family of compact sets $\{g_mK_m\}_m$ is a descending chain of nonempty compact sets, so the finite intersection property for compact groups implies that $\bigcap_m g_mK_m$ is nonempty. Any $g \in G$ in this intersection will map to the sequence (g_m)! ■

2.4.3 The Hydrogen Atom

We now switch topics. Electrons in an atom turn out to not exactly behave as particles (with well-defined position at all times). Instead, they kind of behave like they are in a cloud around the atom, with some probability of being in any particular location. We are interested in calculating the probability distribution $p(x,y,z,t)$ of an electron living in a hydrogen atom, which satisfies

$$\int_{\mathbb{R}^3} p(x,y,z,t)\,dx\,dy\,dz = 1$$

for all t. Schrödinger's quantum mechanics claims that p arises from a more natural complex-valued function $\psi(x,y,z,t)$ from which one recovers p from $p = |\psi|^2$. To calculate ψ, one has the following.

Theorem 2.98 (Schrödinger's equation)**.** The wave function ψ satisfies

$$i\frac{\partial\psi}{\partial t} = H\psi,$$

where H is a quantum Hamiltonian $-\frac{1}{2}\Delta - \frac{1}{r}$.

Remark 2.99. This H arises from quantizing the classical Hamiltonian $\frac{p^2}{2} - \frac{1}{r}$, where the momentum $p = p_x^2 + p_y^2 + p_z^2$ is "quantized" to $\Delta = \partial_x^2 + \partial_y^2 + \partial_z^2$.

Remark 2.100. The previous discussion implies that our wave function ψ is required to satisfy

$$\int_{\mathbb{R}^3} |\psi(x,y,z,t)|^2\,dx\,dy\,dz = 1$$

for all t.

One can solve this equation using separation of variables because the right-hand side does not depend on time. As such, let's quickly review how separation of variables works.

Exercise 2.101. Suppose that we have a Hermitian operator A on a complex finite-dimensional vector space V, and we solve the equation

$$i\frac{\partial v}{\partial t} = Av.$$

Proof. Well, A admits a diagonalization with eigenbasis $\{v_j\}$ and real eigenvalues $\{\lambda_j\}$. Then we may write $v(t) = \sum_j f_j(t)v_j$, so we see that

$$if_j'(t) = \lambda_j f_j(t),$$

which implies that $f_j(t) = f_j(0)e^{-\lambda_j t}$. We thus see that

$$v(t) = \sum_j f_j(0)e^{-i\lambda_j t}v_j,$$

as required. ■

We now attempt to solve Schrödinger's equation. Suppose that we could find a basis $\{\psi_n\}$ of $L^2(\mathbb{R}^3)$ satisfying $H\psi_n = E_n\psi_n$ for each n, where each E_n is real. Then the same argument shows that

$$\psi(x, y, z, t) = \sum_n a_n e^{-iE_n t}\psi_n(x, y, z).$$

Thus, we reduce ourselves to actually finding these functions ψ with $H\psi = E\psi$. Note that we are interested in solving "bounded states" for our electron, which amounts to requiring ψ to stay bounded near the origin, which we can see is equivalent to $E < 0$.

We would like to exploit the symmetry of our system to find these functions ψ. In particular, the form of Δ is invariant under rotations, so we should try to work in spherical coordinates (r, θ, φ). In particular, it turns out that

$$\Delta = \Delta_r + \frac{1}{r^2}\Delta_{\text{sph}},$$

where

$$\Delta_r = \partial_r^2 + \frac{2}{r}\partial_r,$$

and

$$\Delta_{\text{sph}} = \frac{1}{\sin^2\varphi}\partial_\theta^2 + \frac{1}{\sin\varphi}\partial_\varphi \circ \sin\varphi \circ \partial_\varphi,$$

which is the Laplace–Beltrami operator. We are now hunting for functions ψ for which

$$\partial_r^2\psi + \frac{2}{r}\partial_r\psi + \frac{2}{r}\psi + \frac{1}{r^2}\Delta_{\text{sph}}\psi = -2E\psi.$$

We may now apply separation of variables: we may write ψ as a function of $r \in \mathbb{R}^+$ and a unit vector u, and then we will pretend that ψ can be spanned by functions of the form $\psi(r, u) = f(r)\xi(u)$, where ξ is an eigenvector for Δ_{sph} with eigenvalue $-\lambda$. Then we are interested in solving the equations $\Delta_{\text{sph}}\xi = -\lambda\xi$ and

$$f''(r) + \frac{2}{r}f'(r) + \left(\frac{2}{r} - \frac{\lambda}{r^2} + 2E\right)f(r) = 0.$$

We will solve these equations, using the representation theory of $\mathrm{SO}(3)$, next class.

2.5 March 16

Today, we continue with the hydrogen atom.

2.5.1 The Eigenfunctions

Let's start by finding eigenfunctions for the spherical Laplacian.

Lemma 2.102. We compute the eigenvalues λ and functions $\xi \in L^2\left(S^2\right)$ such that

$$\Delta_{\mathrm{sph}}\xi = \lambda\xi.$$

Proof. We are hunting for a function $\xi \in L^2\left(S^2\right)$ and $\lambda \in \mathbb{C}$ for which $\Delta_{\mathrm{sph}}\xi = \lambda\xi$. Note that Δ_{sph} is $\mathrm{SO}(3)$-invariant, so the set of solutions will be a representation of $\mathrm{SO}(3)$. By Theorem 2.43, we know that

$$L^2(\mathrm{SO}(3)) = \widehat{\bigoplus_m} L_{2m} \otimes L_{2m}^*.$$

Here, L_{2m} is the $2m$th symmetric power of the spin representation, and these are all the irreducible representations of $\mathrm{SO}(3)$, which we see because $\mathrm{Spin}(3) = \mathrm{SL}(2)$. Note that $S^2 = \mathrm{SO}(3)/\mathrm{SO}(2)$ as a homogeneous space, so it follows that

$$L^2\left(S^2\right) = \widehat{\bigoplus_m} L_{2m} \otimes \left(L_{2m}^*\right)^{\mathrm{SO}(2)},$$

which one can calculate is

$$L^2\left(S^2\right) = \widehat{\bigoplus_m} L_{2m}.$$

For example, the algebraic functions $L^2_{\mathrm{alg}}\left(S^2\right)$ (namely, the polynomial functions on the sphere) is $\bigoplus_m L_{2m}$. Now, Δ_{sph} can be checked to send trigonometric functions to trigonometric functions, so it descends to an $\mathrm{SO}(3)$-invariant function on $L^2_{\mathrm{alg}}\left(S^2\right)$, which must act by some eigenvalue $-\lambda_m$ on each L_{2m}.

For example, let's hunt for the weight-0 eigenfunctions in L_ℓ. Then they are invariant under $\mathrm{SO}(2)$, so they should depend only on φ, so we can write them as some polynomial $Y_\ell^0 = P_\ell(\cos\varphi)$ of degree ℓ. By orthogonality of the matrix coefficients, one sees that

$$\int_0^\pi P_\ell(\cos\varphi)P_k(\cos\varphi)\sin\varphi\, d\varphi = 0$$

for $\ell \neq k$. (Here, the $\sin\varphi$ comes from our choice of measure.) A change of coordinates makes this

$$\int_{-1}^1 P_\ell(z)P_k(z)\, dz = 0$$

for $\ell \neq k$, so we are looking at the Legendre polynomials! Let's compute the eigenvalues: because P_ℓ does not depend on θ, we see that

$$\Delta_{\mathrm{sph}}P = \partial_z\left(1-z^2\right)\partial_z P_\ell.$$

If the eigenvalue is $-\lambda_\ell$, then a calculation on the leading term reveals that $\lambda_\ell = \ell(\ell+1)$.

In general, for each $L_{2\ell}$, there is an eigenfunction $Y_\ell^m(\varphi,\theta)$ of weight $m \in [-\ell,\ell]$. Accordingly, $e^{-im\theta}Y_\ell^m$ is $\mathrm{SO}(2)$-invariant, so it takes the form $e^{im\theta}P_\ell^m(z)$. For even m, P_ℓ^m turns out to be a polynomial in z; otherwise, it is $\sqrt{1-z^2}$ times a polynomial in z. Our eigenvalue calculation from earlier reveals that

$$\Delta_{\mathrm{sph}}Y_\ell^m = -\ell(\ell+1)Y_\ell^m,$$

so one finds that P_ℓ^m satisfies the Legendre differential equation

$$\partial_z\left(1-z^2\right)\partial_z P - \frac{m^2}{1-z^2}P + \ell(\ell+1)P = 0.$$

One can solve this differential equation. ∎

On the homework, we will solve this differential equation, and we will show that they take the following form.

Definition 2.103 (generalized Legendre polynomials)**.** For a nonnegative integer ℓ and $m \in [-\ell, \ell]$, the *generalized Legendre polynomials* are

$$P_\ell^m(z) = \left(1 - z^2\right)^{m/2} \partial_z^{\ell+m} \left(1 - z^2\right)^\ell.$$

Remark 2.104. Here is a high-level explanation for these eigenvalues. There are angular momentum differential operators L_x, L_y, and L_z, which "quantize" the components of angular momentum (computed as $p \times r$). It turns out that $\{-iL_x, -iL_y, -iL_z\}$ spans $\mathfrak{so}_3(\mathbb{R})$, and it turns out that the Casimir element

$$L_x^2 + L_y^2 + L_z^2$$

is $-\Delta_{\text{sph}}$. However, one can calculate that the action of the Casimir on $L_{2\ell}$ is $\ell(\ell+1)$, which explains why we received those eigenvalues.

We now turn to the part depending on r.

Lemma 2.105. We compute the functions f on $\mathbb{R}$ and energies $E < 0$ for which

$$f''(r) + \frac{2}{r} f'(r) + \left(\frac{2}{r} - \frac{\lambda}{r^2} + 2E\right) f(r) = 0.$$

Proof. It will be convenient to rescale f to a function h satisfying $f(r) = r^\ell e^{-r/n} h(2r/n)$, where n is some positive real number. (It will later turn out that n is a positive integer.) Applying this change of variables yields

$$\rho h''(\rho) + (2\ell + 2 - \rho) h'(\rho) + \left(n - \ell - 1 + \frac{1 + 2En^2}{4}\rho\right) h(\rho) = 0.$$

Thus, it is convenient to set n to be $\sqrt{-1/(2E)}$ so that the equation reads

$$\rho h''(\rho) + (2\ell + 2 - \rho) h'(\rho) + (n - \ell - 1) h(\rho) = 0.$$

This is called the generalized Laguerre equation. Notably, the condition $\|\psi\|^2 < \infty$ translates to

$$\int_{\mathbb{R}^+} \rho^{2\ell+2} e^{-\rho} \left|h(\rho)\right|^2 \, d\rho < \infty.$$

For example, we should check that we do not blow up at zero: if h around zero looks like $\rho^s(1 + o(\rho))$, then we find a characteristic equation $s(s+2\ell+1) = 0$ (from the differential equation). Thus, $s = 0$ or $s = -2\ell - 1$. Let's explain how to throw out the latter case.

- With $\ell > 0$, one finds that $\rho^{2\ell+2} \left|h(\rho)\right|^2 \sim \rho^{-2\ell}$ if $s = -2\ell - 1$ is not integrable at zero, so we may throw out this case.
- If $\ell = 0$, then we can try to take $s = -2\ell - 1 = -1$. Then $\rho^2 \left|h(\rho)\right|^2$ has finite limit at $\rho = 0$, which appears to be integrable. In this case $\ell = 0$, one finds that ξ is constant, so ψ depends only on the radius, and we see $\psi(r) \sim 1/r$ near $r = 0$. But then $H\psi - E\psi$ does not vanish as a distribution because it is roughly δ_0. Thus, ew can throw out $s = 0$ in this case as well.

Thus, for any ℓ, we are required to work with $s = 0$, producing a unique solution h_n satisfying the initial condition $h_n(0) = 1$. We may calculate h_n using a power series. By substituting a power series, we find that

$$h_n(\rho) = \sum_{k=0}^\infty \frac{(1 + \ell - n) \cdots (k + \ell - n)}{(2\ell + 2) \cdots (2\ell + 1 + k))} \cdot \frac{\rho^k}{k!}.$$

The ratio test verifies that this converges for all ρ. If n is not a positive integer, then computing the (large k) power series coefficients finds that $|h(\rho)| \sim e^\rho$, which is not possible by the $\|\psi\|^2 < \infty$ condition. On the other hand, n being a positive integer makes $h(\rho)$ into a polynomial, which is successfully integrable. ■

The moral is that our lemma has found that the solutions h_n are the so-called generalized Laguerre polynomials.

Definition 2.106 (generalized Laguerre polynomial)**.** Choose a positive integer N and real number α. Then the *generalized Laguerre polynomials* is the polynomials

$$L_N^\alpha(\rho) := \sum_{k=0}^{N} (-1)^N \frac{N \cdots (N-k+1)}{(\alpha+1)\cdots(\alpha+k)} \cdot \frac{\rho^k}{k!}.$$

Combined with our discussion last class, we have proven the following theorem.

Theorem 2.107. The bound states of the hydrogen atom are, up to scaling,

$$\psi_{n\ell m} = r^\ell e^{-r/n} L_{n-\ell-1}^{2\ell+1}\left(\frac{2r}{n}\right) Y_\ell^m(\theta,\varphi),$$

where $Y_\ell^m(\theta,\varphi) = e^{im\theta} P_\ell^m(\varphi)$, and n is a positive integer, and ℓ is a nonnegative integer at most $n-1$, and $m \in [-\ell,\ell]$. The energy level E_n of this bound state is $E_n = -\frac{1}{2n^2}$.

Let's name the parameters for our quantum states.

Definition 2.108 (quantum number)**.** The *quantum number* of the bound state $\psi_{n\ell m}$ is the number n. We also call ℓ the *azimuthal quantum number*, and m is the *magnetic quantum number*.

Corollary 2.109. The space W_n of bound states with principal quantum number n is isomorphic to the representation

$$L_0 \oplus L_2 \oplus \cdots \oplus L_{2n}$$

of $\mathrm{SO}(3)$. In particular, it has dimension n^2.

Proof. This follows from the proof of Theorem 2.107. ■

Remark 2.110 (spin)**.** From the periodic table, one expects to have $2n^2$ states for a given n: for example, there are two places the electron can go to in the s orbit (corresponding to the two electrons in the helium atom). It turns out that electrons have an extra degree of freedom, which is called the spin, taking values in $\{+1/2, -1/2\}$. Then the space of states lives in $L^2\left(\mathbb{R}^3\right) \otimes \mathbb{C}^2$. It turns out that $\mathbb{C}^2$ experimentally looks like the two-dimensional representation of $\mathfrak{so}(3)$, so the correct group acting should be $\mathrm{Spin}(3)$ instead of $\mathrm{SO}(3)$. This is not a contradiction in quantum mechanics (though it looks like a contradiction in classical physics) because physical states are only recovered from quantum ones up to a phase factor $e^{i\varepsilon}$, and the extra element in $\mathrm{Spin}(3)$ only adjusts the quantum state up to a sign! Anyway, in the end, we get the correct $2n^2$ states.

2.5.2 Coulomb Waves

The differential operator Δ_{sph} fails to be compact, so we should not expect its spectrum to go to zero. Thus, it is surprising that we only found energy levels going to zero. The solution to this problem is that the bound states $\psi_{n\ell m}$ turn out to not be a basis of $L^2\left(\mathbb{R}^3\right)$. Indeed, let φ be some compactly supported function on $\mathbb{R}^3$ at the origin. Then define $\varphi_s(x) := \varphi(x+sa)$ to be some scaling.

We claim that $(H\varphi_s, \varphi_s)$ is positive for large s: indeed, by integration by parts, one can remove the Laplacian, and we find that

$$(H\varphi_s, H\varphi_s) = \frac{1}{2}\int |\nabla\varphi|^2 \, dV - \int \frac{1}{\|x\|} |\varphi(x+sa)|^2 \, dV.$$

By shifting coordinates, this is

$$(H\varphi_s, H\varphi_s) = \frac{1}{2}\int |\nabla\varphi|^2 \, dV - \int \frac{|\varphi(x+sa)|^2}{\|x-sa\|} \, dV.$$

Now, as $s \to \infty$, the right integral goes to zero, and the left integral is constant and positive, so $(H\varphi_s, \varphi_s)$ is positive! Thus, φ_s cannot be in the span of the $\psi_{n\ell m}$s for sign reasons.

Thus, it is possible to get positive energy levels.

Definition 2.111 (Coulomb waves)**.** The eigenfunctions with positive energy levels in $L^2\left(\mathbb{R}^3\right)$ are called the *Coulomb waves*.

It turns out that the Coulomb waves turn out to have a continuous spectrum, but the core ideas used to understand them are rather similar to the bound states.

2.6 March 18

Here we go.

2.6.1 The Pauli Exclusion Principle

Here is our definition.

Definition 2.112 (Pauli exclusion)**.** If the space of states of a single electron is V, then the space of sates of n electrons is $\wedge^n V$.

Notably, if states were actual vectors in classical mechanics, then we would expect that the collection of spaces of two vectors would be $\operatorname{Sym}^2 V$ because symmetry should be preserved if the electrons are indistinguishable. However, in quantum mechanics, our states only matter up to phase, so it would be okay for permutations to act up to a scalar. There are two such one-dimensional characters of the symmetric group, which correspond to bosons and fermions.

Definition 2.113 (boson)**.** A *boson* is a kind of state in a space V for which the space of n bosons is $\operatorname{Sym}^n V$.

Definition 2.114 (fermion)**.** A *fermion* is a kind of state in a space V for which the space of n bosons is $\wedge^n V$.

Thus, the Pauli exclusion principle is just the physical observation that electrons and fermions.

Example 2.115. The space V_n of electrons at energy level $-\frac{1}{2n^2}$ has dimension $2n^2$. Thus, the space of k electrons is $\wedge^k V_n$; notably, there cannot be more than $2n^2$ electrons at a given level. For example, at the first energy level $-1/2$, we are allowed at most two energy levels, which correspond to the two electrons in helium. The next energy level has eight permissible electrons.

Frequently, the presence of repeated eigenvalues of some matrix is due to the presence of an underlying representation. For example, the adjacency matrix of a graph G will frequently have repeated eigenvalues when G has a large automorphism group. There is a similar notion of "degeneracy" in physics due to our repeated energy levels. However, note that

$$W_n = 2L_1 \oplus 2L_3 \cdots \oplus 2L_{2n-3} \oplus L_{2n-1}$$

as a representation of $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$. The explanation for the many 2s is due to the presence of an additional symmetry of the hydrogen atom: setting $\vec{r} := (x, y, z)$ and $\vec{p} := (-i\partial_x, -i\partial_y, -i\partial_z)$, there is an angular momentum operator $\vec{L} := \vec{r} \times \vec{p}$. The components of $\vec{L}$ define the action of $\mathfrak{so}(3)$, so because $\vec{p}^2 = \vec{p} \cdot \vec{p}$ is the negative Laplacian, which is $\mathfrak{so}(3)$-invariant, we see $[\vec{L}, \vec{p}^2] = 0$.

We now define the second-order differential operator

$$\vec{A}_0 := \frac{1}{2}\left(\vec{p} \times \vec{L} - \vec{L} \times \vec{p}\right),$$

which automatically commutes with $\vec{p}^2$. We would like for $\vec{A}_0$ to also commute with the Hamiltonian $H = -\frac{1}{2}\Delta + U(r) = -\frac{1}{2}p^2 + U(r)$. To this end, we modify $\vec{A}_0$ to be $\vec{A} := \vec{A}_0 + \varphi(r)\vec{r}$, and we would like for $[\vec{A}, H] = 0$. On the homework, we will show that such a φ exists if and only if $U(r) = \frac{C}{r} + D$ (which is the Coulomb potential), and then one can compute φ. The moral is that the components of $\vec{A}$ define a third $\mathfrak{sl}_2$-triple acting on $L^2\left(\mathbb{R}^3\right)$ and commuting with our other two $\mathfrak{sl}_2 = \mathfrak{so}(3)$ acting on W_n.

Remark 2.116. We call this extra $\mathfrak{sl}_2$ a "hidden symmetry." Such things also exist classically: for example, Kepler's laws are periodic only due to the special nature of the potential. Small adjustments to the potential no longer have

It now turns out that $W_n = L_{n-1} \boxtimes L_{n-1}$ as a representation of the combined $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$; notably, it is irreducible, and one finds the decomposition $L_0 \oplus L_2 \oplus \cdots \oplus L_{2n}$ by decomposing the box product against the diagonal $\mathfrak{sl}_2 \subseteq \mathfrak{sl}_2 \oplus \mathfrak{sl}_2$. Accounting for spin, we get $L_{n-1} \boxtimes L_{n-1} \boxtimes L_1$.

2.6.2 The Automorphism Group

We have some time before spring break, so let's say something about our next topic. Let's start by recalling the following facts.

Proposition 2.117. Fix a complex semisimple Lie algebra $\mathfrak{g}$. Then $\mathrm{Aut}(\mathfrak{g})$ is a complex Lie group with Lie algebra $\mathfrak{g}$.

Proof. Note that $\mathrm{Aut}(\mathfrak{g})$ is a closed Lie subgroup of $\mathrm{GL}(\mathfrak{g})$. One can check that the Lie algebra of $\mathrm{Aut}(\mathfrak{g})$ is given by the derivations $\mathfrak{g} \to \mathfrak{g}$. However, a Lie algebra gives a large source of derivations: there is a map $\mathfrak{g} \to \mathrm{Der}(\mathfrak{g})$ given by sending X to the adjoint action. For a semisimple algebra, this map is an isomorphism of Lie algebras, so the result follows. ■

Definition 2.118 (adjoint group)**.** Fix a complex semisimple Lie algebra $\mathfrak{g}$. Then we define the *adjoint group* G_{ad} to be $\mathrm{Aut}(\mathfrak{g})^\circ$.

Example 2.119. For $\mathfrak{g} = \mathfrak{sl}_n$, one can check that G_{ad} is $\mathrm{PGL}_n(\mathbb{C})$.

We also had the following classification of Cartan subalgebras.

Proposition 2.120. Fix a complex semisimple Lie algebra $\mathfrak{g}$. Then all Cartan subalgebras $\mathfrak{h} \subseteq \mathfrak{g}$ are conjugate in the sense that $\mathrm{Aut}(\mathfrak{g})^\circ$ acts transitively on $\mathfrak{g}$.

Proof. Omitted. ■

We are going to get quite a bit of mileage out of a choice of Cartan $\mathfrak{h} \subseteq \mathfrak{g}$ and lift to G_{ad}.

Lemma 2.121. Fix a Cartan subalgebra $\mathfrak{h}$ of a complex semisimple Lie algebra $\mathfrak{g}$. Then $\exp(\mathfrak{h}) \subseteq G_{\mathrm{ad}}$ is a Lie subgroup isomorphic to $(\mathbb{C}^\times)^{\operatorname{rank}\mathfrak{g}}$.

Proof. Let $H \subseteq G$ be the Lie subgroup corresponding to $\mathfrak{h}$. Because $\mathfrak{h}$ is commutative, we see that H is commutative, so $\exp\colon \mathfrak{h} \to H$ is a group homomorphism. Because $\exp$ is also a local diffeomorphism, we conclude that its image contains an open neighborhood of H, so $\exp$ is surjective (because H is connected).

Now, for $X \in \mathfrak{h}$, we see that ad_X acts by 0 on $\mathfrak{h}$ and by $\alpha(X)$ on the root spaces $\mathfrak{g}_\alpha$. This also describes the differential action of $H \subseteq G_{\mathrm{ad}}$ on $\mathfrak{g}$, so we may take the exponential in $G_{\mathrm{ad}} \subseteq \mathrm{GL}(\mathfrak{g})$ to see that $\exp(X)$ must act by 1 on $\mathfrak{h}$ and by $e^{\alpha(X)}$ on $\mathfrak{g}_\alpha$. In particular, the action of H on $\mathfrak{g}$ diagonalizes (via the root space decomposition of $\mathfrak{g}$).

We also see that the kernel of $\exp\colon \mathfrak{h} \to H$ is given by those $X \in \mathfrak{h}$ for which $\alpha(X) \in 2\pi i\mathbb{Z}$ for all roots α, so it follows that H is isomorphic to $\mathfrak{h}/2\pi i P^\vee$. Taking a quotient of a complex vector space by a lattice produces a power of $\mathbb{C}^\times$, so we are done. (Indeed, we merely have to take powers of the isomorphism $\exp\colon \mathbb{C}/2\pi i\mathbb{Z} \to \mathbb{C}^\times$.) ■

Let's use this H for fun and profit.

Proposition 2.122. Fix a Cartan subalgebra $\mathfrak{h}$ of a complex semisimple Lie algebra $\mathfrak{g}$, and set $H := \exp(\mathfrak{h})$ in G_{ad}. Then the normalizer of $N(H)$ of H in G_{ad} coincides with the stabilizer of $\mathfrak{h}$ and contains H as a normal subgroup. In fact, $N(H)/H$ is isomorphic to the Weyl group.

Proof. Set $G := G_{\mathrm{ad}}$ for brevity. Certainly H is normal in $N(H)$ by construction of $N(H)$. Furthermore, recall that G_{ad} acts transitively on the collection of Cartan subalgebras, and $g \in G_{\mathrm{ad}}$ fixes $\mathfrak{h}$ (by adjoint action) if and only if g fixes H (also by adjoint action!) by the uniqueness of the connected Lie subgroups attached to Lie subalgebras. It follows that $N(H)$ is the stabilizer of $\mathfrak{h}$.

Thus, the main content of the proof lies in the last statement. We will now only show the last statement. The general approach will be to first lift the Weyl group to G, and then we will show that H and the lifted Weyl group generate $N(H)$. Let's start with the lifting. For each i, there is a map $\eta_i\colon \mathfrak{sl}_2 \to \mathfrak{g}$ given by sending $e \mapsto e_i$ and $f \mapsto f_i$ and $h \mapsto h_i$. Because $\mathrm{SL}_2(\mathbb{C})$ is simply connected, this lifts (uniquely) to a map $\eta_i\colon \mathrm{SL}_2(\mathbb{C}) \to G_{\mathrm{ad}}$. This η_i let us define

$$w_i := \eta_i\left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}\right).$$

We will use this element as a lift of the simple reflection $s_i \in W$. We can now lift the rest of the Weyl group. Given $w \in W$, we may pick a decomposition $w = s_{i_1} \cdots s_{i_n}$ into simple reflections, and we can define $\widetilde{w} := w_{i_1} \cdots w_{i_n}$.[2] By construction, the adjoint action of w_i on $\mathfrak{h}$ by s_i: indeed, it acts by reflecting α_i's root space.

We emphasize that our lifting is not a group homomorphism: for example, $w_i^2 = \eta_i(-1)$, which may or may not be the identity. On the other hand, note that the differential action of w_i fixes $\mathfrak{h}$, so it lives in $N(H)$. However, it will turn out to be a homomorphism "up to H." Let's make this precise. For any $w, u \in W$, we do know that $\widetilde{wu} = \widetilde{w}\widetilde{u}h$ for some unique $h \in G_{\mathrm{ad}}$; we claim that actually $h \in H$. Well, $\widetilde{wu}$ and $\widetilde{w}\widetilde{u}$ both preserve $\mathfrak{h}$ and even have the same action on the roots, so it follows that h preserves all root spaces. Accordingly, we are granted complex numbers b_j for which $h|_{\mathfrak{g}_{\alpha_j}} = e^{b_j}$. By comparing the action on $\mathfrak{g}$, we conclude that

$$h = e^{\sum_j b_j \omega_j^\vee},$$

which one shows by induction on the height of the root spaces. (Note that the action of h on $\mathfrak{g}$ is enough to know h because $G_{\mathrm{ad}} \subseteq \mathrm{GL}(\mathfrak{g})$.) Thus, $h \in H$.

It follows that the elements $\widetilde{w}_i$ together with H generate a subgroup N of G sitting in a short exact sequence

$$1 \to H \to N \to W \to 1.$$

[2] It turns out that $\widetilde{w}$ does not depend on the choice of decomposition of w, but we will not need this.

It remains to show that $N = N(H)$, which will complete the proof. Well, choose $g \in N(H)$, which we would like to move to N. But then $\{g(\alpha_i)\}_i$ is a collection of positive roots, but all systems of positive roots are conjugate by the Weyl group, so we can adjust g by an element $\widetilde{w} \in N$ so that g acts by a permutation of the simple roots.

We are now in the situation where g acts by an automorphism on the Dynkin diagram. If we could show that this automorphism is trivial, then it would follow that g merely scales all the root spaces (with no permutation), so we are able to conclude that $g \in H$ using the argument above. However, we claim that all of G_{ad} acts trivially on the Dynkin diagram (because the automorphism group of the Dynkin diagram is discrete). It is enough to check that G_{ad} acts trivially on the fundamental representations of $\mathfrak{g}$: indeed, G_{ad} permutes the Verma modules according to its action on $\mathfrak{h}$, which then has quotient to the needed fundamental representations. But G_{ad} needs to act continuously on the characters, which distinguish the fundamental representations, so we conclude that G_{ad} acts trivially on the characters and thus the fundamental representations. ■

Remark 2.123. The extension

$$1 \to H \to N(H) \to W \to 1$$

does not always split. Indeed, this does not split for $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$: one sees that $N(H)$ is generated by the central torus of diagonal elements $\mathrm{diag}(t, 1/t)$ and the element $w = \left[\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right]$, which satisfies $wtw^{-1} = t^{-1}$. Notably, $(wt)^2 = -1$ for any t, but this is not possible for elements in $W \rtimes H$.

We can upgrade this discussion to compute $\mathrm{Aut}(\mathfrak{g})$ in full.

Proposition 2.124. Let $\mathfrak{g}$ be a complex semisimple Lie algebra with Dynkin diagram D. Then there is a split short exact sequence

$$1 \to G_{\mathrm{ad}} \to \mathrm{Aut}(\mathfrak{g}) \to \mathrm{Aut}(D) \to 1.$$

Proof. To begin, note that the automorphisms of the Dynkin diagram embed into $\mathrm{Aut}(\mathfrak{g})$ by using the Serre presentation of $\mathfrak{g}$. Accordingly, there is a map

$$\xi\colon \mathrm{Aut}(D) \ltimes G_{\mathrm{ad}} \to \mathrm{Aut}(\mathfrak{g}).$$

Indeed, certainly G_{ad} is normal in $\mathrm{Aut}(\mathfrak{g})$ because it is the connected component. Also, we may quickly check that ξ is injective: certainly ξ is injective on $\mathrm{Aut}(D)$. Thus, if we have a nontrivial element $x \in \ker \xi$, the image of x in $\mathrm{Aut}(D)$ is not the identity. But then x permutes the fundamental representations nontrivially, which does not happen for elements of G_{ad}.

It remains to show that ξ is surjective. Well, choose $g \in \mathrm{Aut}(\mathfrak{g})$. Because any two Cartan subalgebras are conjugate by G_{ad}, we may assume that g fixes $\mathfrak{h}$. We now redo our previous argument: the collection $\{g(\alpha_i)\}$ is a new collection of simple roots, so there is an automorphism σ of the Dynkin diagram with $g(\alpha_i) = \alpha_{\sigma(i)}$. Adjusting g by σ^{-1}, we may further assume that g fixes the simple roots as well. We conclude that $g \in H$ using the argument from before. ■

Theme 3
Structure Theory

3.1 March 30

Here we go.

3.1.1 Forms of Semisimple Lie Algebras

We will work over a field K of characteristic zero. Recall that we already have a construction of many semisimple Lie algebras.

Definition 3.1 (split form)**.** Fix a field K and a Dynkin diagram D. Then the Serre presentation defines a finite-dimensional Lie algebra $\mathfrak{g}_D$ with Dynkin diagram D over K. We call this the *split form*.

Example 3.2. Over algebraically closed fields, we showed that all Lie algebras with Dynkin diagram D are split forms.

Example 3.3. There are also non-split forms: for example, $\mathfrak{su}_2$ is a real Lie algebra with the same Dynkin diagram as $\mathfrak{sl}_2$, but these are not isomorphic real Lie algebras.

Anyway, the point is that we may classify Lie algebras $\mathfrak{a}$ over a field K of characteristic zero by classifying the Lie algebras $\mathfrak{a} \otimes_K \overline{K}$ over $\overline{K}$ (which is split!) and then classifying the Lie algebras that base-change to the split form. It will be useful to have some language for this situation.

Definition 3.4 (form)**.** Fix a Galois extension L/K of fields, and choose a Dynkin diagram D. Then a Lie algebra $\mathfrak{g}$ over K *splits* over L if and only if $\mathfrak{g}_L$ is isomorphic to the split form. We say that $\mathfrak{g}$ is a *form* of $\mathfrak{g}_D$.

Remark 3.5. Because all forms split over $\overline{K}$, one can show that all forms split over a finite Galois extension.

It will turn out that the Lie algebras $\mathfrak{a}$ over K with $\mathfrak{a} \otimes_K L = \mathfrak{g}_D$ are given by $\mathrm{H}^1(\mathrm{Gal}(L/K); \mathrm{Aut}\, \mathfrak{g}_D)$. Let's recall what H^1 means for our purposes.

Definition 3.6. Fix a group Γ acting by automorphisms on another group G. Then we define $\mathrm{H}^1(\Gamma; G)$ to classify homomorphisms $\Gamma \to \Gamma \rtimes G$ for which the composite back down to Γ is the identity, up to suitable conjugacy.

Remark 3.7. Let's explain how 1-cocycles enter the picture. Such a homomorphism $\varphi\colon \Gamma \to \Gamma \rtimes G$ necessarily has $\varphi(\gamma) = (\eta(\gamma), \gamma)$, where $\eta\colon \Gamma \to G$ is a map which now must satisfy

$$(\eta(\gamma_1\gamma_2), \gamma_1\gamma_2) = (\eta(\gamma_1), \gamma_1)(\eta(\gamma_2), \gamma_2),$$

which is equivalent to $\eta(\gamma_1\gamma_2) = \eta(\gamma_1) \cdot \gamma_1(\eta(\gamma_2))$.

Remark 3.8. Let's explain how 1-coboundaries enter the picture. Well, we are conjugating a map $\gamma \mapsto (\eta(\gamma), \gamma)$ by an element $g \in G$, which corresponds to the map $\gamma \mapsto \big(g\eta(\gamma)\gamma(g)^{-1}, \gamma\big)$. Thus, conjugacy of homomorphisms corresponds to two 1-cocycles being the same up to 1-coboundary.

Let's explain how to relate forms to cohomology. The idea is to note that the Galois group acts on the base-change.

Definition 3.9 (twisted linear)**.** Fix a Galois extension L/K of fields, and let Γ be the Galois group. Then a *twisted linear action* of Γ on a vector space V over L is one which is additive, and

$$\gamma(\ell x) = \gamma(\ell)\gamma(x)$$

for any $\gamma \in \Gamma$ and $\ell \in L$ and $x \in V$.

Example 3.10. For any vector space V over K, there is a trivial twisted linear action on $V \otimes_K L$ given by having Γ act directly on the L-component. By computing a basis, one finds that $V = (V \otimes_K L)^\Gamma$.

Proposition 3.11. Fix a Galois extension L/K of fields, and choose a Dynkin diagram D. Then the pointed set $\mathrm{H}^1(\mathrm{Gal}(L/K); \mathrm{Aut}\,\mathfrak{g}_D)$ classifies Lie algebras $\mathfrak{g}$ over K with Dynkin diagram D which split over L.

Proof. Set $\Gamma := \mathrm{Gal}(L/K)$. For any Lie algebra $\mathfrak{g}$ over K, we see that Γ acts on $\mathfrak{g}_L$ via a twisted linear action, and then one can recover $\mathfrak{g}$ from $\mathfrak{g}_L$ by taking invariants. Conversely, any twisted linear action of Γ on a Lie algebra $\mathfrak{g}_L$ over L can take Γ-invariants to recover a Lie algebra over K.

Now, fix a Lie algebra $\mathfrak{g} := \mathfrak{g}_D$ over K, and let ρ be a twisted linear action of Γ on $\mathfrak{g}_L$. Further, let ρ_0 be the trivial twisted linear action. Then there is $\eta\colon \Gamma \to \mathrm{Aut}(\mathfrak{g})$ for which $\rho(g) = \eta(g)\rho_0(g)$ for all $g \in \Gamma$. Note that ρ is a homomorphism if and only if $\rho(gh) = \rho(g)\rho(h)$ for all g and h, which we can see to be equivalent to having

$$\eta(gh) = \eta(g) \circ g\eta(h),$$

where $g\eta(h) := \rho_0(g)\eta(h)\rho_0(g)^{-1}$. Thus, η is a 1-cocycle in $\mathrm{Z}^1(\Gamma; \mathrm{Aut}(\mathfrak{g}))$.

Conversely, η_1 and η_2 define isomorphic ρ_1 and ρ_2 if and only if there is $a \in \mathrm{Aut}(\mathfrak{g})$ for which $\rho_1(g) \circ a = a \circ \rho_2(g)$ for all g. One can expand that this is equivalent to having $\eta_1(g) = a\eta_2(g)ga^{-1}$ for all g, which is equivalent to η_1 and η_2 being cohomologous. In total, we have obtained a bijection from isomorphism classes of Lie algebras $\mathfrak{g}'$ over D with $\mathfrak{g}'_L = \mathfrak{g}_L$ and $\mathrm{H}^1(\Gamma; \mathrm{Aut}\,\mathfrak{g})$, as required. ∎

Remark 3.12. The same argument works for any linear algebraic structure. Namely, one can replace Lie algebras with vector spaces, quadratic spaces, and so on.

3.1.2 Real Forms

In the sequel, we will focus on the Galois extension $\mathbb{C}/\mathbb{R}$. Let's work out Proposition 3.11 in this case.

Example 3.13. Consider the Galois extension $\mathbb{C}/\mathbb{R}$. In this case, Proposition 3.11 (combined with Proposition 2.124) tells us that real forms of a complex semisimple Lie algebra $\mathfrak{g}_D$ are given by

$$\mathrm{H}^1(\{\pm 1\}; \mathrm{Aut}(D) \ltimes G_{\mathrm{ad}}),$$

where $-1 \in \{\pm 1\}$ acts by complex conjugation. Notably, a 1-cocycle $\eta\colon \{\pm 1\} \to \mathrm{Aut}(\mathfrak{g}_D)$ must have $\eta(1) = \mathrm{id}$, so we only have to worry about the data of $s := \eta(-1)$. The cocycle condition amounts to requiring $s\overline{s} = 1$. Further, note that we only care about η up to conjugation, which means that we are only interested in s up to the conjugation $s \mapsto as\overline{a}^{-1}$, where $a \in \mathrm{Aut}(\mathfrak{g}_D)$. Thus, real forms of G are in bijection with $s \in \mathrm{Aut}(D) \ltimes G_{\mathrm{ad}}$ for which $s\overline{s} = 1$, considered up to twisted conjugation.

Notation 3.14. We work over $\mathbb{R}$. Fix a Dynkin diagram D. For $s \in \mathrm{Aut}(\mathfrak{g}_{D,\mathbb{C}})$ for which $s\overline{s} = 1$, we let $\mathfrak{g}_s$ denote the corresponding real form.

This s is a $\mathbb{C}$-linear transformation, but it will turn out to be convenient to instead consider "antilinear" transformations.

Definition 3.15 (antilinear)**.** Fix a complex vector space V. Then an $\mathbb{R}$-linear map $\sigma\colon V \to V$ is *antilinear* if and only if $\sigma(zv) = \overline{z}\sigma(v)$ for $z \in \mathbb{C}$.

Example 3.16. Suppose that V is a real vector space. Then a linear map $s\colon V_{\mathbb{R}} \to V_{\mathbb{R}}$ induces an antilinear map $\sigma_s\colon V_{\mathbb{C}} \to V_{\mathbb{C}}$ given by composing with complex conjugation.

Remark 3.17. By definition, we have that $\mathfrak{g}_s$ consists of those $X \in \mathfrak{g}_{D,\mathbb{C}}$ for which $\overline{X} = s(X)$, which is equivalent to $X \in \mathfrak{g}^{\sigma_s}$.

Let's now compute some real forms. Note that complex conjugation acts trivially on the Dynkin diagram itself, so our $s \in \mathrm{Aut}(\mathfrak{g}_{D,\mathbb{C}})$ induces an automorphism $s_0 \in \mathrm{Aut}(D)$ with $s_0^2 = 1$. It is not too hard to classify such involutions of the Dynkin diagram. Thus, a semisimple Lie algebra $\mathfrak{g}$ over $\mathbb{R}$ will have $\mathfrak{g}_{\mathbb{C}}$ be given by some Dynkin diagram D with an automorphism s_0; by organizing the Dynkin diagram into s_0-invariant pieces, we see that $\mathfrak{g}$ is a sum of simple Lie algebras.

Further, if $\mathfrak{g}$ is actually simple, then there are two cases.

Example 3.18. It is possible for $\mathfrak{g}_{\mathbb{C}}$ to consists of two copies of a connected Dynkin diagram D_0, meaning that $D = D_0 \sqcup D_0$, and s_0 swaps the two pieces. Thus, we see that $G_{\mathrm{ad}} = K_{\mathrm{ad}} \times K_{\mathrm{ad}}$, where s_0 swaps the two factors. Let's now write out the condition $s\overline{s} = 1$. We know that $s = s_0(g,h)$ where $g, h \in K_{\mathrm{ad}}$. Now, $s\overline{s} = 1$, we see $s_0(g,h) = s_0(\overline{g}\overline{h}) = 1$, which means that $h = \overline{g}^{-1}$. But then $s = s_0\left(g, \overline{g}^{-1}\right) = (1,g)s_0\overline{(1,g)}^{-1}$, so s is cohomologous with s_0!

Thus, we see that we only have a single real form here. Taking s_0-invariants, we see that our Lie algebra $\mathfrak{g}_{s_0}$ is simply given by $\mathfrak{k} = \mathrm{Lie}\, K$. Explicitly, we have taken some simple complex Lie algebra $\mathfrak{k}$ and forgotten the complex structure to produce a simple real Lie algebra.

Alternatively, we could have $\mathfrak{g}_{\mathbb{C}}$ be a simple complex Lie algebra, and s_0 acts nontrivially on the Dynkin diagram. This turns out to require some interesting combinatorics.

3.1.3 Inner Forms

Let's start by organizing our forms by their image in $\mathrm{Aut}\, D$.

Definition 3.19 (inner form)**.** Fix a Dynkin diagram D and a field K. A form $\mathfrak{g}_s$ with Dynkin diagram D is *inner* if and only if the image of s in $\mathrm{Aut}(D)$ is trivial.

Remark 3.20. It is equivalent to ask for s to be in G_{ad}.

Definition 3.21 (inner class)**.** Fix a Dynkin diagram D and a field K. Two forms $\mathfrak{g}_s$ and $\mathfrak{g}_{s'}$ with Dynkin diagram D are in the same *inner class* if and only if s and s' have the same image in $\mathrm{Aut}(D)$.

Remark 3.22. Thus, the inner forms make up the collection of those forms in the same inner class as the split form.

Each inner class admits a distinguished element.

Definition 3.23 (quasisplit)**.** Fix a Dynkin diagram D and a field K. A form $\mathfrak{g}_s$ is *quasisplit* if and only if the image of s in G_{ad} is trivial (up to conjuacy).

Remark 3.24. Each inner class consists of elements of the form $(s_0, g) \in \mathrm{Aut}(\mathfrak{g}) \rtimes G_{\mathrm{ad}}$, where s_0 is fixed. Thus, each inner class admits a unique quasisplit form.

The above discussion works for general fields.

3.1.4 The Compact Form

We now define a special involution of complex Lie algebras.

Definition 3.25 (Cartan involution)**.** Fix a semisimple Lie algebra $\mathfrak{g}$ over $\mathbb{C}$. Then we define the *Cartan involution* $\tau\colon \mathfrak{g} \to \mathfrak{g}$ to be given on the Serre presentation

$$\begin{cases} \tau(h_j) := -h_j, \\ \tau(e_j) := -f_j, \\ \tau(f_j) := -e_j \end{cases}$$

for each j.

Remark 3.26. We will not bother to check that τ preserves the Serre relations, which holds by their symmetry. Note that $\tau\overline{\tau} = 1$.

With an involution, we get a real form.

Definition 3.27 (compact form)**.** Fix a semisimple Lie algebra $\mathfrak{g}$ over $\mathbb{C}$. Then the *compact real form* $\mathfrak{g}^c$ is the real form $\mathfrak{g}_\tau$.

It will turn out that $\mathfrak{g}^c$ corresponds to a compact group, which explains the language. We will prove this shortly, but let's start with an example.

Example 3.28. Consider $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$. Then $\tau(h) = -h$ and $\tau(e) = -f$ and $\tau(f) = -e$, so $\mathfrak{g}^c = \{X \in \mathfrak{g} : \overline{X} = \tau(X)\}$ admits a basis over $\mathbb{R}$ given by the matrices

$$\left\{ \begin{bmatrix} i & \\ & -i \end{bmatrix}, \begin{bmatrix} & 1 \\ -1 & \end{bmatrix}, \begin{bmatrix} & i \\ i & \end{bmatrix} \right\}.$$

Thus, we see that $\mathfrak{sl}_2(\mathbb{C})^c = \mathfrak{su}(2)$.

Theorem 3.29. Fix a complex semisimple Lie algebra $\mathfrak{g}$. Then the Killing form of $\mathfrak{g}^c$ is negative definite.

Proof. The root decomposition allows us to write

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Phi_+} (\mathbb{C}e_\alpha \oplus \mathbb{C}f_\alpha).$$

Now, τ acts on each of the above summands, so we find that

$$\mathfrak{g}^c = \mathfrak{h}^c \oplus \bigoplus_{\alpha \in \Phi_+} (\mathbb{C}e_\alpha \oplus \mathbb{C}f_\alpha)^c.$$

Now, this is an orthogonal decomposition for the Killing form (indeed, this is already true for the root decomposition), so it is enough to show the negative-definiteness on each summand.

- We show that the Killing form is negative-definite on $\mathfrak{h}^c$. Here, τ acts by -1 on $\mathfrak{h}$, we see that $\mathfrak{h}^c$ admits a basis given by the $ih_\bullet$s. On this basis, we choose some $i\sum_j t_j h_j \in \mathfrak{h}^c$, and then we can compute the Killing form by
$$\operatorname{tr}_{\mathfrak{g}} \left(i \sum_j t_j h_j \right)^2 = -\operatorname{tr}_{\mathfrak{g}} \left(\sum_j t_j h_j \right)^2.$$
Now, via the root decomposition, we may compute this trace as $-\sum_\alpha \left(\sum_j t_j \alpha(h_j) \right)^2$, which is negative.

- We show that the Killing form is negative-definite on $(\mathbb{C}e_\alpha \oplus \mathbb{C}f_\alpha)^c$ when α is a simple root α_j. Then by definition of τ, we see that $\mathfrak{g}^c \cap \mathfrak{sl}_{2j}$ is isomorphic to $\mathfrak{su}(2)$, as we just computed in Example 3.28. Here, the Killing form takes the form
$$K(X, Y) = \operatorname{tr}_{\mathfrak{g}}(\rho_{\mathfrak{g}}(X), \rho_{\mathfrak{g}}(Y))$$
for $X, Y \in \mathfrak{su}(2)$. But all representations of $\mathfrak{su}(2)$ lift to the compact group $\mathrm{SU}(2)$ and thus are unitary, so $\rho_{\mathfrak{g}}$ is skew-Hermitian and hence have imaginary eigenvalues, so $\operatorname{tr} \rho_{\mathfrak{g}}(X)^2 < 0$ for nonzero X, so the Killing form is negative-definite!

- It remains to handle the general case for the spaces $(\mathbb{C}e_\alpha \oplus \mathbb{C}f_\alpha)^c$. Well, recall the construction of w_j from Proposition 2.122. Note that $\left[\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right]$ lives in $\mathrm{SU}(2)$. Now, any element G_{ad} preserves the Killing form on its Lie algebra $\mathfrak{g}$, so w_j preserves the Killing form of $\mathfrak{g}^c$. It follows that any $\widetilde{w}$ preserves the Killing form. The point is that any root α can be permuted with W to be simple, and we now know that the Weyl group action preserves the Killing form, so we reduce to the previous case! ■

We may now explain how $\mathfrak{g}^c$ corresponds to a compact real group.

Definition 3.30. Fix a complex semisimple Lie algebra $\mathfrak{g}$. Then we define G_{ad}^c to be the real Lie subgroup of G_{ad} corresponding to the Lie algebra $\mathfrak{g}^c$.

Remark 3.31. By construction, $G_{\mathrm{ad}}^c = \{X \in G_{\mathrm{ad}} : \tau(X) = \overline{X}\}^\circ$, where the action of τ is by conjugation. Indeed, one shows this by taking the differential of the relation $\tau(X) = \overline{X}$. In fact, $\{X \in G_{\mathrm{ad}} : \tau(X) = \overline{X}\}$ is connected, which we will show later.

Remark 3.32. Note that G_{ad}^c acts by negative-definite automorphisms on $\mathfrak{g}$, so G_{ad}^c embeds into an orthogonal group, so it is compact.

Remark 3.33. Later, we will show that the universal cover of G_{ad}^c continues to be compact. Thus, its representation theory is semisimple by Corollary 2.34, so the representation theory of $\mathfrak{g}^c$ is semisimple, so the representation theory of $\mathfrak{g}$ is semisimple!

One can compute this for the classical types without too much effort. We will just give the answers.

Example 3.34. If $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$, then G_{ad}^c turns out to be $\mathrm{PSU}(n)$.

Example 3.35. If $\mathfrak{g} = \mathfrak{so}_n(\mathbb{C})$, then G_{ad}^c is $\mathrm{SO}_n(\mathbb{C})$ if n is odd, and it is $\mathrm{SO}_n(\mathbb{C})/\{\pm 1\}$ if n is even.

Example 3.36. Lastly, if $\mathfrak{g} = \mathfrak{sp}_{2n}(\mathbb{C})$, then G_{ad}^c is $(\mathrm{Sp}_{2n}(\mathbb{C}) \cap \mathrm{U}(2n))/\{\pm 1\}$. Alternatively, one can describe these as the unitary $n \times n$ matrices with coefficients in the quaternions.

Remark 3.37. The compact form is not quasisplit. Indeed, the quasisplit real form has nilpotent elements (it comes from an automorphism of the Dynkin diagram, so it has the element e from the principal $\mathfrak{sl}_2$-triple), so the corresponding real Lie group is not compact.

3.2 April 1

Here we go.

3.2.1 Real Forms of Classical Types

Let's list out some real forms we know.

Exercise 3.38. We list some real forms of $\mathfrak{sl}_n(\mathbb{C})$.

Proof. Here are our forms.

- We have the split real form $\mathfrak{sl}_n(\mathbb{R})$
- One can directly compute that the compact real form is $\mathfrak{su}_n(\mathbb{R})$: one can calculate that τ is given by the Lie algebra involution $\sigma_\tau(X) = -X^\dagger$.
- Because $\operatorname{Aut} A_{n-1}$ is $\mathbb{Z}/2\mathbb{Z}$ (for $n \geq 3$, given by the "flip" of the Dynkin diagram), there is a unique non-inner quasisplit real form. It should be given by the s_0 which sends $e_j \mapsto e_{n-j}$ and $f_j \mapsto f_{n-j}$ and $h_j \mapsto h_{n-j}$. One can check that this automorphism is actually given by $s_0(A) := -JA^\intercal J^{-1}$, where J is the antidiagonal matrix

$$J = \begin{bmatrix} & & & & & -1 \\ & & & & 1 & \\ & & & -1 & & \\ & & 1 & & & \\ & \iddots & & & & \end{bmatrix}.$$

Thus, this quasisplit form is $\mathfrak{su}(p,p)$ if $n = 2p$, and it is $\mathfrak{su}(p,p+1)$ if $n = 2p+1$. Note that comparing τ and s_0 shows that the compact form and this quasisplit form are in the same inner class!

- In general, one has the real forms $\mathfrak{su}(p,q)$ where $n = p+q$. ■

Remark 3.39. For $n = 2$, one has that $\mathfrak{su}(1,1) = \mathfrak{sl}(2)$ because the Dynkin diagram has no nontrivial automorphisms.

Exercise 3.40. We list some real forms of $\mathfrak{so}(2n+1,\mathbb{C})$.

Proof. Here are our forms. Note that all forms are inner because there are no nontrivial automorphism of the Dynkin diagram.

- One has the compact form $\mathfrak{so}(2n+1)$.
- There is also the split form $\mathfrak{so}(n,n+1)$, which one sees from the Serre presentation (or alternatively, the standard form of $\mathfrak{so}(2n+1)$ over $\mathbb{C}$).
- In general, there are forms $\mathfrak{so}(p,q)$ when $p+q = 2n+1$.

Note that when $n = 1$, we have $\mathfrak{so}(1,2) = \mathfrak{sl}(2,\mathbb{R})$. ■

Exercise 3.41. We list some real forms of $\mathfrak{sp}(2n,\mathbb{C})$.

Proof. Note that all forms are inner because the Dynkin diagram has no nontrivial automorphisms. There is the compact form $\mathfrak{su}(n,\mathbb{H})$ and the split form $\mathfrak{sp}(2n,\mathbb{R})$. Note that when $n = 2$, this should agree with $\mathfrak{so}(2n+1)$, so we note that $\mathfrak{su}(2,\mathbb{H}) = \mathfrak{so}(5)$ and $\mathfrak{sp}(4,\mathbb{R}) = \mathfrak{so}(3,2)$ due to the uniqueness of the split and compact forms. ■

Exercise 3.42. We list some real forms of $\mathfrak{so}(2n,\mathbb{C})$.

Proof. There is a compact form. When $n \geq 3$, there are two automorphisms of the Dynkin diagram; even when $n = 2$, there are only two conjugacy classes of automorphisms. Thus, there is a compact form $\mathfrak{so}(2n)$, a split form $\mathfrak{so}(n,n)$ (as in type B), and there is a unique non-inner quasisplit form. By noting that the nontrivial conjugacy class may as well be given by $e_n \mapsto -e_n$ and $e_i \mapsto e_i$ for $i < n$, we may take the quadratic form of the quasisplit form to come from the symmetric matrix

$$\begin{bmatrix} 0 & 0 & 1_{n-1} & 0 \\ 0 & 1 & 0 & 0 \\ 1_{n-1} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

which has signature $(n+1, n-1)$. ■

Example 3.43. For D_2, one can see that the quasisplit inner form recovers $\mathfrak{sl}(2,\mathbb{C})$ because $D_2 = A_2$. Thus, $\mathfrak{sl}(2,\mathbb{C}) \cong \mathfrak{so}(3,1)$ (as real Lie algebras). Note that the corresponding real points of the group $\mathrm{SO}(3,1)$ has two connected components! Indeed, $\mathrm{SO}(3,1)$ acts naturally on the cone cut out by $t^2 - x_1^2 - x_2^2 - x_3^2 = 0$, and there is a real Lie subgroup preserving the points with $t > 0$.

Example 3.44. For $n = 3$, one has $D_3 = A_3$, and one finds that $\mathfrak{so}(6) \cong \mathfrak{su}(4)$ and $\mathfrak{so}(3,3) \cong \mathfrak{sl}(4,\mathbb{R})$ and $\mathfrak{so}(4,2) \cong \mathfrak{su}(2,2)$, again by matching up compact, slit, and quasisplit forms.

One can say something similar for exceptional types, but we will return to them later.

3.2.2 Counting from the Compact Form

For real groups, it turns out to be more convenient to use Galois cohomology to count "from" the compact form instead of the split form.

Proposition 3.45. Fix a complex semisimple Lie algebra $\mathfrak{g}$. Then the real forms of $\mathfrak{g}$ are in bijection with conjugacy classes of involutions in $\operatorname{Aut}\mathfrak{g}^c$.

Proof. Here, we start with the antilinear Cartan involution ω coming from the compact Cartan involution. We now run the same argument as in Proposition 3.11. Now, another real structure on $\mathfrak{g}$ can be described by an antilinear involution, which we may take to have the form $\sigma := \omega \circ g$, where $g \in \operatorname{Aut}(\mathfrak{g})$ is chosen with $\omega(g)g = 1$.

To continue, we claim that

$$\omega(x) \stackrel{?}{=} x^{-\dagger},$$

where $(\cdot)^\dagger$ is induced from the Killing form on $\mathfrak{g}^c$, extended to a Hermitian form. Indeed, to show the claim, it is enough to show that $\omega(X) = -X^\dagger$ on the Lie algebra; by splitting up into Hermitian and skew-Hermitian parts, it is enough to note that $\omega(X) = X$ if X is unitary for the Hermitian form on $\mathfrak{g}$ coming from the Killing form on $\mathfrak{g}^c$ (because ω is negative-definite for the Killing form on $\mathfrak{g}^c$).

The moral is that $\omega(g)g = 1$ implies that

$$g^\dagger g = 1,$$

so g is self-adjoint, so it is self-adjoint with real eigenvalues. Thus, we may decompose

$$\mathfrak{g} = \bigoplus_{\gamma \in \mathbb{R}} \mathfrak{g}[\gamma],$$

where $\mathfrak{g}[\gamma]$ is the γ-eigenspace of $\mathfrak{g}$. Because g is an automorphism of $\mathfrak{g}$, we see that $[\mathfrak{g}[\gamma], \mathfrak{g}[\gamma']] = \mathfrak{g}[\gamma\gamma']$.

We now modify g for convenience. For any t, we define $|g|^t$ to act by $|\gamma|^t$ on $\mathfrak{g}[\gamma]$; note that $|g|^t$ continues to be an automorphism of $\mathfrak{g}$, which one can see by an eigenvalue decomposition. Now, define $\theta := g\,|g|^{-1}$, so it follows that $\theta^2 = 1$ by looking at eigenspace actions. Because $g\omega = \omega g^{-1}$ from earlier, we see that $\theta\omega = \omega\theta^{-1} = \omega\theta$. Additionally, $\theta = |g|^{-1/2}\, g\, |g|^{1/2}$, so θ and g are conjugate, so they define the same real form.

The previous paragraph allows us to replace g with θ, which now squares to 1 and commutes with ω; commuting with ω is enough to view θ as an automorphism of $\mathfrak{g}^c$, so we have at least described how to take a real form and produce some θ. For such θ, we see that another θ' gives the same real form when $\theta' = x\theta\omega(x)^{-1}$ for some automorphism x. To this end, we write $z := \omega(x)^{-1}x$, which is $x^\dagger x$ and thus positive-definite. By construction, we see that $\omega(z) = z^{-1}$ because ω is antilinear with $\omega^2 = 1$, and the construction gives $\theta z = z^{-1}\theta$.

Accordingly, we normalize x with $y := xz^{-1/2}$ so that $\omega(y) = \omega(x)z^{1/2} = xz^{-1/2} = y$. Similarly, we may calculate

$$\begin{aligned}
\theta' &= x\theta\omega(x)^{-1} \\
&= x\theta z x^{-1} \\
&= xz^{-1/2}\theta z^{1/2}x^{-1} \\
&= y\theta y^{-1},
\end{aligned}$$

so θ is actually conjugate by this nice element y, which is ω-invariant! ■

Remark 3.46. Given some θ, the proof shows that our involution for Proposition 3.11 is given by $\omega_\theta := \omega \circ \theta$.

Remark 3.47. Now, given θ, one can split $\mathfrak{g}$ into the complex subspaces $\mathfrak{k} \oplus \mathfrak{p}$, which are the $(+1)$- and (-1)-eigenspaces, respectively. We can see that $\mathfrak{k}$ is a Lie subalgebra, and the definiteness of the Killing form implies that $\mathfrak{k}$ is reductive by Cartan's criterion. Now, this decomposition commutes with ω, so we may write

$$\mathfrak{g}^c = \mathfrak{k}^c \oplus \mathfrak{p}^c.$$

The corresponding real form $\mathfrak{g}_\theta$ can thus be seen to be the ω_θ-conjugate-invariants. Thus, this consists of the invariants in $\mathfrak{k}^c$ (because here θ acts by 1) and consists of $i\mathfrak{p}^c$ (because here θ acts by -1), so $\mathfrak{g}_\theta = \mathfrak{k}^c \oplus i\mathfrak{p}^c$ by gluing these together.

3.2.3 Vogan Diagrams

We will want to use Proposition 3.45 to classify real forms by Dynkin diagrams. Here is a quick coherence check, which allows us to do a root decomposition of $\mathfrak{k}$ and $\mathfrak{g}$ simultaneously.

Proposition 3.48. Fix a real form $\mathfrak{g}_\theta$ of a complex semisimple Lie algebra $\mathfrak{g}$, and set $\mathfrak{k} := \mathfrak{g}^\theta$. Then there is a θ-stable Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ for which $\mathfrak{h} \cap \mathfrak{k}$ is a Cartan subalgebra of $\mathfrak{k}$.

Proof. Fix everything as in Remark 3.47. Choose a generic $t \in \mathfrak{k}^c$, which is known to be regular semisimple, and we let $\mathfrak{h}_+^c$ be its centralizer in $\mathfrak{k}^c$. The complexification is called $\mathfrak{h}_+$, and because t is regular, we see that $\mathfrak{h}_+ \subseteq \mathfrak{h}$ is a Cartan subalgebra of $\mathfrak{k}$.

We now extend $\mathfrak{h}_+$ to $\mathfrak{g}$. Indeed, let $\mathfrak{h}_-^c$ be a maximal subspace of $\mathfrak{p}^c$ so that

$$\mathfrak{h}^c := \mathfrak{h}_+^c \oplus \mathfrak{h}_-^c$$

is an abelian subalgebra of $\mathfrak{g}^c$. We claim that the complexification $\mathfrak{h} \subseteq \mathfrak{g}$ is a Cartan subalgebra, which will complete the proof by construction (note that $\mathfrak{h} \cap \mathfrak{k} = \mathfrak{h}_+$ because $\mathfrak{p} \cap \mathfrak{k} = 0$). Here are our checks.

- Semisimple: note that $\mathfrak{h}^c$ consists of semisimple elements because these elements are anti-Hermitian operators on $\mathfrak{g}^c$.
- Maximal: suppose that z commutes with $\mathfrak{h}$. Then the decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ allows us to decompose $z = z_+ + z_-$. Further, because θ commutes with ω, we see that z_+ must commute with $\mathfrak{h}$, so $z_+ \in \mathfrak{h}_+$ automatically. Thus, we may assume that $z_+ = 0$. Then we write $z_- = x + iy$ where $x, y \in \mathfrak{p}^c$, and both x and y continue to commute with $\mathfrak{h}$ by taking real and complex parts, so $x, y \in \mathfrak{h}_-$ by maximality. Thus, $z_- \in \mathfrak{h}_-$ as well. ■

Remark 3.49. With the above construction, we further claim that $\mathfrak{h}_-$ does not have any coroots of $\mathfrak{g}$. Indeed, suppose that we have a coroot $\alpha^\vee \in \mathfrak{h}_-$. This means that $\theta(\alpha^\vee) = -\alpha^\vee$ because $\mathfrak{h}_- \subseteq \mathfrak{p}$, so $\theta(e_\alpha) = e_{-\alpha}$ and $\theta(e_{-\alpha}) = e_\alpha$ follows. Now, set $x := e_\alpha + e_{-\alpha}$, which we see is θ-stable, so $x \in \mathfrak{k}$. Now, on one hand, $[\mathfrak{h}_+, x] = 0$ because α vanishes on $\mathfrak{h}_+$, but $\mathfrak{h}_+ \subseteq \mathfrak{k}$ is a maximal abelian subalgebra, so it must contain x! On the other hand, the root decomposition means that x is orthogonal to $\mathfrak{h}_+$, so $x = 0$, which is a contradiction.

We now do our root decompositions. Fix a Cartan as in Proposition 3.48. Then we may choose a generic $t \in \mathfrak{h}_+$, and we see that t is regular in $\mathfrak{g}$ because $\mathfrak{h}_+$ has all the roots. Additionally, generic t will have $\operatorname{Re}(t, \alpha^\vee) \neq 0$ for all coroots $\alpha^\vee$, which we use to produce a polarization. By construction, we finally see that θ preserves the set of positive roots (because t is θ-invariant!), so θ preserves the set of simple roots, so θ induces an automorphism of the Dynkin diagram. Let $\{\alpha_i\}$ be the collection of simple roots. There are two cases.

- If $\theta(i) = i$, then $\theta(e_i) = \pm e_i$ and $\theta(f_i) = \pm f_i$ and $\theta(h_i) = \pm h_i$. Indeed, this is required because θ preserves this root space, and $\theta^2 = 1$.

- Otherwise, we can have $\theta(i) \neq i$, but then $\theta(h_i) = h_{\theta(i)}$, and we can normalize everything so that $\theta(e_i) = e_{\theta(i)}$ and $\theta(f_i) = f_{\theta(i)}$.

The moral is that this choice of θ and t can be labeled by a Vogan diagram, which is a decoration of the Dynkin diagram as follows: we choose an automorphism θ of the Dynkin diagram, and we must color the θ-invariant vertices either white or black (corresponding to the sign $+$ or $-$ in the equation $\theta(e_i) = \pm e_i$, respectively).

Remark 3.50. We have shown that every real form $\mathfrak{g}_\theta$ is labeled by at least one Vogan diagram coming from θ and a choice of polarization. But more than one Vogan diagram may produce the same real form!

Example 3.51. For A_1, there are only two Vogan diagrams: we have a single white dot, which is $\mathfrak{sl}(2, \mathbb{R})$; and we have a single black dot, which is $\mathfrak{su}(2)$.

3.2.4 Real Forms of Type A

Let's use Proposition 3.45 for fun and profit.

Exercise 3.52. We classify the real forms of type A_{n-1}.

Proof. Here, $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$, so $\mathfrak{g}^c = \mathfrak{su}(n)$, which has automorphism group $\mathbb{Z}/2\mathbb{Z} \times \mathrm{PU}(n)$; namely, these are the automorphisms of $\mathfrak{g}$ which preserve $\mathfrak{g}^c$. There are thus two cases.

- We have a compact inner class, which comes from those $\theta \in \mathrm{PU}(n)$. Up to conjugacy, we may move such matrices to have p total $(+1)$s and q total (-1)s, where p and q are nonnegative integers with $n = p + q$. We can now directly calculate that $\mathfrak{g}_\theta = \mathfrak{su}(p, q)$. It turns out that $\mathfrak{k} = \mathfrak{sl}(p) \oplus \mathfrak{sl}(q) \oplus \mathbb{R}$.
- We have a split inner class, which corresponds to θ having nontrivial image in $\mathbb{Z}/2\mathbb{Z}$, so θ acts by the flip on the Dynkin diagram. If n is odd, then there is only one Vogan diagram (because θ fixes nothing), so there is only the split form.

  Otherwise, if n is even, then there are two Vogan diagrams (because θ fixes the middle vertex), so we have at most two real forms. Here, one calculates that our real forms are the split form $\mathfrak{sl}(n, \mathbb{R})$ with $\mathfrak{k} = \mathfrak{so}(n)$ (for black vertex) and the additional form $\{X \in \mathfrak{su}(n/2, \mathbb{H}) : \operatorname{Re}\operatorname{tr} X = 0\}$ with $\mathfrak{k} = \mathfrak{sp}(n)$ (with white vertex). ■

Remark 3.53. Already we can see that there are far more Vogan diagrams than real forms in the case where θ acts trivially.

Example 3.54. Consider the following Vogan diagram

$$\circ \text{———} \bullet \text{———} \circ \text{———} \circ \text{———} \bullet \text{———} \circ$$

which gives a form of $\mathfrak{sl}_7(\mathbb{C})$. Here, we see that $\theta(E_{12}) = E_{12}$, $\theta(E_{23}) = -E_{23}$, and so on. Thus, we are looking at the Lie algebra of the special unitary group preserving the bilinear form given by $\theta = \operatorname{diag}(1, 1, -1, -1, -1, 1, 1)$, which we see by examining sign changes. Thus, we have the real form $\mathfrak{su}(4, 3)$! A similar argument shows that each black vertex corresponds to a sign change (whereas a white vertex means staying the same sign), allowing us to calculate the real form for any such Vogan diagram.

3.3 April 6

Today, we write down some Vogan diagrams.

3.3.1 Linear Algebra over the Quaternions

To handle types the other classical types B, C, and D, we need to do a little quaternionic linear algebra. Recall the quaternions $\mathbb{H}$ is a non-commutative quadratic extension of $\mathbb{C}$ by an element j for which $j^2 = -1$ and $ij = -ji$.

Notation 3.55. We define $\mathrm{GL}_n(\mathbb{H})$ to be the group of $\mathbb{R}$-linear automorphisms of $\mathbb{H}^n$.

Remark 3.56. The group $\mathrm{GL}_n(\mathbb{H})$ is a real form of $\mathrm{GL}_{2n}(\mathbb{C})$.

Definition 3.57 (determinant)**.** We define a determinant map $\det\colon \mathrm{GL}_n(\mathbb{H})$ by sending $\alpha \in \mathrm{GL}_n(\mathbb{H})$ to $\det \alpha := \det_{\mathbb{C}}(\alpha; \mathbb{H}^n)$.

Remark 3.58. The determinant turns out to output to $\mathbb{R}^+$. It is enough to check this for diagonal matrices and unipotent upper-triangular matrices (by some row-reduction). There is nothing to say for unipotent matrices, and the case of diagonal matrices reduces to $n = 1$. In this case, we are looking at the determinant map $\det\colon \mathbb{H}^\times \to \mathbb{C}^\times$, which can be computed directly. Alternatively, we can check that $\mathbb{H}^\times \subseteq \mathrm{GL}_2(\mathbb{C})$ turns out to be given by $\mathrm{SU}(2) \times \mathbb{R}^\times$, all of whose determinants go to $\mathbb{R}^+$ because the diagonal $\mathbb{R}^\times$ piece is squared.

The moral of having a determinant is that we can define a special linear group.

Notation 3.59. We define $\mathrm{SL}_n(\mathbb{H})$ to be the subgroup of $\mathrm{GL}_n(\mathbb{H})$ of elements with trivial determinant.

Remark 3.60. By Remark 3.58, there is a short exact sequence

$$1 \to \mathrm{SL}_n(\mathbb{H}) \to \mathrm{GL}_n(\mathbb{H}) \to \mathbb{R}^+ \to 1.$$

The same arguments over $\mathbb{R}$ or $\mathbb{C}$ tell us that we may take the differential to get a short exact sequence

$$1 \to \mathfrak{sl}_n(\mathbb{H}) \to \mathfrak{gl}_n(\mathbb{H}) \to \mathbb{R} \to 0$$

of Lie algebras. One can check that the right-hand map sends X to $\operatorname{Re}\operatorname{tr} X$; for example, this map should vanish on strictly upper-triangular or strictly lower-triangular matrices, so it reduces to a calculation on $n = 1$, where it can be computed directly.

We will also want to have some sesquilinear forms on $\mathbb{H}$.

Definition 3.61 (sesquilinear)**.** Fix a vector space V over $\mathbb{H}$. Then a *sesquilinear* form on V is an $\mathbb{R}$-bilinear map $(-,-)\colon V \times V \to \mathbb{H}$ for which

$$(x\alpha, y\beta) = \overline{\alpha}(x, y)\beta$$

for any $x, y \in V$ and $\alpha, \beta \in \mathbb{H}$. We say that $(-,-)$ is *Hermitian* if and only if $(x, y) = \overline{(y, x)}$, and it is *skew-Hermitian* if and only if $(x, y) = -\overline{(y, x)}$.

One can classify these forms as follows.

Proposition 3.62. Fix a vector space V over $\mathbb{H}$.

(a) Every non-degenerate Hermitian form on V is equivalent to

$$(x, y) = (\overline{x}_1 y_1 + \cdots + \overline{x}_p y_p) - (\overline{x}_{p+1} yp + 1 + \cdots + \overline{x}_{p+q} y_{p+q}),$$

where $p + q = \dim_{\mathbb{H}} V$.

(b) Every non-degenerate skew-Hermitian form is equivalent to

$$(x, y) = \overline{x}_1 j y_1 + \cdots + \overline{x}_n j y_n.$$

Proof. Omitted. This is similar to the argument for complex vector spaces. ■

Definition 3.63 (signature)**.** Fix a non-degenerate Hermitian form B on a quaternionic vector space V. Then its *signature* is the pair (p, q) where B is equivalent to the standard form

$$(x, y) = (\overline{x}_1 y_1 + \cdots + \overline{x}_p y_p) - (\overline{x}_{p+1} yp + 1 + \cdots + \overline{x}_{p+q} y_{p+q}).$$

Remark 3.64. Given a quaternionic sesquilinear form B on V, then we can define $B_1, B_2 \colon V \times V \to \mathbb{C}$ so that $B = B_1 + jB_2$, where each are $\mathbb{R}$-bilinear. In fact, it turns out that B_1 is $\mathbb{C}$-sesquilinear, and B_2 is bilinear, and they satisfy $B_2(x, y) = B_1(xj, y)$. Furthermore, B is Hermitian of signature (p, q) if and only if B_1 is Hermitian of signature $(2p, 2q)$ and B_2 is skew-symmetric. Similarly, B_1 is skew-Hermitian if and only if B_1 is skew-Hermitian of signature (n, n) and B_2 is symmetric.

Our sesquilinear forms now have unitary groups.

Definition 3.65 (unitary group)**.** Fix a sesquilinear form B of a quaternionic vector space V. Then we define the *unitary group* $\mathrm{U}(B)$ to be the group of linear transformations of V which preserve B.

Example 3.66. Suppose that B is Hermitian of signature (p, q). Then Remark 3.64 grants us a decomposition $B = B_1 + jB_2$ where B_1 is Hermitian of signature (p, q) and B_2 is skew-symmetry. It follows that

$$\mathrm{U}(B) = \mathrm{U}(2p, 2q) \cap \mathrm{Sp}(2n).$$

We will denote this group by $\mathrm{U}(p, q, \mathbb{H})$ and its Lie algebra by $\mathfrak{u}(p, q, \mathbb{H})$.

Example 3.67. Suppose that B is skew-Hermitian. Then Remark 3.64 again allows us to decompose B, and we find that

$$\mathrm{U}(B) = \mathrm{U}(n, n) \cap \mathrm{O}(2n).$$

We will denote this group by $\mathrm{O}^*(2n)$ and its Lie algebra by $\mathfrak{so}^*(2n)$.

3.3.2 Real Forms of Classical Types

We are now ready to write down all real forms in classical types. We already handled type A in Exercise 3.52.

Exercise 3.68. We classify the real forms of type B_n.

Proof. Here, $\mathfrak{g} = \mathfrak{so}(2n+1)$. The Dynkin diagram looks like

$$\bullet \text{———} \bullet \text{———} \cdots \text{———} \bullet \Longrightarrow \bullet$$

which has no automorphisms, so all forms are in the same inner class. Now, we are hunting for $\theta^2 = 1$. Because $G_{\mathrm{ad}} = \mathrm{SO}(2n+1)$, we find that θ acts by some matrix which can be taken to commute with our choice of maximal torus. Then θ lives in the corresponding torus of $\mathrm{SO}(2n+1)$ (because it fixes the Dynkin diagram), so it diagonalizes appropriately. By using the Weyl group, we can rearrange the $(+1)$s and the (-1)s in its diagonalization, so we find that θ is given by conjugation by $1_p \oplus (-1)_q$ for some p and q with $p+q=n$. This gives the real form $\mathfrak{so}(2p, 2q+1)$, where $0 \le p \le n$ and q is chosen so that $p+q=n$. Finding θ-stable elements reveals that $\mathfrak{k} = \mathfrak{so}(2p) \oplus \mathfrak{so}(2q+1)$. ■

Exercise 3.69. We classify the real forms of type C_n.

Proof. Once again, the Dynkin diagram has no automorphisms. Here, $G_{\mathrm{ad}} = \mathrm{Sp}_{2n}(\mathbb{C})/\{\pm 1\}$, so $\theta^2 = 1$ has two cases for a lift $\widetilde{\theta}$ to $\mathrm{Sp}_{2n}(\mathbb{C})$. One can always change basis (as in Exercise 3.68) so that $\widetilde{\theta}$ is diagonal and has appropriately arranged diagonal elements.

- If $\widetilde{\theta}^2 = 1$, then we may assume $\widetilde{\theta} = 1_p \oplus (-1_q)$. This gives the real form $\mathfrak{u}(p, q, \mathbb{H})$, which has $\mathfrak{k}^c = \mathfrak{u}(p, \mathbb{H}) \times \mathfrak{u}(p, \mathbb{H})$ so that $\mathfrak{k} = \mathfrak{sp}_{2p}(\mathbb{C}) \times \mathfrak{sp}_{2p}(\mathbb{C})$.

- If $\widetilde{\theta}^2 = -1$, then one can define $V = \mathbb{C}^{2n}$ into two prices $V(i)$ and $V(-i)$, which one can see are both isotropic subspaces. Each must be at most n-dimensional (over $\mathbb{C}$) because this is the largest size of an isotropic subspace, so for dimension reasons, they are both n-dimensional. Thus, $\mathfrak{k} = \mathfrak{gl}_n(\mathbb{C})$, namely fixing the eigenspaces. (The action on $V(-i)$ is determined by the action on $V(i)$ by definition of $\mathfrak{sp}(2n)$.) Note that the real form is $\mathfrak{sp}_{2n}(\mathbb{R})$ by exhaustion. ■

Exercise 3.70. We classify the real forms of type D_n.

Proof. Let's start with the compact inner class. Here, $G_{\mathrm{ad}} = \mathrm{SO}_{2n}(\mathbb{C})/\{\pm 1\}$, so $\theta^2 = 1$ has two cases for a lift $\widetilde{\theta}$ to $\mathrm{SO}_{2n}(\mathbb{C})$. As usual, we may assume that $\widetilde{\theta}$ is given by conjugation by a diagonal element.

- If $\widetilde{\theta}^2 = 1$, then we may assume that $\widetilde{\theta} = 1_p \oplus (-1_q)$. This gives the real form $\mathfrak{so}(2p, 2q)$, which has $\mathfrak{k} = \mathfrak{so}_{2p}(\mathbb{C}) \times \mathfrak{so}_{2q}(\mathbb{C})$.

- If $\widetilde{\theta}^2 = -1$, then we may assume that $\widetilde{\theta} = i1_p \oplus (-i1_q)$. But these eigenspaces subspaces are isotropic, so $p \le n$ and $q \le n$, so $p = q = n$ is forced by $p+q=n$. It follows that $\mathfrak{k} = \mathfrak{gl}_n$ by acting on one of the eigenspaces, and one finds that the real form is $\mathfrak{so}^*(2n)$.

There is also another inner class corresponding to when θ is nontrivial.[1] In this case, we see that θ actually lifts to conjugation by an element of $\mathrm{O}(n)$ with determinant -1. Thus, θ can be taken to be a diagonal matrix $1_p \oplus (-1_q)$ where $p+q=n$ and q is odd. Arguing as in the first point above, we see that we get the real form $\mathfrak{so}(2p+1, 2q-1)$, and $\mathfrak{k} = \mathfrak{so}(2p+1) \times \mathfrak{so}(2q-1)$. ■

Remark 3.71. For D_n at $n = 3$, one can compare our calculation with A_3. By comparing $\mathfrak{k}$s, we find $\mathfrak{so}(6, \mathbb{R}) = \mathfrak{su}(4)$ because are the compact forms. By comparing $\mathfrak{k}$s, we also find that $\mathfrak{so}(4, 2) = \mathfrak{su}(2, 2)$ and $\mathfrak{so}(3, 3) = \mathfrak{sl}(4, \mathbb{R})$ and $\mathfrak{so}(5, 1) = \mathfrak{sl}(2, \mathbb{H})$ and $\mathfrak{so}^*(6) = \mathfrak{su}(3, 1)$.

[1] This inner class is not necessarily the split inner class. Indeed, for D_n, the parity of n determines if the compact inner class is split.

3.4 April 8

Today, we continue playing the Vogan game.

3.4.1 Real Forms of Types F_4 and G_2

We now pass to exceptional types. Linear algebra is no longer so helpful to find our real forms. Instead, we will use Vogan diagrams. We have already seen that many Vogan diagrams can give the same real form, so it will be helpful to have a means to determine when two Vogan diagrams give the same real form. Here is one criterion.

Example 3.72. Choose a Vogan diagram coming from some θ, and choose a vertex i stable under θ. Then θ and $s_i\theta$ give the same real form because they are automatically conjugate. On Vogan diagrams, one sees that

$$\alpha_j(s_i\theta) = s_i\alpha_j(\theta) = (\alpha_j - a_{ij}\alpha_i)(\theta) = \alpha_j(\theta)\alpha_i(\theta)^{-a_{ij}}.$$

Now, $\alpha_j(\theta)$ is $+1$ when j is colored white, and it is -1 when j is colored black. Thus, the Vogan diagram is entirely the same, except when $\alpha_i(\theta) = -1$ and a_{ij} is odd. In practice, this means that we can take a black vertex i in the Vogan diagram, and we may change the colors of all neighbors j with $a_{ij} \in \{-1, -3\}$ to receive a Vogan diagram with an equivalent real form.

The above rule allows us to play the "Vogan game" to try to go between real forms.[2]

Example 3.73. In type G_2, the Dynkin diagram has no nontrivial automorphisms. The Vogan diagram

$$\circ \Lleftarrow \circ$$

gives the compact form; here, $\mathfrak{k} = \mathfrak{g}$. There is also the (non-isomorphic) split form, which must come from the Vogan diagrams

$$\bullet \Lleftarrow \circ \qquad \bullet \Lleftarrow \bullet \qquad \circ \Lleftarrow \bullet$$

which are all equivalent (via Example 3.72). For the split inner form, one can compute as in Theorem 3.29 that the signature of the Killing form on the split form is $(\operatorname{rank}\mathfrak{g} + \#\Phi_+, \#\Phi_+)$, so comparing with the splitting $\mathfrak{g}_\theta = \mathfrak{k}^c \oplus i\mathfrak{p}^c$ shows that the rank of $\mathfrak{k}$ is $\#\Phi_+$. Thus, $\mathfrak{k}$ is some six-dimensional Lie algebra of rank 2; the only possibility is $\mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$.

Example 3.74. We compute the Vogan diagrams in type F_4.

Proof. Before starting, recall that F_4 is a root system in $\frac{1}{2}\mathbb{Z}^4$, given by the roots in B_4 (which are $\{\pm e_i\} \sqcup \{\pm e_i \pm e_j\}$) and adding the roots $\pm_1 e_1 \pm_2 e_2 \pm_3 e_3 \pm_4 e_4$. Thus, F_4 has rank 4 and dimension $4 + 32 + 16 = 52$. The Dynkin diagram has no nontrivial automorphisms. The Vogan diagram

$$\circ \text{———} \circ \Lleftarrow \circ \text{———} \circ$$

gives the compact form; here, $\mathfrak{k} = \mathfrak{g}$.

We now play the Vogan game. By inspection, we see that a_{ij} is odd for all $i \neq j$, except for $a_{23} = -2$. Thus, the right two vertices (which are the long roots) can change the colors of the left two vertices (which are the short roots), but not the other way around. To this end, note that all the diagrams

$$\bullet \text{———} \circ \qquad \bullet \text{———} \bullet \qquad \circ \text{———} \bullet$$

are equivalent. Thus, by iteratively playing the Vogan game on each pair of vertices (first on the long roots and then on the short roots), we get three more Vogan diagrams, as follows.

[2] Because the Weyl group acts transitively on polarizations, this rule should be enough to exhaust the relations in the compact inner class.

- We have the following.

$$\bullet \text{———} \circ \Longleftarrow \circ \text{———} \circ$$

  We may call this real form F_4^1. In this case, one can see by inspection that θ acts by -1 on $\mathfrak{g}_\alpha$ if and only if $\alpha \notin \mathbb{Z}^4$. It follows that $\mathfrak{k} = \mathfrak{g}^\theta$ has exactly the roots from $B_4 = \mathfrak{so}(9)$, so $\mathfrak{k} = \mathfrak{so}(9)$ by comparing ranks. On the remaining roots, one finds that $\mathfrak{p}$ is given by the spin representation by computing its weight decomposition. Note that this is not the split form because $\operatorname{rank} \mathfrak{k} \neq |\Phi_+| = 24$, and it is not the compact form because θ is nontrivial.

- We have the following.

$$\bullet \text{———} \circ \Longleftarrow \circ \text{———} \bullet$$

- We have the following.

$$\circ \text{———} \circ \Longleftarrow \circ \text{———} \bullet$$

  In fact, this is equivalent to the previous diagram: the previous diagram is equivalent to

$$\circ \text{———} \circ \Longleftarrow \bullet \text{———} \bullet$$

  which is equivalent to

$$\circ \text{———} \bullet \Longleftarrow \bullet \text{———} \circ$$

  which is equivalent to

$$\bullet \text{———} \bullet \Longleftarrow \bullet \text{———} \circ$$

  which is equivalent to

$$\circ \text{———} \circ \Longleftarrow \bullet \text{———} \bullet$$

  which is equivalent to

$$\circ \text{———} \circ \Longleftarrow \circ \text{———} \bullet$$

  as needed.

  Now, by exhaustion, we conclude that we must have the split form. Here, we find that θ fixes a root system of type C_3, so $\mathfrak{k}$ contains $\mathfrak{sp}(6)$, which has dimension 21. However, $\mathfrak{k}$ should increase in rank by 1 while increasing in dimension 3, so we are required to have $\mathfrak{k} = \mathfrak{sp}(6) \oplus \mathfrak{sl}(2)$. Indeed, all semisimple algebras strictly containing $\mathfrak{sp}(6)$ have dimension larger than 24, so $\mathfrak{k}$ must have another summand, and the smallest possible summand to increase the rank is either $\mathfrak{sl}(2)$ or $\mathfrak{gl}(1)$, but adding $\mathfrak{gl}(1)$s cannot help us. ■

3.4.2 Real Forms of Type E

Having done everything easy, it remains to do things which are hard. We are now in type E. To start, note that these types have a single inner class, except the Dynkin diagram E_6 admits a nontrivial automorphism. Let's deal with this first.

Exercise 3.75. We compute the real forms in type E_6 which are in the non-compact inner class.

Proof. In type E_6, it is possible to have vertices which are connected in the Vogan diagram: indeed, the split inner class comes from the flip of the Dynkin diagram. In this case, playing the Vogan diagram on the remaining two fixed vertices, we get two real forms, as follows.

- We have the following.



  Here, we see that $\mathfrak{k}$ has rank 4, and by looking at the projection of these roots to $\mathfrak{k}$, we see that the two pairs of associated roots become short, so we $\mathfrak{k}$ has type F_4. This is not the split form for dimension reasons: here, $\dim E_6 = 78$, and the split form has $\dim \mathfrak{k}$ equal to the number of positive roots.

- The Vogan diagram



must correspond to the split form for dimension reasons. The same projection argument shows that $\mathfrak{k}$ contains $\mathfrak{sp}(6)$. One can check that it is $\mathfrak{sp}(8)$, which we will do on the homework. ■

We now turn to the compact inner class. Let's start by making some general comments. The Dynkin diagram has a short leg, a middle leg, and a long leg. For example, E_7 looks like the following.



Once we exclude the compact form, we may assume that there is at least one black vertex. By playing the Vogan game, we may move the black vertex to the end of the short leg, and then we can always use this short leg in order to force the intersection point to be a black vertex. We now do casework on our type.

We now use this to force the long leg of the Dynkin diagram into a particular shape, possibly at the cost of the black vertex at the short leg.

- In type E_6, the short and long leg has two vertices, so there are two possibilities on each leg: $\circ - \circ$ or $\circ - \bullet$. By possibly undoing our work with the nodal vertex, we may force the long leg to be $\bullet - \bullet$.
- In type E_7, we also have to play the Vogan diagram on three vertices. We find that the long leg has $\circ - \circ - \circ$ or $\bullet - \circ - \circ = \bullet - \bullet - \circ = \circ - \bullet - \bullet$ or $\bullet - \circ - \bullet = \bullet - \bullet - \bullet = \circ - \bullet - \circ$. However, using the nodal vertex allows us to see that the rightmost diagrams of the last two classes are equivalent, and the leftmost diagrams of the last two classes are equivalent.
- In type E_8, the long leg has four vertices. One again plays the Vogan game on A_4. We have the following classes.
  - We have $\circ - \circ - \circ - \circ$.
  - We have $\bullet - \circ - \circ - \circ = \bullet - \bullet - \circ - \circ = \circ - \bullet - \bullet - \circ = \circ - \circ - \bullet - \bullet = \circ - \circ - \circ - \bullet$. Note that the nodal vertex allows us to push this to the previous class.
  - We have $\bullet - \circ - \circ - \bullet = \bullet - \circ - \bullet - \bullet = \bullet - \bullet - \bullet - \circ = \circ - \bullet - \circ - \circ$ or $\bullet - \circ - \bullet - \circ = \bullet - \bullet - \bullet - \bullet = \bullet - \bullet - \circ - \bullet = \circ - \bullet - \bullet - \bullet$. Again, the nodal vertex allows us to push this to $\circ - \circ - \circ - \bullet$ and thus to the previous class.

Now, by redoing our nodal vertex, we find that we may assume that the short leg can be assumed to be black, the node can be black, and the long leg can look like $? - \bullet - \cdots - \bullet$. In types E_6 or E_8, one finds that we can even assume that $?$ is black by some combinatorics: in E_6, we are simply saying that $\cdots - \circ - \bullet = \cdots - \bullet - \bullet$. We won't write it our for E_8.

Similarly, the middle leg is a Vogan game on A_2, so it has either $\circ - \circ$ or $\bullet - \bullet = \bullet - \circ = \circ - \bullet$. Thus, E_6 has at most two Vogan diagrams, E_7 has at most four Vogan diagrams, and E_8 has at most two Vogan diagrams. We now work through our types separately.

Exercise 3.76. We compute the real forms in type E_6 in the compact inner class.

Proof. We know that there are at most three Vogan diagrams in the compact inner class, two of which are non-compact. Here are our diagrams.

- We have the following.



Here, we label the black vertex to be the miniscule weight $\alpha_1 = \omega_1$. It follows that every positive root contains α_1 with a coefficient of either 0 or 1 (because it is a fundamental weight), and θ fixes the root if and only if this coefficient is $+1$. We conclude that $\mathfrak{k}$ contains a copy of the rest of the Dynkin diagram, which gives $\mathfrak{so}(5) \subseteq \mathfrak{k}$. However, the rank of $\mathfrak{k}$ should be five, and a dimension argument (with the signature of the Killing form) gives $\mathfrak{k} = \mathfrak{so}(5) \oplus \mathfrak{so}(2)$.

- We have the following.



The same argument shows that $\mathfrak{k}$ contains $\mathfrak{sl}(6)$. The black vertex corresponds to the fundamental weight of the adjoint representation, and one can check that this means that θ will act by $+1$ only on an extra copy of $\mathfrak{sl}(2)$ given by the root spaces of the maximal root. (Notably, $\mathfrak{k}$ exactly contains the torus and the roots where θ acts by $+1$, which can be computed.) ■

Exercise 3.77. We compute the real forms in type E_7 in the compact inner class.

Proof. There are at most four Vogan diagrams which have a black vertex, though we will eventually bring that number down to three. Note $\dim E_7 = 133$, so the number of positive roots is 63.

- We have the following.



Here, $\mathfrak{k}$ has a copy of E_6, which has dimension 78, which is bigger than 63, so $\mathfrak{k}$ is not split. Because the colored root is miniscule, we find that $\mathfrak{k} = E_6 \oplus \mathfrak{so}(2)$ again.

- We have the following.



Here, $\mathfrak{k}$ contains $\mathfrak{so}(12)$ automatically, so dimension reasons make this form not split, and we can see that $\mathfrak{k} = \mathfrak{so}(12) \oplus \mathfrak{sl}(2)$ because the black vertex is maximal, corresponding to the adjoint representation.

- We have the following.



Here, $\mathfrak{k}$ contains $\mathfrak{sl}(7)$. This will turn out to be the split form by exhaustion, so it has dimension

- It remains to show that our last two Vogan diagrams were actually the same. Well,



is



which is



which is



which is



which is



which is among one of our restricted diagrams. ■

Exercise 3.78. We compute the real forms in type E_8 in the compact inner class.

Proof. Note E_8 has 120 positive roots.

- Consider the following Vogan diagram.



  Here, the black vertex is the adjoint representation, so $\mathfrak{k} = E_7 \oplus \mathfrak{sl}(2)$ for rank reasons. This is not the split form because $\dim \mathfrak{k}$ is too large.

- Consider the following Vogan diagram.



  Here, $\mathfrak{k}$ contains $\mathfrak{so}(14)$. This should be the split form by exhaustion. For dimension reasons, this needs to be $\mathfrak{so}(16)$. Also, $\mathfrak{k}$ needs to look like $\mathfrak{so}(14) \oplus R$ where R is some remaining representation of $\mathfrak{so}(14)$ of dimension 29, which we see must be $\mathbb{C}^{14} \oplus \mathbb{C}^{14} \oplus \mathbb{C}$. (This representation needs to be self-dual.) By inspection, we must have $\mathfrak{so}(16)$. While we're here, we remark that $\mathfrak{p}$ turns out to be the spin representation. ■

3.5 April 13

Today we classify connected compact Lie groups.

3.5.1 Classification of Compact Connected Lie Groups

We will show the following on the homework.

Proposition 3.79. Let K be a compact Lie group. Then $\operatorname{Lie} K_{\mathbb{C}}$ is reductive.

Sketch. Using integration, one can find an invariant non-degenerate bilinear form on $\operatorname{Lie} K$, which one can use to show that the adjoint representation is semisimple. It follows that $\operatorname{Lie} K_{\mathbb{C}}$ can be decomposed into a sum of simple Lie algebras. ■

For the rest of the subsection, K^c will be a connected compact Lie group, and its Lie algebra will be denoted $\mathfrak{k}^c$, where $\mathfrak{k} := \mathfrak{k}^c_{\mathbb{C}}$. We recall that the short exact sequence

$$0 \to \mathfrak{z}(\mathfrak{k}) \to \mathfrak{k} \to \mathfrak{k}_{\mathrm{ss}} \to 0$$

splits. Thus, K^c will basically be a product of G^c_{ad} for the semisimple Lie group corresponding to $\mathfrak{k}_{\mathrm{ss}}$ and some S^1s coming from $\mathfrak{z}(\mathfrak{k})$. With this in mind, our classification will be done in two steps.

- We will show that the universal cover G^c of G^c_{ad} is compact.
- Then we will show that K can be realized as a quotient of $G^c \times Z$ by a finite central subgroup, where Z is some finite product of S^1s.

Let's begin with the first point. This will require a finer study of the fundamental group, which comes from the following construction.

Definition 3.80. Let $\mathfrak{g}$ be the complex semisimple Lie algebra, and let $\pi\colon G \to G_{\mathrm{ad}}$ be the universal cover of G_{ad}. For each dominant integral weight λ, we define χ_λ to be the character $\ker \pi \to \mathbb{C}^\times$ giving the action on L_λ. Note that L_λ lifts to a representation of G because G is simply connected.

Remark 3.81. We quickly recall why G is actually a Lie group: the point is that $\pi_1(G)$ is a finitely generated group and hence countable. This holds by finding a countable locally finite cover G with small balls, which shows that G has the same fundamental group as a countable locally finite graph. The fundamental group of such a graph is then finitely generated.

Remark 3.82. Note that $\ker \pi$ is a discrete central subgroup: it is certainly discrete and normal (because it is a kernel). Then it is central because the adjoint action of the total group must act smoothly, so it cannot move around the discrete points.

Remark 3.83. Note that χ_λ is well-defined by Schur's lemma.

Remark 3.84. Because $L_{\lambda+\mu} \subseteq L_\lambda \otimes L_\mu$ (indeed, the latter has a vector of highest weight $\lambda+\mu$), we find that $\chi_{\lambda+\mu} = \chi_\lambda \chi_\mu$. Thus, $\chi_\bullet$ is additive on the dominant integral weights P_+, so we may extend $\chi_\bullet$ to a character

$$\chi_\bullet\colon P \to \mathrm{Hom}(\ker \pi, \mathbb{C}^\times).$$

Remark 3.85. Because the adjoint representation $\mathfrak{g}$ descends to $G_{\mathrm{ad}} \subseteq \mathrm{Aut}(\mathfrak{g})$ for free, we conclude that $\chi_\theta = 1$, where θ is the maximal root.

Proposition 3.86. Let $\mathfrak{g}$ be the complex semisimple Lie algebra, and let $\pi\colon G \to G_{\mathrm{ad}}$ be the universal cover of G_{ad}. Then the highest weight representation L_λ descends to G_{ad} if and only if $\lambda \in Q$.

Proof. We show the two implications separately.

- Suppose $\lambda \in Q$. On the homework, we showed that $L_{\lambda+\alpha} \subseteq \mathfrak{g} \otimes L_\lambda$ for λ large enough (i.e., provided that $(\lambda, \alpha_i^\vee)$ is large enough for i large enough). Thus, $\chi_{\lambda+\alpha} = \chi_\lambda$ for λ large enough, so $\chi_\alpha = 1$ follows. (For example, this recovers the fact that $\chi_\theta = 1$.) Thus, $\chi_\bullet$ descends to a character on P/Q. Now, $\chi\colon P/Q \to \mathrm{Hom}(Z, \mathbb{C}^\times)$ may alternatively be viewed as a character $Z \to (P/Q)^\vee$ by duality, which is also $P^\vee/Q^\vee$.

- For the converse, we see that G_{ad} admits the faithful self-dual representation $\mathfrak{g}$, so L_λ descending to G_{ad} implies that it descends to G_{ad}^c, which in turn is $\otimes$-generated by $\mathfrak{g}$. However, the weights of $\mathfrak{g}$ are in Q, so it follows by taking the character that the weights of L_λ are in Q, so $\lambda \in Q$. ■

Remark 3.87. It follows from the proof that the character $\chi\colon P/Q \to \mathrm{Hom}(Z, \mathbb{C}^\times)$ is injective.

All the same arguments hold for the universal cover $G^c \to G_{\mathrm{ad}}^c$. We are now ready to calculate π_1.

Theorem 3.88. Let $\mathfrak{g}$ be the complex semisimple Lie algebra, and let $\pi\colon G^c \to G_{\mathrm{ad}}^c$ be the universal cover of G_{ad}, and let its kernel be Z^c. Then

$$\chi\colon P/Q \to \mathrm{Hom}(Z^c, \mathbb{C}^\times)$$

is an isomorphism.

Proof. By Remark 3.81, we find that Z^c is a finitely generated abelian group. For a choice of finite-index subgroup Z', we note that G/Z' is a finite cover of G_{ad}^c and therefore a compact Lie group which is a central extension of G_{ad}^c by the center Z^c/Z'.

Notably, the finite-dimensional representations of G^c/Z' are given by the sum L_λs where $\chi_\lambda|_{Z'}$ vanishes; notably, these λs include Q. We then let $P'_+ \subseteq P_+$ be the subset of those dominant integral weights, and we let P' be the induced subgroup. Now, $\chi_\bullet$ descends to an injective homomorphism

$$P'/Q \to\to \mathrm{Hom}(Z^c/Z', \mathbb{C}^\times).$$

Now, G' is compact, so the Peter–Weyl theorem tells us that all characters are in fact achieved, so the χ_λ span a dense subset in $C(G)$, which gives the surjectivity of this map. ■

We can collect all of our work into the following two results.

Corollary 3.89. Let $\mathfrak{g}$ be the complex semisimple Lie algebra.

(a) Then the simply connected Lie group G^c with Lie algebra $\mathfrak{g}^c$ is compact with center $P^\vee/Q^\vee$. Furthermore, $\pi_1(G_{\mathrm{ad}}^c) \cong P^\vee/Q^\vee$.

(b) If $\Gamma \subseteq P^\vee/Q^\vee$ is a subgroup, then the category $\mathrm{Rep}\, G^c/\Gamma$ is semisimple and generated by the highest weight representations L_λ such that $\chi_\lambda|_\Gamma = 1$.

(c) Decompose $\mathfrak{g}$ into a sum $\bigoplus_{i=1}^n \mathfrak{g}_i$ of simple Lie algebras. Then any connected Lie group G with Lie algebra $\mathfrak{g}^c$ is compact and takes the form

$$\left(\prod_{i=1}^n G_{i,\mathrm{ad}}^c\right)/Z^c,$$

where $Z^c = \pi_1(G^c)$. Every semisimple compact connected Lie group is of this form.

Proof. Here, (a) and (b) from Theorem 3.88. It only remains to show the classification parts of (c), which we do in two steps.

- Note that any semisimple compact connected Lie group G will have complex semisimple Lie algebra $\mathfrak{g}_{\mathbb{C}}$. Then $\mathfrak{g}$ must have negative-definite Killing form in order for G to be compact (otherwise, the adjoint group needs to act transitively on a form which is neither positive-definite nor negative-definite!), so $\mathfrak{g} = \mathfrak{g}_{\mathbb{C}}^c$ by our classification of real forms!
- Once G has Lie algebra $\mathfrak{g}^c$, it surjects onto G_{ad}, and this must be a covering map, so G is a quotient of the universal cover G^c. This G^c can be computed on each factor, which proves the first claim. ■

Because it is not much harder, we also do the reductive case. By Corollary 3.89, it is enough to reduce to the semisimple case.

Proposition 3.90. Any connected compact Lie group K is a finite quotient of a Lie group of the form $T \times C$, where T is a product of S^1s, and C is a simply connected semisimple compact Lie group.

Proof. By Proposition 3.79, $\mathfrak{k} := \operatorname{Lie} K$ is reductive. Thus, $\mathfrak{k}$ splits into a sum $\mathfrak{t} \oplus \mathfrak{c}$ where $\mathfrak{t}$ is the center, and $\mathfrak{c}$ is semisimple. It follows that the subgroup $T \subseteq K$ corresponding to $\mathfrak{t}$ is central, and the subgroup $C \subseteq K$ corresponding to $\mathfrak{c}$ is a normal subgroup. Note that C is a closed subset in a compact space, so C is compact.

Now, the image of the homomorphism $T \times C \to K$ is a local diffeomorphism at the identity, so it contains an open neighborhood of the identity, so it is surjective. Thus, the kernel of this map is some discrete subgroup $\Gamma \subseteq T \times C$, so by compactness, Γ is finite, and the result follows! ■

Remark 3.91. The proof also shows that we can take T to be taken to be the connected component of the center of K.

3.5.2 Polar Decomposition

As an aside to our study of real forms, we note that we are able to generalize the decompositions

$$\mathrm{GL}_n(\mathbb{R}) = \mathrm{B}_n(\mathbb{R})\,\mathrm{O}_n(\mathbb{R}),$$

where $\mathrm{B}_n(\mathbb{R})$ is the subgroup of upper-triangular matrices.

To this end, we need to generalize what we mean by B_n and O_n for general real forms. Let's start with the group.

Notation 3.92. Fix a complex semisimple Lie algebra $\mathfrak{g}$. For a Cartan involution θ inducing a real form $\mathfrak{g}_\theta$, we let $G_{\mathrm{ad},\theta} \subseteq G_{\mathrm{ad}}$ be the corresponding subgroup, which are the fixed points of $\omega_\theta := \omega \circ \theta$.

Example 3.93. The group $G_{\mathrm{ad},\theta}$ is not aways connected. For example, for $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$, take θ to induce the split form. Then $G_{\mathrm{ad},\theta} = \mathrm{PSL}_2(\mathbb{R})$, which has two connected components: one can have positive or negative determinant!

Here is our theorem.

Theorem 3.94 (Polar decomposition)**.** Fix a complex semisimple Lie algebra $\mathfrak{g}$. For a Cartan involution θ inducing a real form $\mathfrak{g}_\theta$. Let $K_\theta^c \subseteq G_{\mathrm{ad},\theta}$ be the subgroup of elements which act on $\mathfrak{g}$ by unitary operators, and let P_θ be the image $\exp(i\mathfrak{p}^c)$. Then the multiplication map

$$\mu\colon K^c \times P_\theta \to G_{\mathrm{ad},\theta}$$

is a diffeomorphism.

Remark 3.95. By construction, we see that P_θ is the subgroup of $G_{\mathrm{ad},\theta}$ which acts on $\mathfrak{g}$ by positive Hermitian operators. Thus, there is a $\log\colon P_\theta \to i\mathfrak{p}^c$, which makes P_θ into a manifold.

Proof. The main point is to define an inverse for the multiplication map. We use the polar decomposition for usual matrices: any invertible complex matrix A can be written uniquely as $A = U_A R_A$, where U_A is unitary and R_A is positive; explicitly, one can define $R_A := \sqrt{A^\dagger A}$ and then define $U_A := AR_A^{-1}$.

Now, we may view $g \in G_{\mathrm{ad},\theta}$ as elements of $\mathrm{Aut}(\mathfrak{g})$. For example, $g^\dagger g \in \mathrm{Aut}(\mathfrak{g})$ has positive eigenvalues, and it lies in $G_{\mathrm{ad},\theta}$ because this element commutes with ω_θ. Thus, this element lives in P_θ, and by undoing the exponential, we may take the square root to give $\sqrt{g^\dagger g}$. Thus, we may define U_g and R_g in $G_{\mathrm{ad},\theta}$ using the usual polar decomposition. By construction, $R_g \in P_\theta$, and we can see that U_g acts by a Hermitian operator on $\mathfrak{g}$, so $U_g \in K^c$ as well. Thus, $g \mapsto (U_g, R_g)$ is the required inverse for the multiplication map; note that it is smooth by construction! ■

Remark 3.96. It follows that $G_{\mathrm{ad},\theta}$ is homotopy-equivalent to K^c. Indeed, P_θ is diffeomorphic to a Euclidean space by construction.

Here is an application.

Corollary 3.97. Fix a semisimple complex Lie group G.

(a) The center Z of G is contained in G^c.

(b) Restriction defines an equivalence $\mathrm{Rep}_{\mathbb{C}} G \to \mathrm{Rep}_{\mathbb{C}} G^c$.

Proof. Let's start with (a); (b) then follows by using that the center is the obstruction to the descent of a representation from the simply connected cover, as in Corollary 3.89. By Theorem 3.94, we see that G_{ad} and G^c_{ad} are homotopy-equivalent, so their fundamental groups are the same. This fundamental group embeds into G^c_{ad} as some finite discrete central subgroup, where we see it must also embed into G_{ad}. The case of general G then follows by the classification given by Corollary 3.89. ■

Remark 3.98. The same sort of arguments shows that there is also a polar decomposition to the real form G_θ of any connected complex semisimple group G.

Example 3.99. There is a notable caveat that even if G is simply connected, the real form G_θ need not be connected and can even have infinite fundamental group. Indeed, for $G = \mathrm{SL}_2(\mathbb{C})$, the split real form $\mathrm{SL}_2(\mathbb{R})$ has infinite fundamental group: the polar decomposition gives

$$\mathrm{SL}_2(\mathbb{R}) \cong \mathrm{SO}_2(\mathbb{R}) \times \mathbb{R}^2,$$

and $\mathrm{SO}_2(\mathbb{R}) \cong S^1$ has infinite fundamental group. In general, when $\mathfrak{k} = \mathfrak{g}^\theta$ fails to be semisimple, one finds that G_θ has some torus component, which gives the infinite fundamental group.

Remark 3.100. When G_θ° has infinite fundamental group, it turns out that the nontrivial covers of G_θ° fail to be linear groups. As another example, for the split form $G_\theta = \mathrm{Sp}_{2n}(\mathbb{R})$, one has $\mathfrak{k}_\theta = \mathfrak{u}(n)$, so G_θ is homotopy-equivalent to $\mathrm{U}(n)$, which has fundamental group $\mathbb{Z}$. And indeed, the "generalized metaplectic" covers of G_θ fail to be linear! Roughly speaking, the issue is that $\mathrm{Sp}_{2n}(\mathbb{C})$ has finite fundamental group, so any representation of $\widetilde{\mathrm{Sp}}_{2n}(\mathbb{R})$ descends to the Lie algebra and then lifts to a finite cover of $\mathrm{Sp}_{2n}(\mathbb{C})$, so it cannot be faithful.

3.6 April 15

Here we go.

3.6.1 Reductive and Linear Groups

Intuitively, a reductive group should be a semisimple group times a torus. We take this as a definition.

Definition 3.101 (reductive)**.** A connected complex Lie group G is *reductive* if and only if it takes the form

$$\left((\mathbb{C}^\times)^r \times G_{\mathrm{ss}}\right)/Z,$$

where G_{ss} is a semisimple complex Lie group, and Z is some finite central subgroup.

Remark 3.102. Note that G being reductive implies that $\mathfrak{g}$ is reductive, but the converse does not hold. Indeed, one can show that the Lie group $G = \mathbb{C}$ is not reductive, which has abelian and hence reductive Lie algebra.

These reductive groups have maximal compact subgroups.

Notation 3.103. Let $G = \left((\mathbb{C}^\times)^r \times G_{\mathrm{ss}}\right)/Z$ be a connected complex Lie group. Then we define

$$G^c := \left(\left(S^1\right)^r \times G_{\mathrm{ss}}^c\right)/Z.$$

Remark 3.104. One can show that G^c is independent of the presentation of G, simply by perturbing this presentation accordingly.

Remark 3.105. The restriction functor from holomorphic finite-dimensional representations of G to continuous finite-dimensional representations of G^c is an equivalence. Indeed, one shows this because a holomorphic representation splits as a product of a representation of the torus and a representation of the semisimple group, from which one can use Corollary 3.97 to handle the semisimple part. For the torus, one needs to know that holomorphic representations of $\mathbb{C}^\times$ diagonalize as a direct sum of characters which look like $n\colon \mathbb{C}^\times \to \mathbb{C}^\times$.

Remark 3.106. Choose complex reductive groups G_1 and G_2. From Remark 3.105, it turns out that every homomorphism $G_1^c \to G_2^c$ extends uniquely to a holomorphic homomorphism $G_1 \to G_2$. It turns out that the category of connected compact Lie groups embeds faithfully into the category o connected complex reductive groups, whose inverse on objects is $G \mapsto G^c$.

While we're here, we say something about linear groups.

Definition 3.107 (linear)**.** A connected Lie group G is *linear* if and only if G admits a faithful finite-dimensional representation. Equivalently, G is linear if and only if it can be realized as a Lie subgroup of $\mathrm{GL}_n(\mathbb{R})$ for some n.

Example 3.108. Any compact group admits a faithful representation by Proposition 2.69, so they are linear.

Example 3.109. Any complex reductive group is linear. We don't have to worry about the torus (which can be embedded faithfully separately), and then one can pass to the compact part by Corollary 3.97, so we are done by the previous example.

Non-Example 3.110. Real semisimple groups need not be linear. For example, the universal cover of $\mathrm{SL}_2(\mathbb{R})$ is not linear, by the argument of Remark 3.100.

More generally, we have the following.

Proposition 3.111. Let $\mathfrak{g}_\theta$ be a real form of a complex semisimple Lie algebra $\mathfrak{g}$, and let G be a connected complex Lie group with Lie algebra $\mathfrak{g}$, and set $G_\theta := G^{\omega_\theta}$. Then G_θ° is linear, and every semisimple linear group is of this form.

Proof. We already know that G_θ° is linear because it lands in the semisimple complex group G, which is then reductive. For the converse, we basically repeat the proof of Remark 3.100. The idea is that any faithful representation of G_θ° goes down to the Lie algebra and then lifts to a representation of G^c. ■

3.6.2 Maximal Tori

Here is our definition.

Definition 3.112 (Cartan)**.** A *Cartan subalgebra* $\mathfrak{h}^c$ of $\mathfrak{g}^c$ is a maximal abelian subalgebra of $\mathfrak{g}^c$. We let H^c be the exponential of $\mathfrak{h}^c$ in G^c.

Remark 3.113. We claim that $\mathfrak{h}^c \subseteq \mathfrak{g}^c$ is a Cartan subalgebra if and only if $\mathfrak{h}^c_{\mathbb{C}}$ is a Cartan subalgebra of $\mathfrak{g}$. Indeed, all elements of $\mathfrak{g}^c$ are automatically semisimple, so $\mathfrak{h}^c_{\mathbb{C}}$ succeeds at being a torus in $\mathfrak{g}$, and maximality transfers between the $\mathfrak{g}^c$ and $\mathfrak{g} = \mathfrak{g}^c_{\mathbb{C}}$ because maximal just means that the dimension agrees with the rank.

Remark 3.114. Note that H^c is certainly a torus in G^c, and it is also maximal. In fact, H^c is maximal abelian subgroup: any such commuting element g needs to preserve H^c while also preserving the roots, and in fact, g must act by a scalar in $\mathrm{U}(1)$ on the root spaces. One can then undo the exponential as we did in Proposition 2.122 to show that $g \in H^c$.

Remark 3.115. As a sanity check, we note that $H := \exp(\mathfrak{h}^c_{\mathbb{C}})$ has $H \cap G^c = H^c$. Indeed, $H \cap G^c$ is some abelian subgroup containing H^c.

All these Cartan subalgebras are conjugate.

Lemma 3.116. Any two Cartan subalgebras in $\mathfrak{g}^c$, equipped with a set of positive roots, are conjugate by G^c_{ad}.

Proof. Choose two such pairs $(\mathfrak{h}, \Pi)$ and $(\mathfrak{h}', \Pi')$. Certainly there is some $g \in G_{\mathrm{ad}}$ with $\mathrm{Ad}_g(h, \Pi) = (h', \Pi')$, so it only remains to descend to G^c_{ad}, for which we use the polar decomposition. On one hand, $\omega(g) = \overline{g}$ also satisfies $\mathrm{Ad}_{\overline{g}}(\mathfrak{h}, \Pi) = (\mathfrak{h}', \Pi')$, so $\overline{g}^{-1}g$ commutes with $\mathfrak{h}^c$ while fixing Π. Thus, $\overline{g}^{-1}g$ must live in H, so $\overline{g}h = g$ for some $h \in H$.

On the other hand, Theorem 3.94 lets us write $g = kp$ for $k \in G^c$ and $p \in P$, and $\overline{g} = kp^{-1}$ due to the definition of K and P. It follows that

$$kp^{-1}h = kp,$$

which is then equivalent to $h = p^2$. Thus, p has positive eigenvalues, so h does as well, so we can take a square root of h (in P). Thus, p actually commutes with $\mathfrak{h}^c$ and preserves Π. We conclude that actually $\mathrm{Ad}_k(\mathfrak{h}, \Pi) = (\mathfrak{h}', \Pi')$. ■

Remark 3.117. This conjugacy property is special for the compact. For example, the split form $\mathfrak{sl}_2(\mathbb{R})$ has two different tori, given by $\mathfrak{so}_2(\mathbb{R})$ and the diagonal $\mathfrak{gl}_1(\mathbb{R})$.

Here is a nice structural result.

Theorem 3.118. Fix a connected compact Lie group K. Then every element is contained in a maximal torus, and all maximal tori are conjugate when equipped with a set of positive roots.

Proof. By Proposition 3.90, we may assume that K is semisimple, so K takes the form G^c for some semisimple G. The second statement now follows from Lemma 3.116 once we realize that $\mathfrak{h}^c \mapsto H^c$ is a bijection of Cartan subalgebras with maximal tori. Conversely, any maximal torus $T \subseteq K$ induces a maximal abelian subalgebra $\operatorname{Lie} T \subseteq \operatorname{Lie} K$, and the exponential of $\operatorname{Lie} T$ in K must recover T by Remark 3.114.

It remains to show the first statement. Choose a maximal torus T, and let $\varphi\colon K \times T \to K$ be the action map

$$\varphi(k,t) := ktk^{-1}.$$

Notably, the image of φ consists of those elements of K which can be conjugated into T, which the above paragraph explains is precisely those elements which are contained in a maximal torus. Notably, the source of φ is compact, so the image is closed, so $K \setminus \operatorname{im}\varphi$ is open.

It is thus enough to show that a dense subset of K can be found in a maximal torus. To this end, for $x \in K$, let $P_x(z-1)$ be the characteristic polynomial of Ad_x acting on $\mathfrak{k}$. Then the lowest power of z is the dimension of the centralizer $\mathfrak{z}_x \subseteq \mathfrak{g}$. Now, let $a(x)$ be the coefficient of $z^{\operatorname{rank}\mathfrak{g}}$ in $P_x(z)$. Note that this function is a polynomial in the coordinates of $\operatorname{GL}(\mathfrak{g})$, but a vanishes on $K \setminus \operatorname{im}\varphi$, so either $K \setminus \operatorname{im}\varphi$ is empty or $a = 0$.

Thus, it remains to show that a is nonzero. Well, for generic $X \in \mathfrak{g}^c$, one knows that X is regular, meaning that its centralizer $\mathfrak{z}_X$ has dimension $\operatorname{rank}\mathfrak{g}^c = \operatorname{rank}\mathfrak{g}$; this means that $\mathfrak{z}_X$ is a Cartan subalgebra. Thus, $x = \exp(X)$ commutes with the maximal torus $\exp(\mathfrak{z}_X)$, so $x \in \exp(\mathfrak{z}_X)$. ■

Corollary 3.119. Fix a connected compact Lie group K with Lie algebra $\mathfrak{k}$. Then the exponential map $\exp\colon \mathfrak{k} \to K$ is surjective.

Proof. This follows from Theorem 3.118 because our maximal tori were the image of an exponential by construction. ■

Remark 3.120. This is not true for complex reductive groups. For example, it is not true for $\operatorname{SL}_2(\mathbb{C})$, which we will show on the homework. As an aside, we note that the exponential is surjective for $\operatorname{GL}_n(\mathbb{C})$, which one can show using the Jordan normal form.

While we're here, let's say something about semisimple and unipotent elements.

Definition 3.121 (semisimple, unipotent)**.** Fix a connected complex reductive group G. Then $g \in G$ is *semisimple* (respectively, *unipotent*) if and only if the action of g on any representation is given by a semisimple (respectively, *unipotent*) operator.

Remark 3.122. On the homework, we will show that it is enough to act by a semisimple (respectively, unipotent) operator on a chosen faithful representation.

Remark 3.123. On the homework, we will show that there is a Jordan decomposition: any element $g \in G$ can be written uniquely as a product of a semisimple and a unipotent element.

Remark 3.124. It turns out that the exponential defines a homeomorphism between the nilpotent cone of $\mathfrak{g}$ and the unipotent subvariety of G.

3.6.3 Maximal Abelian Subspaces

We need to start by classifying maximal abelian subspaces of $\mathfrak{p}$.

Proposition 3.125. Fix a real form $\mathfrak{g}_\theta$ of a semisimple complex Lie algebra $\mathfrak{g}$, and let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the usual decomposition into eigenspaces of θ, and similarly $\mathfrak{g}^c = \mathfrak{k}^c \oplus \mathfrak{p}^c$ and $\mathfrak{g}_\theta = \mathfrak{k}^c \oplus \mathfrak{p}_\theta$. Lastly, let K^c be the Lie group of $\mathfrak{k}$ in G^c.

(a) Let $\mathfrak{a}$ be a maximal abelian subspace of $\mathfrak{p}_\theta$. Then the centralizer of $\mathfrak{z}$ of $\mathfrak{a}$ in $\mathfrak{g}^c$ takes the form $\mathfrak{m} \oplus \mathfrak{a}$, where $\mathfrak{m} \subseteq \mathfrak{k}^c$ is reductive. In fact, if $\mathfrak{t} \subseteq \mathfrak{m}$ is a Cartan subalgebra, then $\mathfrak{t} \oplus \mathfrak{a}$ is a Cartan subalgebra of $\mathfrak{g}_\theta$, and $\mathfrak{t} \oplus i\mathfrak{a}$ is a Cartan subalgebra of $\mathfrak{g}^c$.

(b) For generic $a \in \mathfrak{a}$, the centralizer of a in $\mathfrak{p}_\theta$ is $\mathfrak{a}$.

(c) For all $p \in \mathfrak{p}_\theta$, there is $k \in K^c$ for which $\operatorname{Ad}_k p \in \mathfrak{a}$.

(d) All maximal abelian subspace of $\mathfrak{p}_\theta$ are conjugate by K^c.

Proof. Here we go.

(a) Choose $X \in \mathfrak{z}$. Then we may write $X = X_+ \oplus X_-$ where $X_+ \in \mathfrak{k}^c$ and $X_- \in \mathfrak{p}_\theta$. Now, $[X_+, \mathfrak{a}] = 0$ automatically, so it follows that $[X_-, \mathfrak{a}] = 0$ as well, so $X_- \in \mathfrak{a}$ because $\mathfrak{a}$ is maximal. Thus, the collection of X_+ coordinates allow us to decompose $\mathfrak{z}$ into $\mathfrak{m} \oplus \mathfrak{a}$ for some Lie subalgebra $\mathfrak{m} \subseteq \mathfrak{k}^c$. Note that $\mathfrak{m}$ is reductive because the Killing form on $\mathfrak{k}$ restricts to a non-degenerate invariant bilinear form.

Lastly, choosing a Cartan subalgebra $\mathfrak{t} \subseteq \mathfrak{m}$, we see that $\mathfrak{t} \oplus i\mathfrak{a}$ is a maximal subalgebra of in $\mathfrak{g}^c$: indeed, anyone $X_+ + X_-$ commuting with $\mathfrak{t} \oplus i\mathfrak{a}$ has $X_- \in \mathfrak{a}$ as above, so $X_+ \in \mathfrak{m}$, but then $[X_+, \mathfrak{t}] = 0$ implies $X_+ \in \mathfrak{t}$ by its maximality. The argument that $\mathfrak{t} \oplus \mathfrak{a}$ is a Cartan subalgebra of $\mathfrak{g}_\theta$ is similar.

(b) Write $T_\mathfrak{a} := \exp(i\mathfrak{a})$, which is contained in G^c. Then $T_\mathfrak{a}$ is some compact torus, which looks like some power of S^1. For generic a, the one-parameter subgroup in $T_\mathfrak{a}$ is dense. Now, for $p \in \mathfrak{p}_\theta$ which commutes with a, we conclude that p commutes with $T_\mathfrak{a}$. Taking the differential implies that p commutes with $\mathfrak{a}$.

(c) This is the hardest part. Choose $a \in \mathfrak{a}$ generic as in (b). Thus, $\operatorname{Ad}_k(p) \in \mathfrak{a}$ is equivalent to $[\operatorname{Ad}_k(p), a] = 0$. The trick is to consider the function $f\colon K^c \to \mathbb{R}$ defined by

$$f(b) := (\operatorname{Ad}_b p, a).$$

This function is differentiable (because the inner product can be defined by the Killing form, which is algebraic), so we may select k so that f is maximized at $k \in K^c$; set $p' := \operatorname{Ad}_k p$, which we claim will do the trick. Well, the derivative df vanishes at k. On the other hand, for $x \in \mathfrak{k}^c$, this derivative $df_k(x)$ is $([x, p'], a)$, so we see that

$$(x, [p', a]) = 0$$

for all x, so $[p', a] = 0$ by non-degeneracy, so we are done!

(d) Choose maximal abelian subspaces $\mathfrak{a}$ and $\mathfrak{a}'$ of $\mathfrak{p}_\theta$, and choose a generic $p \in \dashv'$ so that $\mathfrak{a}'$ is the centralizer of p. By (c), there is $k \in K^c$ with $\operatorname{Ad}_k p \in \mathfrak{a}$. Then Ad_k sends the centralizer of p to the centralizer of $\operatorname{Ad}_k p$, so it maps $\mathfrak{a}'$ into $\mathfrak{a}$. In other words, $\operatorname{Ad}_k \mathfrak{a}' \subseteq \mathfrak{a}$. For example, this implies that $\dim \mathfrak{a}' \leq \dim \mathfrak{a}$; reversing the argument shows that the dimensions are equal, so we conclude that $\operatorname{Ad}_k \mathfrak{a}' = \mathfrak{a}$. ■

3.7 April 22

Today, we discuss the Cartan decomposition.

3.7.1 The Cartan Decomposition

Here is our statement.

Theorem 3.126. Fix a real form $\mathfrak{g}_\theta$ of a semisimple complex Lie algebra $\mathfrak{g}$, and let $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$ be the decomposition into θ-eigenspaces. We then set $\mathfrak{g}_\theta = \mathfrak{k}^c \oplus \mathfrak{p}_\theta$. Let $G_\theta \subseteq G_{\mathrm{ad}}$ be the invariants, and we define $K^c \subseteq G_\theta$ and $A \subseteq G_\theta$ to be the subgroup corresponding $\mathfrak{k}^c$ and to a maximal abelian subspace $\mathfrak{a} \subseteq \mathfrak{p}_\theta$, respectively. Then

$$G_\theta = K^c A K^c.$$

Proof. The polar decomposition (Theorem 3.94) grants $G_\theta = K^c P_\theta$, so it is enough to show that every element in P_θ can be found in $K^c A K^c$. We will show that every element in P_θ can be conjugated by K^c into A. Well, choose $y \in P_\theta$, which we may write as $\log Y$ for some unique $Y \in \mathfrak{p}_\theta$. By Proposition 3.125, we find that there is $k \in K^c$ with $\mathrm{Ad}_k\, y \in \mathfrak{a}$. Taking the exponential shows that $\mathrm{Ad}_k\, Y \in A$, so we are done. ■

Remark 3.127. There is a generalization to reductive groups, which is no harder to prove.

Remark 3.128. The decomposition is not unique.

Example 3.129. Take $G_\theta = \mathrm{GL}_n(\mathbb{C})$. Then $K = \mathrm{U}(n)$ and A can be taken to be the subgroup of diagonal operators with positive eigenvalues. Thus, we are saying that any $g \in \mathrm{GL}_n(\mathbb{C})$ can be decomposed as $u_1 d u_2$ where $u_1, u_2 \in \mathrm{U}(n)$ and $d \in A$.

Example 3.130. Take $G_\theta = \mathrm{GL}_n(\mathbb{R})$. Then $K = \mathrm{O}(n)$ and A can be taken to be the subgroup of diagonal operators with positive eigenvalues. Thus, we are saying that any $g \in \mathrm{GL}_n(\mathbb{R})$ can be decomposed as $o_1 d o_2$, where $o_2, o_2 \in \mathrm{O}(n)$ and $d \in A$.

3.7.2 Maximal Compact Subgroups

We are now able to prove the most important fact about the K^cs: they are maximal compact subgroups, and with this property, they are unique up to conjugation.

Theorem 3.131 (Cartan)**.** Let G_θ be a real form of a connected semisimple complex Lie group G. Then any compact subgroup $L \subseteq G^\theta$ is conjugate (by G_θ) to a subgroup of K^c.

Note that this gives the following promised corollary.

Corollary 3.132. Let G_θ be a real form of a connected semisimple complex Lie group G. Then all maximal compact subgroups in G_θ are conjugate to K^c.

Proof. This follows from Theorem 3.131: any maximal compact subgroup L is conjugate to a subgroup of K^c, but the conjugate must remain maximal, so L must be conjugate to K^c on the nose. (Note that K^c is compact because it is closed in G^c.) ■

We now prove Theorem 3.131, following Mostow. We start by noticing that it is not actually too hard to show the maximality of K^c.

Theorem 3.133. Let G_θ be a real form of a connected semisimple complex Lie group G. Then K^c is a maximal compact subgroup.

Proof. Suppose that L is a compact subgroup containing K^c. By the polar decomposition, any $k \in K$ can be decomposed into K^c and P_θ, but then the element in P_θ must still be in L. In other words, $L = K^c(P_\theta \cap L)$.

We now aim to show that $P_\theta \cap L$ is trivial, which will complete the proof. Well, choose some $y \in P_\theta \cap L$. Then the eigenvalues of y act on $\mathfrak{g}$ by positive Hermitian operators. Thus, if any of the eigenvalues are not one, then at least one of the eigenvalues is bigger than one, so the eigenvalues of $\{y^n\}_{n\geq 1}$ escape to ∞, so there is no convergent subsequence of the image of these elements in $\mathrm{GL}(\mathfrak{g})$, which contradicts the compactness of L! Thus, $y = 1$, and we are done. ■

Let's now turn to the rest of the proof of Theorem 3.131

Proof of Theorem 3.131. We proceed in steps. Choose a compact subgroup $L \subseteq G_\theta$, and we want to show that it is conjugate to a subgroup of K^c by G_θ. We may as well show that L is conjugate by P_θ by Theorem 3.94. The idea is to construct an L-invariant function $f\colon P_\theta \to \mathbb{R}$ with a unique minimum $\exp(Y) \in P_\theta$, and this minimum will encode the correct conjugate. Indeed, suppose we have such a function f. Then because f is L-invariant, its unique minimum $y \in P_\theta$ must be an L-invariant point. Thus, for all $h \in L$, we have $h\exp(Y) = \exp(Y)\overline{h}$. This rearranges into

$$\exp(-Y/2)h\exp(Y/2) = \exp(Y/2)\overline{h}\exp(-Y/2),$$

which implies that $\exp(-Y/2)h\exp(Y/2)$ is conjugate to itself, meaning that it is in K.

It now remains to hunt for this function f.

1. We set some notation. Let B be the Killing form on $\mathfrak{g}$, restricted to $\mathfrak{g}_\theta$; thus, it is negative-definite on $\mathfrak{k}^c$ and positive-definite on $\mathfrak{p}_\theta$. We then define B_θ by

$$B_\theta(x, y) := -B(x, \theta y),$$

which is positive-definite on $\mathfrak{g}_\theta$ and K^c-invariant (because K^c is θ-invariant). Now, for an operator A on $\mathfrak{g}_\theta$, we let $A^\dagger$ be the adjoint under B_θ. Note the following special cases.

- For $g \in K^c$, the operator Ad_g is orthogonal because it preserves B_θ.
- For $g \in P_\theta$, the operator Ad_g is self-adjoint and has positive eigenvalues.

Accordingly, for $g = kp$ with $k \in K^c$ and $p \in P_\theta$ (given by Theorem 3.94), we find that

$$\begin{aligned}\mathrm{Ad}_g^\dagger &= \mathrm{Ad}_p^\dagger \mathrm{Ad}_k^\dagger \\ &= \mathrm{Ad}_p \mathrm{Ad}_{k^{-1}} \\ &= \mathrm{Ad}_{\overline{g}}^{-1}\end{aligned}$$

by construction of the complex conjugate.

2. We define the function f. To start, we define

$$S := \int_L \mathrm{Ad}_h^\dagger \mathrm{Ad}_h \, dh,$$

which is some endomorphism of $\mathfrak{g}_\theta$. Note that $(\cdot)^\dagger$ is linear on operators, so $S^\dagger = S$. Additionally, S is positive because it is an average of positive operators. Thus, S can be given an orthonormal eigenbasis $\{v_i\}$ with positive eigenvalues $\{\lambda_i\}$. We let λ be the minimal eigenvalue and then define $f\colon P_\theta \to \mathbb{R}$ by

$$f(x) := \mathrm{tr}(\mathrm{Ad}_x \circ S).$$

Note that taking the trace can be computed as

$$f(x) = \sum_i \lambda_i B_\theta(\mathrm{Ad}_x v_i, v_i).$$

Because Ad_x is positive and self-adjoint, this is then bounded below by $\lambda \operatorname{tr} \mathrm{Ad}_x$. While we're here, we also note that f is L-invariant because G_θ acts on P_θ by the twisted adjoint action $g \cdot x = gx\overline{g}^{-1}$.

3. We show that f has a minimum $y \in P_\theta$. Well, for any $R > 0$, note that the set

$$\{A \in M_n(\mathbb{R}) : A = A^\intercal, \det A = 1, \operatorname{tr} A \leq R\}$$

is compact. Indeed, up to the orthogonal sets $\mathrm{O}_n(\mathbb{R})$, any such matrix A can be diagonalized with positive entries by Example 3.130, but then the diagonal entries have positive coefficients and are bounded above by R (with product 1). Thus, the collection of relevant diagonal matrices is compact, so the entire set is compact.

Thus, for large R, we conclude that the set of X with $f(X) \leq R$ is a compact set because they are some self-adjoint operators with bounded trace (and determinant 1 on $\mathfrak{g}$). We conclude that f attains a minimum on P_θ.

4. As an intermission, we discuss a small piece of linear algebra: given symmetric real matrices a and M with a nonzero and M positive-definite. Then we claim that the function

$$\varphi(t) \coloneqq \operatorname{tr}(\exp(ta)M)$$

is strictly convex. Indeed, by changing basis, we may assume that a is a diagonal matrix with diagonal entries $\{a_i\}_i$. Then

$$\operatorname{tr}(\exp(ta)) = \sum_i \exp(ta_i)M_{ii}.$$

Because M is positive-definite, the entries M_{ii} are positive, so this is a positive linear combination of strictly convex functions and hence strictly convex.

5. We now show that f admits a unique minimum $y \in P_\theta$. Suppose that we have another minimum $z \in P_\theta$ of f. Set $Y \coloneqq \log y$ and $Z \coloneqq \log z$, and define $g \coloneqq \exp(Z/2)\exp(-Y/2)$. We would like to show that $g = 1$, which will imply that $Z/2 = Y/2$ by taking logarithms, which then implies $y = z$ by taking exponentials.

To show that $g = 1$, we use the polar decomposition: by Theorem 3.94, we have $k \in K^c$ and $X \in \mathfrak{p}_\theta$ with

$$\exp\left(\frac{Z}{2}\right)\exp\left(-\frac{Y}{2}\right) = k\exp\left(\frac{X}{2}\right).$$

Thus,

$$\exp\left(\frac{X}{2}\right) = k^{-1}\exp\left(\frac{Z}{2}\right)\exp\left(-\frac{Y}{2}\right).$$

Because $\exp(X/2)$ is self-adjoint, we also see that

$$\exp\left(\frac{X}{2}\right) = \exp\left(-\frac{Y}{2}\right)\exp\left(\frac{Z}{2}\right)k,$$

so multiplying gives

$$\exp(X) = \exp\left(-\frac{Y}{2}\right)\exp(Z)\exp\left(-\frac{Y}{2}\right),$$

which rearranges to

$$\exp(Z) = \exp\left(-\frac{Y}{2}\right)\exp(X)\exp\left(-\frac{Y}{2}\right).$$

We now consider the function

$$g(t) := f\left(\exp\left(-\frac{Y}{2}\right)\exp(tX)\exp\left(-\frac{Y}{2}\right)\right).$$

By construction, this function attains minima at $t = 0$ and $t = 1$. On the other hand,

$$g(t) = \operatorname{tr}\left(\operatorname{Ad}_{\exp(-Y/2)} \circ \operatorname{Ad}_{\exp(tX)} \circ \operatorname{Ad}_{\exp(-Y/2)} \circ S\right),$$

which is

$$g(t) = \operatorname{tr}\left(\operatorname{Ad}_{\exp(tX)} \circ \left(\operatorname{Ad}_{\exp(-Y/2)} \circ S \circ \operatorname{Ad}_{\exp(Y/2)}\right)\right)$$

takes the form of the functions in the previous step (recall S is self-adjoint and positive), so it is strictly convex (with a unique minimum) as soon as X is nonzero. Thus, we instead must have $X = 0$, so $g \in K^c$. But then $\exp(Z/2) = k\exp(Y/2)$, so the polar decomposition forces $k = 1$, so we are done. ■

3.7.3 Cartan Subalgebras

We now take a moment to discuss Cartan subalgebras.

Definition 3.134 (Cartan)**.** A *Cartan subalgebra* of a real form $\mathfrak{g}_\theta$ is a maximal abelian subalgebra consisting of semisimple elements.

Remark 3.135. By construction, one finds that $\mathfrak{h}$ is a Cartan subalgebra if and only if $\mathfrak{h}_\mathbb{C}$ is a Cartan subalgebra of $\mathfrak{g}_\theta$. The main point is that any maximal commutative subalgebra $\mathfrak{h}$ with semisimple elements, and we want to show that $\mathfrak{h}_\mathbb{C}$ continues to be maximal. Indeed, if $X \in \mathfrak{g}$ commutes with $\mathfrak{h}_\mathbb{C}$, then so does $\overline{X}$, so the real part $(X + \overline{X})/2$ and the imaginary part $(X - i\overline{X})/2i$ are both in $\mathfrak{h}$, so $X \in \mathfrak{h}_\mathbb{C}$.

Example 3.136. For $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{R})$, choose some nonnegative integer $m \le n/2$. Then there is a Cartan subalgebra $\mathfrak{h}_m$ consisting of the block-diagonal matrices

$$\left\{\operatorname{diag}\left(\begin{bmatrix} a_1 & b_1 \\ -b_1 & a_1 \end{bmatrix}, \dots, \begin{bmatrix} a_m & b_m \\ -b_m & a_m \end{bmatrix}, c_1, \dots, c_{n-2m}\right) : 2(a_1 + \cdots + a_m) + (c_1 + \cdots + c_{n-2m}) = 0\right\}.$$

On the homework, we will show that these Cartan subalgebras exhaust all of them up to isomorphism.

Let's give some invariants to classify Cartan subalgebras.

Definition 3.137 (split)**.** Fix a real form $\mathfrak{g}_\theta$. A semisimple element $X \in \mathfrak{g}_\theta$ is *split* if and only if the adjoint action of X on $\mathfrak{g}_\theta$ has real eigenvalues. A commutative subalgebra $\mathfrak{h}$ of $\mathfrak{g}_\theta$ is *split* if and only if all elements of $\mathfrak{h}$ are split; for any commutative subalgebra $\mathfrak{h}$, we let $\mathfrak{h}^{\mathrm{spl}}$ be the subspace of split elements, and we define $s(\mathfrak{h}) := \dim \mathfrak{h}^{\mathrm{spl}}$.

Remark 3.138. The commutativity implies that $\mathfrak{h}^{\mathrm{spl}}$ is in fact a subspace.

Remark 3.139. Note that $\mathfrak{h}$ is split if and only if $\mathfrak{h}^{\mathrm{spl}} = \mathfrak{h}$, which is equivalent to $s(\mathfrak{h}) = \dim \mathfrak{h}$.

Remark 3.140. One can check that the signature of the Killing form restricted to $\mathfrak{h}$ is $(\dim \mathfrak{h} - s(\mathfrak{h}), s(\mathfrak{h}))$.

Remark 3.141. One can check that $\exp(\mathfrak{h}) \cong \left(S^1\right)^{\dim \mathfrak{h} - s(\mathfrak{h})} \times \left(\mathbb{R}^+\right)^{s(\mathfrak{h})}$.

Remark 3.142. It turns out that $\mathfrak{g}_\theta$ is split if and only if it admits a split Cartan subalgebra. On one hand, being the split form allows the usual $\mathfrak{h}$, which is split. Conversely, having a split Cartan means that its root decomposition can be taken to have real eigenvalues everywhere (by definition) and thus fully defined over $\mathbb{R}$, which allows us to present $\mathfrak{g}_\theta$ as the split real form.

It is interesting to determine what values of $s(\mathfrak{h})$ are possible.

Definition 3.143. Fix a real form $\mathfrak{g}_\theta$. Then $\mathfrak{h}$ is *maximally split* if and only if $s(\mathfrak{h})$ is as large as possible. Similarly, $\mathfrak{h}$ is *maximally compact* if and only if $s(\mathfrak{h})$ is as small as possible.

Example 3.144. For $\mathfrak{sl}_n(\mathbb{R})$, the diagonal torus $\mathfrak{h}_0$ is maximally split, and the diagonal torus $\mathfrak{h}_m$ is maximally compact.

Remark 3.145. Any compact Cartan subalgebra has $s(\mathfrak{h})$ equal to the rank, which is automatically maximally compact.

We will show the following result next time.

Proposition 3.146. Fix a real form $\mathfrak{g}_\theta$, and let $\mathfrak{h}$ be a θ-stable Cartan subalgebra, and let $\mathfrak{g}_\theta = \mathfrak{k}^c \oplus \mathfrak{p}_\theta$ be the polar decomposition.

(a) Then $\mathfrak{h}$ is maximally split if and only if $\mathfrak{h} \cap \mathfrak{p}_\theta$ is a maximal abelian subspace of $\mathfrak{p}_\theta$

(b) Then $\mathfrak{h}$ is maximally compact if and only if $\mathfrak{h} \cap \mathfrak{k}^c$ is a Cartan subalgebra.

(c) Any two maximally split Cartan subalgebras are conjugate.

(d) Any two maximally compact Cartan subalgebras are conjugate.

(e) Any Cartan subalgebra is conjugate to a θ-stable one (by G_θ).

Proof. Omitted. It's present in the notes. ■

Theme 4
Cohomological Methods

4.1 April 27

Here we go.

4.1.1 The Weyl Integration Formula

As an application, we note that we can reformulate the Weyl character formula.

Theorem 4.1 (Weyl integration formula)**.** Fix a connected compact Lie group K with maximal torus $T \subseteq K$. Let Φ be the root system of $\operatorname{Lie} K_{\mathbb{C}}$, and let dx and dt be the probability Haar measures on K and T, respectively. Given a conjugation-invariant continuous function f on K, we have

$$\int_K f(x)\,dx = \frac{1}{|W|}\int_T f(t)\,|\Delta(t)|^2\,dt,$$

where

$$\Delta(t) = \prod_{\alpha\in\Phi_+}\left(\alpha(t)^{1/2} - \alpha(t)^{-1/2}\right).$$

Remark 4.2. Note that $\Delta(t)$ is defined only up to sign (because of the square roots), but we are taking an absolute value later anyway, so this does not matter.

Proof. By the Peter–Weyl theorem, via its application Corollary 2.52, it is enough to check this for characters χ_λ, where λ is dominant integral. We now compute both sides directly.

- On one hand,

$$\int_K \chi_\lambda(x)\,dx = \begin{cases} 0 & \text{if } \chi \text{ is nontrivial,} \\ 1 & \text{if } \chi \text{ is trivial} \end{cases}$$

  by the orthogonality of characters (also proven in Corollary 2.52).

- On the other hand, by the Weyl character formula (Theorem 1.116), we have

$$\frac{1}{|W|}\int_T \chi_\lambda(t)\,|\Delta(t)|^2\,dt = \frac{1}{|W|}\int_T \left(\sum_{w\in W}(-1)^{\ell(w)}e^{w(\lambda+\rho)}\right)\left(\sum_{y\in W}(-1)^{\ell(y)}e^{-y\rho}\right)dt.$$

Here, e^α refers to the character $\alpha\colon T \to \mathbb{C}^\times$. Now, $\int_T e^\gamma\, dt$ vanishes unless $\gamma = 0$, in which case the integral is 1. Now, the only way to receive characters e^0 in this sum is to have $w(\lambda+\rho) = y\rho$, which is only possible when $w = y$ and when $\lambda = 0$. Thus, the integral vanishes when $\lambda \neq 0$, and when $\lambda = 0$, we receive $\frac{1}{|W|} \cdot |W|$, which is 1.

Comparing our two calculations completes the proof. ■

Example 4.3. For $K = \mathrm{U}(n)$, we can set T to be the diagonal torus. We can parameterize the torus as $\mathrm{U}(1)^n$. Then expanding out Δ reveals that

$$\int_{\mathrm{U}(n)} f(x)\, dx = \frac{1}{n!} \int_{\mathrm{U}(1)^n} \left(f(\mathrm{diag}(z_1, \ldots, z_n)) \prod_{k<\ell} (z_k - z_\ell) \right) dz_1 \cdots dz_n.$$

One can write $z_1 = e^{i\theta_1}$, which gives

$$\int_{\mathrm{U}(n)} f(x)\, dx = \frac{1}{(2\pi)^n n!} \int_{(\mathbb{R}/\mathbb{Z})^n} \left(f(\mathrm{diag}(e^{i\theta_1}, \ldots, e^{i\theta_n}) \prod_{k<\ell} 2\sin(\theta_k - \theta_\ell) \right) d\theta_1 \ldots d\theta_n.$$

The factor of 2π in the denominator comes from the probability Haar measure on S^1.

Remark 4.4. By the orthogonality of characters (in Corollary 2.52), we see that

$$\frac{1}{|W|} \int_T \chi_\lambda(t) \overline{\chi_\mu(t)}\, |\Delta(t)|^2\, dt = 1_{\lambda=\mu}.$$

4.1.2 Review of de Rham Cohomology

Recall the following construction.

Definition 4.5 (de Rham complex)**.** Fix a manifold M of dimension n. For each i, we define Ω^i to be the sheaf $\wedge^i \mathrm{T}^* M$. Then we define the *de Rham complex* to be the complex

$$0 \to \Omega^0(M) \to \Omega^1(M) \to \Omega^2(M) \to \cdots \to \Omega^n(M) \to 0.$$

Here, the maps $d\colon \Omega^i(M) \to \Omega^{i+1}(M)$ are given by the de Rham differential, which can be described on pure tensors as $d(f\,\omega) := df \wedge \omega$.

Remark 4.6. One can check that $d^2 = 0$, which explains why this is a complex.

Remark 4.7. The sheaves $\Omega^i(M)$ can be glued into $\Omega^*(M)$, which is a graded commutative ring under the wedge product $\wedge$.

Remark 4.8. One can check that

$$d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^{\deg \alpha} \alpha \wedge d\beta$$

by passing to local coordinates and doing an explicit calculation. Indeed, this is basically the (Leibniz) product rule after keeping track of ordering. Thus, d is a derivation.

Definition 4.9 (de Rham cohomology)**.** Fix a manifold M of dimension n. Then we say that $\omega \in \Omega^i(M)$ is *closed* if and only if $d\omega = 0$ and is *exact* if and only if ω is in the image of d. We then define $\mathrm{H}^i(M;\mathbb{C})$ to be the space of closed forms modulo the space of exact forms.

Remark 4.10. Everything is functorial in the manifold: a smooth map $f\colon M \to N$ produces morphisms $f^*\colon \Omega^i(N) \to \Omega^i(M)$ which commute with the de Rham differential and therefore induce morphisms in de Rham cohomology. We will show on the homework that the induced morphism f^* on cohomology do not depend on the smooth homotopy class of f.

Remark 4.11. It is a theorem that de Rham cohomology agrees (functorially) with singular cohomology.

4.1.3 Group Actions on Cohomology

We will want the following tool.

Definition 4.12 (Lie derivative)**.** Fix a vector field v on a manifold M. Then the *Lie derivative* L_v is the unique degree-zero derivation of $\Omega^*(M)$ for which $[L_v, d] = 0$ and $L_v f$ is the directional derivative of f in the direction v.

Remark 4.13. Note that L_v is uniquely determined by its behavior in degree zero because any differential form is locally exact, so we can always pull a d out of a differential form locally.

Proposition 4.14 (Cartan's magic formula)**.** Fix a vector field v on a manifold M, and let $i_v\colon \Omega^i \to \Omega^{i-1}$ denote the contraction by v. Then

$$L_v = di_v + i_v d.$$

Proof. One can check that i_v is a derivation of degree -1, so both sides are derivations of degree 0. Additionally, both sides commute with d, so they are uniquely determined by their behavior in degree zero. This amounts to checking that $L_v f = i_v df$, which is basically the definition of L_v. ■

Corollary 4.15. Fix a vector field v on a manifold M. Then the induced operation L_v on $\mathrm{H}^i(M;\mathbb{R})$ vanishes.

Proof. Here, the operation L_v on cohomology is well-defined because L_v preserves the complex (and notably commutes with d). Proposition 4.14 tells us that L_v sends closed forms to exact forms. ■

Corollary 4.16. Fix a connected Lie group G acting on a manifold M. Then the action of G on $\mathrm{H}^i(G;M)$ is by the identity.

Proof. Fix $g \in G$, and we would like to show that the action map $g\colon M \to M$ induces the identity on cohomology. In other words, for any closed differential form ω, we would like to check that $g^*\omega - \omega$ is exact. Well, because G is connected, we may find a path $\gamma\colon [0,1] \to G$ with $\gamma(0) = 1$ and $\gamma(1) = g$. Then by Stoke's formula, we have

$$g^*\omega - \omega = \int_0^1 \frac{d}{dt}\gamma(t)^*\omega\, dt,$$

which could alternatively be checked directly in local coordinates. However, by Proposition 4.14, this is

$$g^*\omega - \omega = -\int_0^1 L_{\gamma'(t)\gamma(t)^{-1}}\omega\, dt,$$

and the right-hand side vanishes in cohomology by Corollary 4.15. Here, $\gamma'(t)\gamma(t)$ refers to the invariant vector field of $\gamma'(t)$. ■

Remark 4.17. Intuitively, the previous remark is saying that $g\colon M \to M$ is homotopic to the identity, which would then finish the proof by Remark 4.10. However, this does not really work because the relevant homotopy exists in $\mathrm{Diff}(M)$, which is not actually a manifold.

We can get further mileage from our integration theory in the case where G is compact.

Notation 4.18. Suppose that a compact Lie group G acts on a manifold M. Then we define the operator P on $\Omega^i(M)$ by

$$P\omega := \int_G g^*\omega\, dg,$$

a formula which can be made sense of in local coordinates. (Here, dg is the probability Haar measure.) Lastly, we define $\Omega^i(M)_0$ to be the image of $1 - P$.

Remark 4.19. By construction, $P^2 = P$, and P outputs G-invariant forms while acting by identity on G-invariant forms, so it follows that $\operatorname{im} P = \Omega^i(M)^G$. Because $P^2 = P$, we further receive a decomposition

$$\Omega^i(M) = \Omega^i(M)^G \oplus \Omega^i(M)_0.$$

Remark 4.20. Furthermore, the action of G commutes with d (because G acts by smooth maps), so P commutes with d. Thus, P induces an action on the de Rham complex, and $\Omega^i(M) = \Omega^i(M)^G \oplus \Omega^i(M)_0$ becomes a sum of complexes.

Proposition 4.21. Suppose that a compact Lie group G acts on a manifold M. The complex $\{\Omega^i(M)_0\}_i$ is exact.

Proof. Choose a closed form $\omega \in \Omega^i(M)_0$. Then Corollary 4.16 shows that $[\omega] = g^*[\omega]$, which then equals $[Pw]$ by integration. But $P\omega = 0$, so $[P\omega] = 0$, so $[\omega] = 0$, so $\omega = d\eta$ for some η. Further, because d commutes with P, we find that

$$\begin{aligned}\omega &= (1-P)\omega \\ &= (1-P)d\eta \\ &= d(1-P)\eta,\end{aligned}$$

so ω is in the image of $d\colon \Omega^i(M)_0 \to \Omega^i(M)_0$. ■

Remark 4.22. It follows that just the complex $\left\{\Omega^i(M)^G\right\}_G$ computes the cohomology of M.

4.1.4 The Cohomology of a Lie Group

By Remark 4.22, we can compute the cohomology of a compact connected Lie group G via left-invariant differential forms on G, which are given by $\wedge^\bullet \mathfrak{g}^*$.

Next, we would like to compute the differential on $\wedge^\bullet \mathfrak{g}$ purely algebraically. We will use another tool from differential geometry.

Proposition 4.23 (Cartan differentiation formula)**.** Fix a differential form $\omega \in \Omega^m(V)$ and vector fields $v_0, \ldots, v_m$. Then

$$d\omega(v_0, \ldots, v_m) = \sum_{i=0}^{m} (-1)^i L_v \omega(v_0, \ldots, \widehat{v}_i, \ldots, v_m) + \sum_{0 \le i < j \le m} (-1)^{i+j} \omega([v_i, v_j], v_1, \ldots, \widehat{v}_i, \ldots, \widehat{v}_j, \ldots, v_m)$$

Proof. We may pass to local coordinates. Then the right-hand side is linear in functions on M, so it suffices to compare both sides when we set $v_i = \partial/\partial x_i$ for each i and $\omega = f\, dx_{j_1} \wedge \cdots \wedge dx_{j_m}$. Then one directly computes both sides and concludes. ■

Corollary 4.24. Fix a Lie group G and some G-invariant differential form $\omega \in \Omega^m(G)^G$. Then for all left-invariant vector fields $v_1, \ldots, v_m \in \mathfrak{g}$, we have

$$d\omega(v_0, \ldots, v_m) = \sum_{0 \le i < j \le m} (-1)^{i+j} \omega([v_i, v_j], v_1, \ldots, \widehat{v}_i, \ldots, \widehat{v}_j, \ldots, v_m).$$

Proof. The functions $\omega(v_0, \ldots, \widehat{v}_i, \ldots, v_m)$ on G are G-invariant and thus constant, so their Lie derivatives vanish. Thus, this follows from Proposition 4.23. ■

We are now ready to define cohomology of a Lie algebra.

Definition 4.25 (Chevalley–Eilenberg complex)**.** Fix a Lie algebra $\mathfrak{g}$ over a field F. Then we define the *Chevalley–Eilenberg complex* to be the graded commutative ring $\wedge^\bullet \mathfrak{g}^*$, equipped with the differential

$$d\omega(v_0, \ldots, v_m) = \sum_{0 \le i < j \le m} (-1)^{i+j} \omega([v_i, v_j], v_1, \ldots, \widehat{v}_i, \ldots, \widehat{v}_j, \ldots, v_m).$$

We define $\mathrm{H}^i(\mathfrak{g}; F)$ to be the cohomology spaces associated to this complex.

Remark 4.26. One can check directly with the Jacobi identity that this differential d has $d^2 = 0$. Thus, this is actually a complex.

Example 4.27. Fix a compact connected Lie group G with Lie algebra $\mathfrak{g}$. By Remark 4.22 and corollary 4.24, we find that $\mathrm{H}^i(G; \mathbb{C})$ is isomorphic to $\mathrm{H}^i(\mathfrak{g}; \mathbb{C})$: indeed, these are isomorphic on the level of the complexes (before taking cohomology)!

Example 4.28. Suppose that $G = \mathbb{R}^m$ so that $\mathfrak{g}_\mathbb{C} = \mathbb{C}^m$. Then the differential vanishes, so $\mathrm{H}^i(\mathfrak{g}; \mathbb{C}) = \wedge^i \mathfrak{g}^*$. On the other hand, $\mathrm{H}^i(G; \mathbb{C})$ always vanishes because G is contractible!

Let's use some algebra to compute our cohomology.

Proposition 4.29. Fix a compact connected Lie group G. Then

$$\mathrm{H}^\bullet(G;\mathbb{C}) \cong (\wedge^\bullet \mathfrak{g}^*)^{\mathfrak{g}}.$$

Proof. Note that $G \times G$ acts on G, where the left G acts by left translation, and the right G acts by the adjoint action. By Remark 4.22, we see that we can compute $\mathrm{H}^\bullet(G;\mathbb{C})$ is computed by the complex

$$\Omega^\bullet(G)^{G\times G} = (\wedge^\bullet \mathfrak{g}^*)^G,$$

where this last G acts by the adjoint. Because G is connected, $(\wedge^\bullet \mathfrak{g}^*)^G = (\wedge^\bullet \mathfrak{g}^*)^{\mathfrak{g}}$, so it only remains to compute that the differential on this subcomplex vanishes. Well,

$$d\omega(v_0,\ldots,v_m) = \sum_{0\le i<j\le m} (-1)^{i+j}\omega([v_i,v_j],v_1,\ldots,\widehat{v}_i,\ldots,\widehat{v}_j,\ldots,v_m),$$

which can be decomposed into a sum of terms which look like

$$S_i = \sum_{j\ne i} \omega(v_1,\ldots,[v_i,v_j],\ldots,\widehat{v}_i,\ldots,v_m),$$

up to some signs and a factor of $\frac{1}{2}$, where here $[v_i,v_j]$ lives in coordinate j. However, S_i vanishes because ω is $\mathfrak{g}$-invariant! ■

While we're here, we give another invariant property.

Proposition 4.30. Fix a connected Lie group G, and let $\Gamma \subseteq G$ be a finite subgroup. Then the projection $\pi\colon G \to G/\Gamma$ induces an isomorphism on cohomology.

Proof. To begin, we claim that $\mathrm{H}^i(G/\Gamma;\mathbb{C}) \cong \mathrm{H}^i(G;\mathbb{C})^\Gamma$: indeed, the point is that differential forms on G/Γ are the same as Γ-invariant differential forms on G.[1] However, G acts trivially on its own cohomology, so Γ acts trivially (for example, by Remark 4.22), so we are done. ■

Remark 4.31. This is not true for torsion coefficients.

4.1.5 Using Hopf Algebras

For motivation, we recall the following piece of differential topology.

Theorem 4.32. Fix smooth manifolds M and N. Then the projections define an isomorphism

$$\mathrm{H}^*(M\times N;\mathbb{C}) \to \mathrm{H}^*(M)\otimes \mathrm{H}^*(N).$$

Proof. This is hard, so it is omitted. Unsurprisingly, this is true on the level of complexes. ■

Remark 4.33. Importantly, this is an isomorphism of graded commutative rings, but this is a little twisted on the right-hand side: one has

$$(a\otimes b)(c\otimes d) = (-1)^{\deg(b)\deg(c)}(ac\otimes bd).$$

Intuitively, this is because we have to swap b and c.

[1] Indeed, this identity remains true whenever $\pi\colon M \to M'$ is a finite covering space where Γ is the group of deck transformations.

Corollary 4.34. Fix a connected Lie group G. Then $\mathrm{H}^*(G;\mathbb{C})$ has the structure of a graded commutative Hopf algebra.

Proof. The algebra structure comes from the wedge product in cohomology. The augmentation map ε is the projection onto degree zero (note that G is connected), and the antipode is induced by the inverse map on G. Furthermore, the coproduct Δ is induced by the multiplication $m\colon G\times G\to G$, together with Theorem 4.32; explicitly, Δ is the composite

$$\mathrm{H}^*(G;\mathbb{C}) \xrightarrow{m} \mathrm{H}^*(G\times G;\mathbb{C}) \simeq \mathrm{H}^*(G;\mathbb{C})\otimes \mathrm{H}^*(G;\mathbb{C}).$$

Because these maps make G into a group object in the category of manifolds, they induced pullbacks which show that $\mathrm{H}^*(G;\mathbb{C})$ is a group object in the opposite category of algebras, which is exactly a Hopf algebra. ■

Such Hopf algebras admit a classification.

Theorem 4.35 (Hopf)**.** Fix a field k of characteristic zero. Suppose that R is a finite-dimensional graded commutative bialgebra over k. Then R is free as a k-algebra in generators of odd degree. Explicitly, there is an isomorphism

$$R \cong \wedge^\bullet(\xi_1,\dots,\xi_r),$$

where the right-hand side is the exterior algebra in the generators ξ_i, which all have odd degrees.

Proof. This is on the homework! ■

Remark 4.36. The presence of even-degree elements generate polynomial algebras.

Thus, Theorem 4.35 implies that $\mathrm{H}^*(G;\mathbb{C})$ is some exterior algebra. Let's see what we can say about it.

Proposition 4.37. Fix a compact connected Lie group G with Lie algebra $\mathfrak{g}$. Then $\mathrm{H}^\bullet(G;\mathbb{C})$ is isomorphic (as a $\mathbb{C}$-algebra) to a free exterior algebra on $\mathrm{rank}(\mathfrak{g})$ generators.

Proof. The fact that $\mathrm{H}^\bullet(G;\mathbb{C})$ is isomorphic to a free exterior algebra follows from Theorem 4.35 (together with Corollary 4.34). It remains to compute the number of generators, which we will call r for the time being. Note that

$$2^r = \dim\left(\wedge^\bullet\mathfrak{g}^*\right)^{\mathfrak{g}}$$

by Example 4.27, so it remains to show that the right-hand side equals $2^{\mathrm{rank}(\mathfrak{g})}$.

We will do this using the Weyl character formula. To begin, note that we can explicitly compute the character of $\wedge^\bullet\mathfrak{g} = \wedge^\bullet\mathfrak{g}^*$. Indeed, $\mathfrak{g} = \mathfrak{h}\oplus\bigoplus_\alpha \mathfrak{g}_\alpha$ implies that $\wedge^*\mathfrak{g} = \wedge^*\mathfrak{h}\otimes\bigotimes_\alpha \wedge^*\mathfrak{g}_\alpha$. Then for a maximal torus $T\subseteq G$, we see that $t\in T$ acts on $\wedge^*\mathfrak{h}$ by the identity, and t acts on $\wedge^\bullet\mathfrak{g}_\alpha$ by $(1+\alpha(t))$, so the trace of the action of t is

$$\mathrm{tr}(t;\wedge^*\mathfrak{g}) = 2^{\dim\mathfrak{h}}\prod_\alpha(1+\alpha(t)).$$

By Theorem 4.1 (via Remark 4.4), we conclude that

$$\dim\left(\wedge^\bullet\mathfrak{g}\right)^{\mathfrak{g}} = \frac{2^{\dim\mathfrak{h}}}{|W|} = \int_T \prod_{\alpha\in\Phi}\left(1-e^\alpha\right)\left(1+e^\alpha\right)\,dt.$$

This right-hand side can be collapsed into the product over $\left(1-e^{2\alpha}\right)$. Now, all integrals over nontrivial characters will vanish, so we are just asking for the constant term of this integral. But the constant term for the product $\prod_\alpha\left(1-e^{2\alpha}\right)$ will be the same as the constant term for the product $\prod_\alpha\left(1-e^\alpha\right)$, which is the same as the constant term for Δ. By Theorem 4.1 again, we see that this constant term must be $|W|$, so we are done. ■

4.2 April 29

Here we go.

4.2.1 The Exponents

Let's see what else we can say about $\mathrm{H}^\bullet(G;\mathbb{C})$, where G is a compact simply connected almost simple Lie group.

Notation 4.38. Fix a compact simply connected almost simple Lie group G. Then let $(m_1,\ldots,m_r)$ be the increasing sequence of positive integers for which the generators of $\mathrm{H}^\bullet(G;\mathbb{C})$ as an exterior Hopf algebra have degree $2m_i+1$.

Remark 4.39. In Proposition 4.37, we showed that r is the rank.

Example 4.40. On the homework, we will show that $\left(\wedge^3\mathfrak{g}^*\right)^{\mathfrak{g}}$ is one-dimensional. Thus, $m_1=1$, and $m_i>1$ for all $i\geq 2$.

Remark 4.41. Note that

$$\sum_{i=1}^r(2m_i+1)=\dim\mathfrak{g},$$

so $\sum_i m_i=|\Phi_+|$.

For our next calculations, it will be helpful to have more notation.

Definition 4.42 (Poincaré polynomial)**.** Fix a smooth manifold M. Then we define the *Poincaré polynomial* P_M to be

$$P_M(q):=\sum_{j\geq 0}\dim\mathrm{H}^j(M;\mathbb{C})q^j.$$

Example 4.43. One can calculate that $P\mathrm{SU}(2)(q)=1+q^3$ and $P_{\mathrm{SU}(3)}(q)=\left(1+q^3\right)\left(1+q^5\right)$ and $P_{\mathrm{Sp}_4}(q)=\left(1+q^3\right)\left(1+q^7\right)$ and $P_{G_2}(q)=\left(1+q^3\right)\left(1+q^{11}\right)$.

Remark 4.44. By Poincaré duality, one can find that $m_{r+1-i}=h-m_i$, where h is the Coxeter number. Indeed, the point is that $\dim G=h(r+1)$. This can be used to compute the m_is in rank 3.

In fact, one has the following.

Theorem 4.45. Fix a compact simply connected almost simple Lie group G. Then the numbers $\{m_i\}$ are the exponents of the Lie algebra $\mathfrak{g}$.

We will sketch arguments for classical types. For now, we will content ourselves with arguments in type A, and we will suggest how to argue in the other classical types, leaving details to the homework.

Proposition 4.46. We have

$$P_{\mathrm{U}(n)}(q)=(1+q)\left(1+q^3\right)\cdots\left(1+q^{2n-1}\right).$$

Proof. We use Schur–Weyl duality. Let V be the standard representation. Then $\mathfrak{g}^* = V \otimes V^*$, so by Corollary 1.190, we find that

$$\wedge^\bullet(V \otimes V^*) = \bigoplus_{\lambda=(\lambda_1,\ldots,\lambda_n)} \left(S^\lambda V \otimes S^{\lambda^\dagger} V^*\right).$$

Notably, this has no $\mathfrak{g}$-invariants if $\lambda \neq \lambda^\dagger$ because $S^\lambda V$ is just some irreducible representation. However, even if $\lambda = \lambda^\dagger$, then we only have a one-dimensional space of invariants by Schur's lemma, so it follows that

$$P_{\mathrm{U}(n)}(q) = \sum_{\substack{\lambda=(\lambda_1,\ldots,\lambda_n) \\ \lambda=\lambda^\dagger}} q^{|\lambda|}.$$

Thus, we see that we are hunting for Young diagrams λ which are equal to their transpose. Well, the largest such diagram is the full square of length n, and we see that the desired Young diagrams are simply obtained by removing one of the "hooks" which are a top-left segment. Looping over all decisions to either keep or remove a hook, we find that

$$P_{\mathrm{U}(n)}(q) = (1+q)\left(1+q^3\right)\cdots\left(1+q^{2n-1}\right),$$

as required. ■

We are now ready to prove Theorem 4.45 in type A, which amounts to the following.

Corollary 4.47. We have

$$P_{\mathrm{SU}(n)}(q) = \left(1+q^3\right)\left(1+q^5\right)\cdots\left(1+q^{2n-1}\right).$$

Proof. Note that $\mathrm{U}(n)$ contains the subgroup $\mathrm{SU}(n)$, and it is the kernel of the determinant map. Thus, there is a fiber sequence

$$\mathrm{SU}(n) \to \mathrm{U}(n) \to \mathrm{U}(1).$$

Taking Poincaré polynomials reveals that $P_{\mathrm{U}(n)}(q) = P_{\mathrm{U}(1)}(q)P_{\mathrm{SU}(n)}(q)$, so it follows that

$$P_{\mathrm{SU}(n)} = (1+q)^{-1}P_{\mathrm{U}(n)}(q),$$

so we are done by Proposition 4.46. ■

Here is an application of Theorem 4.45.

Theorem 4.48 (Macdonald's identity)**.** Fix a compact simply connected almost simple Lie group G with Lie algebra $\mathfrak{g}$. Then we have

$$\frac{(1+q)^r}{|W|}\int_T \prod_{\alpha\in\Phi}(1+qe^\alpha)(1-e^\alpha) = \prod_{i=1}^r \frac{1+q^{2m_i+1}}{1+q}.$$

Proof. The root decomposition implies that $\wedge^\bullet\mathfrak{g}^* = \wedge^\bullet\mathfrak{h}^* \otimes \bigotimes_\alpha \wedge^\bullet\mathfrak{g}_\alpha^*$. Now, the character from $\wedge^\bullet\mathfrak{h}^*$ is $(1+q)^r$, and the character from $\wedge^\bullet\mathfrak{g}_\alpha^*$ is $(1+qe^\alpha)$, so we find that our Poincaré polynomial is

$$\operatorname{ch}\wedge^\bullet\mathfrak{g}^* \cdot q^j = (1+q)^r \prod_{\alpha\in\Phi}\left(1+qe^\alpha\right).$$

Thus, arguing exactly as in Proposition 4.37, our Poincaré polynomial is

$$\frac{(1+q)^r}{|W|}\int_T \prod_{\alpha\in\Phi}(1+qe^\alpha)(1-e^\alpha),$$

so Theorem 4.45 is equivalent to

$$\frac{(1+q)^r}{|W|}\int_T \prod_{\alpha\in\Phi}(1+qe^\alpha)(1-e^\alpha) = \prod_{i=1}^r \frac{1+q^{2m_i+1}}{1+q},$$

as desired. ■

Remark 4.49. It is also possible to prove Macdonald's identity directly via some calculations, which then proves Theorem 4.45 by the above argument.

While we're here, we give another approach to Theorem 4.45, which will work for the other classical types. We use cell decompositions.

Proof of Proposition 4.46. Recall that $\mathrm{U}(n)$ acts transitively on S^{2n-1} with stabilizer $\mathrm{U}(n-1)$. On the other hand, S^{2n-1} admits a cell decomposition into cells of dimension 0 and a cell of dimension $2n-1$, which can be described as viewing S^{2n-1} as the one-point compactification of S^{2n-1}. Thus, we can use the fibration to inductively give $\mathrm{U}(n)$ a cell decomposition into 2^n cells.

Thus, we could compute the cohomology of $\mathrm{U}(n)$ using cellular cohomology: indeed, the above cells give a cell decomposition. Observe that if the relevant chain complex had any interesting differentials, then $\dim \mathrm{H}^*(\mathrm{U}(n);\mathbb{C})$ would be smaller than 2^n, which contradicts Proposition 4.37. Thus, we must have an isomorphism of graded abelian groups

$$\mathrm{H}^*(\mathrm{U}(n);\mathbb{Z}) \cong \wedge^\bullet_{\mathbb{Z}}(\xi_1,\ldots,\xi_n),$$

where $\deg \xi_i = 2i-1$. This must also be an isomorphism of rings for degree reasons; namely, we see that we must have a generator in degree one, but then we cannot access degree three, so there must be a generator in degree three, and we can continue this process inductively. ■

Remark 4.50. On the homework, we will repeat this argument for the symplectic groups. Namely,

$$\mathrm{H}^*(\mathrm{U}_n(\mathbb{H});\mathbb{Z}) \cong \wedge^\bullet(\xi_1,\ldots,\xi_n),$$

where $\deg \xi_i = 4i-1$. The moral is that $\mathrm{U}_n(\mathbb{H})/\mathrm{U}_{n-1}(\mathbb{H}) \cong S^{4n-1}$.

Remark 4.51. There is no analogous proof for the orthogonal groups. Indeed, we will need more cells: these cells can be seen in the cohomology of $\mathbb{Z}/2\mathbb{Z}$. Instead, one can use the cohomology of Stieffel manifolds, which will be done on the homework.

4.2.2 The Cohomology of Homogeneous Spaces

Fix a compact connected Lie group G, and let $K \subseteq G$ be some closed Lie subgroup; let $\mathfrak{g}$ and $\mathfrak{k}$ be the corresponding Lie algebras. We are now interested in computing the cohomology of G/K. Because G acts on K, its cohomology is computed by the complex $\Omega^\bullet(G/K)^G$ by Remark 4.22. On the other hand, a G-invariant differential form on G/K can be spread out from the identity uniquely, so it is a differential form on $\mathfrak{g}/\mathfrak{k}$. However, to extend to G/K, it must further be invariant by the stabilizer K at the identity, so we conclude that there is an isomorphism

$$(\wedge^\bullet(\mathfrak{g}/\mathfrak{k})^*)^K \cong \Omega^\bullet(G/K)^G$$

of complexes. Note that taking K-invariants may be done in two steps: we can take $\mathfrak{k}$-invariants (which is algebraic) and then take invariants for the finite group K/K°.

We are now interested in the differential of our complex.

Proposition 4.52. Fix an inclusion $\mathfrak{k} \subseteq \mathfrak{g}$ of Lie algebras. Then there is an inclusion

$$(\wedge^\bullet(\mathfrak{g}/\mathfrak{k})^*)^{\mathfrak{k}} \subseteq \wedge^\bullet \mathfrak{g}^*$$

of complexes.

Proof. This will be shown on the homework. ■

This motivates the following definition.

Definition 4.53 (relative Chevalley–Eilenberg complex)**.** Fix an inclusion $\mathfrak{k} \subseteq \mathfrak{g}$ of Lie algebras. Then the *relative Chevalley–Eilenberg complex* is the subcomplex $(\wedge^\bullet(\mathfrak{g}/\mathfrak{k})^*)^{\mathfrak{k}}$ of $\wedge^\bullet \mathfrak{g}^*$. Its cohomology is denoted $\mathrm{H}^*(\mathfrak{g};\mathfrak{k})$. We denote the relative complex by $\mathrm{CE}^*(\mathfrak{g};\mathfrak{k})$.

Remark 4.54. It turns out that the complex $\Omega^\bullet(G/K)$ consists of the K/K°-invariants of the complex $\mathrm{CE}(\mathfrak{g};\mathfrak{k})$. In particular, the spaces are the same, and the differentials agree, which can be checked via Proposition 4.23. Thus,

$$\mathrm{H}^\bullet(G/K;\mathbb{C}) = \mathrm{H}^\bullet(\mathfrak{g};\mathfrak{k})^{K/K^\circ}.$$

Here, note that K normalizes K°, so K/K° does in fact act on G/K°.

Here is a useful trick, which explains why we insist on allowing K to be disconnected.

Lemma 4.55. Fix a compact connected Lie group G, and let $K \subseteq G$ be a closed Lie subgroups; let the Lie algebras be $\mathfrak{k} \subseteq \mathfrak{g}$. Suppose there is $\varepsilon \in K$ which acts by -1 on $\mathfrak{g}/\mathfrak{k}$. Then the cohomology of G/K is supported in even degrees, and

$$\mathrm{H}^\bullet(G/K;\mathbb{C}) \cong \mathrm{H}^\bullet(\mathfrak{g};\mathfrak{k})^{K/K^\circ}.$$

Proof. We may pass to the relative Chevalley–Eilenberg complex. Having an automorphism ε acting by -1 implies that the complex $(\wedge^\bullet(\mathfrak{g}/\mathfrak{k})^*)^K$ is supported in the even degrees. Thus, the differential vanishes automatically, and the result follows. ■

Exercise 4.56. Fix positive integers m and n. We compute the Poincaré polynomial of the complex Grassmannian $\mathrm{Gr}_{m+n,n}(\mathbb{C})$.

Proof. Fix Hermitian spaces V and W of dimensions m and n, and define $G := \mathrm{U}(V \oplus W)$ and $K := \mathrm{U}(V) \times \mathrm{U}(W)$. Then $G/K = \mathrm{Gr}_{m+n,n}(\mathbb{C})$, which classifies n-dimensional Hermitian subspaces of $\mathbb{C}^{m+n}$. Note that $\varepsilon = \left[\begin{smallmatrix} 1 & \\ & -1 \end{smallmatrix}\right]$ acts by -1 on $\mathfrak{g}/\mathfrak{k}$: indeed, $\mathfrak{g}/\mathfrak{k}$ is given by matrices of the form $\left[\begin{smallmatrix} & B \\ C & \end{smallmatrix}\right]$. Thus, by Lemma 4.55,

$$\mathrm{H}^\bullet(\mathrm{Gr}_{m+n,n}(\mathbb{C});\mathbb{C}) = \wedge^\bullet\left(V \otimes W^* \oplus W \otimes V^*\right)^K.$$

By Corollary 1.190, we find that this is

$$\bigoplus_{\lambda,\mu} S^\mu V \otimes S^{\mu^\dagger} W^* \otimes S^\lambda W \otimes S^{\lambda^\dagger} V^*,$$

where λ has at most n parts, and μ has at most m parts.

Now, taking K-invariants, Schur's lemma tells us that this is $\bigoplus_\lambda \mathbb{C}[2|\lambda|]$, where λ has at most n parts, and $\lambda^\dagger$ has at most m parts. Accordingly, let $N_i(n,m)$ be the number of Young diagrams with i boxes and which fit into the rectangle of height n and length m, so we see that $P_{\mathrm{Gr}_{m+n,n}(\mathbb{C})}(q)$ is the generating function $\sum_i N_i(n,m)q^{2i}$. To this end, we let p_i denote the jump $\lambda_i - \lambda_{i+1}$, where $p_0 = n - \lambda_1$ and $p_n = \lambda_n$. Then the

ps are all nonnegative can be seen to uniquely recover λ, so we may compute with these variables instead: note $|\lambda| = 0p_0 + p_1 + 2p_2 + \cdots + np_n$, so

$$\sum_m f_{n,m}(q) z^m = \sum_{p_0,\ldots,p_m \geq 0} q^{2(0p_0+1p_1+2p_2+\cdots+mp_m)} z^{p_0+\cdots+p_m} = \prod_{j=0}^{n} \frac{1}{1-q^{2j}z},$$

which we call good enough for our calculation. ■

Example 4.57. We find that

$$\sum_m f_{1,m}(q) z^m = \frac{1}{(1-z)(1-q^2 z)}.$$

This recovers the Poincaré polynomial of $\mathbb{CP}^m$, once we isolate the z^m term.

Remark 4.58. In general, one has

$$f_{m,n}(q) = \frac{[m+n]_{q^2}!}{[m]_{q^2}![n]_{q^2}!},$$

where $[n]_{q^2} = \frac{1-q^{2n}}{1-q^2}$ and $[n]_{q^2}! = [1]_{q^2} \cdots [n]_{q^2}$. Indeed, this follows from the q-binomial theorem, which is

$$\sum_{m \geq 0} \frac{[m+n]_{q^2}!}{[m]_{q^2}![n]_{q^2}!} z^m = \prod_{j=0}^{n} \frac{1}{1-q^{2j}z}.$$

Indeed, call the right-hand function $F(z)$, which satisfies

$$F\left(q^2 z\right) = \frac{1-z}{1-q^{2(n+1)}z} \cdot F(z).$$

One can check that the left-hand side also satisfies this, which completes the proof because there is only one power series in z with this property.

It is also possible to compute a cell decomposition of the Grassmannian via Schubert cells. Here is the idea, akin to Lemma 4.55.

Remark 4.59. Suppose that a connected compact manifold X admits a cell decomposition with only even-dimensional cells; for example, such things frequently arise from complex manifolds in algebraic geometry. This means that the differential in the cellular chain complex vanishes for degree reasons, so $\mathrm{H}^\bullet(X;\mathbb{Z})$ admits a basis given by those cells.

Exercise 4.60. Fix positive integers m and n. We compute the integral cohomology of the complex Grassmannian $\mathrm{Gr}_{m+n,m}(\mathbb{C})$.

Proof. Choose a complete flag $\{F_i\}$ of $\mathbb{C}^{m+n}$, such as the standard flag. Now, for any $V \in \mathrm{Gr}_{m+n,m}(\mathbb{C})$, the intersections $\{V \cap F_i\}_i$ have dimensions which increase by at most one and rise from 0 to m. Thus, for each i, we may find the smallest integer ℓ_i for which $\dim(V \cap F_{\ell_i}) = i$; for example, $\ell_0 = 0$. Note that $\{\ell_i\}_i$ is a strictly increasing sequence. Then we may define $\lambda_1 := \ell_m - m$ and $\lambda_2 := \ell_{m-1} - (m-1)$ and so on to $\ell_m := \ell_1 - 1$, and we see that λ is a partition with m parts each of size at most n, meaning that λ fits into the rectangle with m rows and n columns.

Now, for such a partition λ fitting into this rectangle, we define $S_\lambda \subseteq \mathrm{Gr}_{m+n,m}(\mathbb{C})$ to be the subset of subspaces V which induce the partition λ. On the homework, we will show that S_λ is a locally closed embedded complex submanifold of complex dimension $|\lambda|$. Thus, $\mathrm{Gr}_{m+n,m}(\mathbb{C})$ is covered by even cells, so we find that $\mathrm{H}^{2i}(\mathrm{Gr}_{m+n,m}(\mathbb{C});\mathbb{Z})$ is free over $\mathbb{Z}$ with basis given by the Schubert cells. ■

Remark 4.61. For example, it follows that the Grassmannian is simply connected.

Remark 4.62. There is a story for Grassmannians over $\mathbb{R}$, but it is more complicated.

Bibliography

[Shu16] Neal Shusterman. *Scythe*. Arc of a Scythe. Simon & Schuster, 2016.

List of Definitions

adjoint, 11, 63
adjoint group, 78
antilinear, 83

boson, 77
bounded, 62

Cartan, 11, 104, 110
Cartan involution, 84
Cartan matrix, 14
character, 17
Chevalley–Eilenberg complex, 116
commutator, 7
compact, 62, 68
compact form, 84
complex, 49
conjugate, 37
Coulomb waves, 77
Coxeter number, 49

de Rham cohomology, 53, 114
de Rham complex, 113
determinant, 91
differential form, 53
dominant, 14
dual root system, 13
Dynkin diagram, 15

exponent, 48
exponential, 8

fermion, 77
form, 81
fundamental weight, 14

generalized Laguerre polynomial, 76
generalized Legendre polynomials, 75

Haar measure, 57
height, 48
highest weight, 16

inner class, 84
inner form, 84
integral, 55

Killing form, 10

Lie algebra, 7
Lie derivative, 114
Lie group, 5
Lie ideal, 8
Lie subgroup, 6
linear, 103
locally compact, 52
locally finite, 52

maximal root, 46
miniscule, 32

nilpotent, 10

partition of unity, 54
Pauli exclusion, 77
Poincaré polynomial, 119
polynomial, 22
positive, 13
precompact, 62
probability, 68
profinite, 70

quantum number, 76
    azimuthal quantum number, 76
    magnetic quantum number, 76
quasisplit, 84
quaternionic, 49
quotient, 6

radical, 10
real, 49
reducible, 12
reductive, 10, 103
relative Chevalley–Eilenberg complex, 122
representation, 6
root system, 12, 12

Schur algebra, 23
Schur functor, 25
Schur polynomial, 29
semisimple, 10, 105
sesquilinear, 91
signature, 92
simple, 10
solvable, 9
spin, 43
split, 110
split form, 81

twisted linear, 82

unimodular, 57, 67
unipotent, 105
unitary, 7
unitary group, 92
universal enveloping algebra, 9

Verma module, 16

Weyl group, 12